COMMENTARY

Potential Impacts of Global Warming and Climate Change on the Epidemiology of Zoonotic Diseases in Canada

Dominique F. Charron, PhD, DVM

lobal climate change may add to the risk of infectious diseases faced by Canadians. A warmer climate may permit the northward expansion of certain disease reservoir animal hosts and vectors increasing the risk to Canadians from zoonotic diseases, i.e., diseases that are transmissible between animals and humans. A shorter, less severe winter would prolong the transmission season for many existing Canadian zoonoses, and could allow previously exotic zoonoses that enter Canada to become established. Significant challenges to epidemiological forecasting arise from complex interactions among climatic factors, zoonotic disease agents, other infectious agents, underlying human health and immunity, socio-economic factors, and ecological parameters. Public health officials and health care providers need to be kept informed of the changing nature of zoonotic disease risk in order to be vigilant in the early detection of disease, and to plan to respond to new or increased health risks. Surveillance and public education will also help protect Canadians from any additional zoonotic health risks of climate change.

Global warming and climate change

Global warming and associated climate change will be important determinants of future Canadian public health. The International Panel on Climate Change (IPCC) accepts projections of a 1.4 to 5.8°C increase in mean global temperature by 2100 as a result of the accumulation of greenhouse gases (carbon dioxide, ozone, methane, nitrous oxide, and water vapour) in the atmosphere. Achievement of Kyoto emissions reduction targets will reduce the rate of warming but will not prevent it. Climate changes in Canada may include increased temperature over most of Canada, especially in winter and at night; reduced soil moisture and summer drought; increased frequency of extreme precipitation events; more severe weather; and an 11 to 77 cm sea-level rise, by 2100. The greatest changes may affect the Arctic. The sea-level rise is a sea-level rise, by 2100.

Vectorborne zoonoses

Climate change may increase the risk to Canadians from vectorborne diseases (caused by pathogens with an invertebrate vector), including arboviral encephalitis, rickettsial diseases, and Lyme borreliosis. Forecasts under conditions of global warming suggest an increase in arboviral encephalitis in the US.⁴ Increased temperatures and longer summers may extend the survival of vectors in Canada, and allow the northward expansion of the ranges of other vectors.⁵ Sporadic human cases of arboviral encephalitis in Canada have been caused by Western

Centre for Infectious Disease Prevention and Control, Population and Public Health Branch, Health Canada, Guelph, Ontario

Correspondence and reprint requests: Dr. Dominique F. Charron, Centre for Infectious Diseases Prevention and Control, Health Canada, 1 Stone Rd W. 4th Floor, Guelph, ON N1G 2Y4, Tel: 519-826-2173, Fax: 519-826-2244, E-mail: Dominique_charron@hc-sc.gc.ca

The information is extracted from an internal report to the Climate Change and Health Office, Safe Environments Program, Healthy Environments and Consumer Safety Branch, Health Canada, January 2001.

Acknowledgements: The author acknowledges the help of Dieter Riedel in the preparation of this manuscript, and thanks CCHO for supporting this work.

Equine Encephalitis (WEE), Powassan virus and viruses of the California serogroup (including Snowshoe Hare virus). There have also been outbreaks of Eastern Equine Encephalitis (EEE) in horses. ^{6,7} Outbreaks of St. Louis Encephalitis (SLE) are common in the US⁸ and there are American reports of sporadic outbreaks of others, including West Nile Virus (WNV). ⁹ West Nile Virus has been found in bird carcasses from southern Ontario. ¹⁰

Rocky Mountain Spotted fever,⁹ Lyme borreliosis,^{11,12} and ehrlichiosis¹³ are tickborne diseases that may have increased incidence in Canada given milder winters and increased spring precipitation. Warmer temperatures could accelerate pathogen development within the tick,¹² and populations of reservoir mice and deer could expand due to increased food availability and decreased winter die-off. Longer summers might increase the risk of human recreational exposure.

Foodborne and waterborne zoonoses

Every year in Canada, foodborne infections cause significant morbidity and mortality. Amany, including *Salmonella* and *Escherichia coli* O157:H7, have known food animal or wildlife reservoirs. Amanda to increased foodborne disease incidence in the UK. Elevated temperatures in North America may exacerbate food contamination during livestock rearing, food processing, and food preparation. Outbreaks of zoonotic diarrhea in heat-stressed livestock may contribute to this contamination.

Contamination of drinking water in Canada by zoonotic pathogens from livestock has been documented. 19 Linkages between water contamination and precipitation events have been noted in the US,20 and proposed in Canada.¹⁹ Several debilitating outbreaks of waterborne zoonotic diseases have been reported. 19-22 Under conditions of climate change, heavy rain or snowmelt would increase the risk of fecal contamination of surface and ground water, leading to drinking water contamination. In the US, over half the enteric disease outbreaks from 1944 to 1994 were reportedly preceded by an extreme precipitation event.23

Other zoonoses

Changing climate will affect the ecology of wild animals, many of which are reservoirs

of diseases infectious to humans. Longer, warmer summers and milder winters will prolong rodent breeding seasons and reduce mortality. Increased small mammal populations may foster animal epidemics of diseases such as Hantavirus infection.²⁴ Outbreaks of rabies, tuberculosis, and brucellosis, for example, may occur in wild ungulates. If a consistently warmer climate allows more northern agricultural land use, there will be risk of disease spread to livestock and people. Increased recreational activities during longer summers may also bring more Canadians into contact with infected wildlife. Plague is established in rodent populations in several areas in the US, and two areas in southern British Columbia and Alberta.9 Under climate conditions favourable to rodent populations, the likelihood of animal epidemics of plague would increase, and increase the risk of human cases.²⁵ Climate change will likely result in changes to wildlife habitat, range and ecology, changing the risk of human and livestock exposure to zoonoses.26

CONCLUSION

Canadians face new health threats from zoonotic disease from global climate change. Changes are expected in the epidemiology of both existing and exotic infectious diseases. Changes in human behaviour associated with warmer weather and longer summers may increase exposure to infectious diseases, and to zoonoses in particular. Canada's North may face new risks from zoonoses given the degree of climate change expected there, the presence of wildlife reservoirs, the potential for vectors to become established, and the potential for the eventual northward expansion of agriculture. The warming projected for southern Canada will result in rapid and complex changes to ecosystems and to the landscape, bringing unforeseen changes in the ecology of infectious diseases. Surveillance, including animal sentinels, will be key in the early detection of emerging zoonotic disease threats under global climate change. Physicians, other health-care providers, public health workers, indeed, all Canadians need to be made aware of potential zoonotic risks related to climate change to allow for the detection of early cases of unusual or unexpected disease, and to protect Canadian public health. An exhaustive review of the potential impact of global climate change on infectious disease risk to people and animals in Ontario is in preparation for the Ontario Ministry of Natural Resources and the Agricultural Research Institute of Ontario.

REFERENCES

- Duncan K, Guidotti T, Cheng W, et al. Canada country study: Impacts and adaptation - health sector. In: Koshida G, Avis W (Eds.). The Canada Country Study: Climate Impacts and Adaptation - Volume VII, National Sectoral Volume. Ottawa: Environment Canada, 1997.
- Intergovernmental Panel on Climate Change (IPCC). Climate Change 2001: The Scientific Basis. Summary for Policy Makers and Technical Summary of the Working Group I Report. IPCC Secretariat, Geneva, 2001.
- Environment Canada Climate Change website, available at http://climatechange.gc.ca, Accessed November 2000.
- Longsreth J, Wiseman J. The potential impact of climate change on patterns of infectious disease in the United States. In: Smith JB, Tirpak DA, (Eds.). The Potential Impact of Climate Change on the United States. Appendix G. Document 230-05-89-057. Washington: US Environmental Protection Agency, 1989.
- Sellers RF. Seasonal variation in spread of arthropod-borne disease agents of man and animals: Implications for control. *Phil Trans Res Soc London* 1983;302:485-95.
- Sellers RF. Eastern equine encephalitis in Quebec and Connecticut, 1972: Introduction by infected mosquitoes on the wind? Can J Vet Res 1972;53:76-79.
- Sellers RF, Maarouf AR. Impact of climate on Western equine encephalitis in Manitoba, Minnesota, and North Dakota, 1980-1983. Epidemiol Infect 1988;101:511-35.
- 8. Shope RE. Arbovirus-related encephalitis. *Yale J Biol Med* 1980;53:93-99.
- 9. Centers for Disease Control website, available at www.cdc.gov. Accessed November 2000.
- 10. Anonymous. West Nile Virus: Canada. Infectious Diseases News Brief. Division of Disease Surveillance, Population and Public Health Branch, Health Canada. August 24, 2001. Available at http://www.hc-sc.gc.ca/pphbdgspsp/bid-bh/2001/nb3401_e.html. Accessed September 2001.

- 11. Lindsay R, Artsob H, Barker I. Distribution of *Ixodes pacificus* and *Ixodes scapularis* re. concurrent babesiosis and Lyme disease. *Can Commun Dis Rep* 1998;24(15):121-22.
- 12. Barker IK, Lindsay LR. Lyme borreliosis in Ontario: Determining the risks. *CMAJ* 2000;162(11):1573-74.
- Telford SR III, Dawson JE, Katavolos P, et al. Perpetuation of the agent of human granulocytic ehrlichiosis in a deer-tick-rodent cycle. *Proc Natl Acad Sci* 1996;93:6209-14.
- Khakhria R, Woodward D, Johnson WM, Poppe C. Salmonella isolated from humans, animals and other sources in Canada, 1983-92. *Epidemiol Infect* 1997;119(1):15-23.
- Van Donkersgoed J, Graham T, Gannon V. The prevalence of verotoxins, *Escherichia coli* 0157:H7, and Salmonella in the feces and rumen of cattle at processing. *Can Vet J* 1999;40(5):332-38
- Letellier A, Messier S, Pare J, et al. Distribution of Salmonella in swine herds in Quebec. Vet Microbiol 1999;67(4):299-306.
- Mikaelian I, Daignault D, Duval MC, Martineau D. Salmonella infection in wild birds from Quebec. Can Vet J 1997;38(6):385.
- 18. Bentham G, Langford I. Climate change and the incidence of food poisoning in England and Wales. *Int J Biometeorol* 1995;39:81-86.
- 19. McQuigge M. The Investigative Report on the Walkerton Outbreak of Waterborne Gastroenteritis, May-June 2000. Unpublished report prepared by Bruce-Grey-Owen Sound Health Unit, Ontario, October 2000.
- MacKenzie WR, Hoxie NJ, Proctor ME. A massive outbreak in Milwaukee of *Cryptosporidium* infection transmitted through the public water supply. N Engl J Med 1994;331:161-67.
- Anonymous. Outbreak of toxoplasmosis associated with municipal drinking water British Columbia. Can Commun Dis Rep 1995;21:161-64.
- denHollander N, Notenboom R. *Toxoplasma gondii* in Ontario and waterborne toxoplasmosis in Victoria, BC. *PHERO* 1996;7:28-31.
- Curriero FC, Patz JA, Rose JB, Lele S. The association between extreme precipitation and water-borne disease outbreaks in the United States, 1948-1994. Am J Public Health 2001;91(8):1194-99.
- 24. Hjelle B, Glass GE. Outbreak of hantavirus infection in the Four Corners region of the United States in the wake of the 1997/1998 El Nino-southern oscillation. *J Infect Dis* 2000;187(5):1569-73.
- 25. Gubler DJ, Reitler P, Ebi KL, Yap W, Nasci R, Patz JA. Climate variability and change in the United States: Potential impacts on vector- and rodent-borne diseases. *Environ Health Perspect* 2001;109(supp2):223-33.
- 26. Intergovernmental Panel on Climate Change (IPCC). Climate Change 2001: Impacts, Adaptation, and Vulnerability Contribution of Working Group II to the IPCC Third Assessment Report. McCarthy JJ, Canziani OF, Leary NA, Dokken DJ, White KS (Eds.). Geneva: IPCC Secretariat, 2001.

Received: October 3, 2001 Accepted: May 21, 2002