

How will climate change affect the temporal and spatial distributions of a reservoir host, the Indian gerbil (*Tatera indica*), and the spread of zoonotic diseases that it carries?

Kordiyeh Hamidi¹, Saeed Mohammadi² and Naeimeh Eskandarzadeh³

¹Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran,
²Department of Environmental Sciences, Faculty of Natural Resources, University of Zabol, Zabol, Iran
and ³Young Researchers and Elite Club, Islamic Azad University, Shirvan Branch, Shirvan, Iran

ABSTRACT

Background: The Indian gerbil (*Tatera indica*) is a main reservoir host of cutaneous leishmaniasis, a great public health problem in many rural areas of Iran.

Questions: How do climatic variables affect the habitat suitability and distribution of *T. indica*? How will changes in climatic variables affect the spatial distribution of *T. indica* across Iran? Will those changes influence the outbreak regions of zoonotic cutaneous leishmaniasis?

Organism: The Indian gerbil, *T. indica*, a rodent.

Analytical methods: Maximum entropy modelling (MaxEnt) to predict suitable regions and the potential distribution of this gerbil in the present and future in Iran.

Results: Species distribution models revealed the four variables most effective in determining Indian gerbil occurrence: the mean precipitation of the year's driest month; the seasonality of precipitation; the mean temperature of the warmest quarter of the year; and the mean temperature of the wettest quarter. According to our model, the southern parts of Iran have the most suitable habitat for *T. indica*. With global climate change, suitable habitats for the gerbil will increase considerably in Iran spreading outwards toward the southwest, centrally, and the northeast.

Conclusions: Our results may be used to estimate outbreaks and prevalence of cutaneous leishmaniasis (as well as other infectious diseases which this gerbil carries).

Keywords: climate change, ectoparasites, Indian gerbil, leishmaniasis, species distribution projections.

INTRODUCTION

All wildlife can carry pathogens (bacteria, viruses, fungi, and parasites) that may cause infectious diseases in humans. These diseases are called zoonotic diseases. Because of this,

Correspondence: K. Hamidi, Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran. email: kordiyeh.hamidi@yahoo.com; kordiyeh.hamidiloyen@mail.um.ac.ir
Consult the copyright statement on the inside front cover for non-commercial copying policies.

small mammals (mainly rodents) play a major role in the outbreak of zoonoses (Krauss *et al.*, 2003). Mammalian reservoirs of zoonoses may be arthropods (fleas, lice, mites, ticks, flies) or they be transmitted directly from animals to humans (mainly viral diseases such as hantaviruses) (Smith *et al.*, 2008; Montoya *et al.*, 2010).

In Iran, both cutaneous and visceral leishmaniasis are significant arthropod-borne diseases. Zoonotic cutaneous leishmaniasis (ZCL) is a tropical disease caused by different species of the genus *Leishmania*. *Leishmania* is a kinetoplastid (Order: Trypanosomatida), a group of flagellated protozoans (which include free-living microorganisms and parasites of various invertebrate, vertebrate, and plant species). *Leishmania* is spread by sandflies of the genera *Phlebotomus* and *Lutzomyia* in the Old World and New World, respectively (de Vries *et al.*, 2015), whose primary hosts are vertebrates such as canids, rodents, and humans (Myler and Fasel, 2008). *Leishmania* occurs widely in rodent populations of savanna and arid regions. It is also a major public health problem in many rural areas of Iran (Khoobdel *et al.*, 2003; Akhavan *et al.*, 2010).

Several species of sandflies belonging to the genus *Phlebotomus* (Insecta; Diptera), including *P. alexandri*, *P. ansarii*, *P. caucasicus*, *P. mongolensis*, *P. papatasi*, and *P. salehi*, act as vectors in the transmission of ZCL between rodents. The most significant parasite carried by these sandflies is *Leishmania major* (Yaghoobi-Ershadi and Akhavan, 1999; Yaghoobi-Ershadi *et al.*, 2005). A few wild rodent species are known to be the principal reservoir hosts of ZCL in Iran (Bates *et al.*, 2015) – these are gerbilline rodents including jirds (*Meriones crassus*, *M. hurrianae*, and *M. libycus*) and gerbils (such as *Rhombomys opimus* and *Tatera indica*) (Rassi *et al.*, 2001, 2006; Hepburn, 2003; Gramiccia and Gradoni, 2005; Gholamrezaei *et al.*, 2016).

Tatera indica is found in Afghanistan, Pakistan, India, Iran, Iraq, Kuwait, Turkey, Syria, Nepal, and Sri Lanka (Harrison and Bates, 1991). Lay (1967) reported this rodent throughout the southern half of Iran. It is locally abundant and presumed to have a large population (Yiöit *et al.*, 2001). *Tatera indica* is the principal reservoir rodent in the western border foci of ZCL (e.g. Khuzestan province – mainly in Dezful and Shush – and Ilam province) (Mohammadi *et al.*, 2017), the southern foci (southern part of Fars province), and the southeastern foci (Sistan and Baluchestan province, especially Chabahar) in Iran (Mohebbi *et al.*, 2004; Oshaghi *et al.*, 2009). The presence of numerous rodent burrows in an area portends a high risk of contracting the disease.

Distribution and ecological niche modelling is widely used to predict the potential ecological distribution of a species and to increase our knowledge about its distribution while identifying the habitats in which it may be undocumented. In the case of species that may carry diseases, ecological niche modelling may permit us to anticipate those areas at high risk for disease occurrence (Gonzalez *et al.*, 2010; Karimi *et al.*, 2014). Ecological niche modelling is based on the idea that the known occurrence of an organism can be related to raster geographic datasets summarizing the factors that likely influence the suitability of the environment for the species across the occurrence landscapes (Phillips and Dudík, 2008; Elith and Leathwick, 2009). So we conducted species distribution modelling to predict the distribution of *T. indica*, to reveal the most important abiotic factors influencing its distribution, and to assess the effects of future climate changes on that distribution. Thus we might estimate the outbreak and prevalence of cutaneous leishmaniasis in Iran.

METHODS

Occurrence records and environmental data

For mapping the distribution range and habitat suitability of *T. indica*, presence data (39 georeferenced distribution records) of the species were assembled during field expeditions using custom-made mesh live traps in North and South Khorasan provinces between the years 2017 and 2018, and also personal collections and available literature (e.g. Darvish and Rastegar-Pouyani, 2012; Hamidi *et al.*, 2015, 2016; Amirafzali *et al.*, 2017; author's unpublished data, and data available on the internet: GBIF at <https://www.gbif.org> and VerNet at <http://www.vernet.org>). We assembled 19 bioclimatic variables as well as elevation with 30 arc-seconds (~1 km) resolution, obtained from the WorldClim database (<http://www.worldclim.org/>). Environmental layers and records were applied by open Modeler v.1.0.7 (de Souza Muñoz *et al.*, 2011) to determine the relevant grid values for each environmental layer. The extracted values were tested by SPSS v.16.0 to determine the Pearson correlation coefficient. We then approximated a set of independent layers by choosing layers with correlations less than 0.75 to run the final model.

Modelling of species distribution

We used MaxEnt (maximum entropy machine learning algorithm) v.3.4.1k (Phillips *et al.*, 2017) with 15 replications to produce the final model. MaxEnt uses 'presence only' records to achieve high-performance model predictions even with low occurrence data; several ecologists believe it to be the most efficient method of ecological niche modelling (Hernandez *et al.*, 2006; Phillips *et al.*, 2006; Elith *et al.*, 2011; Merow *et al.*, 2013). Finally, we considered the area under the receiver operating characteristic curve (AUC) as the criterion for the training data and act as a measure for the model's ability to discriminate 'present' from 'absent' at each point. The value of AUC ranges between 0 and 1. A model with no ability to discriminate should return an AUC of 0.5; an AUC greater than 0.5 indicates that the model is better than random; and a value close to 1 indicates near perfect accuracy of the model (Phillips and Dudik, 2008). These threshold values produced a potential species distribution map that we regrouped into two classes: high potential (>0.5) and low potential (<0.005) (Abolmaali *et al.*, 2018).

Climate data under the present and future change scenarios

To evaluate the potential future distribution of species for the year 2050 (average for 2046–2065), we projected the model using MaxEnt (Morueta-Holme *et al.*, 2010). Global coupled Atmospheric-Ocean General Circulation Models (GCMs) are essential tools in theoretical investigations of climatic change mechanisms (Covey *et al.*, 2003). By using GCMs from <http://www.ccafs-climate.org/>, we can simulate the present and project future climatic changes under different scenarios. We used the CCSM 4.0 (Community Climate System Model) simulations for the Intergovernmental Panel on Climate Change (IPCC) Fourth Assessment Report (AR4) in 44 CC (Climate Change) scenarios to predict habitat suitability of species in Iran (Pendergrass and Hartmann, 2012). All layers were downloaded in 30 arc-second resolutions and cropped for Iranian boundaries using ArcGIS v.10.3. Based on different inputs of greenhouse gas emission drivers (i.e. population and economic growth, and technological choices), land use changes, environmental policy options,

and adaptation processes (Solomon *et al.*, 2007), the IPCC recommended four representative concentration pathways (RCPs) (Miao *et al.*, 2014). The RCPs are consistent with a wide range of possible changes in future anthropogenic, greenhouse gas (GHG) emissions, and aim to represent the atmospheric concentrations. We considered RCPs v.2.6 and 6, which described a possible future range of energy states for the earth on the basis of different trends in climate change drivers.

RESULTS

Based on the Pearson correlation coefficient results, we ran the MaxEnt model in seven layers. Table 1 records the layers we selected. The ecological niche modelling output based on the model calibration results for testing data was 0.7862 (standard deviation = 0.1032). The following had the most marked effect on the model's output:

- mean temperature of warmest quarter (percentage contribution value (PC): 54.12% and permutation importance value (PIMP): 7.92%);
- precipitation of driest month (PC: 13.90%, PIMP: 1.38%);
- precipitation seasonality (PC: 10.02%, PIMP: 27.88%);
- mean temperature of wettest quarter (PC: 9.53%, PIMP: 40.38%).

Environments with high temperatures and low rainfall were most likely to harbour the species (Fig. 1). The Khuzestan plain in southwest Iran, the coastal area of southeast Iran, southeast Dasht-e Kavir (the Great Salt Desert) in Kerman province, as well as central and northeastern areas in Sistan and Baluchestan province were the most suitable habitats for *T. indica* (Fig. 2). Figure 3 shows the potential distribution map of the MaxEnt model by CCSM4 and RCP45. The potential distribution map of *T. indica* in the future shows a remarkable expansion outwards from currently suitable areas, with shifts into the south-west, central, and northeast of the country. Meanwhile, the south of Iran will remain suitable.

Table 1. Bioclimatic variables used to develop the distribution model in MaxEnt. Contribution ratio (%) and permutation importance value for each layer are also shown

Variable	Percentage contribution	Permutation importance
Bio10 (mean temperature of warmest quarter)	54.12	7.92
Bio14 (precipitation of driest month)	13.90	1.38
Bio15 (precipitation seasonality; coefficient of variation)	10.02	27.88
Bio8 (mean temperature of wettest quarter)	9.53	40.38
Bio3 (isothermality)	5.36	3.14
Bio12 (annual precipitation)	4.27	8.64
Bio2 (mean diurnal range; mean of monthly [maximum – minimum] temperature)	2.79	10.67




Fig. 1. The response curves of four variables with the greatest effect on the output of the model of current conditions. (A) Bio8: mean temperature of wettest quarter (in $^{\circ}\text{C} \times 10$); (B) Bio10: mean temperature of warmest quarter (in $^{\circ}\text{C} \times 10$); (C) Bio14: precipitation of driest month (in mm); (D) Bio15: precipitation seasonality (coefficient of variation) (%).




Fig. 2. Potential distribution map of the Indian gerbil, *Tatera indica*, and its classified map in the present. Black dots indicate the localities.




Fig. 3. Potential distribution map of the Indian gerbil, *Tatera indica*, and its classified map in the future. Black dots indicate the localities.

DISCUSSION

Species distribution patterns are among the most prominent issues in biodiversity management (Tittensor *et al.*, 2014). Distribution models have provided a new tool to explore diverse questions about species distributions (Nally and Fleishman, 2004). More specifically, investigating these patterns for host reservoir species helps us to use them for public health purposes. In spite of the growing number of GIS-based studies of leishmaniasis vectors, especially wild rodent hosts, only a few studies have been conducted in Iran (Ghatee *et al.*, 2013; Mollalo *et al.*, 2015; Fakhar *et al.*, 2017).

Zoonotic cutaneous leishmaniasis is an endemic zoonosis in many rural districts of Iran. The Indian gerbil, *Tatera indica*, together with sandflies of the genus *Phlebotomus* play a major role in the transmission of the pathogen (Mehrabani *et al.*, 2007; Ghatee *et al.*, 2017). Published literature shows that *T. indica*, in southern and southwestern Iran, is one of the main reservoir hosts (Gholamrezaei *et al.*, 2016). Our own output distribution map shows that the more southern parts of Iran have the largest amount of suitable habitat for *T. indica*. According to the response curves, increases in rainfall and decreases in temperature will result in the significant decline of its presence probability (Fig. 1).

Epidemiological research has shown that temperature and precipitation affect the distribution of tick-borne diseases (Gage *et al.*, 2008). Recent studies have illustrated that the geographical distribution of vectors and reservoirs of leishmaniasis can be affected by environmental variables (weather parameters, vegetation and topography changes) (Patz and Olson, 2006) and man-made disturbances (e.g. urbanization). Another important factor in

the distribution of *T. indica* is seasonality of precipitation (coefficient of variation). This variation stems mainly from the fact that these areas are near the sea and show coastal climate conditions – mild climate with hot summers and cool winters, high seasonal precipitation, and temperatures averaging about 50°C (Islamic Republic of Iran Meteorological Organization Site, at <http://www.irimo.ir/>). In agreement with Molur *et al.* (2005), we confirm that this species lives in a range of dry or arid habitats with sparse or low density of vegetation, from low elevations in the south and centre to the northern part of Iran. When we overlaid our predicted distribution map on the map of Iran's biotopes, we could also see that *T. indica* prefers 'hot dry desert', 'hot semi-desert', and 'coastal dry' climatic zones (<http://www.irimo.ir/>). The average elevation of the observed localities in our study is 755 m a.s.l. (maximum elevation 2800 m a.s.l., only two points above 2000 m, and a minimum value of 4 m a.s.l.). The Indian gerbil is a warm area species that penetrated Torbat-e Jam in the northeast of Iran (Razavi Khorasan province) (Darvish and Rastegar-Pouyani, 2012).

A half century ago, the northeasternmost part of the range of *T. indica* was the Tabas region (South Khorasan province) (Misonne, 1959). But recent data indicate that *T. indica* is expanding further northeast, probably due to the effects of global warming and drought. The climate changes in Iran have increased attention towards the impact of climate change on the incidence and prevalence of cutaneous leishmaniasis in different provinces of Iran (Alvar *et al.*, 2012; Mollalo *et al.*, 2015; Ramezankhani *et al.*, 2017).

Climate change scenarios help us predict future conditions, prepare to face them, and plan better so that we may prevent disease. The future distribution showed that in Iran suitable habitats for the Indian gerbil will increase considerably outwards toward the south-west, centre, and northeast. The impact of climatic factors on the outbreak of leishmaniasis in the west of Iran showed that there was a significant negative correlation between the temperature and disease incidence, with leishmaniasis peaking in October and November. However, there is no significant correlation between disease outbreak and rainfall or relative humidity (Yazdanpanah *et al.*, 2013).

In a similar study, Akbari *et al.* (2014) showed that the highest incidence of the disease happened during the second half of the year (especially in the autumn; between October to December in Razavi Khorasan, northeast Iran). Moreover, they noted a weak positive correlation with relative humidity and rainfall, and a weak inverse correlation with sunshine and temperature. However, Entezari and Eskandari (2014) showed that zoonotic cutaneous leishmaniasis had the highest incidence and prevalence in January and February in Fars province. In analysing bioclimatic factors epidemiologically, Githeko *et al.* (2000) showed that temperature and hours of sunshine must play a fundamental role in the transmission of vector-borne diseases; these show a direct relationship with the prevalence of disease. Moreover, rainfall and relative humidity had an inverse correlation. On the other hand, the impact of vegetation cover on the incidence of leishmaniasis and the life cycle of sandflies was studied in Yazd-Ardakan Plain. Comparing spatial models of human disease with vegetation cover showed that the highest incidence of cutaneous leishmaniasis was concentrated in areas with sparse or low density of vegetation (Mozaffari *et al.*, 2012). Bavia *et al.* (2005) showed similar results in correlation analysis between sandflies and vegetation density of mosquitoes in Brazil.

Generally, the Indian gerbil lives in colonies. Its external morphology overlaps that of many species of *Meriones* (especially with Sundevall's jird, *M. crassus*, with which it co-occurs). These several species constitute alternative reservoirs for zoonotic diseases such as leishmaniasis (Fig. 4).




Fig. 4. (A) Habitat and vegetation cover of the Indian gerbil (*Tatera indica*) and Sundevall's jird (*Meriones crassus*), two gerbilline rodents that often co-occur. Both are reservoirs for leishmaniasis. Ferdows county, South Khorasan province, Iran. (B) Burrow entrances of a colony of *T. indica*. (C) Comparison of external morphology of *T. indica* and *M. crassus*. The main diagnostic character is the dual-coloured tail (dark colour above and below, pale at the sides) of *T. indica*.

The main diagnostic characters which distinguish *T. indica* from other gerbilline rodents are its tail colour (dark above and below, but pale at the sides); its tail sometimes being slightly shorter than its head and body length; and its hind feet having naked soles (Baramaki Yazdi *et al.*, 2011). So it can be difficult to discriminate from jirds in the field. And accurate identification is made no easier when we consider the great overlap of the habitats of different species of jirds and gerbils. Perhaps these similarities have even resulted in misidentification of the species. This, along with inadequate information about the ecological distribution of *T. indica* populations and its habitat structure (and also those of other gerbillines) are among the factors that may result in difficulties in the study of *Leishmania* outbreaks, and programmes to manage it by controlling its reservoir hosts. However, undertaking such studies is crucial to public health.

CONCLUSION

The Indian gerbil, *Tatera indica*, harbours several ectoparasites such as mites (e.g. *Dermanyssus americanus*, *Echinolaelaps echidninus*, *Haemolaelaps glasgowi*, *Laelaps nuttalli*, and *Ornithonyssus bacoti*), fleas (mainly *Xenopsylla buxtoni*), and lice species (such as *Polyplax gerbilli* and *Rhipicephalus* sp.) (Hanafi-Bojd *et al.*, 2007; Hamidi *et al.*, 2015, 2016). Reasonably accurate prediction of its distribution as well as the role of bioclimatic factors in its presence will be valuable for preventing zoonoses and is vital to programmes attempting to control zoonotic arthropod-borne diseases.

Some zoonotic infections have no vaccine and lack specific drugs for their treatment. Hence it is better to prevent the outbreak of a disease than to try to get rid of it once it has begun (Oglesbee, 2011). In a world with unusually rapid climatic change, our results can be used to improve prevention by anticipating which places will become havens for the rodent and so require enhanced public health defences.

ACKNOWLEDGEMENTS

The authors wish to express their sincere thanks to H. Mozaffari, A. Hamidi, and E. Ebrahimi for their kind support, and Dr. M. Nassirkhani and Professor M.L. Rosenzweig for their critical reviews of the manuscript.

REFERENCES

- Abdolmaali, S.M.R., Tarkesh, M. and Bashari, H. 2018. MaxEnt modeling for predicting suitable habitats and identifying the effects of climate change on a threatened species, *Daphne mucronata*, in central Iran. *Ecol. Inform.*, **43**: 116–123.
- Akbari, E., Mayvaneh, F., Entezari, A.R. and Nazari, M. 2014. Survey of the role of bioclimatic factors in the outbreak of cutaneous leishmaniasis. *Iran. J. Epidemiol.*, **10** (3): 65–74 [in Persian with English abstract].
- Akhavan, A.A., Yaghoobi-Ershadi, M.R., Khamesipour, A., Mirhendi, H., Alimohammadian, M.H., Rassi, Y. *et al.* 2010. Dynamics of *Leishmania* infection rates in *Rhombomys opimus* (Rodentia: Gerbillinae) population of an endemic focus of zoonotic cutaneous leishmaniasis in Iran. *Bull. Soc. Pathol. Exot.*, **103**: 84–89.
- Alvar, J., Vélez, I.D., Bern, C., Herrero, M., Desjeux, P., Cano, J. *et al.* 2012. Leishmaniasis worldwide and global estimates of its incidence. *PLoS One*, **7** (5): e35671.
- Amirafzali, Y., Darvish, J. and Yazdani-Moghaddam, F. 2017. Study of rodents' fauna of the Jiroft, Kerman Province in southeast of Iran. *Iran. J. Anim. Biosyst.*, **13**: 119–129.
- Baramaki Yazdi, R., Shaghoozaie, M., Galdavi, S., Yousef Elahi, M., Shahriar Moghadam, M., Shokri Bousjei, N. *et al.* 2011. Identification of a new morphotype of the Indian Gerbil, *Tatera indica* Hardwicke, 1807 (Muridae, Rodentia), from Iran. *Zool. Middle East*, **52**: 111–113.
- Bates, P.A., Depaquit, J., Galati, E.A., Kamhawi, S., Maroli, M., McDowell, M.A. *et al.* 2015. Recent advances in phlebotomine sand fly research related to leishmaniasis control. *Parasit. Vectors*, **8**: 131.
- Bavia, M.E., Carneiro, D.D., Gurgel Had, C., Madureira Filho, C. and Barbosa, M.G. 2005. Remote sensing and geographic information systems and risk of American visceral leishmaniasis in Bahia, Brazil. *Parassitologia*, **47**: 165–169.
- Covey, C., AchutaRao, K.M., Cubasch, U., Jones, P., Lambert, S.J., Mann, M.E. *et al.* 2003. An overview of results from the Coupled Model Intercomparison Project. *Glob. Planet. Change*, **37**: 103–133.

- Darvish, J. and Rastegar-Pouyani, E. 2012. Biodiversity conservation of reptiles and mammals in the Khorasan provinces, northeast of Iran. *Prog. Biol. Sci.*, **2** (1): 95–109.
- De Souza Muñoz, M.E., De Giovanni, R., De Siqueira, M.F., Sutton, T., Brewer, P., Pereira, R.S. *et al.* 2011. OpenModeller: a generic approach to species' potential distribution modelling. *GeoInformatica*, **15**: 111–135.
- de Vries, H.J.C., Reedijk, S.H. and Schallig, H.D.F.H. 2015. Cutaneous leishmaniasis: recent developments in diagnosis and management. *Am. J. Clin. Dermatol.*, **16**: 99–109.
- Elith, J. and Leathwick, J.R. 2009. Species distribution models: ecological explanation and prediction across space and time. *Annu. Rev. Ecol. Evol. Syst.*, **40**: 677–697.
- Elith, J., Phillips, S.J., Hastie, T., Dudík, M., Chee, Y.E. and Yates, C.J. 2011. A statistical explanation of MaxEnt for ecologists. *Divers. Distrib.*, **17**: 43–57.
- Entezari, M. and Eskandari, F. 2014. Relationship between climatic factors and the prevalence of cutaneous leishmaniasis in Larestan City. *J. Mil. Med.*, **16** (2): 99–104 [in Persian with English abstract].
- Fakhar, M., Karamian, M., Ghatee, M.A., Taylor, W.R., Pazoki Ghohe, H. and Rasooli, S.A. 2017. Distribution pattern of anthroponotic cutaneous leishmaniasis caused by *Leishmania tropica* in Western Afghanistan during 2013–2014. *Acta Trop.*, **176**: 22–28.
- Gage, K.L., Burkot, T.R., Eisen, R.J. and Hayes, E.B. 2008. Climate and vectorborne diseases. *Am. J. Prev. Med.*, **35**: 436–450.
- Ghatee, M.A., Sharifi, I., Haghdooost, A.A., Kanannejad, Z., Taabody, Z., Hatam, G. *et al.* 2013. Spatial correlations of population and ecological factors with distribution of visceral leishmaniasis cases in southwestern Iran. *J. Vector Borne. Dis.*, **50**: 179–187.
- Ghatee, M.A., Haghdooost, A.A., Kooreshnia, F., Kanannejad, Z., Parisaie, Z., Karamian, M. *et al.* 2017. Role of environmental, climatic risk factors and livestock animals on the occurrence of cutaneous leishmaniasis in newly emerging focus in Iran. *J. Infect. Public Health* [<https://doi.org/10.1016/j.jiph.2017.12.004>].
- Gholamrezaei, M., Mohebbali, M., Hanafi-Bojd, A.A., Sedaghat, M.M. and Shirzadi, M.R. 2016. Ecological niche modeling of main reservoir hosts of zoonotic cutaneous leishmaniasis in Iran. *Acta Trop.*, **160**: 44–52.
- Githeko, A.K., Lindsay, S.W., Confalonieri, U.E. and Patz, J.A. 2000. Climate change and vector-borne diseases: a regional analysis. *Bull. World Health Organ.*, **78**: 1136–1147.
- Gonzalez, C., Wang, O., Strutz, S.E., Gonzalez-Salazar, C., Sanchez-Cordero, V. and Sarkar, S. 2010. Climate change and risk of leishmaniasis in North America: predictions from ecological niche models of vector and reservoir species. *PLoS Neglect. Trop. Dis.*, **4**: e585.
- Gramiccia, M. and Gradoni, L. 2005. The current status of zoonotic leishmaniasis and approaches to disease control. *Int. J. Parasitol.*, **35**: 1169–1180.
- Hamidi, K., Nourani, L. and Moravvej, G.H. 2015. The relationship of ectoparasite prevalence to the capturing season, locality and species of the murine rodent hosts in Iran. *Persian J. Acarol.*, **4**: 409–423.
- Hamidi, K., Nourani, L. and Moravvej, G.H. 2016. New rodents' hosts of sucking lice, fleas (Insecta: Anoplura, Siphonaptera) and hard ticks (Acari: Ixodida) from Iran. *Persian J. Acarol.*, **5**: 85–88.
- Hanafi-Bojd, A.A., Shahi, M., Baghaii, M., Shayeghi, M., Razmand, N. and Pakari, A. 2007. A study on rodent ectoparasites in Bandar Abbas: the main economic southern seaport of Iran. *Iran. J. Environ. Health Sci. Eng.*, **4** (3): 173–176.
- Harrison, D.L. and Bates, P.J.J. 1991. *The Mammals of Arabia*, 2nd edn. Sevenoaks: Harrison Zoological Museum.
- Hepburn, N.C. 2003. Cutaneous leishmaniasis: current and future management. *Expert Rev. Anti-infect. Ther.*, **1**: 563–570.
- Hernandez, P.A., Graham, C.H., Master, L.L. and Albert, D.L. 2006. The effect of sample size and species characteristics on performance of different species distribution modelling methods. *Ecography*, **29**: 773–785.

- Karimi, A., Hanafi-Bojd, A.A., Yaghoobi-Ershadi, M.R., Akhavan, A.A. and Ghezalbashi, Z. 2014. Spatial and temporal distributions of phlebotomine sand flies (Diptera: Psychodidae), vectors of leishmaniasis, in Iran. *Acta Trop.*, **132**: 131–139.
- Khoobdel, M., Fajrak, H., Ladonni, H., Shayeghi, M. and Asadzadeh, R. 2003. A new method for military personnel protection against insects. *J. Mil. Med.*, **5** (2): 147–155.
- Krauss, H., Weber, A., Appel, M., Enders, B., Isenburg, H.D., Schiefer, H.G. *et al.* 2003. *Zoonoses: Infectious Diseases Transmissible from Animals to Humans*. Washington, DC: ASM Press.
- Lay, D.M. 1967. A study of the mammals of Iran resulting from the street expedition of 1962–63. *Fieldiana: Zool.*, **54**: 1–282.
- Mehrabani, D., Motazedian, M., Oryan, A., Asgari, Q., Hatam, G. and Karamian, M. 2007. A search for the rodent hosts of *Leishmania major* in the Larestan region of southern Iran: demonstration of the parasite in *Tatera indica* and *Gerbillus* sp., by microscopy, culture and PCR. *Ann. Trop. Med. Parasitol.*, **101**: 315–22.
- Merow, C., Smith, M.J. and Silander J.A., Jr. 2013. A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography*, **36**: 1058–1069.
- Miao, C., Duan, Q., Sun, Q., Huang, Y., Kong, D., Yang, T. *et al.* 2014. Assessment of CMIP5 climate models and projected temperature changes over Northern Eurasia. *Environ. Res. Lett.*, **9**: 055007.
- Misonne, X. 1959. *Zoogeography the mammifere de l'Iran*. Bruxelles: Institut Royal des Science Naturelles de Belgique. Memoires, deuxième série, fasc. 59, 160.
- Mohammadi, S., Vazirianzadeh, B., Fotouhi-Ardakani, R., Alaee Novin, E., Amirkhani, A., Samii, J. *et al.* 2017. *Tatera indica* (Rodentia: Muridae) as the prior concern and the main reservoir host of zoonotic cutaneous leishmaniasis on the border of Iran and Iraq. *Jundishapur J. Microbiol.*, **10** (1): e42452.
- Mohebbi, M., Javadian, E., Yaghoobi-Ershadi, M.R., Akhavan, A.A., Hajjarian, H. and Abaei, M.R. 2004. Characterization of *Leishmania* infection in rodents from endemic areas of the Islamic Republic of Iran. *East Mediterr. Health J.*, **10**: 591–599.
- Mollalo, A., Alimohammadi, A., Shirzadi, M.R. and Malek, M.R. 2015. Geographic information system-based analysis of the spatial and spatio-temporal distribution of zoonotic cutaneous leishmaniasis in Golestan Province, north-east of Iran. *Zoonoses Public Health*, **62**: 18–28.
- Molur, S., Srinivasulu, C., Srinivasulu, B., Walker, S., Nameer, P.O. and Ravikumar, L., eds. 2005. *Status of South Asian Non-volant Small Mammals: Conservation Assessment and Management Plan (C.A.M.P.)*. Workshop Report. Coimbatore, India: Zoo Outreach Organization/ CBSG-South Asia.
- Montoya, J.G., Boothroyd, J.C. and Kovacs, J.A. 2010. *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*, 7th edn. Philadelphia, PA: Elsevier.
- Moruela-Holme, N., Fløjgaard, C. and Svenning, J.C. 2010. Climate change risks and conservation implications for a threatened small-range mammal species. *PLoS One*, **5** (4): e10360.
- Mozaffari, G.H., Bakhshizadeh, F. and Gheibi, M. 2012. Analysis relationship between vegetation cover and salak skin disease in Yazd-Ardakan Plain. *Geogr. Environ. Plan.*, **22** (4): 47–50.
- Myler, P. and Fasel, N. 2008. *Leishmania: After the Genome*. Poole: Caister Academic Press.
- Nally, R. and Fleishman, E. 2004. Influence of temporal scale of sampling on detection of relationships between invasive plants and the diversity patterns of plants and butterflies. *Conserv. Biol.*, **18**: 1525–1532.
- Oglesbee, B.L. 2011. *Blackwell's Five-Minute Veterinary Consult: Small Mammal*, 2nd edn. New York: Wiley.
- Oshaghi, M.A., Maleki Ravasan, N., Javadian, E., Mohebbi, M., Hajjarian, H., Zare, Z. *et al.* 2009. Vector incrimination of sand flies in the most important visceral leishmaniasis focus in Iran. *Am. J. Trop. Med. Hyg.*, **81**: 572–577.

- Patz, J.A. and Olson, S.H. 2006. Climate change and health: global to local influences on disease risk. *Ann. Trop. Med. Parasitol.*, **100**: 535–549.
- Pendergrass, A.G. and Hartmann, D.L. 2012. Global-mean precipitation and black carbon in AR4 simulations. *Geophys. Res. Lett.*, **39** (1): L01703.
- Phillips, S.J. and Dudik, M. 2008. Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography*, **31**: 161–175.
- Phillips, S.J., Anderson, R.P. and Schapire, R.E. 2006. Maximum entropy modelling of species geographic distributions. *Ecol. Model.*, **190**: 231–259.
- Phillips, S.J., Anderson, R.P., Dudik, M., Schapire, R.E. and Blair, M. 2017. Opening the black box: an open-source release of Maxent. *Ecography*, **40**: 887–893.
- Ramezankhani, R., Sajjadi, N., Nezakati Esmailzadeh, R., Jozi, S.A. and Shirzadi, M.R. 2017. Spatial analysis of cutaneous leishmaniasis in an endemic area of Iran based on environmental factors. *Geospat. Health*, **12** (2): 578.
- Rassi, Y., Jalali, M., Javadian, E. and Moatazedian, M.H. 2001. Confirmation of *Meriones libycus* (Rodentia: Gerbillidae) as the main reservoir host of zoonotic cutaneous leishmaniasis in Arsanjan, Fars Province, south of Iran (1999–2000). *Iran. J. Public Health*, **30**: 143–144.
- Rassi, Y., Javadian, E., Amin, M., Rafizadeh, S., Vatandoost, H. and Motazedian, H. 2006. *Meriones libycus* is the main reservoir zoonotic cutaneous leishmaniasis in south Islamic Republic of Iran. *East Mediterr. Health J.*, **12**: 474–477.
- Smith, V.S., Light, J.E. and Durden, L.A. 2008. Rodent louse diversity, phylogeny, and cospeciation in the Manu Biosphere Reserve, Peru. *Biol. J. Linn. Soc.*, **95**: 598–610.
- Solomon, I.S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B. *et al.* 2007. *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group 1 to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press.
- Tittensor, D.P., Walpole, M., Hill, S.L., Boyce, D.G., Britten, G.L., Burgess, N.D. *et al.* 2014. A mid-term analysis of progress toward international biodiversity targets. *Science*, **346** (6206): 241–244.
- Yaghoobi-Ershadi, M.R. and Akhavan, A.A. 1999. Entomological survey of sand flies (Diptera: Psychodidae) in a new focus of zoonotic cutaneous leishmaniosis in Iran. *Acta Trop.*, **73**: 321–326.
- Yaghoobi-Ershadi, M.R., Akhavan, A.A., Zahraei-Ramazani, A.R., Jalali-Zand, A.R. and Piazak, N. 2005. Bionomics of *Phlebotomus papatasi* (Diptera: Psychodidae) in an endemic focus of zoonotic cutaneous leishmaniasis in central Iran. *J. Vector. Ecol.*, **30**: 115–118.
- Yazdanpanah, H., Baratian, A. and Karimi, S. 2013. Study the relationship of the effective climatic factors on the outbreak of leishmaniasis in Qasr-e qand county. *Spatial Plan.*, **3** (3): 69–86 [in Persian].
- Yiđit, N., Colak, E., Verimli, R., Ozkurt, S. and Sozen, M. 2001. A study on the distribution morphology and karyology of *Tatera indica* (Hardwicke, 1807) (Mammalia: Rodentia) in Turkey. *Turk. J. Zool.*, **25**: 65–67.