

Ocular rhinosporidiosis in Tamil Nadu, India

J.S. Moses¹ & C. Balachandran², S. Sandhanam & N. Ratnasamy, S. Thanappan, Johnson Rajaswar & Dinakar Moses³

Madras Veterinary College, Madras-600 007, Ophthalmologists, Govt. Headquarter's Hospital, Nagercoil-629 001, Veterinary Doctors, Nagercoil-629 001

Received 11 December 1989; accepted 23 December 1989

Key words: Rhinosporidiosis, ocular, Tamil Nadu, India

Abstract

A high incidence of ocular rhinosporidiosis in Kanyakumari district of Tamil Nadu, India is reported. Among the four taluks (sub-division for administration), highest occurrence was found in Agastheeswaram taluk (51.2%) followed by Kalkulam (24.4%), Thovalai (22%) and Vilavancode (2.4%). The disease occurred among both sexes equally and preponderance of a particular sex was not observed. The young adolescents were found to be more susceptible. The disease was found in all socio-economic strata and among all communities and persons belonging to different religious groups. Most of the patients gave a history of bathing in muddy stagnant pools of water.

Introduction

Rhinosporidiosis, a mycotic infection, is caused by *Rhinosporidium seeberi*, producing polyps mostly on the mucous membranes of the nostrils and throat, less frequently in the eye and rarely in the bronchus, genitalia, skin and bone. Rhinosporidiosis of the eye has been reported from various parts of the world and in India [1–27]. The elusive nature of the epidemiology of rhinosporidiosis makes it as one of the most unique and interesting diseases. This paper is about the high incidence of ocular rhinosporidiosis (OR) in Kanyakumari (KK) district of Tamil Nadu, India.

Materials and methods

Two hundred and nineteen cases of primary OR recorded between August, 1984 and November, 1988 at Government Headquarters Hospital, Nagercoil were analysed. Tissues preserved in 10% formalin were processed, stained by haematoxylin and eosin and examined. The growth varied in size from a pin head to a polyp measuring 2 × 3 cm (Figs. 1–3).

¹ Retired Professor of Microbiology and Project Leader University Scheme on 'Electronmicroscopic and Ultrastructural studies on rhinosporidiosis in nasal polyps of animals and man'. *Present address:* Emeritus Professor of Indian Council of Agricultural Research at Madras Veterinary College, Madras-600 007, India.

² Then Assistant Professor of the Scheme. *Present address:* Associate Professor of Pathology, Veterinary College and Research Institute, Namakkal-637 002, India.

³ E.N.T. specialist, Madras-600 007.




Fig. 1.



Fig. 2.



Fig. 3.

Results and discussion

Histopathological examination of the ocular growths were positive for rhinosporidiosis. The histological features agree with earlier findings [9, 10].

The age and sex distribution and various anatomic sites involved are given in Table 1 and 2 respectively. The youngest was one year old and the oldest 79 years old. Of 219 patients, 119 were male and 100 were female.

Moral and Gracia [14] stated that out of 510 cases of OR, 233 were from India. Shukla *et al.* [24] reported 119 cases from Raipur during a 13 year period. Six out of 272 cases attended at 3 medical colleges were OR 926. Billore [4] found

a prevalence rate of 1% in urban area and 0.22% in rural area of Raipur. This report of 219 cases in a period of 4 years and 3 months, clearly indicates a high incidence of the disease in KK district. Among the four taluks, the highest occurrence was found in Agastheeswaram (51.2%) followed by Kalkulam (24.4%), Thovalai (22%) and Vilavancode (2.4%). A high incidence of nasal rhinosporidiosis in Agastheeswaram taluk has been reported previously [5, 13, 15]. Even in a sample survey conducted in a high school located in this area, one case of OR was seen out of 4 rhinosporidiosis cases [15].

The male to female ratio in this study was 1.19 : 1.00 as against 2 : 1 [24]. In the age group of 11 to 20 the number of females even though

Table 1. Age and sex distribution of ocular rhinosporidiosis

Age group	Male	Female	Total
0-10	32	26	58
11-20	51	48	99
21-30	12	4	16
Above 30	24	22	46
Total	119	100	219

arithmetically more, was not statistically significant. A preponderance of rhinosporidiosis in the male [3, 24] or the female [5, 13] has been reported. But, in this study almost an equal distribution among both sexes were observed.

The growths were more common in upper tarsal conjunctiva (62.5%) than in the bulbar conjunctiva (7.3%) and very rare in lacrimal sac (1.4%). An increased incidence in the upper tarsal conjunctiva may be due to the presence of sulcus tarsalis, a linear groove, where the spore of foreign body gets lodged and is not washed off easily. Rippon [21] also states that in 90% of ocular infection, the palpebral conjunctiva is involved.

Shukla *et al.* [24] has categorically stated that all the patients belong to the lower economic strata. However, in this study the OR was found to occur in people belonging to different strata of socio-economic order and there was no preponderance of the disease in any community belonging to a particular caste or religion. Kannankutty *et al.* [8] stated that a member of one community is a much vulnerable to this disease as a member of another community.

The disease may occur after dust storms [11] and is common in dry dusty areas [21]. But, in this study, most of the patients hailed from wet areas. The KK district gets the benefit of both southwest and northeast monsoon. Average rainfall is more than 140 cm. Further, most of the patients gave a history of taking baths in local ponds. This habit has been attributed to be a primary factor in the causation of disease by several authors. But, so far, the method of transmission and the occurrence of the infective stage in the ponds have not been known.

Table 2. Anatomic sites of ocular rhinosporidiosis

Sites	No. of cases
Upper tarsal conjunctiva	137
Lower tarsal conjunctiva	63
Bulbar conjunctiva	16
Lacrimal sac	3
Total	219

Hence, it can be concluded that KK district, a hyperendemic area for nasal rhinosporidiosis [15] is also highly endemic for OR, mostly affecting young adolescents, frequently involving upper tarsal conjunctiva, equally affecting both sexes belonging to various socio-economic and religious groups.

Acknowledgement

The authors are thankful to the District Medical Officer, Staff of Animal Husbandry Department and Livestock Research and Development Centre, Nagercoil, Kanyakumari District for their help and facilities provided to collect the specimens. They also thank the Tamil Nadu Agricultural University, Coimbatore for the funds and facilities provided.

References

1. Agarwal PL, Bhargava S, Nagar CK, Nepali LK. Ocular rhinosporidiosis. *J Indian Med Ass* 1975; 64: 90-100.
2. Allen FRWK, Dave ML. The treatment of rhinosporidiosis in man based on the study of sixty cases. *Indian Med Gaz* 1936; 71: 376-95.
3. Bhaskar Reddi D. Rhinosporidiosis: a review based on the study of 156 biopsies. *Antiseptic* 1954; 51: 702-11.
4. Billore OP. Epidemiological survey of rhinosporidiosis in India. Paper presented in Asia-Pacific Congress on Ophthalmology. New Delhi Feb'85.
5. Chitravel V, Subramaniam S, Sundaram BH, Thirunellakantan S. Rhinosporidiosis in a South Indian village. *Sabouraudia* 1980; 18: 241.
6. Gupta K, Gupta AK, Jain SB. Rhinosporidiosis of Conjunctiva (a report of a case from Patiala) *J Postgr Med* 1984; 30: 119-20.

7. Jain SC, Darbari BS, Bhatnagar BS. Ocular rhinosporidiosis ENT Monthly 1968; 47: 380-2.
8. Kannankutty M, Sreedharan T, Mathew KT. Some observation on rhinosporidiosis. Am J Med Sci 1963; 246: 695-701.
9. Kameswaran S. Rhinosporidiosis in 'ENT Diseases in a tropical environment'. Higginbothams Ltd. Madras 1975; p. 53-70.
10. Karunaratnae WAE. The pathology of rhinosporidiosis. J Path Bact 1936, 42: 193-202.
11. Kaye H. A case of rhinosporidiosis on the eye. Br J Ophthalmol 1938; 22: 447-55.
12. Kirpatrick H. Rhinosporidiosis of lacrimal sac. Ophthalmoscope 1916; 14: 177.
13. Kumaresam M. Study of rhinosporidiosis in Tamil Nadu and Kerala. 'Stanscope'. Stanley Medical College, Madras 1982; p. 35-47.
14. Moral-Gracia JL. Rhinosporidiosis of conjunctiva. Indian Med Gaz 1954; 57: 81.
15. Moses JS, Shanmugham A. Epidemiological survey of rhinosporidiosis in man. A sample survey in a high school located in a hyperendemic area. Indian Vet J 1987; 64: 34-8.
16. Moses JS, Shanmugham A, Kingsly N, Vijayam JC, Balachandran C, Venkateswaran, Albert A. Epidemiological survey of rhinosporidiosis in Kanyakumari district of Tamil Nadu 1988; Mycopathologia 101: 177-9.
17. Natarajan S, Biswas J, Moses JS, Badrinath SS. Scleral staphyloma due to rhinosporidiosis of the conjunctiva. Nethralaya Insight 1985; 2: 8-9.
18. Newmayor TC. Bilateral rhinosporidiosis of the conjunctiva. Arch Ophthalmol 1964; 71: 379-81.
19. Peter HJ, DeBelly CG. Conjunctival polyps caused by *Rhinosporidium seeberi*. Report of a case. Am J Clin Path 1969; 51: 256-9.
20. Purandare NH. Rhinosporidiosis in Bombay. Indian J Med Sci 1953; 7: 602.
21. Rippon JW. 'Medical Mycology'. The pathogenic fungi and pathogenic actinomycetes. II Ed. WB Saunders Co London 1982.
22. Robertson MC. Conjunctival rhinosporidium. Annal Ophthalmol 1985; 17: 262-3.
23. Savino DF, Maryo CE. Conjunctival rhinosporidiosis. Light and electronmicroscopic study. Ophthalmol 1983; 90: 1482-9.
24. Shukla JM, Arora MM, Darbari BS, Arora MP. Ocular rhinosporidiosis. Proc All India Ophthalmol Soc 1970; 31: 133-7.
25. Suseela V, Subramaniam KS. Rhinosporidiosis and the eye: a study of 49 cases. Indian J Ophthalmol 1975; 23: 1-4.
26. Taralakshmi VV, Pankajalakshmi VV, Arumugam SS, Subramaniam S. Rhinosporidiosis: some observation. J Lab Sci 1977; 2: 46-51.
27. Wright RE. *Rhinosporidium kinealvi* of conjunctiva. Indian Med Gaz 1922; 57: 81.