



[Trends Parasitol.](#) Author manuscript; available in PMC 2019 Apr 1.

PMCID: PMC5879000

Published in final edited form as:

NIHMSID: NIHMS929580

[Trends Parasitol.](#) 2018 Apr; 34(4): 310–321.

PMID: [29305089](#)

Published online 2018 Jan 2. doi: [10.1016/j.pt.2017.12.004](#)

How do virus-mosquito interactions lead to viral emergence?

[Claudia Rückert](#) and [Gregory D. Ebel](#)

Department of Microbiology, Immunology and Pathology, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO 80523, USA

*Correspondence: claudia.rueckert@Colostate.edu and gregory.ebel@colostate.edu

[Copyright notice](#)

[Publisher's Disclaimer](#)

Abstract

Arboviruses such as West Nile, Zika, chikungunya, dengue and yellow fever viruses have become highly significant global pathogens through unexpected, explosive outbreaks. While the rapid progression and frequency of recent arbovirus outbreaks is associated with long-term changes in human behavior (globalization, urbanization, climate change), there are direct mosquito-virus interactions which drive shifts in host range and alter virus transmission. This review summarizes how virus-mosquito interactions are critical to their ability to become global pathogens at molecular, physiological, evolutionary and epidemiological scales. Integrated proactive approaches are required in order to effectively manage the emergence of mosquito-borne arboviruses, which appears likely to continue into the indefinite future.

Keywords: mosquitoes, arboviruses, virus emergence, virus evolution, virus-mosquito, interactions

Mosquito-borne viruses as the new global pathogens

Viral pathogens are major causes of morbidity and mortality among humans and animals. Efficient transmission of viruses between susceptible hosts is required in order for these agents to persist in nature and ultimately cause disease. Several mechanisms for this exist, and include direct contact, aerosol and sexual transmission, among others. A subset of viruses, termed arthropod-borne viruses (arboviruses) requires hematophagous arthropods, mainly mosquitoes and ticks, for transmission between vertebrates. In general, perpetuation of arboviruses requires vertebrate viremia so that arthropods acquire infectious virus along with nutrient-containing blood during feeding. Transmission of virus to a new host by an arthropod infected in this manner requires that this arthropod be a competent vector. In public health entomology, the term “vector competence” refers to the inherent ability of a particular arthropod to transmit a particular virus. In competent vectors, virus is acquired during feeding, undergoes replication in gut tissue, disseminates to secondary sites of replication, including the salivary glands and is ultimately released into the arthropod's salivary secretions, where it may be inoculated into the skin and cutaneous vasculature of the host during subsequent feeding ([Figure 1](#)). Arboviruses, therefore, are those viruses that have evolved an intimate association with both a vertebrate and arthropod host in order to perpetuate in nature.




Figure 1

Different tissue barriers determine vector competence in mosquitoes


An arbovirus is taken up by the mosquito during an infectious bloodmeal. The virus infects the midgut epithelium and replicates before it passes the basal lamina into the hemolymph and disseminates throughout the mosquito body. In order to be transmitted to the next host, the virus has to infect the salivary glands from where it can be released into the saliva and transmitted to the next host. Virus population genetic diversity is reduced stochastically as viruses pass through anatomical barriers to transmission, such as midgut infection and escape barriers, and salivary gland infection and escape barriers. Potential changes in virus populations that have passed through such bottlenecks are depicted as a change in color (increasingly dark blue). Through this genetic drift as well as positive selection in the mosquito, new genotypes may emerge.

Mosquitoes are the most important vectors of arboviruses[1], although many are maintained by ticks[2], phlebotomines[3] and other arthropods[4]. The global health burden of mosquito-borne viruses is immense. It is commonly estimated that 50 to 100 million cases of infection by dengue virus (DENV) serotypes 1-4 occur per year. Recent estimates placed the burden of DENV at 1.14 million disability-adjusted life-years in 2013[5]. Most of the individuals who are at risk of DENV infection are at risk of other arboviruses including yellow fever (YFV), chikungunya (CHIKV) and Zika (ZIKV) viruses which share the same mosquito vector, *Aedes aegypti*[6]. Additional arboviruses that burden the health of individuals living in or traveling to the tropics include Japanese encephalitis virus (JEV), Venezuelan equine encephalitis virus (VEEV), Mayaro virus (MAYV), o'nyong nyong virus (ONNV) and many others. Temperate regions also experience seasonal epidemics of arboviral disease caused by West Nile virus (WNV), La Crosse virus (LACV), eastern equine encephalitis virus (EEEV), Jamestown Canyon virus (JCV) and related viruses. Although quantitative estimates of the collective burden of mosquito-borne arboviruses on human health worldwide do not currently exist, it is clear that their burden is enormous, and increasing[5, 7, 8].

The geographic distribution of many arboviruses has expanded in recent decades[6, 9], resulting in infection of naïve populations and providing opportunities for new host-virus relationships to develop. For example, after incursions into Europe in the 1990s, WNV (genus *Flavivirus*) was introduced into the Americas in 1999 and rapidly spread from a small focus near New York City throughout the New World. Similarly, CHIKV (genus *Alphavirus*) spread from an African focus into Asia during the mid-2000s and was introduced into the Caribbean region in 2013[10, 11]. CHIKV is now endemic in the

Americas and has caused over one million infections, many of which result in debilitating arthralgia[10]. ZIKV (genus *Flavivirus*) has also emerged in recent years[12]. Following an expansion from an African focus into the Pacific islands, the virus was introduced into South America and has spread throughout most of the range of its *Ae. aegypti* vector[8]. ZIKV has caused notable disease among developing fetuses and unexpected neurological disease among adults[13]. WNV, CHIKV and ZIKV, along with DENV and YFV, underscore the emergence of mosquito-borne viruses as truly global pathogens. The combination of increased travel and trade has resulted in frequent exchange of pathogens and vectors across continents, such that the notion of “geographic diseases” is increasingly irrelevant. Coupled with the rapid growth of tropical megacities, these exchanges continue to result in explosive epidemics of pathogens transmitted by mosquito vectors that require the human footprint on the environment in order to survive, such as *Ae. aegypti* and *Culex quinquefasciatus*. The ongoing emergence of mosquito-borne viruses is occurring on a scale (geographic, economic and human) that is without precedent in human history[7, 8, 14].

How, then, do mosquito-virus relationships lead to the emergence of these global pathogens (Figure 2, Key Figure)? This review will examine the ways that mosquitoes influence the emergence of mosquito-borne viruses in order to provide perspectives on the history and future of this phenomenon. The first section examines two central concepts in public health entomology, vector competence and vectorial capacity, which outline basic mosquito-virus interactions and are key in understanding how mosquitoes impact virus emergence. These concepts are also required for readers to develop a basic understanding of the biology and epidemiology of arboviruses. The second section deals with new knowledge of the evolutionary relationships between viruses and their arthropod hosts. This section illuminates the complexities of arthropod-host interactions and how these can influence virus population biology and phenotype, sometimes leading to emergence. The third section provides a historical perspective on how mosquitoes influence arbovirus emergence by examining the cases of WNV, CHIKV and ZIKV. Finally, we provide perspectives on the future emergence of mosquito-borne viruses, highlighting emerging mosquito-borne viruses that have yet to capture the attention of the general population.



[Figure 2](#)

Key Figure. Summary of factors influencing arbovirus emergence

Cellular and molecular interactions such as RNAi drive virus diversification in the mosquito. Differences in mosquito vector competence and bottlenecks that the virus encounters during dissemination within a mosquito can result in further divergence of the virus population and drive virus evolution and emergence. Due to urbanization and deforestation humans and livestock are frequently in close proximity to mosquito vectors and vertebrate hosts maintaining viruses in sylvatic/enzootic transmission cycles. Human settlements may also bring along anthropophilic mosquitoes such as *Aedes aegypti* and *Aedes albopictus*, which may encounter viremic reservoir hosts, such as primates. Interactions between new vector mosquitoes and vertebrate hosts can drive arbovirus evolution and emergence. Due to intense travel and ubiquitous distribution of *Aedes* spp. mosquitoes, such spillover events may easily lead to the outbreak of a new global pathogen.

Understanding vector competence and vectorial capacity

Arbovirus emergence is driven by the hematophagous behavior of their arthropod vectors. This unique mode of transmission has important consequences for the ecology of arboviruses. There are two concepts that are central to our understanding of arbovirus transmission and epidemiology: **vector competence** (see Glossary) and **vectorial capacity**.

As described above, vector competence defines the ability of a particular arthropod to transmit a given virus. Vector competence has been studied extensively in mosquitoes, and it is determined by both genetic and non-genetic (e.g. environmental) factors. It may vary depending on mosquito species, local mosquito populations and even individual mosquitoes. Importantly, vector competence is a quantitative rather than a qualitative quantity: rarely do all mosquitoes of a given species or population transmit a given virus. Thus, vector competence is usually expressed as a proportion (e.g. 34% of *Culex tarsalis* mosquitoes transmit WNV after 10 days incubation). Vector competence of a specific mosquito population is also dependent on the virus, and even different isolates of the same virus species may result in changes in vector competence of a mosquito population [15, 16]. Generally, there are four major barriers that the virus must cross within the mosquito in order to be transmitted (reviewed in [17]). First, when a mosquito ingests an infectious bloodmeal, the virus must successfully infect and replicate in the midgut epithelial cells (Figure 1). The midgut is the first tissue in which virus-mosquito cell interactions occur that can shape the outcome of infection. Mosquitoes in which the virus cannot establish an infection have a midgut infection barrier (MIB). This can occur due to genetic factors, such as lacking expression of receptors on the cell surface, or non-genetic determinants such as microbiome density and composition [18-20]. However, once a virus has established midgut infection it must cross the basal lamina surrounding the midgut epithelium in order to disseminate throughout the rest of the

mosquito ([Figure 1](#)). When virus replication is limited to the midgut and dissemination does not occur, mosquitoes are said to have a strong midgut escape barrier (MEB). The basal lamina of the midgut presents a physical barrier for the virus and the thickness of the basal lamina has previously been linked to decreased dissemination of DENV-1 in different *Aedes albopictus* populations[21]. Furthermore, it has been shown that smaller, nutritionally deprived *Aedes triseriatus* mosquitoes have increased LACV dissemination rates compared to larger mosquitoes reared on a normal or rich larval diet[22, 23], which may also correlate with the development of a thick basal lamina. However, it has recently been shown that the uptake of a bloodmeal alters expression of specific enzymes in the mosquito midgut, including collagenases, which results in transient degradation and increased permissibility of the basal lamina, allowing CHIKV to disseminate[24, 25]. Viruses with slower replication rates such as DENV may not benefit as much from early transient degradation of the basal lamina following a bloodmeal. Another possible midgut escape route for arboviruses may be via tracheal or neuronal cells. Dong and colleagues[24] recently showed that CHIKV can infect tracheal cells connected to *Ae. aegypti* midguts. When the virus has disseminated from the midgut it replicates in other mosquito tissues including the fat body, hemocytes, nerve tissue and muscle tissue (depending on the virus) ultimately reaching the salivary glands, the next crucial anatomical barrier to infection, the ‘salivary gland infection barrier’ (SGIB). Upon salivary gland infection, the virus replicates and is deposited in the apical cavities of acinar cells in order to be expectorated with saliva ([Figure 1](#)). However, not all mosquitoes will be able to expectorate virus (for reasons yet unknown) and thus have a ‘salivary gland escape barrier’ (SGBE).

Other important mechanisms to limit virus replication throughout the mosquito body are mosquito antiviral immune responses (reviewed in [26, 27]). The most specific and potent mosquito antiviral defense is RNA interference (RNAi). During infection of a mosquito cell, viral dsRNA intermediates are recognized by the endonuclease Dicer2 and cleaved into 21 nt virus-derived small RNAs. These small RNAs are integrated into the RNA-induced silencing complex (RISC) and can target viral RNA for degradation. This sequence-specific response can be very efficient at controlling viral replication and many viruses have evolved mechanisms to antagonize or evade the RNAi response[28]. All mosquito antiviral responses may pose selective pressures on the virus, but RNAi is unique in its sequence specificity, which poses a direct evolutionary pressure on the viral genome. Mosquito barriers and antiviral responses together contribute to the overall phenotype of vector competence.

Vectorial capacity is a second critical concept that describes the basic reproductive rate of a vector-borne pathogen by a particular vector species/population ([Figure 3](#)), and highlights the power of mosquitoes as drivers of virus emergence. Factors which influence vectorial capacity are vector density with respect to host (m), the daily probability of the host being fed upon (a), vector competence (VC), the probability of daily survival (p) and the extrinsic incubation period (n). The factors that influence vectorial capacity and lead to arbovirus emergence have been reviewed and discussed extensively [29, 30]. Briefly, the most influential variables in the formula are the probability of daily mosquito survival and the extrinsic incubation period (EIP). The EIP refers to the amount of time that it takes for a virus to infect the midgut and disseminate to the saliva (i.e. the time between uptake of virus and ability to transmit). If the probability of mosquito daily survival is low and the EIP long, the likelihood of transmission is low. Extensive mosquito control programs that shorten the lifespan can thus efficiently reduce transmission. However, the EIP is affected both by environmental conditions such as temperature[31-33] as well as genetic factors[34-36] influencing vector competence of the mosquito population. Viruses may adapt to faster dissemination in susceptible mosquito species and thus shorten the EIP[37].




Figure 3

Vectorial capacity

The vectorial capacity formula describes the total number of future infectious bites arising from mosquitoes biting an individual infectious host on a single day. It consists of five factors: vector density with respect to host (*m*), the daily probability of the host being fed upon (*a*), vector competence of the mosquito population (*VC*), the probability of daily survival (*p*) and the extrinsic incubation period (*n*). None of these factors are constants, but variables which depend on both environmental influences as well as specific virus-mosquito interactions as indicated.

Another important variable is the probability of a particular host being fed upon, or the degree of host focus. This variable, *a* in the vectorial capacity equation, is a proportion (e.g. 80%, or 0.8 of mosquito X bloodmeals are taken from host Y) and is raised to the second power to reflect the need for susceptible hosts to be bitten twice in order for perpetuation to occur. Different mosquito species vary in their blood-feeding behavior and host preference. The mosquito species *Ae. aegypti*, for example, has adapted to life around humans, in particular urban areas of the tropics and subtropics, and mosquitoes of this species feed almost exclusively on human hosts and frequently feed indoors[38]. This behavior makes them extremely efficient vectors for viruses such as DENV, CHIKV and ZIKV, which replicate to high titers in human hosts. In contrast, *Ae. albopictus*, which can also serve as a vector for DENV, CHIKV and ZIKV, is more likely to feed outside, and while it is still anthropophilic, it is more of an opportunistic feeder[39]. Other mosquitoes, such as several species within the genus *Culex*, may feed on a human host, but tend to prefer birds[40]. This makes them efficient vectors for WNV and other zoonotic arboviruses. WNV infection of human or horse hosts is incidental and results in a ‘dead end’ for the virus due to inefficient replication and consequent low viremia. Thus, the competence of a given mosquito vector for an arbovirus may be irrelevant if that mosquito feeds only infrequently on susceptible vertebrate hosts.

Virus adaptation to new mosquito vectors clearly shapes the patterns and dynamics of arbovirus emergence. Generally, viruses with an *Ae. aegypti*-driven urban life cycle emerged from enzootic progenitors that circulated between non-human primates and sylvatic mosquito species such as *Aedes africanus* or *Aedes furcifer*[8, 41]. Adaptation to transmission by *Ae. aegypti* mosquitoes provided access to a new and abundant vertebrate host, humans. This has resulted in explosive outbreaks as seen before for ZIKV, which originated in a sylvatic cycle in Africa. Similarly, there are likely numerous viruses in current sylvatic viruses that may cause disease outbreaks in the future, some of which we already know, many of which we may be unaware of.

Mosquitoes as sources of virus genetic diversity

Arbovirus transmission requires active virus replication both in the arthropod vector and the vertebrate host. Arboviruses are thus subject to selective pressures from two evolutionary distant hosts. In vertebrates, where purifying selection is strong[42, 43], these pressures include innate and adaptive immune responses. Mosquitoes lack a classic adaptive immune system and do not produce interferon. However, Jak/STAT signaling and other signaling pathways, such as Toll, have been identified as antiviral responses against DENV[44, 45], WNV[46], Semliki Forest virus[47] and ONNV[18]. RNAi is the main antiviral defense in mosquitoes[27], which has direct consequences for virus intrahost evolution in mosquitoes (which can significantly impact virus emergence). Virus-derived small RNAs inhibit virus replication and translation by directly binding to complementary viral RNA[27]. The generation and perpetuation of novel sequences of viral RNA (i.e. containing mutations) may thus be beneficial for the virus within the mosquito because small RNAs will not match with perfect complementarity. Experimental evolution studies have confirmed the hypothesis that RNAi targeting by the arthropod vector leads to diversification of arbovirus genomes[48-50]. Rapid evolution and generation of a complex virus population could be particularly beneficial for the virus in order to reduce complementary binding, and because small RNAs have been shown to spread from cell-to-cell and may thus prepare neighboring cells for imminent virus infection[51]. However, these rapidly generated complex viral populations encounter bottlenecks within the mosquito when crossing the MEB, the SGIB, as well as during transmission of saliva itself[52, 53]. These bottlenecks vary in their size between different mosquito species and most likely virus-vector combinations. In *Cx. quinquefasciatus*, for example, WNV diversification is high, purifying selection is relatively weak, and bottlenecks are significant, resulting in high divergence of WNV populations in the saliva compared to the virus contained in the bloodmeal[52]. Different mosquito species may thus alter the virus population that is ultimately transmitted. Additionally, it has been shown that *Culex* mosquitoes may transmit unique WNV populations during each feeding episode[54], which is a further example of how mosquito infection promotes virus diversification. However, it is not known if the same phenomenon occurs with other viruses and mosquito species.

Rapid intrahost evolution of a mosquito-borne virus in the mosquito may result in the emergence of a new variant with a replication advantage in the vertebrate host or a new tissue tropism resulting in increased disease severity. For instance, VEEV emerges frequently from **enzootic** to **epizootic** transmission cycles with horses serving as amplifying hosts causing large outbreaks in both humans and horses. These epizootic strains appear to have only a few amino acid changes in the **envelope glycoprotein E2** in common compared to the enzootic strains[55]. There are many VEEV strains currently being transmitted in enzootic cycles, posing a constant threat in South and Central America[56].

While virus adaptation to replication in mosquitoes may reduce the ability of the virus to replicate to high titers in the mammalian host, such adaptations can also affect pathogenesis in undesirable ways. This can be seen in EEEV, for example, which does not reach high viremia in humans, but can cause severe to fatal encephalitis. It has recently been shown that the virus cannot replicate in myeloid cells of mice due to a miRNA target site in the 3'UTR for miRNA 142-3p, a myeloid cell specific miRNA[57]. However, while mutating this target sequence led to an increase in viremia and replication in peripheral tissues in mice, it resulted in a decrease in replication in mosquito cells and in infectivity of mosquitoes, suggesting that this sequence is important for replication in mosquitoes. This is one example of how adaptation to the vector may interfere with replication in a mammalian host.

Conversely, insect-specific viruses may have at some point developed mechanisms to inhibit mosquito antiviral defenses, which may also help the virus replicate efficiently in mammalian cells. One such mechanism is the generation of subgenomic flaviviral RNA (sfRNA) during replication. Flaviviruses have complex secondary structures in the 3'UTR, which stall the 3'→5' exonuclease XRN1, resulting

in the generation of sfRNAs. These sfRNAs inhibit mRNA degradation by sequestering XRN1, both in mosquito and mammalian cells. sfRNAs also suppress interferon signaling in vertebrate cells and are important for cytopathicity as well as pathogenicity in mice[58]. Production of sfRNAs is also important for replication and dissemination of flaviviruses in mosquitoes[59, 60], which may be due to both repression of XRN1 as well as their other function as suppressors of the mosquito RNAi machinery[61, 62]. However, it has recently been shown that DENV sfRNA patterns differ when virus is passaged in mosquito cells compared to mammalian cells[63], suggesting that while sfRNA generation is important in both cell types it is not optimized for one over the other. Overall, sfRNAs may have evolved in mosquitoes (similar yet less complex 3'UTR structures exist in insect-specific flaviviruses), but have an important role in vertebrate cells that is related to, but distinct from, its role in mosquitoes.

Virus-mosquito interactions have previously shaped virus emergence

In recent decades, WNV, CHIKV and ZIKV have emerged as global pathogens in explosive outbreaks that have caused significant human morbidity and mortality. The general phenomenon of formerly geographically restricted arboviruses emerging on a global scale has become common enough that it is perhaps best considered a new *status quo*. Several factors that have led to this phenomenon have been extensively discussed, including in this review. These include global increases in travel and trade, the rise of tropical megacities and the decline of public health programs to manage vector mosquitoes[9, 64, 65]. Further, as we have seen above, arboviruses have the capacity for explosive outbreaks due to their coadaptation to mosquitoes, which carry these agents between individuals with direction (i.e. as a vector). However, in each case (WNV, CHIKV and ZIKV) the virus adapted to local conditions during global spread, which maximized transmission potential (and health burden). These changes highlight the ways that the association of viruses with mosquitoes can lead to their emergence.

WNV was introduced into the Americas in 1999, most likely from the Middle East[66]. Molecular epidemiologic studies confirmed that the virus remained fairly homogeneous until approximately 2001, when a new virus genotype emerged in the central US. During the next two years, this genotype (somewhat erroneously called the “WN02” lineage because it was first recognized during 2002) outcompeted the introduced “NY99” genotype to the extent that the NY99 genotype seems to have become extinct. The most notable difference between this genotype and the introduced genotype was a mutation that resulted in a conservative Valine to Alanine change on the exposed surface of the WNV envelope glycoprotein[67, 68]. This mutation conferred faster transmission by *Culex pipiens* mosquitoes, important vectors of WNV in North America [37]. The reduced EIP of the WN02 genotype enhances vectorial capacity (see above) because mortality has less time to act on WN02-infected mosquitoes before they can transmit virus to a new host. This enhanced vectorial capacity of the WN02 genotype coincided with a massive WNV outbreak in N. America in 2002. Thus, specific virus-mosquito interactions between WNV and *Cx. pipiens* led to epidemiological changes (via vectorial capacity) that contributed to a significant and at the time unprecedented arbovirus outbreak in the US.

In contrast, CHIKV provides evidence of how an arbovirus can acquire the capacity for efficient transmission by a new vector mosquito species with disastrous consequences. The current global CHIKV epidemic began when the virus emerged in coastal Kenya around 2004. From there, it spread throughout the Indian Ocean region. During this process of emergence, the virus acquired a mutation to the coding sequence of the **envelope glycoprotein E1** that resulted in the substitution of a Valine for an Alanine at position 226 (A226V) of the protein[69]. This mutation rendered the virus more transmissible by the highly abundant *Ae. albopictus* mosquito[70] (i.e. increased the competence of *Ae. albopictus*). Since the competence of any given vector (again, see above) clearly impacts its reproductive rate, the acquisition of the A226V mutation conferred an advantage and spread rapidly

[70]. Initial adaptation to *Ae. albopictus* later allowed for mutations in CHIKV E2 to develop, which further enhanced infection of *Ae. albopictus* midguts[71]. Interestingly, the CHIKV that was introduced into the Caribbean in 2013 lacked the A226V mutation, and this mutation has not yet been detected in CHIKV from the Americas.

Most recently, ZIKV has emerged as a global pathogen in a manner similar to WNV and CHIKV before it. The consequences of ZIKV infection were not fully understood prior to this emergence, and are now notorious: microcephaly in developing fetuses and Guillain-barré syndrome in some infected adults[13]. Further, it is now clear that ZIKV may be transmitted sexually[72, 73]. Several mutations have been described in the literature that impact ZIKV phenotype in potentially significant ways. A mutation that changes an alanine at position 188 of the ZIKV NS1 protein to a valine has been associated with enhanced infectivity to *Ae. aegypti* mosquitoes via increased NS1 **antigenemia**[74]. Similarly, the serine to asparagine mutation at amino acid 139 (S139N) of the prM protein that appears to have arisen in around 2013 has been shown to contribute to pathogenesis in developing fetuses[75]. It is not yet clear what specific interactions could have produced either of these changes and how they may impact other important ZIKV phenotypes. This promises to be a fruitful area for future research.

Future arbovirus threats

The recent history of mosquito-borne virus emergence, coupled with what we know about the molecular and ecological interactions that facilitate their transmission, indicate that new explosive outbreaks of arbovirus disease are likely to occur in the future. Several arboviruses ([Table 1](#)) currently circulate at low levels in geographically limited settings that result in few human cases. While all of these viruses have the (theoretical) capacity to spread rapidly and cause outbreaks, some of them seem more likely than others to cause large scale epidemics in the near future.

Table 1

Selected arboviruses with currently limited geographic distribution and disease incidence

Genus	Virus	Suspected mosquito vector	Geographic distribution	Disease
<i>Alphavirus</i>	Mayaro virus (MAYV)	<i>Haemagogus</i> spp.	South and Central America	Febrile illness, arthralgia, myalgia
	Una virus (UNAV)	<i>Psorophora</i> spp.	South America	Febrile illness, arthralgia, myalgia
	Pixuna virus (PIXV)	unclear	South America	Febrile illness,
	Rio Negro virus (RNV)	unclear	South America	Febrile illness, myalgia
	Tonate virus (TONV)	unknown	French Guiana	Febrile illness, encephalitis
	Everglades virus (EVEV)	<i>Culex cedecei</i>	Florida	Fever, headache, myalgia
	Mucambo virus (MUCV)	<i>Culex</i> spp.	South America	Febrile illness, arthralgia, myalgia
	Trocara virus (TROCV)	<i>Aedes serratus</i>	South America	unknown
	o'nyong nyong virus (ONNV)	<i>Anopheles</i> spp.	sub-Saharan Africa	Febrile illness, arthralgia, myalgia
	Spondweni virus (SPONV)	<i>Aedes circumluteolus</i>	sub-Saharan Africa	Febrile illness, arthralgia, myalgia
<i>Flavivirus</i>	dengue virus type 5 (DENV-5)	<i>Aedes</i> spp.	South East Asia	High fever, arthralgia, myalgia
	Rocio virus (ROCV)	<i>Aedes</i> spp., <i>Psorophora</i> spp.	South America	Febrile illness, encephalitis
<i>Orthobunyavirus</i>	Oropouche virus (OROV)	mainly <i>Culicoides</i> biting midges*	South America	High fever, arthralgia, myalgia
<i>Phlebovirus</i>	Rift Valley fever virus (RVFV)	>40 species of mosquitoes	sub-Saharan Africa	Febrile illness, arthralgia, myalgia

[Open in a separate window](#)

*Oropouche virus is possibly also transmitted by *Aedes* spp. and *Coquillettidia* spp. mosquitoes

One of these viruses is MAYV, which has been circulating in parts of South and Central America since at least 1954 when it was first isolated in Trinidad[76]. Disease symptoms are similar to those of CHIKV infection, including fever, rash, myalgia and arthralgia. MAYV is suspected to be transmitted between non-human primates by vector mosquitoes of the genus *Haemagogus*; however, *Ae. aegypti*

and *Ae. albopictus* are capable of transmission in an experimental setting[77, 78]. While few human infections have been reported overall, a relatively large outbreak of 77 cases occurred in 2010 in Venezuela[79] and MAYV received some media attention due to a human case in Haiti in 2015[80]. Historically, MAYV infections occurred predominantly in northern South America, in regions bordering the Amazon Basin[81]. Since there was no history of travel in the Haitian case, infection probably occurred through local mosquito transmission. The patient was also coinfecting with DENV-1, possibly suggesting that infection occurred through the bite of an *Ae. aegypti* mosquito. Since there are no non-human primates on Hispaniola, this may indicate the presence of a local human-mosquito transmission cycle. However, MAYV has previously been isolated from a migrating bird in Colorado[82], suggesting another potential route of MAYV introduction into Haiti (and source of infection of local mosquitoes). Whether continuous local transmission was established remains unknown. While reports of potential recent recombinant MAYV strains[83] are of some concern, recombination of alphaviruses is likely a rare event. The only recombinant alphaviruses described are the three members of the Western equine encephalitis virus serocomplex, which are recombinants of a strain of EEEV and a Sindbis-like virus[84]. However, with a multitude of alphaviruses circulating in South and Central America and the recent addition of CHIKV, a small chance of recombination events remains. This could result in rapid emergence of new viruses with altered transmission phenotypes and pathogenesis. The risk of MAYV emergence has been discussed in detail in several recent reviews[85-87].

In addition, a little-known flavivirus of some concern is Spondweni virus (SPONV). SPONV is currently circulating solely in sub-Saharan Africa in sylvatic cycles involving zoophilic *Aedes* mosquitoes. Peri-domestic mosquitoes such as *Ae. aegypti*, *Ae. albopictus* and *Cx. quinquefasciatus* do not appear to be competent vectors[88]. However, as previously discussed, adaptation to an urban transmission cycle may be achieved by only one or a few mutations in the virus genome that increase the infection potential for *Ae. aegypti* or *Ae. albopictus* mosquitoes. Current research on SPONV is limited, but recent evidence shows its potential to be sexually transmitted among mice, albeit inefficiently[89]. Our current lack of understanding of SPONV pathogenesis and transmission warrants further investigation. The zoonotic potential of arboviruses currently circulating in Africa has been reviewed elsewhere[90].

Concluding remarks

Over the last decades, arboviruses have truly emerged as global pathogens. Viruses previously existing only in local transmission cycles in rural tropical settings are now distributed worldwide causing devastating disease outbreaks. This is largely due to the now global distribution of mosquito species with high vector potential, such as *Ae. aegypti* and *Ae. albopictus*, as well as human travel and population density in tropical megacities. Virus-mosquito interactions and mosquito adaptation to humans contribute significantly to arboviral emergence. Future research should be aimed at increasing our understanding of neglected arboviruses, novel surveillance methods and implementation of surveillance programs to recognize spillover events and arbovirus outbreaks early on (see Outstanding Questions). Both surveillance for arboviruses as well as invasive mosquito species can help with implementation of rapid vector control responses. As with all arboviruses, vector control will be crucial for the prevention and containment of future arbovirus emergence. The development of simple inexpensive diagnostic tests for a large selection of pathogens may also help identify and contain an outbreak. Moreover, the mosquito-virus interaction has facilitated the rapid emergence of several arboviruses, a pattern that is likely to continue, requiring an integrated approach to outbreak management.

Trends Box

- Arboviruses such as West Nile, Zika, chikungunya, dengue and yellow fever viruses have become highly significant global pathogens through unexpected, explosive outbreaks
- Recent advances in next generation sequencing have allowed rapid detection and identification of viruses, as well as tracing virus movement using phylogenetic analysis
- Evolutionary and modeling approaches, combined with experiments and new surveillance modalities are attempting to predict virus emergence
- Vector competence and virus evolution studies using selected mosquito species and neglected arboviruses are helping identify which virus may cause the next outbreak
- Integrated proactive approaches are required in order to effectively manage the emergence of mosquito-borne arboviruses, which appears likely to continue into the indefinite future.

Outstanding Questions Box

- Can the emergence of mosquito-borne arboviruses be predicted and preemptively managed?
- Which sylvatic/enzootic arboviruses can be maintained in a human/primate to mosquito transmission cycle?
- What vector control mechanisms and other preventative measures can specifically help prevent or contain future outbreaks?
- What are molecular mechanisms that underpin virus host shifts?
- To what extent can evolutionary approaches contribute to predictive efforts? Defining transmission and/or pathogenesis-altering mutations to a viral genome using experimental studies is, in some cases, highly tractable. However, frequently these studies may result in a so-called “gain-of-function” variants. Are the benefits of understanding the plasticity of viral genotypes and phenotypes worth the risk of generating gain-of-function variants?
- Is our society willing to invest in determining what arboviruses may emerge next?

Systems to monitor arbovirus emergence exist, but are difficult to implement at an appropriate scale. Further, many modern systems for monitoring health “events” such as arbovirus emergence rely on passive reporting and social media. Appropriate surveillance systems should focus on detecting agents that are early in the process of emergence and contain mechanisms to support critical research on an ongoing basis. Doing so requires financial commitment and political will.

Glossary

Antigenemia	describes levels of antigen in the blood. In the particular case of increased NS1 antigenemia mentioned here, it means that the more recent isolates of ZIKV result in increased NS1 levels in the blood of infected individuals
E1 and E2 glycoproteins	these are structural proteins incorporated into the envelope of the alphaviruses, such as CHIKV. E1 and E2 are important for attachment, entry and fusion events of the virus
Enzootic transmission cycle	describes the natural transmission cycle of a pathogen between wild animals. The term ‘enzootic’ is equivalent to the term ‘endemic’ as used for human diseases
Epizootic transmission cycle	refers to disease outbreaks among animal populations. The term ‘epizootic’ is equivalent to the term ‘epidemic’ as used for human diseases
Vector competence	describes the ability of a particular arthropod to transmit a specific pathogen
Vectorial capacity	describes the basic reproductive rate of a vector-borne pathogen by a particular vector species/population

Footnotes

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

References

1. Conway MJ, et al. Role of the Vector in Arbovirus Transmission. *Annu Rev Virol.* 2014;1:71–88. [[PubMed](#)] [[Google Scholar](#)]
2. Mansfield KL, et al. Emerging Tick-Borne Viruses in the Twenty-First Century. *Frontiers in cellular and infection microbiology.* 2017;7:298. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
3. Alkan C, et al. Sandfly-borne phleboviruses of Eurasia and Africa: epidemiology, genetic diversity, geographic range, control measures. *Antiviral Res.* 2013;100:54–74. [[PubMed](#)] [[Google Scholar](#)]
4. Carpenter S, et al. *Culicoides* biting midges, arboviruses and public health in Europe. *Antiviral Res.* 2013;100:102–113. [[PubMed](#)] [[Google Scholar](#)]
5. Stanaway JD, et al. The global burden of dengue: an analysis from the Global Burden of Disease Study 2013. *Lancet Infect Dis.* 2016;16:712–723. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
6. Leta S, et al. Global risk mapping for major diseases transmitted by *Aedes aegypti* and *Aedes albopictus*. *Int J Infect Dis* 2017 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
7. Gould E, et al. Emerging arboviruses: Why today? *One Health.* 2017;4:1–13. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

8. Mayer SV, et al. The emergence of arthropod-borne viral diseases: A global prospective on dengue, chikungunya and zika fevers. *Acta Trop.* 2017;166:155–163. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
9. Weaver SC. Urbanization and geographic expansion of zoonotic arboviral diseases: mechanisms and potential strategies for prevention. *Trends in microbiology.* 2013;21:360–363. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
10. Burt FJ, et al. Chikungunya virus: an update on the biology and pathogenesis of this emerging pathogen. *Lancet Infect Dis.* 2017;17:e107–e117. [[PubMed](#)] [[Google Scholar](#)]
11. Silva LA, Dermody TS. Chikungunya virus: epidemiology, replication, disease mechanisms, and prospective intervention strategies. *The Journal of clinical investigation.* 2017;127:737–749. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
12. Aliota MT, et al. Zika in the Americas, year 2: What have we learned? What gaps remain? A report from the Global Virus Network. *Antiviral Res.* 2017;144:223–246. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
13. Miner JJ, Diamond MS. Zika Virus Pathogenesis and Tissue Tropism. *Cell host & microbe.* 2017;21:134–142. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
14. Powers AM, Waterman SH. A decade of arboviral activity-Lessons learned from the trenches. *PLoS neglected tropical diseases.* 2017;11:e0005421. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
15. Weger-Lucarelli J, et al. Vector Competence of American Mosquitoes for Three Strains of Zika Virus. *PLoS neglected tropical diseases.* 2016;10:e0005101. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
16. Azar SR, et al. Differential Vector Competency of *Aedes albopictus* Populations from the Americas for Zika Virus. *The American journal of tropical medicine and hygiene.* 2017;97:330–339. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
17. Franz AW, et al. Tissue Barriers to Arbovirus Infection in Mosquitoes. *Viruses.* 2015;7:3741–3767. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
18. Carissimo G, et al. Antiviral immunity of *Anopheles gambiae* is highly compartmentalized, with distinct roles for RNA interference and gut microbiota. *Proceedings of the National Academy of Sciences of the United States of America.* 2015;112:E176–185. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
19. Hegde S, et al. The microbiome modulates arbovirus transmission in mosquitoes. *Curr Opin Virol.* 2015;15:97–102. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
20. Terradas G, McGraw EA. Wolbachia-mediated virus blocking in the mosquito vector *Aedes aegypti*. *Current opinion in insect science.* 2017;22:37–44. [[PubMed](#)] [[Google Scholar](#)]
21. Thomas RE, et al. Midgut basal lamina thickness and dengue-1 virus dissemination rates in laboratory strains of *Aedes albopictus* (*Diptera: Culicidae*) *Journal of medical entomology.* 1993;30:326–331. [[PubMed](#)] [[Google Scholar](#)]
22. Paulson SL, Hawley WA. Effect of body size on the vector competence of field and laboratory populations of *Aedes triseriatus* for La Crosse virus. *Journal of the American Mosquito Control Association.* 1991;7:170–175. [[PubMed](#)] [[Google Scholar](#)]

23. Grimstad PR, Walker ED. *Aedes triseriatus* (Diptera: Culicidae) and La Crosse virus. IV. Nutritional deprivation of larvae affects the adult barriers to infection and transmission. Journal of medical entomology. 1991;28:378–386. [[PubMed](#)] [[Google Scholar](#)]
24. Dong S, et al. Chikungunya virus dissemination from the midgut of *Aedes aegypti* is associated with temporal basal lamina degradation during bloodmeal digestion. PLoS neglected tropical diseases. 2017;11:e0005976. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
25. Dong S, et al. The midgut transcriptome of *Aedes aegypti* fed with saline or protein meals containing chikungunya virus reveals genes potentially involved in viral midgut escape. BMC genomics. 2017;18:382. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
26. Cheng G, et al. Mosquito Defense Strategies against Viral Infection. Trends Parasitol. 2016;32:177–186. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
27. Blair CD, Olson KE. The role of RNA interference (RNAi) in arbovirus-vector interactions. Viruses. 2015;7:820–843. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
28. O'Neal ST, et al. Mosquito-borne viruses and suppressors of invertebrate antiviral RNA silencing. Viruses. 2014;6:4314–4331. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
29. Kramer LD, Ciota AT. Dissecting vectorial capacity for mosquito-borne viruses. Curr Opin Virol. 2015;15:112–118. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
30. Liu-Helmerson J, et al. Vectorial capacity of *Aedes aegypti*: effects of temperature and implications for global dengue epidemic potential. PloS one. 2014;9:e89783. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
31. Danforth ME, et al. The Impact of Cycling Temperature on the Transmission of West Nile Virus. Journal of medical entomology. 2016;53:681–686. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
32. Mbaika S, et al. Vector competence of *Aedes aegypti* in transmitting Chikungunya virus: effects and implications of extrinsic incubation temperature on dissemination and infection rates. Virol J. 2016;13:114. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
33. Xiao FZ, et al. The effect of temperature on the extrinsic incubation period and infection rate of dengue virus serotype 2 infection in *Aedes albopictus*. Archives of virology. 2014;159:3053–3057. [[PubMed](#)] [[Google Scholar](#)]
34. Mercado-Curiel RF, et al. A dengue receptor as possible genetic marker of vector competence in *Aedes aegypti*. BMC microbiology. 2008;8:118. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
35. Bennett KE, et al. Quantitative trait loci that control dengue-2 virus dissemination in the mosquito *Aedes aegypti*. Genetics. 2005;170:185–194. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
36. Fansiri T, et al. Genetic mapping of specific interactions between *Aedes aegypti* mosquitoes and dengue viruses. PLoS genetics. 2013;9:e1003621. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
37. Moudy RM, et al. A newly emergent genotype of West Nile virus is transmitted earlier and more efficiently by *Culex* mosquitoes. The American journal of tropical medicine and hygiene. 2007;77:365–370. [[PubMed](#)] [[Google Scholar](#)]
38. Scott TW, Takken W. Feeding strategies of anthropophilic mosquitoes result in increased risk of pathogen transmission. Trends Parasitol. 2012;28:114–121. [[PubMed](#)] [[Google Scholar](#)]
39. Delatte H, et al. Blood-feeding behavior of *Aedes albopictus*, a vector of Chikungunya on La Reunion. Vector borne and zoonotic diseases. 2010;10:249–258. [[PubMed](#)] [[Google Scholar](#)]

40. Farajollahi A, et al. “Bird biting” mosquitoes and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology. Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in infectious diseases. 2011;11:1577–1585. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
41. Eastwood G, et al. Enzootic Circulation of Chikungunya Virus in East Africa: Serological Evidence in Nonhuman Kenyan Primates. The American journal of tropical medicine and hygiene 2017 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
42. Deardorff ER, et al. West Nile virus experimental evolution in vivo and the trade-off hypothesis. PLoS pathogens. 2011;7:e1002335. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
43. Coffey LL, et al. Arbovirus evolution in vivo is constrained by host alternation. Proceedings of the National Academy of Sciences of the United States of America. 2008;105:6970–6975. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
44. Ramirez JL, Dimopoulos G. The Toll immune signaling pathway control conserved anti-dengue defenses across diverse *Ae. aegypti* strains and against multiple dengue virus serotypes. Dev Comp Immunol. 2010;34:625–629. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
45. Souza-Neto JA, et al. An evolutionary conserved function of the JAK-STAT pathway in anti-dengue defense. Proceedings of the National Academy of Sciences of the United States of America. 2009;106:17841–17846. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
46. Paradkar PN, et al. Secreted Vago restricts West Nile virus infection in *Culex* mosquito cells by activating the Jak-STAT pathway. Proceedings of the National Academy of Sciences of the United States of America. 2012;109:18915–18920. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
47. Fragkoudis R, et al. Semliki Forest virus strongly reduces mosquito host defence signaling. Insect molecular biology. 2008;17:647–656. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
48. Brackney DE, et al. RNAi targeting of West Nile virus in mosquito midguts promotes virus diversification. PLoS pathogens. 2009;5:e1000502. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
49. Brackney DE, et al. Modulation of flavivirus population diversity by RNA interference. Journal of virology. 2015;89:4035–4039. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
50. Grubaugh ND, et al. Transmission bottlenecks and RNAi collectively influence tick-borne flavivirus evolution. Virus evolution. 2016;2:vev033. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
51. Attarzadeh-Yazdi G, et al. Cell-to-cell spread of the RNA interference response suppresses Semliki Forest virus (SFV) infection of mosquito cell cultures and cannot be antagonized by SFV. Journal of virology. 2009;83:5735–5748. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
52. Grubaugh ND, et al. Genetic Drift during Systemic Arbovirus Infection of Mosquito Vectors Leads to Decreased Relative Fitness during Host Switching. Cell host & microbe. 2016;19:481–492. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
53. Lequime S, et al. Genetic Drift, Purifying Selection and Vector Genotype Shape Dengue Virus Intra-host Genetic Diversity in Mosquitoes. PLoS genetics. 2016;12:e1006111. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
54. Grubaugh ND, et al. Mosquitoes Transmit Unique West Nile Virus Populations during Each Feeding Episode. Cell Rep. 2017;19:709–718. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

55. Brault AC, et al. Positively charged amino acid substitutions in the e2 envelope glycoprotein are associated with the emergence of venezuelan equine encephalitis virus. *Journal of virology*. 2002;76:1718–1730. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
56. Aguilar PV, et al. Endemic Venezuelan equine encephalitis in the Americas: hidden under the dengue umbrella. *Future virology*. 2011;6:721–740. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
57. Trobaugh DW, et al. RNA viruses can hijack vertebrate microRNAs to suppress innate immunity. *Nature*. 2014;506:245–248. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
58. Roby JA, et al. Noncoding subgenomic flavivirus RNA: multiple functions in West Nile virus pathogenesis and modulation of host responses. *Viruses*. 2014;6:404–427. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
59. Goertz GP, et al. Noncoding Subgenomic Flavivirus RNA Is Processed by the Mosquito RNA Interference Machinery and Determines West Nile Virus Transmission by *Culex pipiens* Mosquitoes. *Journal of virology*. 2016;90:10145–10159. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
60. Pompon J, et al. Dengue subgenomic flaviviral RNA disrupts immunity in mosquito salivary glands to increase virus transmission. *PLoS pathogens*. 2017;13:e1006535. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
61. Moon SL, et al. Flavivirus sfRNA suppresses antiviral RNA interference in cultured cells and mosquitoes and directly interacts with the RNAi machinery. *Virology*. 2015;485:322–329. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
62. Schnettler E, et al. Noncoding flavivirus RNA displays RNA interference suppressor activity in insect and mammalian cells. *Journal of virology*. 2012;86:13486–13500. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
63. Filomatori CV, et al. Dengue virus genomic variation associated with mosquito adaptation defines the pattern of viral non-coding RNAs and fitness in human cells. *PLoS pathogens*. 2017;13:e1006265. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
64. Zellweger RM, et al. Socioeconomic and environmental determinants of dengue transmission in an urban setting: An ecological study in Noumea, New Caledonia. *PLoS neglected tropical diseases*. 2017;11:e0005471. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
65. Matysiak A, Roess A. Interrelationship between Climatic, Ecologic, Social, and Cultural Determinants Affecting Dengue Emergence and Transmission in Puerto Rico and Their Implications for Zika Response. *Journal of tropical medicine*. 2017;2017:8947067. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
66. Lanciotti RS, et al. Origin of the West Nile virus responsible for an outbreak of encephalitis in the northeastern United States. *Science*. 1999;286:2333–2337. [[PubMed](#)] [[Google Scholar](#)]
67. Ebel GD, et al. Genetic and phenotypic variation of West Nile virus in New York, 2000-2003. *The American journal of tropical medicine and hygiene*. 2004;71:493–500. [[PubMed](#)] [[Google Scholar](#)]
68. Davis CT, et al. Phylogenetic analysis of North American West Nile virus isolates, 2001-2004: evidence for the emergence of a dominant genotype. *Virology*. 2005;342:252–265. [[PubMed](#)] [[Google Scholar](#)]
69. Schuffenecker I, et al. Genome microevolution of chikungunya viruses causing the Indian Ocean outbreak. *PLoS Med*. 2006;3:e263. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

70. Tsetsarkin KA, et al. A single mutation in chikungunya virus affects vector specificity and epidemic potential. *PLoS pathogens*. 2007;3:e201. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
71. Tsetsarkin KA, Weaver SC. Sequential adaptive mutations enhance efficient vector switching by Chikungunya virus and its epidemic emergence. *PLoS pathogens*. 2011;7:e1002412. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
72. Foy BD, et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA. *Emerging infectious diseases*. 2011;17:880–882. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
73. Magalhaes T, et al. Mosquito-borne and sexual transmission of Zika virus: Recent developments and future directions. *Virus Res* 2017 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
74. Liu Y, et al. Evolutionary enhancement of Zika virus infectivity in *Aedes aegypti* mosquitoes. *Nature*. 2017;545:482–486. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
75. Yuan L, et al. A single mutation in the prM protein of Zika virus contributes to fetal microcephaly. *Science* 2017 [[PubMed](#)] [[Google Scholar](#)]
76. Anderson CR, et al. Mayaro virus: a new human disease agent. II. Isolation from blood of patients in Trinidad, B.W.I. *The American journal of tropical medicine and hygiene*. 1957;6:1012–1016. [[PubMed](#)] [[Google Scholar](#)]
77. Long KC, et al. Experimental transmission of Mayaro virus by *Aedes aegypti*. *The American journal of tropical medicine and hygiene*. 2011;85:750–757. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
78. Smith GC, Francy DB. Laboratory studies of a Brazilian strain of *Aedes albopictus* as a potential vector of Mayaro and Oropouche viruses. *Journal of the American Mosquito Control Association*. 1991;7:89–93. [[PubMed](#)] [[Google Scholar](#)]
79. Auguste AJ, et al. Evolutionary and Ecological Characterization of Mayaro Virus Strains Isolated during an Outbreak, Venezuela, 2010. *Emerging infectious diseases*. 2015;21:1742–1750. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
80. Lednicky J, et al. Mayaro Virus in Child with Acute Febrile Illness, Haiti, 2015. *Emerging infectious diseases*. 2016;22:2000–2002. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
81. Tesh RB, et al. Mayaro virus disease: an emerging mosquito-borne zoonosis in tropical South America. *Clin Infect Dis*. 1999;28:67–73. [[PubMed](#)] [[Google Scholar](#)]
82. Calisher CH, et al. Isolation of Mayaro virus from a migrating bird captured in Louisiana in 1967. *Bull Pan Am Health Organ*. 1974;8:243–248. [[PubMed](#)] [[Google Scholar](#)]
83. Mavian C, et al. Emergence of recombinant Mayaro virus strains from the Amazon basin. *Sci Rep*. 2017;7:8718. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
84. Allison AB, et al. Evolutionary genetics and vector adaptation of recombinant viruses of the western equine encephalitis antigenic complex provides new insights into alphavirus diversity and host switching. *Virology*. 2015;474:154–162. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
85. Hotez PJ, Murray KO. Dengue, West Nile virus, chikungunya, Zika-and now Mayaro? *PLoS neglected tropical diseases*. 2017;11:e0005462. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
86. Esposito DLA, Fonseca B. Will Mayaro virus be responsible for the next outbreak of an arthropod-borne virus in Brazil? *Braz J Infect Dis*. 2017;21:540–544. [[PubMed](#)] [[Google Scholar](#)]

87. Mackay IM, Arden KE. Mayaro virus: a forest virus primed for a trip to the city? *Microbes Infect.* 2016;18:724–734. [[PubMed](#)] [[Google Scholar](#)]
88. Haddow AD, et al. Genetic Characterization of Spondweni and Zika Viruses and Susceptibility of Geographically Distinct Strains of *Aedes aegypti*, *Aedes albopictus* and *Culex quinquefasciatus* (*Diptera: Culicidae*) to Spondweni Virus. *PLoS neglected tropical diseases.* 2016;10:e0005083. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
89. McDonald EM, et al. Pathogenesis and sexual transmission of Spondweni and Zika viruses. *PLoS neglected tropical diseases.* 2017;11:e0005990. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
90. Venter M. Assessing the zoonotic potential of arboviruses of African origin. *Curr Opin Virol.* 2017;28:74–84. [[PubMed](#)] [[Google Scholar](#)]