

Evolutionary ecology of plant–microbe interactions: soil microbial structure alters selection on plant traits

Jennifer A. Lau^{1,2} and Jay T. Lennon^{1,3}

¹W.K. Kellogg Biological Station Michigan State University, 3700 East Gull Lake Drive, Hickory Corners, MI 49060, USA; ²Department of Plant Biology, Michigan State University, 166 Plant Biology Building, East Lansing, MI 48824, USA; ³Department of Microbiology and Molecular Genetics, Michigan State University, 2215 Biomedical Physical Sciences, East Lansing, MI 48824, USA

Summary

Author for correspondence:

Jennifer Lau

Tel: +1 269 671 2107

Email: jenlau@msu.edu

Received: 15 March 2011

Accepted: 8 May 2011

New Phytologist (2011) **192**: 215–224

doi: 10.1111/j.1469-8137.2011.03790.x

Key words: abiotic stress, *Brassica rapa*, drought stress, microbial ecology, natural selection, phenotypic selection analysis, plant–microbe interaction, species interactions.

- Below-ground microbial communities influence plant diversity, plant productivity, and plant community composition. Given these strong ecological effects, are interactions with below-ground microbes also important for understanding natural selection on plant traits?

- Here, we manipulated below-ground microbial communities and the soil moisture environment on replicated populations of *Brassica rapa* to examine how microbial community structure influences selection on plant traits and mediates plant responses to abiotic environmental stress.

- In soils with experimentally simplified microbial communities, plants were smaller, had reduced chlorophyll content, produced fewer flowers, and were less fecund when compared with plant populations grown in association with more complex soil microbial communities. Selection on plant growth and phenological traits also was stronger when plants were grown in simplified, less diverse soil microbial communities, and these effects typically were consistent across soil moisture treatments.

- Our results suggest that microbial community structure affects patterns of natural selection on plant traits. Thus, the below-ground microbial community can influence evolutionary processes, just as recent studies have demonstrated that microbial diversity can influence plant community and ecosystem processes.

Introduction

Below-ground microbial communities play a key role in determining the productivity, diversity and composition of plant communities (van der Heijden *et al.*, 2008; Kulmatiski *et al.*, 2008). For example, empirical studies have demonstrated that below-ground microbial diversity influences plant growth and productivity (Marschner & Rumberger, 2004; Hol *et al.*, 2010), nutrient availability (Bonkowski & Roy, 2005), and ecosystem functioning (Degens, 1998; Bradford *et al.*, 2002; Bell *et al.*, 2005). Additionally, many other studies illustrate the importance of plant–soil feedbacks, whereby shifts in microbial community composition feedback to affect plant coexistence and community composition (Bever, 2003; Reynolds *et al.*, 2003). While the effects of plant–soil microbe interactions on plant ecological dynamics appear to be widespread, little research has addressed how below-ground microbial communities influence plant evolutionary processes.

One way in which microbes could alter patterns of natural selection on plant populations is by ameliorating plant responses to abiotic environmental stress. Microbes have relatively short generation times, a high degree of genetic diversity, and the ability to evolve on ecologically relevant timescales. These attributes may allow for rapid changes in microbial community structure, which in turn may shape the way that plant populations respond to novel selective pressures in their environment. Several recent studies have demonstrated that local adaptation in plants is driven by genetic differentiation in closely associated microbes (reviewed in Rodriguez & Redman, 2008). For example, plant tolerance to high temperatures, salinity, and drought has been attributed to colonization by fungal endophytes (Rodriguez *et al.*, 2008), and heavy-metal tolerant strains of mycorrhiza have been shown to facilitate plant colonization of contaminated mine tailings (Adriaensen *et al.*, 2005). Similarly, plant growth-promoting bacteria can increase drought tolerance of their plant hosts by producing

hormones that induce the expression of plant genes associated with drought tolerance (e.g. Timmusk & Wagner, 1999; reviewed in Yang *et al.*, 2009).

These examples highlight the important role that a few key microbial taxa can have in promoting plant adaptation to stress. More often, however, plants form associations with complex, highly diverse, and potentially coevolved microbial communities. The net effect of such plant–microbe interactions can have implications for plant growth and fitness, carbon sequestration, and nutrient cycling (van der Heijden *et al.*, 2006, 2008; Kulmatiski *et al.*, 2008), and may influence plant community responses to global environmental changes (van der Putten *et al.*, 2009). Specifically, diverse microbial communities improve plant growth and vigor (Marschner & Rumberger, 2004) and help to maintain ecosystem functioning in stressful environments (Griffiths *et al.*, 2000). Therefore, diverse microbial communities may protect plants from the negative consequences of stress more effectively than less diverse microbial communities. For example, plant stress tolerance was heightened in the presence of multiple strains of plant growth-promoting bacteria, suggesting potential benefits of diverse microbial communities (Figueiredo *et al.*, 2008). However, soil food webs are notoriously complex, making it difficult to predict how changes in microbial community structure affect ecological processes. As a result, Bradford *et al.* (2002) advocate for empirical approaches for predicting how shifts in soil communities affect ecological functions. For similar reasons, empirical approaches may also be necessary for predicting effects of soil communities on plant evolutionary processes.

Here, we examine how below-ground microbial community structure influences selection on plant traits. We imposed treatments that altered the diversity and composition of the below-ground microbial community associated with replicated *Brassica rapa* populations growing in two soil moisture environments. These experiments allowed us to explore how biotic and abiotic factors influence plant phenotypes and selection on plant traits. We were particularly interested in determining whether below-ground microbes alter selection on plant traits by mitigating the effects of abiotic environmental stress on plant populations. Specifically, we tested whether more complex microbial communities reduce the negative consequences of drought stress on plant growth and reproduction; variation in microbial community complexity alters selection on plant growth and physiological traits; and more complex microbial communities minimize the selective impacts of drought stress on plant traits.

Materials and Methods

Experimental design

We conducted a factorial experiment to assess how differences in microbial community structure (simplified vs

complex) influence plant phenotypes, selection on plant traits, and plant responses to soil moisture (low vs high). We sowed 125 *Brassica rapa* L. (Brassicaceae) seeds into each of 16 mesocosms at 4 cm intervals on 30 May 2008. Seeds used in this experiment were obtained by haphazardly pollinating 200 *B. rapa* Wisconsin Fast Plants™ (standard stock lines, Wisconsin Fast Plants Program, University of Wisconsin, Madison, WI, USA). We used Wisconsin Fast Plants because this study is part of a multigeneration experiment that required plant taxa with short generation times. *B. rapa*, like most members of the Brassicaceae, is non-mycorrhizal; however, *B. rapa* still associates with a diverse rhizosphere community, and prior studies have shown that microbial community composition influences *Brassica oleraceae* growth and nutrient availability (e.g. Hol *et al.*, 2010). The mesocosms consisted of 76 l plastic pots (58 cm diameter, 34 cm height) filled with an autoclaved soil medium (1 part Baccto High Porosity Mix: 1 part perlite: 1 part vermiculate; 121°C, 15 PSI (pounds per square inch), 16 h). Each of the 16 populations was then randomly assigned to one of the replicate ($n = 4$) microbe \times soil moisture treatment combinations. In addition, we exposed eight mesocosms without plants to the soil moisture and microbe manipulations (i.e. microbe \times soil moisture) to test for the direct effects of moisture on microbial structure. Together, this resulted in a total of 24 experimental units.

We manipulated soil microbial community structure by inoculating each mesocosm with 3 l of either autoclaved (two cycles at 121°C, 15 PSI, 45 min) or unautoclaved field soil. The field soil was collected from an early successional field (Oshtemo Sandy Loam, heavily disturbed *c.* 25 yr ago (Burbank *et al.*, 1992)) dominated by many exotic grasses and forbs, including several species in the Brassicaceae, at the Kellogg Biological Station (KBS), Hickory Corners, MI, USA in May 2008. Although autoclaving can alter soil organic matter chemistry and the availability of plant nutrients (McNamara *et al.*, 2003), the inoculum was added to a large quantity of autoclaved potting media and comprised < 5% of the total mesocosm volume. Because the bulk of the soil used in each treatment experienced identical steam-sterilization regimes, we attribute any differences between microbe treatments to the addition of inocula that contained either live or dead soil microorganisms. Owing to the rapid growth rate of many microbial populations, the inoculum size in our study was sufficient to create microbial communities that differed in complexity (primarily diversity) by the end of the experiment (see the 'Results' section). Moreover, when sampled at the end of the experiment, estimates of bacterial richness in the treatments with live inoculum were similar to that observed in natural Michigan field soils (using identical fingerprinting methods) (mean \pm 1 SE = 58 ± 3.8 operational taxonomic units (OTUs), this study; 57.5 ± 2.3 , field study; JT Lennon, unpublished).

Bacterial and fungal communities in mesocosms receiving autoclaved inocula were less diverse compared with mesocosms receiving nonautoclaved inocula (see the 'Results' section). As such, we refer to the mesocosms with autoclaved inocula as 'simplified' and mesocosms with nonautoclaved inocula as 'complex'. Quantitative PCR (see the 'Microbial Structure' section) indicated that autoclaving the inocula did not reduce the total abundance of bacteria or fungi sampled at the end of the experiment (mean \pm 1 SE: \log_{10} bacteria – 'complex' = 6.62 ± 0.06 , 'simplified' = 6.50 ± 0.10 , $F_{1,17} = 1.14$, $P = 0.30$; \log_{10} fungi – 'complex' = 5.77 ± 0.11 , 'simplified' = 5.96 ± 0.18 , $F_{1,17} = 0.90$, $P = 0.36$), presumably because weedy, fast-growing microorganisms were able to recolonize the mesocosms. Such effects are probably common, even though investigators rarely characterize the effects of autoclaving on the abundance or composition of microbial communities.

All mesocosms were initially watered with reverse-osmosis (RO) water to stimulate germination of seeds. Shortly after emergence (*c.* 2 d), we ceased watering mesocosms in the low moisture treatment. Mesocosms in the high soil moisture treatment received enough water to ensure that the soil was consistently moist; this resulted in the application of 1000–1500 ml of RO water every other day for mesocosms with plants, and 500–1000 ml of RO water every other day for mesocosms without plants. On a single sampling date halfway through the experiment, we measured gravimetric water content from the surface soils as the loss of water mass after drying soil at 100°C for 24 h. We also measured volumetric soil moisture continuously in one mesocosm per treatment using Decagon (EC-TM) soil moisture probes with accompanying data loggers (EM50 ECH20) (Decagon Devices, Inc., Pullman, WA, USA).

Microbial structure

We characterized microbial community structure from mesocosm soils using culture-independent molecular methods. We collected and pooled five subsamples of soil from the top 5 cm of each mesocosm (approximate rooting depth) after plants had set seed. These soil samples were immediately stored at –20°C until later processing. We extracted genomic DNA from 1 g of soil sample using an UltraClean™ Soil DNA Isolation Kit, which was subsequently quantified using a Nanodrop spectrophotometer. We used this DNA to evaluate how microbial structure changed in response to our soil moisture and microbe treatments using quantitative PCR (qPCR) and terminal restriction fragment length polymorphism (T-RFLP).

First, we estimated the abundance of fungi and bacteria in our samples using qPCR. Briefly, qPCR assays consisted of 30 μ l reactions containing 1 μ l of DNA template, 0.5 μ l of each primer (10 μ mol l^{–1}), 14.5 μ l of DNase- and RNase-free H₂O, and 13.5 μ l of 5 Prime 2.5x Real-

MasterMix SYBR ROX (5 Prime, Inc. Gaithersburg, MD, USA). For fungi we used ITS1f (forward) and 5.8S (reverse) primers, and for bacteria we used Eub338 (forward) and Eub518 (reverse) primers (Fierer *et al.*, 2005). qPCR assays were performed with an Eppendorf Mastercycler realplex² system using previously reported thermal cycle conditions (Fierer *et al.*, 2005). We generated qPCR standards from a bacterial (*Micrococcus* sp.) and fungal (*Trichosporon* sp.) isolate using the TOPO TA Cloning Kit (Invitrogen). We extracted plasmids from transformed cells (Sambrook & Russell, 2001), and used M13 forward and reverse primers from the cloning kit to generate PCR products that we used for our standard curve, which captured a range of 10²–10⁷ copies μ l^{–1}. The coefficients of determination (r^2) for our assays ranged from 0.96 and 0.99, while amplification efficiencies fell between 0.93 and 0.99. Based on melting curve analyses, we found no evidence for primer dimers. We estimated fungal and bacterial abundance based on the estimated gene copy number from their respective standard curves and used these values to calculate fungal to bacteria ratios (F : B).

Secondly, we fingerprinted the soil microbial community using T-RFLP. For fungi, we PCR-amplified DNA using a fluorescently (FAM-6) labeled ITS1-F forward primer, an unlabeled ITS4 reverse primer, and the thermal cycler pattern described by Avis *et al.* (2006). For bacteria, we amplified DNA using a fluorescently (FAM-6) labeled 8F forward primer, an unlabeled 1492R reverse primer, and the thermal cycler pattern described in Lennon & Martiny (2008). We then digested 20–30 ng of the fluorescently labeled product for 24 h with five units of *Hae*III restriction enzyme. Lastly, we used a Qiagen nucleotide removal kit to clean up the enzyme-digested product. The T-RFLP samples were analyzed with an ABI Prism 3100 Genetic Analyzer at the Research Technology Support Facility (RTSF) at Michigan State University. We quantified the size of fluorescently labeled fragments in our samples by comparison to an internal ROX-labeled size standard (50–2000 base-pairs). From the resulting fragment profiles, we differentiated signal peaks from background noise using the methods of Jones & McMahon (2009). We then estimated the OTU richness for bacterial and fungal samples by calculating the sum of peaks that were present in the fragment profiles.

Plant measurements

We censused the central 64 plants in each mesocosm every other day and recorded flowering day and counted the number of leaves at flowering. After most plants had begun to flower, we measured chlorophyll content of the second or third leaf using a chlorophyll meter (SPAD 502; Spectrum Technologies, Inc., Plainfield, IL, USA) as an indicator of leaf nitrogen content and general plant vigor (Bullock & Anderson, 1998; Swiader & Moore, 2002). We estimated

specific leaf area (SLA = leaf area/dry mass) by weighing a leaf disk (0.28 cm^2) collected from the second true leaf produced on each plant (*c.* 3 wk post-germination) with a microbalance after the disk had been dried for 2 d at 65°C . This technique likely overestimated SLA because structural masses (e.g. midveins), which may increase leaf weights, were avoided; however, the SLA estimation technique should be consistent across treatments. This technique was necessary because plants produced less than four leaves on average and removing a whole leaf likely would have had severe effects on plant fitness. All open flowers in each mesocosm were hand-pollinated by other open flowers in that same mesocosm with a soft paintbrush three times per wk (brushes were cleaned with 30% isopropyl alcohol between mesocosms to prevent gene flow). When most plants in each mesocosm had ceased flowering and senesced, we harvested each individual, counted the number of seeds produced, and estimated total flower production by counting the number of flower scars on each plant. After removing seeds, we dried the above-ground biomass at 65°C for 2 d and weighed each individual to estimate above-ground biomass.

Statistical analyses

Microbial structure We used general linear models to determine the significance of main effects (soil moisture, plant presence, and microbe) and interactions (soil moisture \times plant presence and soil moisture \times microbe) on F : B ratios and OTU richness of the bacterial and fungal communities. When necessary, we used \log_{10} transformations to help meet the assumption of normality and equal variance. To analyze microbial composition data, we used the relative fluorescence output from the fragment profiles to generate separate sample \times OTU matrices for fungi and bacteria. We then tested for the main effects and interactions of our experimental manipulations with *adonis* (a permutation-based multivariate analysis of variance) with Bray–Curtis distance matrices using the *vegan* package in R statistical software (R Development Core Team, 2009).

Plant ecological and evolutionary effects We employed three methods to test how the soil moisture and microbe manipulations influenced plant populations: ANOVA to test for effects of microbe and soil moisture on the expression of plant traits; phenotypic selection analyses (Lande & Arnold, 1983) to estimate selection on each plant trait in each soil moisture \times microbe environment; and ANCOVA to test whether microbe and soil moisture treatments altered selection on plant traits.

We tested for the effects of microbe and soil moisture treatments on plant growth, phenological, and fitness traits with mixed-model ANOVA (PROC MIXED, SAS

Institute 2001). The microbe treatment, soil moisture treatment, and their interaction (microbe \times soil moisture) were included as fixed factors. Mesocosm nested within the microbe \times soil moisture interaction was included as a random factor. Response variables included flowering day (d since sowing), leaf number at flowering, chlorophyll content, SLA, above-ground biomass, lifetime flower production, and seed production. All response variables were natural log-transformed to help meet the assumptions of normality and equal variance.

We used phenotypic selection analyses (Lande & Arnold, 1983) to estimate selection differentials and gradients on flowering day, SLA, and biomass in each microbe \times soil moisture treatment. Phenotypic selection analyses use regression or multiple regression to quantify the relationship between traits and relative fitness. In all selection analyses, we used seed number as our estimate of lifetime female fitness. Selection differentials, which include both direct selection acting on the trait and indirect selection resulting from selection acting on tightly correlated traits, were estimated by performing separate linear regressions for each plant trait. Selection gradients estimate only direct selection on each trait by accounting for correlations with other traits included in the model and were estimated by performing multiple regressions that included all focal traits (flowering day, SLA and final biomass) as predictor variables and relative fitness (seed number) as the response variable. Traits were standardized within each mesocosm to a mean of zero and a variance of one (results from analyses of nonstandardized traits were qualitatively similar), and relative fitness was calculated based on mean seed number for each mesocosm.

We used analysis of covariance (ANCOVA) to determine whether the direction and/or intensity of selection differed between treatments (Wade & Kalisz, 1990). In the univariate (selection differential) analyses, each model included microbe treatment, soil moisture treatment, the standardized trait value, and all interactions as predictors of relative fitness (seed number). Mesocosm and the mesocosm \times standardized trait value interaction were included as random factors. Significant interactions between the microbe or soil moisture treatments and the plant trait indicate that selection differentials differed between treatments. A similar multiple regression analysis that included all traits and all interactions between traits and the microbe and soil moisture treatments as predictor variables was used to test for effects of microbe or soil moisture treatments on selection gradients. Additionally, because the residuals for seed number were not normally distributed (Shapiro–Wilk statistics ranged from 0.86 to 0.92, all $P < 0.0001$), we also used more conservative bootstrapping methods to estimate 95% confidence intervals for each selection differential and gradient in each treatment (%Boot Macro, SAS Institute, 2001).

Results

Soil moisture

Gravimetric water content (θ_d) was more than fourfold higher in the high moisture treatment than the low moisture treatment (1.16 ± 0.070 vs 0.27 ± 0.070 g H₂O/g soil; $F_{1,12} = 74.5$, $P < 0.0001$). Continuous measurements of soil moisture in a subset of mesocosms ($n = 1$ per treatment) revealed that these patterns were consistent throughout the duration of the experiment (Supporting Information, Fig. S1).

Microbial structure

Our experimental manipulations had strong effects on the structure of soil microbial communities. The mesocosms inoculated with autoclaved field soil ('simplified' microbe treatment) had microbial communities that were less diverse than the mesocosms that were inoculated with intact field soil ('complex' microbe treatment). OTU richness was significantly higher in the complex than in the simplified microbe treatment for both bacteria (ANOVA, $F_{1,16} = 17.3$, $P = 0.0007$, 58 ± 3.8 vs 45 ± 4.7 OTUs) and fungi (ANOVA, $F_{1,18} = 7.05$, $P = 0.016$, 24 ± 1.4 vs 16 ± 2.4 OTUs). In addition, the microbe manipulation altered bacterial composition (*adonis*, $F_{1,18} = 2.55$, $P = 0.003$), but when averaged across the plant presence and soil moisture treatments, we found no difference between fungal composition in the simplified and complex microbe treatments (*adonis*, $F_{1,18} = 1.22$, $P = 0.28$). Inoculation with autoclaved field soil tended to reduce the F : B ratio compared with inoculation with intact field soil, although this effect was not statistically significant (ANOVA, $F_{1,17} = 3.23$, $P = 0.09$; simplified = 0.16 ± 0.165 , complex = 0.34 ± 0.156).

The plant and soil moisture manipulations also affected soil microbial structure, and in some cases interacted with the microbial treatment. For example, fungal OTU richness was significantly greater in the high soil moisture treatments, but only in mesocosms containing *Brassica* (Fig. 1a) and in mesocosms assigned to the complex microbe treatment (Fig. 1b). The soil moisture manipulation also had a strong effect on fungal composition, but this effect was influenced by the presence of *Brassica* (plant presence \times soil moisture, *adonis*, $F_{1,18} = 3.64$, $P = 0.008$) and by the complexity of the microbial community (soil moisture \times microbe, *adonis*, $F_{1,18} = 2.45$, $P = 0.025$). By contrast, bacterial communities were less responsive to the soil moisture and plant presence manipulations. Both bacterial OTU richness (Fig. 1c) and bacterial composition were unaffected by plant presence (*adonis*, $F_{1,18} = 0.64$, $P = 0.93$) and soil moisture manipulations (*adonis*, $F_{1,18} = 1.25$, $P = 0.18$). The F : B ratio was significantly higher under low soil moisture than under high soil moisture (ANOVA,




Fig. 1 Interactive effects of soil moisture and plant (a, c) or microbe treatments (b, d) on soil microbial structure. Fungal operational taxonomic unit (OTU) richness in low (open bars) and high (closed bars) soil moisture mesocosms, with and without plants (i.e. *Brassica rapa*) (a) or in the simplified vs complex microbial communities (b). Bacterial OTU richness in mesocosms with or without plants (c) and with simplified vs complex microbial communities (d). Significant interactions between soil moisture and the plant (a, $F_{1,18} = 18.19$, $P = 0.0004$) and microbe manipulations (b, $F_{1,18} = 12.75$, $P = 0.0022$) were detected on fungal OTU richness. The values are marginal means and standard errors derived from ANOVA models.

$F_{1,17} = 5.80$, $P = 0.028$; low soil moisture = 0.530 ± 0.107 , high soil moisture = 0.214 ± 0.115).

Plant ecological effects

The microbe and soil moisture treatments both affected plant traits. On average, plants growing in soils with a complex microbial community were larger (e.g. greater leaf numbers at flowering and above-ground biomass), produced more flowers, and had a higher chlorophyll content than plants growing in the simplified microbe treatment (Table 1). Plants in the low soil moisture treatment had accelerated flowering and marginally reduced chlorophyll concentration (Table 1). Neither the soil moisture nor microbe treatment significantly affected SLA (Table 1).

We did not detect any statistically significant interactions between the soil moisture and microbe treatments on plant traits, but did observe a marginally nonsignificant microbe \times soil moisture interaction on seed production ($F_{1,12} = 3.50$, $P = 0.09$; Table 1). Specifically, simplification of the microbial community resulted in a 50% reduction in seed production for plants grown in low soil moisture treatments, but the microbe treatment had minimal effects on seed production for plants grown in high soil moisture treatments (Table 1).

Table 1 Influence of abiotic (soil moisture) and biotic (microbial complexity) manipulations on *Brassica rapa* plant traits

	Low moisture		High moisture		F-statistics			
	Simple microbial community	Complex microbial community	Simple microbial community	Complex microbial community	Soil moisture (S)	Microbe (M)	S × M	Mesocosm
Flowering day	20.70 ± 0.49	20.70 ± 0.49	21.86 ± 0.49	21.80 ± 0.49	5.37*	0.00	0.00	$\chi^2 = 27.2^{***}$
Leaf number	3.30 ± 0.12	3.68 ± 0.12	3.57 ± 0.12	3.71 ± 0.12	0.96	5.48*	0.85	$\chi^2 = 21.6^{***}$
SLA (cm ² g ⁻¹)	0.73 ± 0.03	0.70 ± 0.03	0.76 ± 0.03	0.74 ± 0.03	1.53	0.52	0.02	$\chi^2 = 33.1^{***}$
Chlorophyll content	9.68 ± 1.60	16.16 ± 1.53	13.88 ± 1.55	18.51 ± 1.53	3.39 ⁺	12.27**	1.00	$\chi^2 = 53.7^{***}$
Above-ground biomass (mg)	59.42 ± 14.00	98.41 ± 14.00	73.97 ± 14.00	113.6 ± 14.00	0.06	9.68**	0.02	$\chi^2 = 76.6^{***}$
Flower number	7.87 ± 1.22	8.60 ± 1.22	11.68 ± 1.22	12.12 ± 1.22	0.01	11.72**	0.01	$\chi^2 = 24.7^{***}$
Seed number	10.66 ± 4.32	23.16 ± 4.32	17.62 ± 4.32	14.82 ± 4.32	0.20	2.61	3.50 ⁺	$\chi^2 = 52.2^{***}$

SLA, specific leaf area.

Least-squares mean ± 1 SE for each trait in each treatment and F-statistics and chi-square values showing the effects of microbe (complex/simple) and soil moisture treatments on plant phenotypes. Values shown are per plant means. Values in bold are statistically significant at $P < 0.05$.

***, $P < 0.0001$; **, $P < 0.01$; *, $P < 0.05$; +, $P < 0.10$.

Selective effects of microbes and soil moisture

Selection favored plants that flowered earlier and accumulated more biomass in all treatments; however, below-ground microbial communities significantly altered the strength of selection on plant traits (Tables 2, 3). Selection for increased biomass was significantly stronger and selection for earlier flowering tended to be stronger for plants grown in simplified than in complex microbial communities (Table 2). We also detected evidence that soil moisture and microbe treatments interacted to affect selection on SLA. Low soil moisture resulted in significant selection for increased SLA, but only in the simplified soil microbe treatment (Table 2). Similar results were observed with the more conservative bootstrapping approach (Table 2). The selection gradient analysis revealed similar patterns (Table 3), although selection gradients on flowering time tended to be weaker than selection differentials, suggesting that a portion

of the observed selection on flowering time was likely indirect and the result of selection on correlated traits.

By measuring selection on replicated plant populations in each microbe × soil moisture treatment, we can definitively attribute observed differences in selection to differences in microbial community structure. Selection differentials were similar across mesocosms in a given treatment (coefficient of variation ranged from 0.24 to 0.45), and an analysis in which the selection differentials of each mesocosm were treated as independent response variables also revealed that selection on both flowering day and biomass tended to be weaker on populations grown with complex microbial communities compared with populations grown with simplified microbial communities (e.g. LSMean selection differential ± 1 SE: flowering day – complex -0.26 ± 0.05 , simplified -0.39 ± 0.05 , $F_{1,12} = 3.01$, $P = 0.11$; biomass – complex 0.59 ± 0.12 , simplified 0.97 ± 0.12 , $F_{1,12} = 5.35$, $P = 0.04$).

Table 2 Influence of abiotic (soil moisture) and biotic (microbial complexity) manipulations on selection differentials (univariate analyses) of *Brassica rapa*

	Low moisture		High moisture		F-statistics			
	Simple microbial community	Complex microbial community	Simple microbial community	Complex microbial community	Soil moisture (S)	Microbe (M)	S × M	Mesocosm
Flowering day	-0.33 (-0.43, -0.22)	-0.20 (-0.28, -0.11)	-0.44 (-0.59, -0.32)	-0.32 (-0.44, -0.21)	2.16	2.57	0.00	$\chi^2 = 0.1$
SLA	0.19 (0.04, 0.33)	-0.05 (-0.16, 0.07)	-0.08 (-0.27, 0.12)	0.15 (-0.02, 0.40)	0.16	0.00	5.83*	$\chi^2 = 0.5$
Above-ground biomass	0.83 (0.62, 0.97)	0.63 (0.50, 0.74)	1.08 (0.92, 1.27)	0.56 (0.36, 0.78)	0.44	7.58*	1.25	$\chi^2 = 33.2^{***}$

SLA, specific leaf area.

Values in brackets are bootstrapped 95% confidence intervals. Selection differentials that are significantly different from zero are shown in bold. F-statistics from the ANCOVA analyses testing the effects of soil moisture, microbe and microbe × soil moisture interactions on selection differentials also are shown.

***, $P < 0.0001$; *, $P < 0.05$.

Table 3 Influence of abiotic (soil moisture) and biotic (microbial complexity) manipulations on selection gradients (multivariate analysis) of *Brassica rapa*

	Low soil moisture		High soil moisture		F-statistics			
	Simple microbial community	Complex microbial community	Simple microbial community	Complex microbial community	Soil moisture (S)	Microbe (M)	S × M	Mesocosm
Flowering day	-0.13 (-0.25, -0.01)	-0.13 (-0.22, -0.04)	-0.14 (-0.33, 0.04)	-0.26 (-0.40, -0.10)	0.38	0.48	0.84	$\chi^2 = 0.3$
SLA	0.18 (0.07, 0.28)	0.01 (-0.09, 0.12)	0.13 (0.00, 0.26)	0.21 (0.05, 0.48)	1.26	0.56	3.95*	$\chi^2 = 0.0$
Above-ground biomass	0.78 (0.52, 0.95)	0.60 (0.47, 0.74)	1.06 (0.84, 1.30)	0.56 (0.38, 0.77)	0.80	5.92*	1.77	$\chi^2 = 21.9^{***}$

SLA, specific leaf area.

Values in parentheses are bootstrapped 95% confidence intervals. Selection gradients that are significantly different from zero are shown in bold. F-statistics from the ANCOVA analyses testing the effects of soil moisture, microbe and microbe × soil moisture interactions on selection gradients also are shown.

***, $P < 0.0001$; *, $P < 0.05$; +, $P < 0.10$.

Discussion

In this study, we quantified how below-ground microbial structure influenced plant growth and fitness traits; plant responses to abiotic stress (soil moisture); and selection on plant traits. Microbes did not modify plant phenotypic responses to soil moisture; however, microbial community structure altered plant growth and influenced the strength of selection on plant traits (above-ground biomass and SLA) and tended to alter selection on flowering day. Selection for increased above-ground biomass was 50% stronger in mesocosms containing a simplified microbial community compared with a more complex microbial community. Contrary to our predictions, we detected little evidence that microbes influenced selection by ameliorating drought stress. Instead, the influence of our microbial manipulation on selection on plant growth and phenological traits typically was consistent across soil moisture treatments, suggesting that soil communities can alter the strength of selection on plant traits independent of abiotic environmental conditions. Just as recent findings have shown that microbial diversity is important to plant productivity and ecosystem functioning (Marschner & Rumberger, 2004; Bell *et al.*, 2005), our study suggests that aspects of microbial diversity affect plant evolutionary processes. If our system is typical, effects of variation in below-ground microbial community composition on natural selection may be common and strong. Furthermore, although few studies manipulate both abiotic and biotic environmental variables, our study suggests that biotic selection agents can be equally or more important drivers of natural selection than abiotic selection agents.

Ecological consequences of plant–microbe interactions

Simplification of the microbial community led to reductions in below-ground bacterial and fungal richness, which

ultimately reduced plant growth, chlorophyll content (a trait correlated with plant nutrient status), and seed production. While several recent studies have shown how the presence, abundance, or diversity of specific classes of microbes, such as mycorrhizal mutualists or rhizobium symbionts, may influence plant productivity (van der Heijden *et al.*, 2006; Maherali & Klironomos, 2007; reviewed in van der Heijden *et al.*, 2008), very few studies have examined how the complexity of microbial communities influences above-ground productivity (but see Marschner & Rumberger, 2004; Hol *et al.*, 2010). In our study, the positive effects of complex microbial communities on plant performance may be the result of effects of microbial diversity on resource availability (Griffiths *et al.*, 2000; Loreau, 2001) or through the suppression of pathogenic microbes (Garbeva *et al.*, 2004). For example, in subsequent plant generations, we noted high incidences of disease in the simplified mesocosms and were able to recover viable populations of *Fusarium* sp. (a commonly pathogenic fungus) from mesocosms in the simplified microbe treatment, but not from the more diverse microbial communities in the complex microbe treatment. Interestingly, recent studies in other systems have convincingly shown how manipulation of the soil environment in ways that favor bacterial growth can limit the growth and abundance of *Fusarium*, presumably because of increases in *Fusarium*-antagonists, including antibiotic-producing bacteria (Perez *et al.*, 2008).

Compared with the microbe manipulation, soil moisture had rather limited effects on plant phenotypes. The only significant effect we detected was that plants in the low soil moisture treatment flowered *c.* 1 d earlier than plants in the high soil moisture mesocosms, which is consistent with other studies showing accelerated phenologies in a variety of plant taxa under drought stress (Aronson *et al.*, 1992; Wu *et al.*, 2010). Thus, both abiotic and biotic factors can influence the expression of plant phenotypes, but, perhaps not surprisingly, abiotic and biotic factors affected different

traits. In addition, we detected limited evidence that microbial community composition influenced plant responses to soil moisture; no statistically significant interactions between soil moisture and microbe treatments on any plant traits were detected.

While plant phenotypes showed minimal responses to the soil moisture treatments, the below-ground fungal community responded strongly to this manipulation. Culture-independent PCR-based techniques indicated that high soil moisture treatments decreased F : B ratios, increased fungal richness, and altered fungal community composition (relative abundances of OTUs). These results are consistent with several studies demonstrating that soil moisture can alter both total microbial biomass and microbial community structure (Schimel *et al.*, 1999, 2007; Gleeson *et al.*, 2008; see Blankinship *et al.*, 2011 for review). Bacteria community composition was less responsive to our soil moisture treatments, similar to findings from glasshouse experiments (Kassem *et al.*, 2008) and natural surveys (Clark *et al.*, 2009) but inconsistent with a recent field experiment documenting large negative effects of precipitation reductions on bacterial communities, but limited effects on fungi (Yuste *et al.*, 2011).

Natural selection and plant–microbe interactions

Microbial community structure altered selection on plant traits. The general trend was for the simplified microbial community to increase selection on plant growth and phenological traits compared with the complex microbial community (Tables 2, 3). This finding may be driven by reductions in fungal diversity in the simplified microbial treatments. Fungal OTU richness was typically negatively correlated with the strength of selection (selection differentials for flowering time, $r = -0.55$, $P < 0.03$; SLA, $r = -0.30$, $P = 0.26$; and biomass, $r = -0.47$, $P < 0.07$). Although it is possible that microbial diversity directly affects patterns of natural selection, given that plants were significantly larger when grown in the presence of a more complex microbial community, the observed microbe effects on selection may result from indirect effects that occur because microbial diversity influences nutrient availability. Regardless of the proximate mechanism, however, our results suggest that microbial diversity can affect selection on plant traits, just as recent studies have demonstrated that fungal diversity can affect plant productivity, plant diversity, and ecosystem functions (van der Heijden *et al.*, 1998; Jonsson *et al.*, 2001; Maherali & Klironomos, 2007; reviewed in van der Heijden *et al.*, 2008).

Given the lack of strong evidence that variation in microbial community composition mediates plant fitness responses to soil moisture, it is perhaps not surprising, that the effects of microbial communities on selection on plant growth and phenological traits typically were consistent

across soil moisture treatments. In this system, the effects of microbes on patterns of selection do not appear to result from microbes ameliorating selection imposed by drought stress.

The relative importance of biotic vs abiotic factors

Biotic agents are recognized as important evolutionary forces that contribute to the generation and maintenance of biodiversity (Ehrlich & Raven, 1964; Kursar *et al.*, 2009; see also Benton, 2010). Our results revealed that biotic agents (i.e. soil microbes) affected the expression of plant traits and patterns of selection more so than abiotic agents (i.e. soil moisture). Relatively few studies of natural selection simultaneously manipulate abiotic and biotic variables, so it is difficult to ascertain whether our results are part of a more general pattern. In fact, two recent studies show contrasting results: Stanton *et al.* (2004) found that patterns of selection on *Sinapis arvensis* differed more between two light environments (abiotic variable) than between two competition treatments (biotic), while Lau *et al.* (2010) detected virtually no effect of elevated atmospheric CO₂ concentrations (abiotic variable), but strong effects of competitors (biotic) on selection on *Arabidopsis thaliana*. We suspect that the relative importance of abiotic vs biotic factors on selection varies from system to system and likely depends on both the intensity of the biological interaction or abiotic stressor and the amount of genetic variation in traits mediating the interaction or the abiotic stress response.

Indirect effects and the potential for co-adaptation to a common abiotic environment

Given that both plants and below-ground microbial communities responded to the soil moisture treatments in this experiment and given that variation in microbial community composition can alter selection on plant traits, it is interesting to consider how plants and their below-ground microbes might respond in a tightly linked manner to variation in the abiotic environment. Feedbacks between above-ground plant and below-ground microbial communities are well documented (Bever, 2003; Kulmatiski *et al.*, 2008), and because multiple generations of both plants and microbes may face similar environmental conditions, plants and below-ground microbes could potentially co-adapt to soil moisture environments or other stressors. Although few studies have investigated evolutionary interactions between plants and their total below-ground community, Pregitzer *et al.* (2010) observed plant local adaptation to soils that differed in microbial structure and soil effects on the heritability of plant traits. By contrast, Wagner *et al.* (2011) found no evidence that plant populations were locally adapted to soil biota. In more tightly linked, pairwise symbiotic relationships, microbially mediated adaptive changes

have been documented for a wide variety of systems from aphids and their *Buchnera* endosymbionts (Zilber-Rosenberg & Rosenberg, 2008) to plants and their mycorrhiza mutualists (Johnson *et al.*, 2010). Moreover, recent studies on plant–rhizobia mutualisms suggest that specific plant–mutualist combinations are necessary for high fitness (Heath & Tiffin, 2007) and that genome \times genome epistatic interactions (*sensu* Wade, 2007) may be common. As a result, changes in both plants and their associated microbes may be necessary for strong adaptive responses to environmental change. Understanding how a focal plant population changes in response to abiotic environmental variation, therefore, also requires understanding how closely associated community members, such as below-ground microbes, respond to the same environmental variable and how these biotic interactors influence plant evolutionary processes.

Acknowledgements

We thank Sandy Erwin, Mark Hammond, Robert Snyder and Brent Lehmkuhl for substantial help with glasshouse and laboratory work, and K. Heath, J. Hoeksema, R. Lankau, S. Jones, S. Strauss and two anonymous reviewers for providing insightful comments on earlier versions of this manuscript. This is KBS publication #1562. We acknowledge support from the Rackham Research Endowment and the Michigan Agricultural Experiment Station (MAES). In addition, this project was supported by National Research Initiative grants (2006-35107-16725 (J.T.L.) and 2008-35107-04481 (J.T.L.)) from the USDA National Institute of Food and Agriculture.

References

- Adriaenssens K, Vralstad T, Noben JP, Vangronsveld J, Colpaert JV. 2005. Copper-adapted *Suillus luteus*, a symbiotic solution for pines colonizing Cu mine spoils. *Applied and Environmental Microbiology* 71: 7279–7284.
- Aronson J, Kigel J, Shmida A, Klein J. 1992. Adaptive phenology of desert and Mediterranean populations of annual plants grown with and without water-stress. *Oecologia* 89: 17–26.
- Avis PG, Dickie IA, Mueller GM. 2006. A 'dirty' business: testing the limitations of terminal restriction fragment length polymorphism (trFLP) analysis of soil fungi. *Molecular Ecology* 15: 873–882.
- Bell T, Newman JA, Silverman BW, Turner SL, Lilley AK. 2005. The contribution of species richness and composition to bacterial services. *Nature* 436: 1157–1160.
- Benton MJ. 2010. Evolutionary biology: new take on the red queen. *Nature* 463: 306–307.
- Bever JD. 2003. Soil community feedback and the coexistence of competitors: conceptual frameworks and empirical tests. *New Phytologist* 157: 465–473.
- Blankinship JC, Niklaus PA, Hungate BA. 2011. A meta-analysis of responses of soil biota to global change. *Oecologia* 165: 553–565.
- Bonkowski M, Roy J. 2005. Soil microbial diversity and soil functioning affect competition among grasses in experimental microcosms. *Oecologia* 143: 232–240.
- Bradford MA, Jones TH, Bardgett RD, Black HJ, Boag B, Bonkowski M, Cook R, Eggers T, Gange AC, Grayston SJ *et al.* 2002. Impacts of soil faunal community composition on model grassland ecosystems. *Science* 298: 615–618.
- Bullock DG, Anderson DS. 1998. Evaluation of the Minolta SPAD-502 chlorophyll meter for nitrogen management in corn. *Journal of Plant Nutrition* 21: 741–755.
- Burbank DH, Pregitzer KS, Gross KL. 1992. *Vegetation of the W.K. Kellogg biological station*. Michigan State University, Agricultural Experiment Station Research Report 510.
- Clark JS, Campbell JH, Grizzle H, Acosta-Martinez V, Zak JC. 2009. Soil microbial community response to drought and precipitation variability in the Chihuahuan Desert. *Microbial Ecology* 57: 248–260.
- Degens BP. 1998. Decreases in microbial functional diversity do not result in corresponding changes in decomposition under different moisture conditions. *Soil Biology & Biochemistry* 30: 1989–2000.
- Ehrlich PR, Raven PH. 1964. Butterflies and plants: a study in coevolution. *Evolution* 18: 586–608.
- Fierer N, Jackson JA, Vilgalys R, Jackson RB. 2005. Assessment of soil microbial community structure by use of taxon-specific quantitative PCR assays. *Applied and Environmental Microbiology* 71: 4117–4120.
- Figueiredo MVB, Burity HA, Martinez CR, Chanway CP. 2008. Alleviation of drought stress in the common bean (*Phaseolus vulgaris* L.) by co-inoculation with *Paenibacillus polymyxa* and *Rhizobium tropici*. *Applied Soil Ecology* 40: 182–188.
- Garbeva P, van Veen JA, van Elsas JD. 2004. Microbial diversity in soil: selection of microbial populations by plant and soil type and implications for disease suppressiveness. *Annual Review of Phytopathology* 42: 243–270.
- Gleeson DB, Herrmann AM, Livesley SJ, Murphy DV. 2008. Influence of water potential on nitrification and structure of nitrifying bacterial communities in semiarid soils. *Applied Soil Ecology* 40: 189–194.
- Griffiths BS, Ritz K, Bardgett RD, Cook R, Christensen S, Ekelund F, Sørensen SJ, Bååth E, Bloem J, de Ruiter PC *et al.* 2000. Ecosystem response of pasture soil communities to fumigation-induced microbial diversity reductions: an examination of the biodiversity–ecosystem function relationship. *Oikos* 90: 279–294.
- Heath KD, Tiffin P. 2007. Context dependence in the coevolution of plant and rhizobial mutualists. *Proceedings of the Royal Society B-Biological Sciences* 274: 1905–1912.
- van der Heijden MGA, Bakker R, Verwaal J, Scheublin TR, Rutten M, van Logtestijn R, Staehelin C. 2006. Symbiotic bacteria as a determinant of plant community structure and plant productivity in dune grassland. *FEMS Microbiology Ecology* 56: 178–187.
- van der Heijden MGA, Bardgett RD, van Straalen NM. 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology Letters* 11: 296–310.
- van der Heijden MGA, Klironomos JN, Ursic M, Moutoglou P, Streitwolf-Engel R, Boller T, Wiemken A, Sanders IR. 1998. Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* 396: 69–72.
- Hol WHG, de Boer W, Termorshuizen AJ, Meyer KM, Schneider JHM, van Dam NM, van Veen JA, van der Putten WH. 2010. Reduction of rare soil microbes modifies plant–herbivore interactions. *Ecology Letters* 13: 292–301.
- Johnson NC, Wilson GWT, Bowker MA, Wilson JA, Miller RM. 2010. Resource limitation is a driver of local adaptation in mycorrhizal symbioses. *Proceedings of the National Academy of Sciences, USA* 107: 2093–3098.
- Jones SE, McMahon KD. 2009. Species-sorting may explain an apparent minimal effect of immigration on freshwater bacterial community dynamics. *Environmental Microbiology* 11: 905–913.
- Jonsson LM, Nilsson MC, Wardle DA, Zackrisson O. 2001. Context dependent effects of ectomycorrhizal species richness on tree seedling productivity. *Oikos* 93: 353–364.

- Kassem II, Joshi P, Sigler V, Heckathorn S, Wang Q. 2008. Effect of elevated CO₂ and drought on soil microbial communities associated with *Andropogon gerardii*. *Journal of Integrative Plant Biology* 50: 1406–1415.
- Kulmatiski A, Beard KH, Stevens JR, Cobbold SM. 2008. Plant–soil feedbacks: a meta-analytical review. *Ecology Letters* 11: 980–992.
- Kursar TA, Dexter KG, Lokvam J, Pennington RT, Richardson JE, Weber MG, Murakami ET, Drake C, McGregor R, Coley PD. 2009. The evolution of antiherbivore defenses and their contribution to species coexistence in the tropical tree genus *Inga*. *Proceedings of the National Academy of Sciences, USA* 106: 18073–18078.
- Lande R, Arnold SJ. 1983. The measurement of selection on correlated characters. *Evolution* 37: 1210–1226.
- Lau JA, Shaw RG, Reich PB, Tiffin P. 2010. Species interactions in a changing environment: elevated CO₂ alters the ecological and potential evolutionary consequences of competition. *Evolutionary Ecology Research* 12: 435–455.
- Lennon JT, Martiny JBH. 2008. Rapid evolution buffers ecosystem impacts of viruses in a microbial food web. *Ecology Letters* 11: 1178–1188.
- Loreau M. 2001. Microbial diversity, producer–decomposer interactions and ecosystem processes: a theoretical model. *Proceedings of the Royal Society of London Series B-Biological Sciences* 268: 303–309.
- Maherali H, Klironomos JN. 2007. Influence of phylogeny on fungal community assembly and ecosystem functioning. *Science* 316: 1746–1748.
- Marschner P, Rumberger A. 2004. Rapid changes in the rhizosphere bacterial community structure during re-colonization of sterilized soil. *Biology and Fertility of Soils* 40: 1–6.
- McNamara NP, Black HIJ, Beresford NA, Parekh NR. 2003. Effects of acute gamma irradiation on chemical, physical and biological properties of soils. *Applied Soil Ecology* 24: 117–132.
- Perez C, Dill-Macky R, Kinkel LL. 2008. Management of soil microbial communities to enhance populations of *Fusarium graminearum*-antagonists in soil. *Plant and Soil* 302: 53–69.
- Pregitzer CC, Bailey JK, Hart SC, Schweitzer JA. 2010. Soils as agents of selection: feedbacks between plants and soils alter seedling survival and performance. *Evolutionary Ecology* 24: 1045–1059.
- van der Putten WH, Bardgett RD, de Ruiter PC, Hol WHG, Meyer KM, Bezemer TM, Bradford MA, Christensen S, Eppinga MB, Fukami T *et al.* 2009. Empirical and theoretical challenges in aboveground-belowground ecology. *Oecologia* 161: 1–14.
- R Development Core Team. 2009. *R: A language and environment for statistical computing, reference index version 2.8.1*. Vienna, Austria: R Foundation for Statistical Computing. URL <http://R-project.org>.
- Reynolds HL, Packer A, Bever JD, Clay K. 2003. Grassroots ecology: plant–microbe–soil interactions as drivers of plant community structure and dynamics. *Ecology* 84: 2281–2291.
- Rodriguez R, Redman R. 2008. More than 400 million years of evolution and some plants still can't make it on their own: plant stress tolerance via fungal symbiosis. *Journal of Experimental Botany* 59: 1109–1114.
- Rodriguez RJ, Henson J, van Volkenburgh E, Hoy M, Wright L, Beckwith F, Kim YO, Redman RS. 2008. Stress tolerance in plants via habitat-adapted symbiosis. *ISME Journal* 2: 404–416.
- Sambrook J, Russell DW. 2001. *Molecular cloning: a laboratory manual*. Cold Spring Harbor, NY, USA: Cold Spring Harbor Laboratory Press.
- SAS Institute. 2001. *SAS version 8.1*. SAS Institute: Cary, NC, USA.
- Schimel J, Balser TC, Wallenstein M. 2007. Microbial stress-response physiology and its implications for ecosystem function. *Ecology* 88: 1386–1394.
- Schimel JP, Gulledge JM, Clein-Curley JS, Lindstrom JE, Braddock JF. 1999. Moisture effects on microbial activity and community structure in decomposing birch litter in the Alaskan taiga. *Soil Biology & Biochemistry* 31: 831–838.
- Stanton ML, Thiede DA, Roy BA. 2004. Consequences of intraspecific competition and environmental variation for selection in the mustard *Sinapis arvensis*: contrasting ecological and evolutionary perspectives. *American Naturalist* 164: 736–752.
- Swader JM, Moore A. 2002. Spad-chlorophyll response to nitrogen fertilization and evaluation of nitrogen status in dryland and irrigated pumpkins. *Journal of Plant Nutrition* 25: 1089–1100.
- Timmusk S, Wagner EGH. 1999. The plant-growth-promoting rhizobacterium *Paenibacillus polymyxa* induces changes in *Arabidopsis thaliana* gene expression: a possible connection between biotic and abiotic stress responses. *Molecular Plant-Microbe Interactions* 12: 951–959.
- Wade MJ. 2007. The co-evolutionary genetics of ecological communities. *Nature Reviews Genetics* 8: 185–195.
- Wade MJ, Kalisz S. 1990. The causes of natural selection. *Evolution* 44: 1947–1955.
- Wagner V, Antunes PM, Ristow M, Lechner U, Hensen I. 2011. Prevailing negative soil biota effect and not evidence for local adaptation in a widespread Eurasian grass. *PLoS ONE* 6: e17580.
- Wu CA, Lowry DB, Nutter LI, Willis JH. 2010. Natural variation for drought-response traits in the *Mimulus guttatus* species complex. *Oecologia* 162: 23–33.
- Yang J, Kloepper JW, Ryu C-M. 2009. Rhizosphere bacteria help plants tolerate abiotic stress. *Trends in Plant Science* 14: 1–4.
- Yuste JC, Peñuelas J, Estiarte M, Garcia-Mas J, Mattana S, Ogaya R, Pujol M, Sardans J. 2011. Drought-resistant fungi control soil organic matter decomposition and its response to temperature. *Global Change Biology* 17: 1475–1486.
- Zilber-Rosenberg I, Rosenberg E. 2008. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiology Reviews* 32: 723–735.

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Time series showing volumetric water content in one representative mesocosm of each soil moisture treatment over the duration of the experiment.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.