

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/285929094>

Vectors and vector-borne diseases in Turkey

Article · January 2013

CITATIONS

27

READS

845

12 authors, including:



[Abdullah Inci](#)

Erciyes Üniversitesi

111 PUBLICATIONS 983 CITATIONS

[SEE PROFILE](#)



[Aydin Tuncbilek](#)

Erciyes Üniversitesi

54 PUBLICATIONS 403 CITATIONS

[SEE PROFILE](#)



[Ramazan Canhilal](#)

Erciyes Üniversitesi

41 PUBLICATIONS 316 CITATIONS

[SEE PROFILE](#)



[Mehmet Doganay](#)

Erciyes Üniversitesi

181 PUBLICATIONS 3,287 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Pesticide Residues in Turkish Foods and Agriculture, Uluslararası Atom Enerjisi Ajansı Teknik Yardım Projesi, IAEA-TC,TUR/5/015 [View project](#)



Ecology [View project](#)

Review / Derleme

Vectors and Vector-Borne Diseases in Turkey

Abdullah INCI, Suleyman YAZAR, Aydın S. TUNCBILEK, Ramazan CANHILAL, Mehmet DOGANAY, Levent AYDIN, Munir AKTAS, Zati VATANSEVER, Aykut OZDARENDELİ, Yusuf OZBEL, Alparslan YILDIRIM, Onder DUZLU

Vectors and Vector-borne Diseases Implementation and Research Center, Erciyes University, Kayseri, TURKEY.

Summary: In Turkey, arthropod-borne diseases such as human babesiosis, dirofilariasis, leishmaniasis, malaria, lymphatic filariasis, onchocerciasis, anthrax, bartonellosis, louse-borne relapsing fever, lyme disease, plague, trench fever, tularemia, epidemic typhus, tick-borne typhus, Crimean-Congo hemorrhagic fever, phlebovirus infections, tick-borne encephalitis and west nile virus in humans; animal trypanosomiasis, avian haemosporidiasis, babesiosis, canine hepatozoonosis, canine leishmaniasis, dipetalonemiasis, dirofilariasis, habronemiasis, onchocerciasis, parafilariasis, setariasis, theileriosis, thelaziasis, anaplasmosis, ehrlichiosis, feline infectious anemia, akabane, bluetongue, bovine ephemeral fever, equine infectious anemia, louping ill, west nile virus, colony collapse disorders (CCD) were reported in domestic animals and *Wolbachia* endobacteria infection was also reported in mosquitos so far. On the other hand, aphids (Aphididae), leafhoppers (Cicadellidae), white flies (Aleyrodidae), beetles (Coleoptera), trips (Thysanoptera), gal mites (Eriophyidae), and nematodes are also very important vectors of plant diseases in Turkey. Some of the most harmful plant viruses in the world such as tristeza, plum pox virus, and tomato ringspot viruses are also transmitted by arthropod vectors. In recent years, in spite of changes in ecological balance caused by abandoning agricultural land, use of natural areas to farm, release of rabbits, preventing water flooding, leaving the fields uncultivated, migration from urban to rural areas, hunting of wild animals, the dramatic drop in the number of farm animals, global warming and the strengthening of wild life was probably responsible for the increase of tick populations. The construction of ponds and dams, changes in riverbeds, drying out of swamps and marshes, dense ad-hoc and haphazard construction, changes in picnicking habits, wild irrigation in agriculture, and the use of unsuitable control measures by local authorities, resulted in an increase of flies. This led to an increase in infected vector populations and vector-borne diseases and a decrease in the potential for economic development. In conclusion, it is very important to initiate and develop integrated control programs targeting arthropod vectors, disease agents of human, animal and plants and natural reservoirs. These diseases have been caused high economic losses in Turkey, and therefore they should be further investigated.

Key words: Animal, human, plant, Turkey, vectors, vector-borne diseases,.

Türkiye’de Vektörler ve Vektörlerle Bulaştırılan Hastalıklar

Özet: Türkiye’de insanlarda babesiosis, dirofilariasis, şark çıbanı, sıtma, lenfatik filariasis, onchocerciasis, şarbon, bartonellosis, bitle bulaşan dönemli ateş (louse-borne relapsing fever), lyme, veba, siper ateşi, tularemia, epidemik tifüs, tick-borne tifüs, Kırım-Kongo kanamalı ateşi, phlebovirus enfeksiyonu, tick-borne encephalitis ve batı nil virüsü; evcil hayvanlarda trypanosomiasis, kanatlı haemosporidiasisi, babesiosis, köpek hepatozoonosisi, köpek leishmaniosis, dipetalonemiasis, dirofilariasis, yaz yarası, onchocerciasis, yaz kanaması, setariasis, theileriosis, thelaziasis, anaplasmosis, ehrlichiosis, kedi enfeksiyöz anemisi, akabane, mavidil, üçgün hastalığı, loupingill, enfeksiyöz at anemisi, arıların colony collapse hastalığı (CCD) ve sivrisineklerde *Wolbachia* endobakteri enfeksiyonu şimdiye kadar rapor edilen artropod kaynaklı hastalıklardır. Diğer taraftan, Türkiye’de yaprak bitleri (Aphididae), yaprak zararlıları (Cicadellidae), beyazsinekler (Aleyrodidae), kın kanatlılar (Coleoptera), kirpik kanatlılar (Thysanoptera), gal akarları (Eriophyidae) ve nematodlar da bitki hastalıklarının önemli vektörleridir. Tristeza, şarka, domates halkalı leke virüsleri gibi dünyadaki en zararlı bitki patojenlerinin de uygun vektörleri olmadan epidemi yapamadıkları bilinmektedir. Tarım alanlarının terk edilmesi, doğal alanların tarıma açılması, doğaya tavşan salınması, su taşkınlarının önlenmesi, yayla yasağı, av yasağı, tarlaların işlenmemesi, kırsal hayattan şehirlere göç, yırtıcı hayvanların öldürülmesi, çiftlik hayvanlarının sayısındaki dramatik düşüşe karşın yaban hayatının güçlenmesi ve küresel ısınma gibi son yıllarda ekolojik dengelerin değişmesine yol açan faktörler kene sayısında artışa yol açan olası nedenlerdir. Fizibilite yapılmadan gölet ve baraj inşası, nehir yataklarının değiştirilmesi, bataklık ve sazlıkların kurutulması, plansız ve gelişigüzel yoğun yapılaşma, piknik kültüründeki değişiklik, tarımda vahşi sulama, yerel yöneticilerin bilimsel yaklaşımdan yoksun, bilinçsiz vektör mücadelesi de sinek populasyonlarında artışa sebep olmuştur. Bu durum da vektör populasyonunda, vektörlerle bulaştırılan hastalıklarda artışa ve dolayısıyla ekonomik gelişmelerde azalmalara veya yavaşlamalara yol açmaktadır. Sonuç olarak, Türkiye’de insan, hayvan ve bitki sağlığını tehdit ederek ekonomik kayıplara yol açan vektörlere ve bu vektörlerin bulaştırdıkları hastalıklara karşı mücadele programları geliştirmek ve başlatmak, bu konuda daha ileri araştırmalar yapmak oldukça önemlidir.

Anahtar sözcükler: Bitki, hayvan, insan, Türkiye, vektörler, vektörlerle bulaşan hastalıklar.

Introduction

The majority of the described infections are vector-borne diseases and most of them are described as emerging, reemerging and zoonotic diseases in The World (92). In Turkey, several vector-borne diseases including equine infectious anemia have already been investigated until today. Currently, presence of these diseases such as human babesiosis, leishmaniasis, malaria, lymphatic filariasis, onchocerciasis, anthrax, bartonellosis, louse-borne relapsing fever, Lyme disease, plague, trench fever, tularemia, epidemic typhus, tick-borne typhus, Crimean-Congo hemorrhagic fever, phlebovirus Infections, tick-borne encephalitis and West Nile Virus in humans; animal trypanosomiasis, avian haemosporidiosis, babesiosis, canine hepatzoonosis, canine leishmaniasis, dipetalonemiasis, dirofilariosis, habronemiasis, onchocerciasis, parafilariosis, setariosis, theileriosis, thelaziasis, anaplasmosis, ehrlichiosis, feline infectious anemia, akabane, bluetongue, bovine ephemeral fever, equine infectious anemia, louping ill and West Nile virus in domestic animals and colony collapse disorders (CCD) in honeybee, *Wolbachia* endobacteria infection in mosquitoes have been recorded in Turkey. On the other hand, Aphids (Aphididae), leafhoppers (Cicadellidae), white flies (Aleyrodidae), beetles (Coleoptera), trips (Thysanoptera), gal mites (Eriophyidae), and nematodes have been also described as important vectors of plant diseases in Turkey. Some of the most harmful plant viruses in the world such as Tristeza, plum pox virus, tomato ringspot viruses are transmitted by vectors. In this article it was aimed to review vector-borne diseases which have already been investigated and described in Turkey.

Human babesiosis (HB): HB is an infectious disease caused by protozoa of the genus *Babesia*. *Babesia* species are transmitted by ticks. *Babesia divergens* and *Babesia microti* are the most significant etiologic agents of human babesiosis cases in Europe and in U.S.A. respectively. In Turkey, the seropositivity of *B. microti*, *B. divergens* and *B. bovis* was reported as 6.23% (149), 8% (65) and 18% (106) in human populations respectively.

Leishmaniasis: Zoonotic visceral leishmaniasis (ZVL) caused by *Leishmania infantum* and cutaneous leishmaniasis (CL) caused by *L. tropica* and *L. infantum* were reported in Turkey. These diseases are notifiable diseases in Turkey since long time. The ZVL has been seen in Aegean and Mediterranean regions endemically and reported sporadically in other regions. It was reported from 38 provinces and the population under the risk has been estimated around 45 million. The official number of cases was reported between 50 and 60 annually but most cases are underreported. It is consistent with Mediterranean type and mostly seen in infants. *Leishmania* strains were isolated from human/dogs in different regions and identified as *L.*

infantum MON-1 and MON-98 by multilocus enzyme electrophoresis (MLEE) also known as zymodeme analysis. Antimonials and Liposomal Amphotericin B are being used for the treatment as first choice (136, 142). CL was characterized by long-term nodulo-ulcerative lesions healing spontaneously with scarring and the disease has been seen in southeastern and East Mediterranean Regions endemically/epidemicly. It has been spreading from endemic regions to other regions because of the different epidemiological factors. CL with different clinical types was reported in 41 provinces but more than 90% of the total cases were concentrated in 5 provinces (Sanliurfa, Adana, Osmaniye, Mersin, Aydin) located in southeastern, east Mediterranean and western regions of Turkey. According to Ministry of Health's official records, 46.003 new cases were reported between 1990 and 2010. Among those cases, 96% of them were reported from Şanlıurfa, Adana, Osmaniye, Hatay, Diyarbakır, İçel and Kahramanmaraş provinces. Although 45% of the cases were notified from Şanlıurfa in the last 20 years, its ratio is currently decreasing while other regions' ratios have been showing an increasing trend. Antimonials (intralesional or systemic) are the first line drugs for treatment (70).

Human Malaria: In recent years, the number of malaria cases in Turkey and their active foci has decreased significantly. Although more than two million patients were treated in 1945, only four cases of *Plasmodium vivax* (relapsing cases) were reported in 2011. The first malaria control program was launched in 1925. Since 1971, the number of malaria cases in the Çukurova and Amikova plains began to increase, reaching epidemic proportions in 1976 and 1977, when 37,320 and 115,512 cases were reported, respectively. The situation deteriorated again, however, with over 34,000 cases in 1980 and 66,673 in 1983. Between 1990 and 1996 years, the malaria situation remained critical. The cases peaked to 84,321 in 1994. The situation began to improve only in 1997, when case numbers were cut in half from the previous year. Incidence of the disease was increased particularly in areas where the Southeastern Anatolia Project (GAP) irrigation program was being implemented (14).

Lymphatic Filariasis (Elephantiasis): Elephantiasis is caused by the nematodes such as *Wuchereria bancrofti*, *Brugia malayi* and *B. timori*. The adults of these round worms lodge in the lymphatic system and disrupt the immune system of the host. They live for 6-8 years and, during their life time, produce so many of microfilariae circulating system of the host. The disease is transmitted by mosquitoes belonging to the genera *Culex*, *Anopheles* and *Aedes*. Currently, about 1.3 billion people are under the risk potential in several countries and approximately 65% of those live in the South-East Asia Region, 30% in the African Region, and the

remainder in other tropical areas according to WHO records. In Turkey, a case was recorded from an 11-year-old girl presented with swelling in both legs from Kozan in 2006 (35).

Anthrax: Anthrax is an infection of humans and other mammals caused by the bacterium *Bacillus anthracis*. Although the vast majority of human cases are related directly from infected carcasses or the handling of contaminated products from morbid animals, the transmission of the disease has been demonstrated with wide variety of tabanid and mosquito species, and with stable-flies (*Stomoxys* spp) (44). The disease has wide distribution in the World, and is also an endemic zoonosis in Turkey. A total of 967 (464 from animals and 503 from humans) anthrax cases were reported from Eastern Turkey between 1992 and 2004 years (146). On the other hand only human anthrax cases have been recorded as 926 in Turkey between 1990 and 2007 (42).

Bartonellosis: Bartonellosis is caused by different *Bartonella* species, and is usually transmitted by hematophagous insects, such as sandflies (*Lutzomyia* spp), fleas, and lice and also by animal scratches and bites (120). Although the reports about this disease are limited in Turkey, the seropositivity rate of *Bartonella henselae* was recorded as 18.6% in cats (33), 6% in human blood donors (195) and 16.9% in kidney transplant patients (111), 22.2% in cattle breeders and veterinarians (161) while the seroprevalence of *B. vinsonii* subsp. *berkhoffii* was recorded as 6.6% in dogs (34) in Turkey.

Louse-Borne Relapsing Fever: Louse-borne relapsing fever is a disease of human caused by spirochaetes, *Borrelia recurrentis*, and can be disseminated by war, famine and other disasters. It was reported that during the first half of the 20th century, there were at least 50 million people with the infection and 10 million of them died in Europe, the Middle East and northern Africa. After the 1st World War, the disease spread and the epidemics extended notably into Turkey and Persia in 1920-1921 (183). After the 2nd World War, the infection rose again into the Middle East and it has continuing potential for epidemic spread of the infection from its current foci in some parts of Africa (181).

Lyme Disease: The disease is caused by *Borrelia burgdorferi* and transmitted via ixodid ticks especially *Ixodes ricinus*. The distribution of the disease is correlated the prevalence of hard ticks. The vector tick mainly occurs in moist habitats like woodlands, urban parks and also pastures in maritime climates. In Turkey, the seroprevalence of *B. burgdorferi* was found 17% in human in rural area of Samsun province (18) and 23.2% in dogs and 6% in horses (25).

Plague (Black Death): The disease is zoonotic and transmitted from rodent to rodent and from rodent to man via biting of fleas. Man can also be infected by direct

contact with infected animal tissues. Pneumonic plague can result from direct human to human transmission. The causative agent is the bacterium, *Yersinia pestis*. Urban plague describes the situation where plague circulating among wild rodents. The infection is maintained in the rat population by fleas, such as *Xenopsylla cheopis* (Asia, Africa Europe and the Americas), *Xenopsylla astia* (South-East Asia) and *Xenopsylla brasiliensis* (Africa, India and South America). Rarely, plague is spread directly from person to person by fleas, such as *Xenopsylla* species and so-called human flea, *Pulex irritans* (130). The first recorded appearance of the plague in Europe was at Messina, Sicily in Middle Ages (October, 1347). It was estimated that it arrived on trading ships that very likely came from the Black Sea, past Istanbul and through the Mediterranean. At that time, it was estimated that a quarter people of Europe killed from the Black Death. Within the last decade, human plague cases have been reported from countries in Africa, America and Asia. During the decade of the 1990s through 1996, there were 16,005 cases of plague and 1214 deaths (7.6%) reported to the World Health Organization (30). There is no official record about this disease in Turkey.

Trench Fever: The disease is caused by *Bartonella Quintana*, and was first detected during the First World War and the Second World War years and then it disappeared. Later it has been reported in homeless people in Mexico, North America and Europe. The agent of the infection is transmitted by human body louse, *P. humanus* (73). The disease has not been reported from Turkey by this date.

Tularemia: Tularemia is also another important vector-borne zoonotic disease in Turkey. The pathogenic agent of the disease is Gram-negative and facultative intracellular bacterium *Francisella tularensis*. The transmission of the infection can be disseminated by contact with infected animals or blood-sucking vectors such as ticks, mosquitos, and flies. Although many human cases of tularemia have been reported each year in Turkey, there is no record that reflects the status of this disease in animals and as well in the vectors. This disease has been initially reemerged in 1988 in Marmara Region of Turkey. Then, the infection has been spread all over the Turkey. A total of 4824 human cases with tularemia were recorded by Ministry of Health in Turkey between 2005 and 2011 years (43). Tularemia outbreak was initially observed in 2005. First case was diagnosed at the Department of Infectious Disease, Erciyes University Hospital in 2005. The patient came to the hospital from Sariz town which is from east part and 120 km far from the center of Kayseri. Following years, the cases spread all villages located at the northeast, east and southeast part of Kayseri. Tularemia is still an endemic disease in Kayseri and around (108).

Epidemic Typhus (louse-borne typhus): Epidemic typhus is a louse-borne rickettsial disease caused by *Rickettsia prowazekii* in humans. In the past, reported epidemic typhus cases were associated with wars and also human disasters. It was estimated that over 20 million people were infected *R. prowazekii* after the First World War (1914-1918) and almost 20% of the infected people died in Eastern Europe from the infection. It was also estimated that 7 million people died from epidemic typhus after the Second World War (1939-1945) and typhus cases still occur in war-torn countries today (74). Only one epidemic typhus case was reported from Turkey (170). After that there is no information about presence of the infection in Turkey by this time.

Tick-borne Typhus (Mediterranean Spotted Fever, MSF): The disease is caused by *Rickettsia conorii* and transmitted by dog tick *Rhipicephalus sanguineus*. Until today, a limited number tick typhus cases were reported from Turkey (127, 162, 186).

Crimean-Congo Hemorrhagic Fever (CCHF): CCHF is an acute viral and contagious infection in human (140). Wild and livestock animals serve as amplifiers for CCHF virus in field conditions. The infection is mainly transmitted by ixodid ticks especially *Hyalomma marginatum* in endemic areas. CCHF virus can also be transmitted to human through contact with the infected blood of viremic man and as well livestock animals. In Turkey, since 2002 about 6000 people became infected with the Crimean-Congo hemorrhagic fever and nearly 400 of them have died. Ticks like *H. marginatum*, which are common in Turkey, have been identified as the vectors of Crimean Congo hemorrhagic fever to humans and animals in Kayseri region. The disease is spreading throughout Turkey (180); however the conditions in Kayseri province and its surroundings are particularly favorable for its development and spread.

Phlebovirus Infections: Four serotypes of phleboviruses, sand fly Sicilian virus (SFSV), sand fly Cyprus virus (SFCV), sand fly Naples virus (SFNV), and Toscana virus (TOSV), have been circulating in the Mediterranean Basin including Turkey. Sand flies are the biological vector for phleboviruses and the virus can also be transmitted transovarially (39, 122). SFSV, SFNV, and other related viruses cause sand fly fever, also known as 'three-day fever' or 'pappataci fever' in human. TOSV also causes neurological disorders, such as meningitis, encephalitis and meningoencephalitis (39, 41). In Turkey, first reports were published in 1975 and 1976. The seroprevalence of SFSV was reported as 5.2% in Aegean region and 22% in Mediterranean region while SFNV was reported as 28.3% in Aegean region and 62% in Mediterranean Region (172). Later studies were also showed the activity of SFNV and TOSV in different areas (139, 163). Toscana virus exposure was also detected in blood donors in Central, North and South/Southeast Anatolia of Turkey (55) and its

association with central nervous system infections (54). Recently, the outbreaks in 3 different areas (Izmir, Ankara and Adana provinces) were happened in Turkey and new phlebovirus, Sand fly Fever Turkey Virus, was described (32). In 2011, phlebovirus RNA was also detected in 3 *Phlebotomus major s.l.* female specimens collected from Ankara province and characterized as Sand fly Fever Turkey Viruses (56).

Tick-borne Encephalitis (TBE): Tick-borne encephalitis is most important tick-transmitted disease in man and occurs in Europe and Asia. The pathogenic agent of the disease is tick-borne encephalitis virus, a member of the genus *Flavivirus* in the family *Flaviviridae*. Although TBE reports are very limited, the seropositivity of the disease was found as 10.5% in southeastern (52) and 1.43% in Central Anatolia (179) in Turkey.

West Nile Virus (WNV): WNV infection has a worldwide distribution. The infection is mainly transmitted by mosquitoes. The virus has been isolated from several mosquito species. Especially *Culex* species have been found infected with the virus. In Turkey, a few seroepidemiological studies about WNV infection have been carried out in humans and also in animal. The results of serological investigations were reported as 0.99% (53) and 16% (141) in man and between 1-37.7% in domestic animals (143). Recently some molecular epidemiological studies have been also performed on WNV infection and the first case was detected in horse in Central Anatolia, Turkey (144). Additionally, the infection was also investigated in mosquitos using molecular techniques (141).

Trypanosomiasis (Animal Trypanosomiasis): The disease is caused by belonging to the genus *Trypanosoma*. Although some *Trypanosoma* species (e.g. *Try. equiperdum*) are transmitted directly between hosts, the transmission of the infection is possible by blood-sucking insects. The disease principally affects domestic livestock animals in Africa, and is associated with severe production losses (102). In Turkey, this disease has not been reported except one sub clinical case *Trypanosoma theileri* infection associated with *Theileria annulata* infection in cattle in Ankara (81) by this date.

Avian Haemosporidiasis (Avian malaria): The disease is a mosquito-borne protozoan infection of wild birds and fowl. The infections are caused by a complex of more than fifty species of *Plasmodium*. In addition to *Plasmodium* species, infections with *Haemoproteus* and *Leucocytozoon* are also responsible for the avian haemosporidian diseases. Species belonging to *Culicoides* and *Simulium* are the known vectors of the latest species, respectively. The occurrence of the infections is differed depend on host range, geographical distribution, vector species and the virulence of the parasite. Avian haemosporidiasis begins when infective sporozoites are injected by a dipteran insect and are carried around the

body until they invade cells of the lymphoid-macrophage system in brain, spleen, kidney, lung and liver tissue where they undergo a phase of asexual multiplication (exoerythrocytic schizogony) resulting in the production of many uninucleate merozoites. The life cycle involves the sexual process and sporogony which occurs in the invertebrate host (vector) and the merogony and development of gametocytes which takes place in avian hosts (20, 75, 99, 134). Recently, presence of partial *cytb* gene of avian haemosporidian mt-DNA from *Culex pipiens*, *Aedes vexans*, *Culex theileri* and *Culiseta annulata* collected from Kayseri province has been reported for the first time in Turkey. In addition among phylogenetically analyzed isolates, 11 lineages belonging to *Plasmodium* were characterized in four phylogenetic groups whereas two lineages belonging to *Haemoproteus* were characterized in one phylogenetic group in the same study. On the other hand *Cx. pipiens* was found to be the major vector of avian *Plasmodium* in the study area (94). Regarding to *Haemoproteus* infections the first molecular detection and characterization of *Haemoproteus* spp. lineage in an owl from Mugla region was also reported in Turkey (192).

Babesiosis: Babesiosis [bovine babesiosis (BB, Redwater, tick fever), ovine babesiosis, equine babesiosis, canine babesiosis] is very important tick-borne disease of domesticated animals such as cattle, sheep, horses, dogs, and is caused by intraerythrocytic parasite of the genus *Babesia*. The main causative agents of BB are *Babesia bigemina* and *B. bovis* and the principle tick vectors of the disease are *Boophilus* species in tropical and subtropical climates, and the disease causes magnitude economical losses in livestock industry in the endemic areas. BB is also very important cattle disease, and causes great economic losses in Turkey as well as in the World (47). BB was first defined by microscopically in 1890 in Turkey (91). Following, some studies on BB were carried out by microscopic (50, 61, 77, 165), serological (36, 80, 85) and molecular studies (46-48, 90, 97, 187) in Turkey. The causative agents of BB were detected in ticks which were collected from cattle in Kayseri region (76). On the other hand some field studies were also performed on ovine babesiosis (37, 40, 83, 87, 156), equine babesiosis (82, 112) and canine babesiosis (96, 178).

Canine Hepatozoonosis: The disease is a tick-borne and protozoal disease of dogs. The causative agent is *Hepatozoon canis* which primarily infects haemolymphatic tissues and causes anemia and lethargy, and *Hepatozoon americanum* which localizes in muscular tissues and leads to severe myositis and lameness in dogs. Old World canine hepatozoonosis caused by *H. canis* is transmitted by *Rhipicephalus sanguineus* whereas American canine hepatozoonosis caused by *H. americanum* is transmitted by *Amblyoma maculatum*. The transmission of the both

infections to dogs occurs by oral ingestion of the infected ticks with the mature oocysts of protozoan parasites (23). *H. canis* was first detected in India in 1905 and then it has been reported from Europe, the Middle East, Africa and Far East while *H. americanum* was recorded from Texas in 1978 and since this date some cases associated with *H. americanum* have been reported from several states in U.S.A. In Turkey, *H. canis* infection was first detected in 1933 (174) and in last decade a few *H. canis* cases were also reported from several parts of Turkey (104, 110, 166).

Canine Leishmaniasis (CanL): CanL is another serious vector-borne disease of domesticated dogs and wild canids. Dogs can be infected with several *Leishmania* species, but the most important clinical picture of canine leishmaniasis is viscerocutaneous leishmaniasis, caused by *Leishmania infantum*. CanL mainly occurs in the Mediterranean Basin. The infection is also widespread in some countries in South and Central America. The infection (all forms) is transmitted by sand flies in the endemic regions (17). The infection is also prevalent in dogs in Mediterranean and Aegean Sea regions of Turkey. The epidemiology of leishmaniasis was first investigated in the Mediterranean Region (137). Following, some studies were also carried out for CanL using serological tests (38, 138) and molecular techniques (78, 131, 138) in various regions in Turkey.

Dipetalonemiasis (Acanthocheilonemiasis): Dipetalonemiasis is another arthropod-borne and helminthic infection of humans and animals caused by genus *Dipetalonema*. Although the disease is described as a rare disease, it has been recorded in some parts of the World and recently a clinical dipetalonemiasis case in a dromedary camel in Pakistan was reported (129). In Turkey, *D. reconditum* was first reported in 1966 in dogs in various parts of Turkey (51). Afterwards, canine filariasis was investigated around Istanbul and the microfilariae of *D. reconditum* was detected as an agent of canine filariasis in the research area (173).

Dirofilariosis (Canine Heartworm): Dirofilariosis is another vector-borne disease of dogs, caused by most important nematode parasite *Dirofilaria immitis*, and has a worldwide distribution in tropical, subtropical and temperate zones. *D. immitis* is transmitted by mosquito species including genera of *Culex*, *Aedes*, *Ochlerotatus*, *Anopheles*, *Armigeres* and *Mansonia* (15, 31, 147). Transmission of dirofilariosis depends on the presence of sufficient numbers of infected, microfilaremic dogs, susceptible mosquitoes, and a suitable climate to permit extrinsic incubation of parasite in the mosquito vector (123). In Turkey, the canine heartworm is widespread and was reported from dogs in several provinces. The prevalence of disease has been reported between 0.2% and 30.0% from several regions of Turkey (27, 79, 189, 190). Currently, the risk potential of *D. immitis* as a

problem for Turkey was also investigated using molecular analyzing techniques in potential mosquito vectors in Central Anatolia, according the results of the research, *Ae. vexans* was determined as the main potential vector of *D. immitis* in Kayseri (191). On the other hand, some dirofilariasis cases were also reported as orbital dirofilariasis (24, 66, 167), premaseteric (117) and foot nodules (116) in man.

Habronemiasis (Summer Sores): Habronemiasis is a dipteran- borne nematode infection of horses, caused by *Habronema muscae*, *H. majus* (synonym *H. microstoma*) and *Draschia megastoma* (formerly *H. megastoma*) species. The disease causes gastric, pulmonary, cutaneous lesions called "summer sores" (118) and ophthalmic (185) habronemiasis forms in infected horses. The infection is ubiquitous and it was reported from several countries in The World and the prevalence of habronemiasis in horses was recorded as 0.8% according to the necropsy findings (8) in Turkey.

Onchocerciasis: Onchocerciasis is very serious simuliid-borne disease of humans and animals which has three principal manifestations: subcutaneous nodule formation, dermatitis, and blindness. About 27 species of *Onchocerca* infect animals and onchocercal infections occur worldwide and very common in cattle, horses and donkeys. The accidental infection of humans by animal *Onchocerca* species (probably *O. gutturosa*) has been reported from North America, Europe and Japan (152). The causative agents of the disease are transmitted by black-flies of the genus *Simulium*. *Simulium* species feed on a variety of mammals and birds, and their painful bites cause considerable annoyance. The transmission of the disease associated with afflictions; first microfilariae ingested during feeding migrate to the flight muscles of the fly and moult twice over one week. They then migrate to the proboscis and develop into infective third-stage larvae which are transmitted to vertebrate hosts during feeding. In Turkey, one human onchocercosis case caused *O. caecutienuis* was reported for first time in 1976 (196). Following, bovine onchocercosis caused *O. armillata* was recorded in the southern part of Turkey, and the prevalence of the disease was also reported as 86% in over 1 year old and 18% in under 1 year old cattle (9). Recently, ocular onchocercosis case in a human was reported from Edirne province of Turkey and the nematode was identified as *O. lupi* based on its morphology and molecular phylogenetic analysis of partial *cox 1* and 12S ribosomal DNA genes (155). Recently, the risk potential of simuliid infestation as a vector for transmission of *Onchocerca* species to humans and also livestock animals in Central Kizilirmak basin (including along the river of Nevsehir area) in Turkey was also investigated using TaqMan probe based Real Time PCR assay. According the results of the study, 2 of the examined pools which were consisted from adult

simuliid specimens found to be positive for *Onchocerca* spp (98).

Parafilariasis (Summer Bleeding): Parafilariasis is another insect-borne infection of horses and cattle caused by nematods of the genus *Parafilaria*. The causative agents of the disease are *P. bovicola* and *P. multipapillosa* which live in subcutaneous tissues of infected animals and pierce the skin to release eggs in blood. The vectors of the infections are muscid flies, *Musca autumnalis*, *M. lusoria* and *M. xanthomelis* and the biting dipteran *Haemotobia atripalpis*. The transmission of nematod parasites are disseminated by vector insects while feeding blood from the lesions. Vector flies ingest eggs of the parasites together with blood, then the eggs hatch, larvae invade the tissues of the infected fly and develop to third-stage larvae (L3). L3 move to the mouthparts of the fly and enter the host at the refeeding time on a lesion or on the orbit of another host. Parafilariasis has a widespread distribution in the World (12). In Turkey, some clinical parafilariosis cases in cattle have been reported from Malatya area (169) until today.

Setariasis: Setariasis is a mosquito-borne disease of ungulates with the filarial worms of the genus *Setaria*. It was reported that, of the livestock bovinds and horses are natural hosts for *S. digitata* and *S. equine*, respectively whereas sheep, goats and horses are abnormal hosts for *S. digitata*. *S. digitata* is considered to be non-pathogenic in natural hosts while the transmission of the infective stage larvae to abnormal hosts (e.g. sheep) causes severe neuropathological defects (184). The causative agents of the infections in abnormal hosts are transmitted by mosquito species belonging to the genera *Aedes*, *Anopheles*, *Armigeres* and *Culex* in different regions of the World. When the infected mosquito feeds, the infective larvae of the parasite actively pass down the proboscis to be deposited on the host's skin, and then enter the host through the puncture caused by the mosquito bite. The infections occur often in Far East and Asia. For example in China, the prevalence of the disease in sheep and goats was reported as many as 30-40% (184). In Turkey, several *S. equina* infections in horses were reported (8, 29, 67). Recently the prevalence of *S. equina* was recorded as 15% in equids in Ankara (135).

Theileriosis: Theileriosis is a tick-borne protozoan disease of ruminants, equids and felids. The causative agents of infection are *Theileria* species. The natural transport of the infection from an infected animal to other susceptible animal is possible with trans-stadial transmission, depending on gametocytes in the blood of infected animals being ingested by the larvae or nymphs of the particular species of two or three host ixodid ticks (150, 154). The disease occurs in two forms such as malignant theileriosis and benign theileriosis in infected animals (95). The malignant theileriosis of cattle and

sheep causes morbidity and mortality in susceptible animals. European cattle breeds and their cross-bred are very susceptible to the malign bovine theileriosis and so considerable loss of productivity is seen due to infection in the endemic areas. Tropical theileriosis (Mediterranean coast fever, bovine malaria) is one of the malignant cattle theileriosis is caused *Theileria annulata*. The agent of the disease is transmitted by *Hyalomma* ticks (2, 159), and is widespread throughout Turkey (2, 21, 88, 158). The epidemiology of the disease has been also investigated deeply using microscopic (83), serologic (160), biochemical (22) and molecular (10, 45, 76, 97) methods in Turkey. The infection causes heavy damage to animals in the Develi, Yesilhisar and Incesu districts of Kayseri province as well as in the other endemic regions of Turkey (84, 86). In this region only, economic losses due to tropical theileriosis in cattle were estimated to be approximately USD 600,000 for two years (89). On the other hand ovine theileriosis was also researched by field observations and laboratory analysis (3, 11, 88, 93) but malign theileriosis of sheep has not been reported in Turkey up today. Meanwhile recently, a few molecular researches were also carried out on *Theileria equi* which causes equine theileriosis (112). Feline theileriosis has not been documented until today in Turkey.

Thelaziasis: Thelaziasis is a dipteran-borne and ocular infection of animals and humans, caused by spirurid nematods belonging to the genus *Thelazia* commonly known as eyeworms. The most pathogenic species of these eyeworms is *T. rhodesii* that infects cattle in many regions of The World (119). In Turkey, bovine thelaziasis caused by *T. rhodesii* reported as widespread and the prevalence of the parasite was reported as 22% (69) and 5.5% (171) in cattle and 1,2% in buffalo (68).

Anaplasmosis (Gall Sickness, Tropical Bovine Ehrlichiosis): Anaplasmosis or gall sickness is principally a tick-borne disease of cattle, sheep and wild ruminants that caused by intra-erythrocytic *Anaplasma* species. In addition the transmission of the disease is also possible iatrogenic and mechanical by blood sucking insects such as *Tabanus* spp. and *Stomoxys* spp. The disease has a worldwide distribution in tropical and sub tropical regions. Causative organisms of gall sickness are *Anaplasma marginale*, *A. centrale* and *A. caudatum* while the agent of anaplasmosis of leucocytes (Tropical Bovine Ehrlichiosis or mononuclear/agranulocytic anaplasmosis) is *Anaplasma (Ehrlichia) bovis*. *Anaplasma* species recently have been classified within the genogroup II of ehrlichial complex based on analyzing of 16S ribosomal RNA gene sequences (113, 153). In Turkey, bovine anaplasmosis cases have been reported clinically (145), microscopically (26), serologically (49) and molecularly (4). The molecular prevalence of causative agents of bovine anaplasmosis

was detected in the ticks which were collected from the cattle in Black Sea Cost of Turkey (5, 6). On the other hand additionally, blood parasites of dogs were investigated to determine the prevalence with molar techniques in Kayseri area. In the study, the prevalence of *A. phagocytophilum* was found to be 8% according to results detected by real time PCR analyses (96).

Ehrlichiosis: Ehrlichiosis is a tick-borne disease caused by *Ehrlichia* species in tribe Ehrlichieae. The causative agents of the disease are obligate intracellular Rickettsiales and infect granulocytes, agranulocytes or platelets of human, domesticated animals and also wild animals. Ehrlichiosis was recognized as a problem in the Old World before its presence in the New World was discovered. Dogs in Africa were to be found infected in 1930s, and tick-borne fever (TBF) was recognized in ruminants in Europe and the UK around 1950 (59). The causative agent of TBF, now known as *Ehrlichia phagocytophila*, was found to be transmitted by the tick *Ixodes ricinus*. Following the discovery of *E. chaffeensis* as a zoonotic agent, another human pathogen was recognized in North America. *Ehrlichia canis* is widely distributed in domestic dogs in The World, and is transmitted by *Rhipicephalus sanguineus*, called dog tick (153). In Turkey, limited studies were carried out for canine ehrlichiosis. In a study in Ankara, 12 dog blood samples were examined for *E. canis* infection using *E. canis*-specific PCR and three of them were found to be positive by PCR using *E. canis*-specific primers. One of the positive isolates (*E. canis* strain, Kutahya) was found identical to Venezuelan Dog Ehrlichia (VDE) and was closely related (99.9%) to strain of *E. canis*, Oklahoma (177). Following clinically first ehrlichiosis case was reported from a dog in Aydin province in 2007 (176). On the other hand the seroprevalence of *Anaplasma phagocytophilum* in humans was reported as 10.62% in Sinop and 5.77% in Tokat (64). In addition very soon the molecular prevalence of blood parasites of dogs in Kayseri region was investigated by highly specific TaqMan Real Time PCR technique for the first time. In the study, totally 224 (56.0%) out of the 400 dog blood samples were found to be infected with at least one parasite species in Kayseri province. *Ehrlichia canis* had the highest prevalence among with the rate of 15.0% according to results detected by real time PCR analyses. This rate was followed by *B. canis canis* (12%), *B. gibsoni* (9%), *A. phagocytophilum* (8%) and *N. caninum* (6%) (96).

Feline Infectious Anemia: Feline infectious anemia (FIA, haemobartonellosis) is one of most imported vector-borne diseases of cats. The infection is caused by *Mycoplasma haemofelis* when combined with feline leukemia virus. Cats become infected via biting of infected flea. Some clinical cases about the infection were reported in Turkey (19, 115, 175).

Akabane: Akabane (AKA) virus is a mosquito-borne and infects pregnant ruminants such as cattle, sheep and goats. The virus belongs to Bunyaviridae family and has a wide distribution in the World (125). In Turkey, the seroprevalence of AKA has been reported as 0.5% in sheep, 22% in cattle in Black Sea Region (Samsun, Sinop, Ordu, Amasya and Tokat provinces) (7), as 0.14% in cattle in Thrace (105) in Turkey.

Bluetongue Virus: Disease with bluetongue virus (BTV) is an insect-borne disease and causes non contagious infections in domestic ruminants particularly in sheep. The virus has worldwide distribution. Several bluetongue outbreaks were reported from Aegean, Mediterranean Sea and Thrace regions of Turkey between 1977 and 2000 years; BTV-4, BTV-9 and BTV-16 serotypes of Bluetongue virus were isolated; BTV-4 of them was attenuated and then used successfully as vaccine to control of the disease in vaccination programs in the country (58). In this period, some epidemiological investigations were also performed on seroprevalence of the disease and on the vector of the agent. The epidemiological studies showed that the vector of the BTV is *Cluicoides imicola* (100); the seroprevalence of disease were recorded as 73.54% cattle in Thrace (105), 3% in sheep, 11% in cattle in Middle Black Sea Region (7) in Turkey.

Bovine Ephemeral Fever (BEF): Bovine ephemeral fever is another vector-borne disease of cattle and water buffalo. The infection is caused by BEF virus and occurs in virtually all of Africa, the Middle East and Asia. BEF epizootics occur in summer and autumn and have a strong association with recent rainfall and BEF virus has been isolated from a pool of *Culicoides* species in several countries (126). In Turkey, the seroprevalence of Ephemeral fever virus in cattle was reported between 2.5% and 15.3% in Thrace (105) and 13.5% in Middle Black Sea Region (7).

Equine Infectious Anemia (Swamp Fever, EIA): EIA is a viral infection of equids. The causative agent is a lentivirus, and transmitted by biting flies. Although a lot of seroepidemiological studies were carried out for EIA in the World, the seropositivity has not been reported in equids in Turkey until today (121).

Louping-ill (Infectious Encephalomyelitis of Sheep, LI): LI is a tick-transmitted and acute viral disease of mainly sheep/goats, but it can be recognized rarely in cattle and horses. LI virus characteristically causes an encephalomyelitic disease of sheep. The occurrence of LI infection is closely related to the distribution of the primary vector tick, *Ixodes ricinus*. A specific acute encephalomyelitis was recognized in sheep in north western part (Gebze and Kırklareli areas) of Turkey and the isolated agent was defined as member of the Louping ill/Russian Spring Summer Virus complex and then the isolated virus was described as Turkish sheep

encephalitis (TSE) virus and coded as TTE80 strain (71). Later TTE80 strain was also characterized as a subgroup in tick-borne viruses of Flaviviridae family by antigenic, pathogenic and molecular analyses (60).

Colony Collapse Disorders (CCD): Honey bees (*Apis mellifera*) are the most important insects both economically valuable and as a pollinators of agricultural crops worldwide. These economically valuable insects are affected from so many pathogens such as bacterium, virus, fungus, parasite and also some predators including man. The most of these pathogens cause severe diseases and very serious damages in honey bee colonies. CCD is associated several diseases. It was characterized with the severity and unusual colony declines in honeybee colonies, and was first described in 2006 in U.S.A. Recently, CCD was also reported as the main factor of the huge economic losses in beekeeping units all around the World (101). In Turkey, a few studies were carried out on the role of the viruses [e.g. Deformed Wing Virus (DWV)] which cause CCD and their transmission by "Varroa Mites", *Varroa destructor*. The vectoral role of *V. destructor* for DWV was determined in the honey bee colonies with CCD in Hatay. The vectoral role of varroa for deformed wing virus was determined in the CCD colonies of Hatay (132).

Wolbachia Endobacteria: *Wolbachia* Endobacteria is maternally inherited endosymbiont in so many kinds of insects (e.g. mosquito species), arachnids, crustaceans and some filarial nematod species such as *Dirofilaria* spp. It is estimated that 65% of insect species are infected with *Wolbachia*. In arthropods, *Wolbachia* is responsible for inducing a number of reproductive modifications that enable its spread and maintenance in natural populations (28). Recently considerable researches have been performed on *Wolbachia* that related to the interaction of *Wolbachia* and its host and impacts on parasite transmission (28). The results of the study have prompted researchers to propose strategies that utilize *Wolbachia* infections that limit parasite proliferation in the insect host to aid in the control of disease transmission and so an important step has been taken for *Wolbachia*-based control strategies, the release of *Wolbachia*-infected mosquitoes in a natural population with the ultimate goal of aiding in the control of mosquito-transmitted disease (28). In Turkey, a recent study was carried out to investigate the presence of *Wolbachia* Endobacteria in *Culex pipiens* specimens collected from Kayseri province of Turkey. The minimum infection rate of *Wolbachia* spp. in the totally analyzed 118 *C. pipiens* specimens was determined as 5.08% in the study. Furthermore one of the isolates (WolKys1) was also determined in *Wolbachia* Group B and wPIP subgroup according to the phylogenetic analyses of the *Wolbachia* surface protein (wsp) gene (GenBank accession no: JX474753). In conclusion, this study reports the first molecular

detection and characterization of *Wolbachia* endobacteria in *C. pipiens* populations in Turkey (193).

Vector-borne diseases of plants: In addition, there are many harmful arthropods and nematodes causing loss in quality and quantity of crop production in Turkey. These organisms can directly and sometimes indirectly harm plants (103). Aphids (Aphididae), leafhoppers (Cicadellidae), whiteflies (Aleyrodidae), beetles (Coleoptera), trips (Thysanoptera), gal mites (Eriophyidae), fruit flies (Tephritidae), ash-gray leaf bugs (Piesmatidae), bees and ants (Apidae and Formicidae) and nematodes are important vectors of plant diseases. Some of the most harmful plant viruses in the world such as tristeza, plum pox virus, tomato ringspot viruses are transmitted by vectors (128). The study of vector behavior and their control are of paramount importance to diminish damage to crops. The citrus tristeza virus spread by aphids brings about epidemics worldwide, causing the death of millions of citrus trees. This virus is also of extreme importance for the economy of Turkey (13, 16, 57). Plant and animal viruses have some very different constraints regarding the transmission step. While animal hosts are mobile and often come into physical contact with each other plant hosts are immobile and close contact is rare. Furthermore, plant cells are surrounded by cell walls, thick, solid barriers, which a virus cannot penetrate unless a crack is created either by mechanical stress or by plant-feeding organisms. This apparently explains why nearly all plant viruses are exclusively vector transmitted in the real world, whereas many animal viruses rely partly or completely on non-vector transmission (114). As a consequence, a diverse range of specific and intricate interaction strategies has evolved, and these strategies have been the subject of considerable research efforts in the plant virus field. Over the last half century, the classification of these plant-virus-vector interactions has been frequently modified, complemented and updated (72, 107, 148, 168, 182). Vector transmission of animal viruses was originally thought to be of two possible sorts, either 'mechanical' or 'biological'. The former is generally associated with non-specific inoculation of viruses by contaminated mouthparts or body of the vector, while the latter applies to specific associations and requires that the virus can propagate within the vector. Although this simple dichotomy is still in use today (62, 124, 133), it may not reflect the actual diversity of strategies that viruses can develop, particularly for so-called 'mechanical' transmission. Because the classification system developed for the transmission of plant viruses is more detailed, it will be used in the rest of this chapter as a common system applying to both plant and animal viruses.

In recent years, in spite of changes in ecological balance caused by abandoning agricultural land, use of

natural areas to farm, release of rabbits, preventing water flooding, leaving the fields uncultivated, migration from urban to rural areas, hunting of wild animals, the dramatic drop in the number of farm animals, climate change and the strengthening of wild life was probably responsible in the increase of tick populations (151, 164). The construction of ponds and dams, changes in riverbeds, drying out of swamps and marshes, dense ad-hoc and haphazard construction, changes in picnicking habits, wild irrigation in agriculture, and the use of unsuitable control measures by local authorities, resulted in an increase of flies (1, 109, 188, 194). This leads to an increase in infected vector populations, and also vector-borne diseases and a decrease in the potential for economic development (63, 157).

In conclusion, it is very significant to initiate and develop struggle programs against to vectors and vector-borne diseases which threaten human, animal and plant health and cause high economic losses in Turkey. Therefore they should be further investigated and monitorized regularly by using new and specific diagnostic tools.

Acknowledgements

The authors are grateful to the foundation of Erciyes University for supporting the study with project number TSS-12-4118. This article was presented at 1st National Symposium on Vectors and Vector-Borne with International Participation, 9-10 September, 2012, Avanos, Cappadocia, Nevsehir, Turkey.

Conflict of interest statement: None of the authors of this paper has a financial or personel relationship with other people or organizations that could inappropriately influence or bias the content of the paper.

References

1. **Adler PH** (2012): *World Perspective on Black Flies as Pests and Vectors*. 29-31. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
2. **Aktas M, Dumanli N, Angin M** (2004): *Cattle infestation by Hyalomma ticks and prevalence of Theileria in Hyalomma species in the east of Turkey*. Vet Parasitol, **119**, 1-8.
3. **Aktas M, Altay K, Dumanli N** (2005): *Survey of Theileria parasites of sheep in eastern Turkey using polymerase chain reaction*. Small Ruminant Research, **60**, 289-293.
4. **Aktas M, Vatansever Z, Altay K, Aydin MF, Dumanli N** (2010): *Molecular evidence for Anaplasma phagocytophilum in Ixodes ricinus from Turkey*. Trans R Soc Trop Med Hyg, **104**, 10-15.
5. **Aktas M, Altay K, Dumanli N** (2011): *Molecular detection and identification of Anaplasma and Ehrlichia species in cattle from Turkey*. Ticks Tick Borne Dis, **2**, 62-65.

6. **Aktas M, Altay K, Ozubek S, Dumanli N** (2012): *A survey of ixodid ticks feeding on cattle and prevalence of tick-borne pathogens in the Black Sea region of Turkey*. Vet Parasitol, **187**, 567-571.
7. **Albayrak H, Ozan E** (2010): *Seroprevalence of some arboviral infections transported blood sucking insects in ruminants and equids in Middle Blacksea Region in Turkey*. Kafkas Univ Vet Fak Derg, **16**, 33-36.
8. **Alibaşoğlu M, Yalçın S** (1965): *1933-1961 yılları arasında Ankara ve yöresinde atlarda görülen hastalıklara toplu bakış*. Ankara Üniv Vet Fak Derg, **12**, 98-111.
9. **Alibaşoğlu M, Göksu K, Ertürk E, Güler S** (1969): *Türkiye'de sığırlarda görülen onchocercosis olayları (Onchocerca armillata Railliet ve Henry, 1909)*. Ankara Üniv Vet Fak Derg, **16**, 50-60.
10. **Altay K, Aydın MF, Dumanli N, Aktas M** (2008): *Molecular detection of Theileria and Babesia infections in cattle*. Vet Parasitol, **158**, 295-301.
11. **Altay K, Aktas M, Dumanli N, Aydın MF** (2012): *PCR-Based Detection of Theileria sp. MK in Naturally Infected Sheep and Goats*. 110-111. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevşehir, Turkey.
12. **Anderson RC** (2006): *Parafilaria*. 394-395. In: Service MV, (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
13. **Anonymous** (2008): *Turuncgil tristesza virüsü*. 47-51. In: Tarım ve Köyişleri Bakanlığı Tarımsal Araştırmalar Genel Müdürlüğü Zirai Mücadele Teknik Talimatları **5**.
14. **Anonymous** (2012): *World Health Organization Regional Office for Europe report*. <http://www.euro.who.int/en/home>. 15.01.2013.
15. **Anyanwu IN, Agbede RI, Ajanusi OJ, Umoh JU, Ibrahim ND** (2000): *The incrimination of Aedes (stegomyia) aegypti as the vector of Dirofilaria repens in Nigeria*. Vet Parasitol, **92**, 319-327.
16. **Arsalan M, Uremis I, Sertkaya E, Sertkaya G, Caliskan S** (2012): *Phyllody Disease and Leafhopper Vectors of Sesame in Amik Plain-Turkey*. 105-106. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevşehir, Turkey.
17. **Ashford RW** (2006): *Leishmaniasis*. 269-284. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
18. **Aslan Başbulut E, Gözalan A, Sönmez C, Cöplü N, Körhasan B, Esen B, Akin L, Ertek M** (2012): *Samsun kırsalında Borrelia burgdorferi ve kene ensefaliti virusu seroprevalansının araştırılması*. Mikrobiyol Bul, **46**, 247-256.
19. **Atalay O, İca A, Cam Y, Kibar M** (2008): *A cat haemobartonellosis case in Kayseri*. J Fac Vet Med Univ Erciyes, **7**, 131-135.
20. **Atkinson CT** (2008): *Avian Malaria*. 39. In: Atkinson CT, Thomas NJ, Hunter DB (Eds), Parasitic Diseases of Wild Birds. Wiley-Blackwell, USA.
21. **Aydın L** (2012): *Geographical Distribution of Ticks in Turkey*. 91-94. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevşehir, Turkey.
22. **Ayerdem B, İnci A, Uyanık F, İca A, Çakmak A, Yildirim A** (2006): *Sığırlarda doğal Theileria annulata enfeksiyonlarında monosit nitrik oksit düzeyleri*. ERÜ Sağlık Bil Derg, **15**, 116-121.
23. **Baneth G** (2006): *Hepatozoonosis, canine*. 215-220. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
24. **Beden U, Hokelek M, Acici M, Umur S, Gungor I, Sullu Y** (2007): *A case of orbital dirofilariasis in Northern Turkey*. Ophthal Plast Reconstr Surg, **23**, 329-331.
25. **Bhide M, Yilmaz Z, Golcu Z, Torun S, Mikula I** (2008): *Seroprevalence of anti-Borrelia burgdorferi antibodies in dogs and horses in Turkey*. Ann Agric Environ Med, **15**, 85-90.
26. **Birdane FM, Sevinc F, Derinbay O** (2006): *Anaplasma marginale infections in dairy cattle: clinical disease with high seroprevalence*. Bull Vet Inst Pulawy, **50**, 467-470.
27. **Bişkin Z, Düzlü Ö, Yıldırım A, İnci A** (2010): *Kayseri'nin Felahiye Yöresinde Dirofilaria immitis'in vektör sivrisineklerde moleküler biyolojik tanısı*. Türkiye Parazitoloj Derg, **34**, 200-205.
28. **Brelsfoard CL, Dobson SL** (2011): *Short note: An update on the utility of Wolbachia for controlling insect vectors and disease transmission*. Asia Pac J Mol Biol Biotechnol, **19**, 85- 92.
29. **Burgu A, Doğanay A, Oge H, Oge S, Pişkin C** (1995): *Atlarda bulunann helmint türleri*. Ankara Üniv Vet Fak Derg, **42**, 193-205.
30. **Butler T** (2006): *Plauge*. 401-408. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections, CABI Publishing, Oxfordshire, UK.
31. **Cancrini G, Pietrobello M, Frangipane di Regalbano A, Tampieri MP, Della Torre A** (1995): *Development of Dirofilaria and Setaria nematodes in Aedes albopictus*. Parassitologia, **37**, 141-145.
32. **Carhan A, Uyar Y, Ozkaya E, Ertek M, Dobler G, Dilcher M, Wang Y, Spiegel M, Hufert F, Weidmann M** (2010): *Characterization of a new phlebovirus related to Sandfly fever Sicilian virus isolated during a Sandfly fever epidemic in Turkey*. J Clin Virol, **48**, 264-269.
33. **Celebi B, Kilic S, Aydın N, Tarhan G, Carhan A, Babur C** (2007): *Investigation of Bartonella henselae in cats in Ankara, Turkey*. Zoonoses Public Health, **56**, 169-175.
34. **Celebi B, Ozkan Taylan A, Kilic S, Akca A, Koenhems L, Pasa S, Yildiz K, Mamak N, Guzel M** (2010): *Seroprevalence of Bartonella vinsonii subsp. berkhoffi in urban and rural dogs in Turkey*. J Vet Med Sci, **72**, 1491-1494.
35. **Cengiz N, Savas L, Uslu Y, Anarat A** (2006): *Filariasis in a child from southern Turkey: a case report*. Turk J Pediatr, **48**, 152-154.
36. **Çakmak A** (1987): *Untersuchungen zur inzidenz von haemoparasiten in der provinz Ankara*. Tierarztl Hochsch Diss, Hannover.
37. **Çakmak A, İnci A, Karaer Z** (1998): *Çankırı yöresinde koyun ve keçilerde subklinik Babesia ovis enfeksiyonlarının indirekt floresan antikor testiyle saptanması*. Türkiye Parazitoloj Derg, **22**, 73-76.
38. **Çoşkun Ş, Batmaz H, Aydın L, Yılmaz F** (1997): *Türkiye'nin batısında köpeklerde Leishmania infantum*

- enfeksiyonunun seroprevalansı. Türkiye Parasitol Derg, **21**, 287-291.
39. **Depaquit J, Grandadam M, Fouque F, Andry PE, Peyrefitte C** (2010): *Arthropod-borne viruses transmitted by Phlebotomine sandflies in Europe: a review*. Euro Surveill, **15**, 19507.
 40. **Dinçer Ş, Sayın F, Karaer Z, Çakmak A, Friedhoff KT, Muller I, İnci A, Yukarı BA, Eren H** (1991): *Karadeniz Bölgesi sığırlarında bulunan kan parazitlerinin sero-insidensi üzerine araştırmalar*. Ankara Üniv Vet Fak Derg, **38**, 206-226.
 41. **Dionisio D, Esperti F, Vivarelli A, Valassina M** (2003): Epidemiological, clinical and laboratory aspects of sandfly fever. Curr Opin Infect Dis, **16**, 383-388.
 42. **Doganay M, Metan G** (2009): *Human anthrax in Turkey from 1990 to 2007*. Vector Borne Zoonotic Dis, **9**, 131-140.
 43. **Doganay M** (2012): *Tularaemia: Re-Emerging Disease*. 62-64. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 44. **Dragon DC** (2006): *Anthrax*. 46-48. In: Service MV (Ed), *The Encyclopedia of Arthropod-Transmitted Infections*. CABI Publishing, Oxfordshire, UK.
 45. **Dumanlı N, Aktas M, Cetinkaya B, Cakmak A, Koroglu E, Saki CE, Erdogmus Z, Nalbantoglu S, Ongor H, Simsek S, Karahan M, Altay K** (2005): *Prevalence and distribution of tropical theileriosis in Eastern Turkey*. Vet Parasitol, **127**, 9-15.
 46. **Düzlü Ö, İnci A, Yıldırım A** (2011): *Karadeniz bölgesindeki sığırlardan elde edilen Babesia bovis suşlarının moleküler karakterizasyonu*. ERÜ Sağlık Bilimleri Derg, **20**, 18-28.
 47. **Duzlu O, Yıldırım A, İnci A, Avcioglu H, Balkaya I, Biskin Z** (2012): *The Investigation of Babesia bovis and Babesia bigemina by Real Time PCR in Cattle in Erzurum Province of Eastern Part of Turkey*. 119-121. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 48. **Düzlü Ö, İnci A, Yıldırım A** (2012): *Türkiye’de evcil ruminantlarda babesiosis*. Türkiye Klinikleri J Vet Sci, **3**, 27-34.
 49. **Ekici OD, Sevinc F** (2011): *Comparison of cELISA and IFA tests in the serodiagnosis of anaplasmosis in cattle*. African Journal of Microbiology Research, **5**, 1188-1191.
 50. **Ekrem İ** (1931): *Piroplasmoslar hakkında en yeni malumat*. Türk Baytarlar Mecmuası, **6**, 65-73.
 51. **Erdil N** (1966): *Köpeklerde Dirofilaria immitis ve Dipetalonema reconditum üzerinde araştırma (A study on Dirofilaria immitis and Dipetalonema reconditum in dogs)*. KKK As Vet Akademisi İç Hastalıklar Şube ve Kliniği Çalışmaları, Ankara.
 52. **Ergunay K, Ozer N, Us D, Ozkul A, Simsek F, Kaynas S, Ustacelebi S** (2007): *Seroprevalence of West Nile virus and tick-borne encephalitis virus in southeastern Turkey: first evidence for tick-borne encephalitis virus infections*. Vector Borne Zoonotic Dis, **7**, 157-61.
 53. **Ergunay K, Saygan MB, Aydoğan S, Menemenlioglu D, Turan HM, Ozkul A, Us D** (2010): *West Nile virus seroprevalence in blood donors from Central Anatolia, Turkey*. Vector Borne Zoonotic Dis, **10**, 771-775.
 54. **Ergunay K, Sayiner AA, Litzba N, Lederer S, Charrel R, Kreher P, Us D, Niedrig M, Ozkul A, Hascelik G** (2012): *Multicentre evaluation of central nervous system infections due to Flavi and Phleboviruses in Turkey*. J Infect, **65**, 343-349.
 55. **Ergunay K, Aydoğan S, İlhami Özcebe O, Cilek EE, Hacıoglu S, Karakaya J, Ozkul A, Us D** (2012): *Toscana Virus (TOSV) exposure is confirmed in blood donors from Central, North and South/Southeast Anatolia, Turkey*. Zoonoses and Public Health, **59**, 148-154.
 56. **Ergunay K, Erisoz Kasap O, Kocak Tufan Z, Turan MH, Ozkul A, Alten B** (2012): *Molecular evidence indicates that Phlebotomus major sensu lato (Diptera: Psychodidae) is the vector species of the recently-identified sandfly fever Sicilian virus variant: sandfly fever turkey virus*. Vector Borne Zoonotic Dis, **12**, 690-698.
 57. **Ertunc F** (2012): *A New Threat for Turkish Horticulture: Phytoplasma Diseases and Their Vectors*. 77-84. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 58. **Erturk A, Tatar N, Kabaklı O, Incoglu S, Cizmeci SG, Barut FM** (2004): *The current situation of bluetongue in Turkey*. Vet Ital, **40**, 137-140.
 59. **Ewing SA** (2006): *Ehrlichiosis*. 159-162. In: Service MV (Ed), *The Encyclopedia of Arthropod-Transmitted Infections*. CABI Publishing, Oxfordshire, UK.
 60. **Gao GF, Hussain MH, Reid HW, Gould EA** (1993): *Classification of a new member of the TBE flavivirus subgroup by its immunological, pathogenetic and molecular characteristics: identification of subgroup-specific pentapeptides*. Virus Res, **30**, 129-144.
 61. **Göksu K** (1970): *Yurdumuzun çeşitli bölgelerinde sığırlarda Piroplasmida enfeksiyonları (piroplasmossis, babesiosis, theileriosis ve anaplasmosis’in yayılış durumları)*. Türk Vet Hek Dern Derg, **40**, 29-39.
 62. **Gray SM, Banerjee N** (1999): *Mechanisms of arthropod transmission of plant and animal viruses*. Microbiol Mol Biol Rev, **63**, 128-148.
 63. **Gray EW, Noblet R, Iburg JP, Wyatt RD** (2012): *Standardized Methods for Black Fly Rearing and Product Development for the Operational Control of Black Flies*. 41-43. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 64. **Gunes T, Poyraz O, Atas M, Turgut NH** (2011): *The seroprevalence of Anaplasma phagocytophilum in humans from two different climatic regions of Turkey and its co-seroprevalence rate with Borrelia burgdorferi*. Turk J Med Sci, **41**, 903-908.
 65. **Gün H, Tanyüksel M, Yukarı BA, Çakmak A, Karaer Z** (1996): *Türkiye’de babesiosis’in ilk insan serodiagnozu*. Türkiye Parazitoloj Derg, **20**, 1-7.
 66. **Gungel H, Kara N, Pinarci EY, Albayrak S, Baylancicek DO, Uysal HK** (2009): *An uncommon case with intravitreal worm. Intravitreal Dirofilaria infection*. Br J Ophthalmol, **93**, 573-574.
 67. **Güralp N, Doğru C** (1968): *Türkiye’de ehli hayvanlarda setariose*. Ankara Üniv Vet Fak Derg, **15**, 29-40.
 68. **Güralp N, Oğuz T** (1970): *Türkiye’de mandalarda (Bubalus bubalis) thelaziose*. Ankara Üniv Vet Fak Derg, **17**, 109-113.

69. **Güralp N** (1981): *Helmintholoji*. 599. In: Ankara Üniversitesi Veteriner fakültesi yayınları: 368, ders Kitabı No: 266, Ankara Üniversitesi Basımevi, Ankara.
70. **Gurel MS, Olgen MK, Yesilova Y, Ozbel Y** (2012): *Cutaneous leishmaniasis in Turkey*. *Turkiye Parazitoloj Derg*, **36**, 121-129.
71. **Hardly WJ, Martin WB, Hakioglu F, Chifney STE** (1969): *A viral encephalitis of sheep in Turkey*. *Pendik Institute J*, **1**, 89-100.
72. **Harris KF** (1977): *An ingestion-egestion hypothesis of non circulative virus transmission*. 166-208. In: Harris KF, Maramorosch K (Eds), *Aphids as Virus Vectors*, Academic Press, New York.
73. **Hechemy KE, Anderson BE** (2006): *Trench Fever*. 71-72. In: Service MV (Ed), *The Encyclopedia of Arthropod-Transmitted Infections*. CABI Publishing, Oxfordshire, UK.
74. **Hechemy KE, Azad AF** (2006): *Epidemic typhus*. 170-174. In: Service MV (Ed), *The Encyclopedia of Arthropod-Transmitted Infections*. CABI Publishing, Oxfordshire, UK.
75. **Hillyer JF, Barreau C, Vernick KD** (2007): *Efficiency of salivary gland invasion by malaria sporozoites is controlled by rapid sporozoite destruction in the mosquito haemocoel*. *Int J Parasitol*, **37**, 673-681.
76. **Ica A, Vatansever Z, Yildirim A, Duzlu O, İnci A** (2007): *Detection of Theileria and Babesia species in ticks collected from cattle*. *Vet Parasitol*, **148**, 156-160.
77. **Ica A, İnci A, Yildirim A** (2007): *Parasitological and molecular prevalence of bovine Theileria and Babesia species in the vicinity of Kayseri*. *Turk J Vet Anim Sci*, **31**, 33-38.
78. **İça A, İnci A, Yildirim A, Aslan Ö, Düzlü Ö** (2008): *Kayseri ve civarında köpeklerde leishmaniasisin Nested-PCR ile araştırılması*. *Turkiye Parazitoloj Derg*, **32**, 187-191.
79. **Icen H, Sekin S, Simsek A, Kochan A, Celik OY, Altas MG** (2011): *Prevalence of Dirofilaria immitis, Ehrlichia canis, Borrelia burgdorferi infection in dogs from Diyarbakir in Turkey*. *Asian J Anim Vet Adv*, **6**, 371-378.
80. **İnci A** (1992): *Ankara'nın Çubuk ilçesinde sığırlarda babesiosis'in seroinsidensi üzerine araştırmalar*. *Ankara Üniv Vet Fak Derg*, **39**, 153-167.
81. **İnci A, Sayın F** (1995): *Türkiye'de sığırlardan Trypanosoma theileri (Laveran, 1902)'nin izolasyonu ve kültürasyonu*. *Ankara Üniv Vet Fak Derg*, **42**, 97-103.
82. **İnci A** (1997): *Gemlik askeri harası atlarında Babesia caballi (Nuttal, 1910) ve Babesia equi (Laveran, 1901)'nin mikroskopik muayeneyle saptanması*. *T J Vet and Anim Sci*, **21**, 43-46.
83. **İnci A, Yukari BA, Sayın F** (1998): *Çankırı yöresinde bazı koyun ve keçi sürülerinde babesiosis ve theileriosis etkenlerinin mikroskopik kan muayenesiyle araştırılması*. *Ankara Üniv Vet Fak Derg*, **45**, 105-113.
84. **İnci A, Çakmak A, Çam Y, Karaer Z, Atasever A, İça A** (2002): *Kayseri yöresinde tropikal theileriosis'e bağlı ekonomik kayıplar*. *Turkiye Parazitoloj Derg*, **26**, 156-160.
85. **İnci A, Çakmak A, Karaer Z, Dinçer Ş, Sayın F, İça A** (2002): *Kayseri yöresinde sığırlarda babesiosisin seroprevalansı*. *Turk J Vet Anim Sci*, **26**, 1345-1350.
86. **İnci A, Çakmak A, İça A, Günay O** (2002): *Kayseri yöresinde tropikal theileriosis'in istatistiksel analizi*. *Turkiye Parazitoloj Derg*, **26**, 38-41.
87. **İnci A, Karaer Z, İça A** (2002): *Kayseri yöresinde koyun ve keçilerde babesiosis*. *F Ü Sağ Bil Derg*, **16**, 79-83.
88. **İnci A, Nalbantoglu S, Çam Y, Atasever A, Karaer Z, Çakmak A, Sayın F, Yukarı B, İça A, Deniz A** (2003): *Kayseri yöresinde koyun ve keçilerde theileriosis ve kene enfestasyonları*. *Turk J Vet Anim Sci*, **27**, 57-60.
89. **İnci A, Ica A, Yildirim A, Vatansever Z, Cakmak A, Albasan H, Cam Y, Atasever A, Sariozkan S, Duzlu O** (2007): *Economical impact of tropical theileriosis in the Cappadocia Region of Turkey*. *Parasitol Res*, **101**, 171-174.
90. **İnci A, İça A, Florin-Christensen M, Vatansever Z, Yildirim A, Düzlü Ö, Mosqueda J, Yavuz A** (2007): *Türkiye'nin çeşitli yörelerinden elde edilen Babesia bovis ve Babesia bigemina'nın moleküler karakterizasyonu*. 163-164. In: XV. Ulusal Parazitoloji Kongresi, 18-23 Kasım, Kayseri ve Ürgüp.
91. **İnci A, İça A, Albasan H** (2007): *Babesia Enfeksiyonlarında İmmunité*. 508-515. In: Özcel MA, İnci A, Turgay N, Köroğlu E (Eds), *Tıbbi ve Veteriner İmmunoparazitoloji*. Parazitoloji Derneği Yayınları, No: 21, İzmir, Türkiye.
92. **İnci A, Düzlü Ö** (2009): *Vektörler ve vektörlerle bulaşan hastalıklar*. *ERÜ Vet Fak Derg*, **6**, 53-63.
93. **İnci A, Ica A, Yildirim A, Düzlü O** (2010): *Identification of Babesia and Theileria species in small ruminants in Central Anatolia (Turkey) via reverse line blotting*. *Turk J Vet Anim Sci*, **34**, 205-210.
94. **İnci A, Yildirim A, Njabo KY, Duzlu O, Biskin Z, Ciloglu A** (2012): *Detection and molecular characterization of avian Plasmodium from mosquitoes in central Turkey*. *Vet Parasitol*, **188**, 179-184.
95. **İnci A, Yavuz A, Yildirim A, Düzlü O, Bişkin Z** (2012): *Sığır theileriosis'inde metastaz*. *Erciyes Üniv Vet Fak Derg*, **9**, 113-122.
96. **İnci A, Yildirim A, Duzlu O, Ciloglu A, Biskin Z** (2012): *The investigation of canine blood protozoon infections by taqman real time PCR in Kayseri province of Central Turkey*. 102-104. In: *Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation*, 9-10 September, Avanos, Cappadocia, Nevşehir, Turkey.
97. **İnci A, Duzlu O, Yildirim A** (2012): *Molecular prevalence of babesiosis and theileriosis in cattle in Turkey*. 95-96. In: *Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation*, 9-10 September, Avanos, Cappadocia, Nevşehir, Turkey.
98. **İnci A, Yildirim A, Düzlü O** (2013): *Orta Kızılırmak Havzası'nda sorun oluşturan karasinek (Diptera: Simuliidae) türlerinin moleküler klasifikasyonu ve vektörlük potansiyellerinin real time pcr ile araştırılması*. In: TÜBİTAK 111O426 Numaralı Araştırma Projesi, 2. Gelişme Raporu.
99. **Ishtiaq F, Guillaumot L, Clegg SM, Phillimore AB, Black RA, Owens IP, Mundy NI, Sheldon BC** (2008): *Avian haematozoan parasites and their associations with mosquitoes across Southwest Pacific Islands*. *Mol Ecol*, **17**, 4545-4555.
100. **Jennings M, Boorman JPT, Ergun H** (1983): *Culicoides from Western Turkey in relation to bluetongue disease of sheep and cattle*. *Rev Elev Med Vet Pays Trop*, **36**, 67-70.

101. **Johnson R** (2010): *Honey bee colony collapse disorder*. In: CRS Report for Congress, Congressional Research Service, 7-5700, January 7, 2010. <http://www.fas.org/sgp/crs/misc/RL33938.pdf>. 15.01.2013.
102. **Jones TW** (2006): *Animal Trypanosomiasis*. 33-46. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
103. **Kansu İA** (1994): Genel Entomoloji. Kıvanç Basımevi, Ankara.
104. **Karagenc TI, Pasa S, Kirli G, Hosgor M, Bilgic HB, Ozon YH, Atasoy A, Eren H** (2006): *A parasitological, molecular and serological survey of Hepatozoon canis infection in dogs around the Aegean coast of Turkey*. Vet Parasitol, **135**, 113-119.
105. **Karaoglu T, Ozgunluk I, Demir B, Ozkul A, Burgu I** (2007): *Seroprevalence of culicoides-borne disease in cattle in Eurpoean Turkey*. Ankara Univ Vet Fak Derg, **54**, 121-125.
106. **Kaya M** (2011): *Investigation of Frequency of Babesiosis among Human who Have Tick Bite History Living in Tatvan Region*. Thesis of Master, Erciyes University, Graduate School of Health Sciences, Department of Parasitology, Kayseri.
107. **Kennedy JS, Day MF, Eastop VF** (1962): *A Conspectus of Aphids as Vectors of Plant Viruses*. Commonwealth Institute of Entomology, London.
108. **Kilic S** (2010): *Francisella tularensis ve Türkiye’de tularemii epidemiyolojisine genel bir bakış*. Flora, **15**, 37-58.
109. **Kilic E, Canhilal R** (2012): *Entomopathogens for Controlling of Urban Pests*. 122-124. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
110. **Kiral F, Karagenc T, Pasa S, Yenisey C, Seyrek K** (2005): *Dogs with Hepatozoon canis respond to the oxidative stress by increased production of glutathione and nitric oxide*. Vet Parasitol, **131**, 15-21.
111. **Kiriş Satılmış O, Akkaya Y, Ergin C, Kaleli I, Dursun B, Aydın C** (2012): *Investigation of Bartonella henselae antibodies in serum and plasma samples of kidney transplant patients*. Mikrobiyol Bul, **46**, 568-574.
112. **Kızılaslan F** (2012): *Bursa Karacabey Yöresinde Atlarda Theileria equi Ve Babesia caballi'nin Real Time PCR ile Araştırılması ve Moleküler Karakterizasyonu*. ERÜ Sağlık Bilimleri Enstitüsü, Doktora Tezi, Kayseri.
113. **Kocan KM** (2006): *Anaplasmosis*. 28-33. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
114. **Kuno G** (2001): *Transmission of arboviruses without involvement of arthropod vectors*. Acta Virol, **45**, 139-150.
115. **Kurtdede A, Ural K** (2004): *Haemobartonellosis in cats in Ankara, Turkey*. Acta Vet Brno, **73**, 507-512.
116. **Küçükaydin M, Sahin I** (1986): *The third case of dirofilariasis conjunctivae in Turkey*. Mikrobiyol Bul, **20**, 25-28.
117. **Latifoglu O, Ozmen S, Sezer C, Yavuzer R, Altintas K, Uluoglu O** (2002). *Dirofilaria repens presenting as a premaseteric nodule*. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. **94**, 217-20.
118. **Lyons ET** (2006): *Habronemiasis of Horses*. 198-200. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
119. **Lyons ET** (2006): *Thelaziasis*. 502-504. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
120. **Maguiña C, Guerra H, Ventosilla P** (2009): *Bartonellosis*. Clinics in Dermatology, **27**, 271-280.
121. **Marenzoni ML, Cuteri V, De Parri F, Danzetta ML, Yilmaz Z, Yaramis CP, Kennerman E, Or Me, Marchi S, Casciari C, De Mia Gm, Valente C, Costarelli S** (2013): *A pilot study on the epidemiological status of equine infectious anaemia, equine viral arteritis, glanders, and dourine in Turkey*. Turk J Vet Anim Sci, **37**, doi:10.3906/vet-1110-46.
122. **Maroli M, Feliciangeli MD, Bichaud L, Charrel RN, Gradoni L** (2012): *Phlebotomine sandflies and the spreading of leishmaniasis and other diseases of public health concern*. Med Vet Entomol, DOI: 10.1111/j.1365-2915.2012.01034.
123. **Medlock JM, Barras I, Kerrod E, Taylor MA, Leach S** (2007): *Analysis of climatic predictions for extrinsic incubation of Dirofilaria in the United Kingdom*. Vector Borne Zoonotic Dis, **7**, 4-14.
124. **Mellor PS** (2000): *Replication of arboviruses in insect vectors*. J Comp Pathol, **123**, 231-247.
125. **Mellor PS** (2006): *Akabane virus*. 25-27. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
126. **Mellor PS** (2006): *Bovine Ephemeral Fever*. 87-91. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
127. **Mert A, Ozaras R, Tabak F, Bilir M, Ozturk R, Aktuglu Y** (2003): *Akdeniz benekli ateşi: 14 vakanın değerlendirilmesi*. Flora, **8**, 158-162.
128. **Meyer JR** (2003): *Insect vectors of plant pathogens*. Department of Entomology, North Carolina State University. <http://www.cals.ncsu.edu/course/ent425/text18/plantvectors.html#wasps>.
129. **Muhammad SA, Farooq MS, Akhtar MS, Hayat CS** (2004): *Dipetalonemiasis in a dromedary camel and its treatment*. Pakistan Vet J, **24**, 205-206.
130. **Mumcuoglu KY** (2012): *The Vectorial Capacity of the Human Body Louse, Pediculus humanus humanus*. 35-37. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
131. **Muz MN, Yaman M, Karakavuk M, Dik B** (2012): *Role of Phlebotomines as Vector of Leishmania sp. in Hatay, Turkey. Preliminary Report for Sandfly-borne Diseases and DNA barcoding*. 117-118. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
132. **Muz MN, Muz D** (2012): *Vectorial Role of Mites on CCD in Turkey, Varroa destructor as a Vector of Deformed Wing Virus in Honeybee Colony Collapse Disorders*. 97-98. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International

- Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
133. **Nault LR** (1997): *Arthropod transmission of plant viruses: a new synthesis*. Ann Entomol Soc Am, **90**, 521-541.
 134. **Njabo KY, Smith TB** (2012): *Vectors of Avian Malaria in Africa and Their Blood Feeding Habits*. 32-34. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 135. **Oge S, Oge H, Yildirim A, Kircali F** (2003): *Setaria equine infection of Turkish equines: estimates of prevalence based on necropsy and detection of micrflaraemia*. Ann Trop Med Parasitol, **97**, 403-409.
 136. **Ok UZ, Balcioglu IC, Taylan Ozkan A, Ozensoy S, Ozbel Y** (2002): *Leishmaniasis in Turkey*. Acta Tropica, **84**, 43-49.
 137. **Ozbel Y, Turgay N, Ozensoy S, Ozbilgin A, Alkan MZ, Ozel MA, Jaffe CL, Schnur L, Oskam L, Abranches P** (1995): *Epidemiology, diagnosis and control of leishmaniasis in Mediterranean region*. Ann Trop Med Parasitol, **89**, 89-93.
 138. **Ozbel Y, Oskam L, Ozensoy S, Turgay N, Alkan MZ, Jaffe CL, Ozel MA** (2000): *A survey on canine leishmaniasis in western Turkey by parasite, DNA and antibody detection assays*. Acta Tropica, **74**, 1-6.
 139. **Ozbel Y, Ertabaklar H, Ciufolini MG, Marks A, Fiorentino C, Erensoy S, Ozacar T, Rastgeldi S, Nicoletti N** (2003): *Sand fly Fever viruses (Phleboviruses) transmitted by Phlebotomus in Turkey*. Microbiologica Balkanica 3rd Balkan Conference of Microbiology Proceedings, 1152-1155.
 140. **Ozdarendeli A** (2012): *Development of an Inactivated Vaccine against Crimean-Congo Hemorrhagic Fever Virus*. 60-61. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 141. **Ozer N, Ergünay K, Simsek F, Kaynas S, Alten B, Caglar SS, Ustacelebi S** (2007): *West Nile virus studies in the Sanliurfa Province of Turkey*. J Vector Ecol, **32**, 202-206.
 142. **Ozensoy Toz S, Nasereddin A, Ozbel Y, Ertabaklar H, Culha G, Sevil N, Alkan MZ, Jaffe CL** (2009): *Leishmaniasis in Turkey: molecular characterization of Leishmania from human and canine clinical samples*. Trop Med Int Health, **14**, 1401-1406.
 143. **Ozkul A, Yildirim Y, Pinar D, Akcali AA, Yilmaz V, Colak D** (2006): *Serological evidence of West Nile Virus (WNV) in mammalian species in Turkey*. Epidemiol Infect, **134**, 826-829.
 144. **Ozkul A, Ergunay K, Koysuren A, Alkan F, Arsava EM, Tezcan S, Emekdas G, Hacıoglu S, Turan M, Us D** (2013): *Concurrent occurrence of human and equine West Nile Virus infection in Central Anatolia, Turkey: The first evidence for circulation of lineage I viruses*. IJID, (DOI: 10.1016/j.ijid.2013.02.005).
 145. **Ozlem MB, Karaer Z, Turgut K, Eren H, Irmak K, Inci A** (1988): *Efficacy of long-acting oxytetracycline on bovine anaplasmosis*. Ank Üniv Vet Fak Derg, **35**, 1-5.
 146. **Ozkurt Z, Parlak M, Tastan R, Dinler U, Saglam YS, Ozyurek F** (2005): *Anthrax in Eastern Turkey, 1992-2004*. Emerg Infect Dis, **11**, 1939-1941.
 147. **Pampiglione S, Rivasi F** (2000): *Human dirafilariosis due to Dirofilaria (Nochtiella) repens: an update of world literature from 1995 to 2000*. Parassitologia, **42**, 231-254.
 148. **Pirone T, Blanc S** (1996): *Helper-dependent vector transmission of plant viruses*. Annu Rev Phytopathol, **34**, 227-247.
 149. **Poyraz O, Gunes T** (2010): *Seroprevalance of Babesia microti in Humans Living in Rural Areas of the Sinop Region*. Turkiye Parazit Derg, **34**, 81-85.
 150. **Preston PM** (2006): *Theileriosis*. 487-502. In: Service MV (Ed), The Encyclopedia of Artropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
 151. **Randdolph S** (2008): *To what extent has climate change contributed to the recent epidemiology of tick-borne diseases?* 2-3. In: Proceeding VI International Conferece on Ticks and Tick-Borne Pathogens. 21-26 September, Buenos Aires, Argentina.
 152. **Renz A** (2006): *Onchocerciasis, Animal*. 375-381. In: Service MV (Ed), The Encyclopedia of Artropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
 153. **Rikihisa Y** (2012): *Ehrlichiae of Public Health and Veterinary Importance: How Can They Infect Phagocytes?* 38-40. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 154. **Robinson PM** (1982): *Theileria annulata and its transmission-A review*. Trop Anim Health Prod, **14**, 3-12.
 155. **Sakru N, Testini G, Gurlu VP, Yakar K, Lia RP, Dantas-Tores F, Bain O** (2012): *Case Report: First evidence of human zoonotic infection by Onchocerca lupi (Spirurida, Onchocercidae)*. Am J Trop Ed Hyg, **84**, 55-58.
 156. **Saraylı H, İnci A, İça A, Yildirim A, Düzlü Ö** (2006): *Yeşilhisar yöresindeki koyun ve keçilerde Babesia etkenlerinin Reverse Line Blotting (RLB) yöntemiyle araştırılması*. ERÜ Sağlık Bil Derg, **15**, 181-188.
 157. **Sariozkan S, Inci A** (2012): *An economic perspective on Simulium Infestation in Cappadocia Region of Turkey*. 86-87. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
 158. **Sayin F, Dincer S, Karaer Z, Dumanli N, Cakmak A, Inci A, Yukari BA, Vatansever Z** (1997): *Status of tick infestation of sheep and goats in Turkey*. Parassitologia, **39**, 145-152.
 159. **Sayin F, Karaer Z, Dincer S, Cakmak A, Inci A, Yukari BA, Eren H, Vatansever Z, Nalbantoglu S, Melrose T** (2003): *A Comparison of susceptibilities to Infection of four species of Hyalomma ticks with Theileria annulata*. Vet Parasitol, **113**, 115-121.
 160. **Sayin F, Dincer S, Karaer Z, Cakmak A, Inci A, Yukari BA, Eren H, Vatansever Z, Nalbantoglu S** (2003): *Studies on the epidemiology of tropical theileriosis (Theileria annulata lfection) in cattle in Central Anatolia, Turkey*. Trop Anim Health Prod, **35**, 521-539.
 161. **Sayin-Kutlu S, Ergin C, Kutlu M, Akkaya Y, Akalin S** (2012): *Bartonella henselae seroprevalence in cattle breeders and veterinarians in the rural areas of Aydin and Denizli, Turkey*. Zoonoses Public Health, **59**, 445-449.

162. **Serdar ZA, Yasar S, Inan A, Doner N, Erfan G** (2010): *Kene tifüsü: 2 olgu sunumu*. Türk Aile Hek Derg, **14**, 46-49.
163. **Serter D** (1980): *Present status of arbovirus sero-epidemiology in the Aegean region of Turkey*. 155-161. In: Vesenjak-Hirjan J, Calisher C (Eds), Arboviruses in the Mediterranean Countries, Gustav Fischer Verlag, Stuttgart.
164. **Service MW** (2006): *Ticks (Ixodida)*. 524-528. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
165. **Sevinç F, Özdemir Ö, Coşkun A** (2005): *On bir günlük bir buzağıda akut babesiosis*. ERU Vet Fak Derg, **2**, 131-135.
166. **Seyrek K, Karagenc T, Pasa S, Kiral F, Atasoy A** (2009): *Serum zinc, iron and copper concentrations in dogs infected with Hepatozoon canis*. Acta Vet Brno, **78**, 471-475.
167. **Soylu M, Ozcan K, Yalaz M, Varinli S, Slem G** (1993): *Dirofilariasis: an uncommon parasitosis of the eye*. Br J Ophthalmol, **77**, 602-603.
168. **Sylvester ES** (1956): *Transmission of Beet yellows virus by the green peach aphid*. J Econ Entomol, **49**, 789-800.
169. **Şaki CE, Sevgili M, Özer E** (1999): *Malatya ve yöresi sığırlarında parafilariosis*. Turk J Vet Anim Sci, **23**, 541-545.
170. **Tarizzo ML** (1973): *Geographic distribution of louse-borne diseases*. 50-60. In: Horwitz A (Ed), Proceeding of the International Symposium on the control of lice and louse-borne diseases. Vol **263**. Pan American Health Organization, Washington, D.C., U.S.A.
171. **Taşcı S, Toparlak M, Yılmaz H** (1989): *Van mezbahasında kesilen sığırlarda thelaziose'un yayılışı*. Ankara Üniv Vet Fak Derg, **36**, 352-357.
172. **Tesh RB, Saidi S, Gajdamovic SJ, Rodhain F, Vesenjak-Hirjan J** (1976): *Serological studies on the epidemiology of sandfly fever in the Old World*. Bull World Health Organ, **54**, 663-674.
173. **Toparlak M, Gargili A, Ulutas Esatgil M, Cetinkaya H** (2005): *Canine filariosis around Istanbul, Turkey Employing Naphthol AS-TRm phosphate*. Acta Vet Brno, **74**, 233-236.
174. **Tüzdil AN** (1933): *Bizde ilk defa görülen bir Hepatozoon canis vak'ası*. Türk Baytarlar Cemiyeti Mecmuası, **13**, 35.
175. **Tuzer E, Goksu K, Bilal T, Yesildere T** (1993): *A case of haemobartonellosis in a cat in Istanbul*. J Protozool Res, **3**, 69-70.
176. **Ulutas B, Bayramlı G, Karagenc T** (2007): *First case of Anaplasma (Ehrlichia) platys infection in a dog in Turkey*. Turk J Vet Animal Sci, **31**, 279-282.
177. **Unver A, Rikihisa Y, Borku K, Ozkanlar Y, Hanedan B** (2005): *Molecular detection and characterization of Ehrlichia canis from dogs in Turkey*. Berl Munch Tierarztl Wochenschr, **118**, 300-304.
178. **Ural K, Aysul N, Tuna GE, Atasoy A, Ulutas B** (2012): *Abdominal Compartment Syndrome in a Babesia canis Infected Dog*. 138-139. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
179. **Uyar Y, Akcali A, Carhan A, Ozkaya E, Ertek M** (2007): *Türkiye'de kene ısırığı öykülü olgularda tick-borne encephalitis virüsünün seroprevalansı*. Turk Hij Den Biyol Derg, **64**, 21-25.
180. **Vatansever Z** (2012): *Ecology of Hyalomma marginatum and Crimean-Congo hemorrhagic fever*. 44-59. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
181. **Warnel DA** (2006): *Louse-borne relapsing fever*. 295-299. In: Horwitz A (Ed), Proceeding of the International Symposium on the control of lice and louse-borne diseases. Vol **263**. Pan American Health Organization, Washington, D.C., U.S.A.
182. **Watson MA, Roberts FM** (1939): *A comparative study of the transmission of Hyoscyamus Virus 3, potato virus Y and cucumber virus 1 by the vectors Myzus persicae (Sulz), M. circumflexus (Buckton), and Macrosiphum gei (Koch)*. Proc R Soc Lond B, **127**, 543-576.
183. **Willcox WH** (1920): *Typhus and relapsing fever in Mesopotamia and Northern Persia*. Proc R Soc Med, **13** (Med Sect), 59-81.
184. **Wjesundera WSS** (2006): *Setariosis*. 465-468. In: Service MV (Ed), The Encyclopedia of Arthropod-Transmitted Infections. CABI Publishing, Oxfordshire, UK.
185. **Yarmut Y, Brommer H, Weisler S, Shelah M, Komarovskiy O, Steinman A** (2008): *Ophthalmic and cutaneous habronemiasis in a horse: case report and review of the literature*. Isr J Vet Med, **63**, 87-90.
186. **Yasa O, Erguven M, Karaca Atakan S, Emmungil S** (2010): *Ateş ve döküntü birlikteliğinin ender nedeni: Bir riketsiyöz vakası*. Çocuk Derg, **10**, 90-93.
187. **Yavuz A, İnci A, Düzlü Ö, Bişkin Z, Yıldırım A** (2011): *Babesia bovis'in msa-2c Geninin Moleküler Karakterizasyonu*. Türkiye Parazitoloj Derg, **35**, 140-144.
188. **Yılmaz A, İnci A, Tunçbilek A, Yeşilöz H, Koçak O, Şirin Ü, İça A, Yıldırım A, Demircioğlu A, Düzlü Ö** (2007): *Orta Kızılırmak Havzasında Karasinek (Simulium (Wilhelmia) Lineatum) (Diptera: Simuliidae) İstilası*. ERU Üniv Vet Fak Derg, **4**, 91-95.
189. **Yıldırım A, İça A, Atalay Ö, Düzlü Ö, İnci A** (2007): *Kayseri Yöresi Köpeklerinde Dirofilaria immitis'in Membran Filtrasyon-Asit Fosfat Histokimyasal Boyama, Antijen ELISA ve PCR Yöntemleri ile Araştırılması*. 140-141. In: XV. Ulusal Parazitoloji Kongresi, 18-23 Kasım, Kayseri ve Ürgüp.
190. **Yıldırım A, İca A, Atalay O, Duzlu O, İnci A** (2007): *Prevalence and epidemiological aspects of Dirofilaria immitis in dogs from Kayseri province, Turkey*. Res Vet Sci, **82**, 358-363.
191. **Yıldırım A, İnci A, Duzlu O, Biskin Z, İca A, Şahin İ** (2011): *Aedes vexans and Culex pipiens as the potential vectors of Dirofilaria immitis in Central Turkey*. Vet Parasitol, **178**, 143-147.
192. **Yıldırım A, Aysul N, Bayramlı G, İnci A, Eren H, Aypak S, Duzlu O** (2012): *Detection and Molecular Characterization of a Haemoproteus lineage in an owl in Turkey*. 88-90. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
193. **Yıldırım A, İnci A, Duzlu O, Biskin Z, Ciloglu A** (2012): *Detection and Molecular Characterization of the Wolbachia Endobacteria in the Culex pipiens (Diptera: Culicidae) Specimens Collected from Kayseri Province of*

- Turkey*. 99-101. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
194. **Yilmaz A, Yesiloz H, Demircioglu A** (2012): *Control Programs of the Simulium Species in the Center Kizilirmak Basin of Turkey*. 65-67. In: Proceedings of 1st National Symposium on Vectors and Vector Borne Diseases with International Participation, 9-10 September, Avanos, Cappadocia, Nevsehir, Turkey.
195. **Yilmaz C, Ergin C, Kaleli I** (2009): *Investigation of Bartonella henselae seroprevalence and related risk factors in blood donors admitted to Pamukkale University blood center*. Mikrobiyol Bul, **43**, 391-401.
196. **Yüce G, Girgin H** (1967): *Bir onchocerciasis olayı*. Türk Vet Hek Dern Derg, **37**, 18-21.
- Geliş tarihi: 11.01.2013 / Kabul tarihi: 22.03.2013*

Address for correspondence:

Prof. Dr. Abdullah Inci
Erciyes University,
Vectors and Vector-Borne Diseases Implementation and
Research Center,
Melikgazi , 38038, Kayseri-Turkey.
e-mail: ainci@erciyes.edu.tr