

## Floods increase similarity among aquatic habitats in river-floodplain systems

Sidinei M. Thomaz · Luis Mauricio Bini ·  
Reinaldo Luiz Bozelli

Received: 30 March 2006 / Revised: 30 June 2006 / Accepted: 30 July 2006 / Published online: 21 November 2006  
© Springer Science+Business Media B.V. 2006

**Abstract** During low water levels, habitats in river-floodplain systems are isolated from each other and from the main river. Oppositely, floods tend to connect water bodies with distinct hydrological characteristics and, as a result, ecological processes and biological communities tend to be more similar among the distinct habitats that comprise a river-floodplain system. Based on a literature review and also using unpublished data obtained in tropical floodplains, the aim of this paper is to highlight the effects of floods as a process that reduce spatial variability. The usual negative relationship between the coefficient of variation of any ecological indicator (e.g., chlorophyll-*a* or total phosphorus) and water level is the main result demonstrating a reduction in

spatial variability due to floods. Considering physical, chemical or biotic data gathered in distinct habitats within the floodplain, this pattern was found in temperate and tropical regions, subjected to distinct levels of anthropogenic impacts, and at different spatial extents. The main mechanism that accounts for this pattern may be stated as follow. During low water level, the biological communities of each habitat in the floodplain (e.g., lagoons, backwater, streams) follow distinct temporal trajectories due to the effects of local driving forces (e.g., an efficient predator trapped in a lagoon but not in another). Management plans and biodiversity conservation in river floodplain systems will benefit by considering the effects of flood homogenization and increased connectivity peculiar to these unique ecosystems.

Handling editor: K. Martens

S. M. Thomaz (✉)  
Universidade Estadual de Maringá, Nupelia,  
Av. Colombo, 5790 Maringá, PR, Brasil  
e-mail: smthomaz@nupelia.uem.br

L. M. Bini  
Universidade Federal de Goiás, ICB, DBG, 74001-  
970 Goiânia, GO, Brasil  
e-mail: bini@icb.ufg.br

R. L. Bozelli  
Departamento de Ecologia, Universidade Federal do  
Rio de Janeiro, 21941-540 Rio de Janeiro, RJ, Brasil  
e-mail: bozelli@biologia.ufrj.br

**Keywords** Flood pulse · River-floodplain systems · Biodiversity

### Introduction

Hydrological regime is considered the key factor driving ecological functioning and biodiversity patterns in river floodplain systems (RFSs) (Junk et al., 1989; Neiff, 1990; Bunn & Arthington, 2002). Due to the dramatic rates of river regulation and habitat degradation worldwide (e.g., Agostinho

et al., 2000; Tockner et al., 2000a, b), investigations concerning how hydrological regimes affect attributes of floodplain systems, especially biodiversity, have become a central issue for river management in the last years.

Many RFSs have been studied intensively in the last two decades (see, for example Gopal et al., 2000; 2001) and different causal relationships between hydrology and ecosystem attributes have been proposed. Nevertheless, few patterns, that can be generalized for a wide variety of systems, at different latitudes and scales, were proposed (e.g., Carvalho et al., 2001; Ward & Tockner, 2001). In addition, the processes regulating biodiversity in these systems are not still well understood (Neiff, 2001).

The relationship between water level and landscape heterogeneity were studied in Neo-tropical (Furch & Junk, 1985; Hamilton & Lewis, 1990; Bozelli, 1992; Thomaz et al., 1997) and north temperate floodplains (Tockner & Ward, 1999; Tockner et al., 2000a, b; Ward & Tockner, 2001; Amoros & Bornette, 2002). In both regions, the among habitats (i.e., river channels, secondary channels and lagoons) variability of limnological variables (e.g., nitrate, conductivity, nutrients and chlorophyll-*a*) decrease with increasing water levels, suggesting that floods increase similarity among RFSs habitats. We consider this a good candidate to be a general pattern for RFSs.

In this paper we show an increase in homogeneity between river channel and a diverse array of floodplain aquatic habitats during floods through analyses of physical, chemical and biological data from Neo-tropical and temperate floodplains. Both empirical data and literature values are part of this analysis. Our goal is to show that the increase in habitat similarity during floods (or “flood homogenization”) can be considered a general pattern in RFSs. In addition, we consider the probable causes of this pattern and its consequences on attributes of RFSs, especially biodiversity maintenance. We believe that our theoretical framework will have high potential for the management and conservation of RFSs integrity.

### The pattern

RFSs are characterized by high habitat heterogeneity (e.g., Junk et al., 1989; Tockner et al., 2000a, b;

Drago et al., 2004). Besides several ecotonal habitats, which are found in the aquatic terrestrial transition zone (ATTZ) (Junk et al., 1989), RFSs have a wide variety of aquatic habitats, such as temporal and permanent lagoons, channels, backwaters, shallow lakes, sloughs, and the main river channel. The diversity of those habitats is of key importance for maintenance of high biological diversity in the system as a whole (i.e., gamma diversity).

Higher similarities among aquatic habitats during floods than during non-flood periods were found in several floodplains. In its most simple form, this homogeneity was first described as the existence of high similarity in physical, chemical or biological features between the main river and the backwaters of its floodplain (e.g., Furch & Junk, 1985, in the Amazonian floodplain; Hamilton and Lewis, 1990, in the Orinoco River floodplain; and Bozelli, 1992, in the Trombetas River floodplain). It was also explicitly quantified by the first time in the Upper Paraná River floodplain, when Thomaz et al. (1997) and later Agostinho et al. (2000), recognized the “homogenization effect of floods” using multivariate analyses.

In a non-exhaustive survey in the literature (Table 1) we found several other examples that illustrate the pattern. These examples show that it was consistently repeated in different spatial extents in several ecosystems.

A first group of factors, which tend to be more similar during the flood events, are the physical and chemical factors. Among these factors, nutrients (both nitrogen and phosphorus) and conductivity have been considered in several studies (Table 1). Similarity of chlorophyll-*a* concentrations of lagoons and the river channel also increases during flood periods (Table 1).

In addition to the limnological characteristics cited previously, floods seem to decrease community turnover or beta diversity. The composition of phytoplankton, zooplankton, fish and macrophytes are less similar among habitats during low water, but they are more similar during floods. Spatial variability in planktonic richness and abundance is also lessened by flood events (Table 1).

Studies indicate that the homogenization effect of floods operate at different spatial extents. For example, at the scale of whole waterbody, different

**Table 1** Data confirming the increase of similarity within and between habitats during floods recorded in several floodplains

Floodplain	Factors studied	Ecosystem/community responses
Amazon (Brazil) (pristine)	Abiotic	DOC, inorganic carbon and electrical conductivity values were more heterogeneous among stations of a same lake and between the lake and the river during low water (1)
Orinoco (Venezuela) (pristine)	Abiotic and biotic	Chemical characteristics, suspended load and plankton composition are more dissimilar among lakes during low water (2); structure of fish assemblage in lakes tend to converge as flood water enters lakes (3)
Trombetas (Brazil) (subject to impact by bauxite tailings)	Abiotic and biotic	Limnological abiotic factors, such as phosphorus, nitrogen and chlorophyll- <i>a</i> concentrations (4), phytoplankton (5) and zooplankton (6) densities are more similar between sampling stations of a same lake and between the lake and Trombetas River during high water
Pantanal – Piraim River (Brazil) (pristine)	Abiotic	The coefficient of variation of limnological factors, such as phosphate and ammonium, obtained in different sampling stations of a inundation field are lower during inundations (7)
Pantanal – Paraguay River (Brazil) (pristine)	Abiotic	Electrical conductivity and oxygen are more similar between a lake and the river after connection (8)
Upper Paraná (Brazil) (subject to water level control by reservoirs' operation)	Abiotic and biotic	Limnological abiotic factors (e.g., phosphorus, nitrogen, chlorophyll, oxygen etc.), summarized through principal component analyses (9), and the structure of zooplankton (10) and fish (11) assemblages are more similar among lagoons and between lagoons and Paraná main channel during high waters
Middle Paraná (Argentina) (subject to low degree of water level control)	Biotic	Composition of aquatic macrophyte assemblage in different lagoons are more dissimilar during low waters (12)
Mogi Guaçu River (Brazil) (impacted by pollution and deforestation)	Abiotic and biotic	During drought period the littoral region of a lagoon differed more from the other sites, indicating a higher within lagoon heterogeneity during this period (13); similarity of the fish assemblages between lagoons were higher at the wet season (14);
Acre River (Brazil) (pristine)	Abiotic	8 limnological parameters (temperature, conductivity, total seston, ammonium, nitrate, phosphate, silicate and total pigments) out of 11 analyzed in three sampling stations of Amapá lake had higher coefficient of variation during low water period (15);
Danube (Austria) (subject to high degree of water level control)	Abiotic	Coefficient of variation of nitrate and chlorophyll obtained in different sites (between 35 and 49) are lower during high water (16)
Illinois (USA) (subject to high degree of water level control)	Biotic	Nitrification and denitrification are more variable among sites during low water periods (17)
Mississippi (USA)	Biotic	Annual floods homogenize the floodplain and promote connectivity to various degrees, allowing for fish exchanges between river and floodplain that directly affect community composition (18)

(1) Funk & Junk (1985); (2) Hamilton & Lewis (1990); (3) Lewis et al. (2000); (4) Panosso and Kubrusly (2000); (5) Huszar (2000); (6) Bozelli (1992, 2000); (7) David (2001); (8) Calheiros & Hamilton (1998); (9) Thomaz et al. (1997); (10) Bonecker et al. (1998); (11) Verissimo (1994); (12) Neiff (1979); (13) Taniguchi et al. (2004); (14) Meschiatti et al. (2000); (15) Almeida (2000); (16) Tockner et al. (2000a, b); (17) M. Lemke (unpublished data); (18) Miranda (2005)


sampling sites at the same lagoon tend to be more similar during periods of high water (e.g., in the Pantanal and Batata Lake – Table 1). At a broader spatial extent, different lagoons of a same

floodplain and the lagoons of a floodplain and their main river (e.g., Orinoco, Paraná, Amazon, Danube – Table 1) tend to be more similar during high water periods.

To illustrate the occurrence of this pattern across different spatial extents we used the coefficient of variation (CV) in limnological data obtained in (i) three sampling stations at a single lake (Lake Batata) every three months during 8 years and (ii) in ten lakes and two sampling sites of Trombetas River obtained in a single year at the high and low water levels. All lakes, including Lake Batata, are in the same region (Amazon) and they are connected with the Trombetas River.

For Lake Batata, a significant negative relationship ( $P < 0.05$ ) was observed between water level and CV values of water temperature, suspended matter, site depth, conductivity, pH and chlorophyll-*a* (Fig. 1). These results show that high water levels create more similar sites within this lake.

The same pattern was observed at a broader spatial extent, considering the data for the 10 lakes plus Trombetas River sites. Coefficients of variation estimated for dissolved organic carbon,



**Fig. 1** Coefficients of variation (CV) estimated for the limnological variables obtained in three sampling sites of Lake Batata (Trombetas River floodplain, Amazon)

plotted against water level. Each point represents the spatial CV estimated with the data obtained in these three sampling sites in the same day

dissolved phosphorus, dissolved nitrogen, conductivity, turbidity, depth, Secchi disk and suspended matter were higher during low water period (Fig. 2). Thus, lakes are more similar among themselves and with Trombetas River when they are connected.

The pattern seems to occur also in RFSs subjected to different degrees of conservation (Table 1). For example, the Pantanal, Amazon and Orinoco are representative of relatively pristine RFSs, whereas Paraná and Danube are subject to impacts associated with water level control. The homogenization effect was also observed in lakes subject to other types of impact, as is the case of Lake Batata (Trombetas River), which is highly impacted by bauxite tailings (Roland, 2000; Guenther & Bozelli, 2004).

### Main mechanisms


Trying to untangle the ecological complexity of RFSs and the main mechanisms that produce the patterns found in nature are not easy tasks. Nevertheless, we present empirical evidence from several sources that suggest some explanations as

more plausible than others. We first hypothesize that these patterns are due to (i) increasing dissimilarities during low water levels caused by local driving forces and (ii) increasing connectivity during high water periods. We will discuss these factors separately below.

#### The influence of local driving forces during low waters

The floodplain aquatic habitats are isolated from each other and subject to local driving forces during low water periods (Thomaz et al., 1997; Camargo & Esteves, 1995; Tockner et al., 1999; Lewis Jr. et al., 2000; Carvalho et al., 2001). These driving forces inducing heterogeneity include: (i) water inputs from lateral tributaries and/or seepage leading to localized physical and chemical characteristics that are basin-specific; (ii) wind and animal induced sediment re-suspension, which affects water bodies according to their morphometry, and (iii) differences in ecological succession. These local forces act with different intensities in the floodplain landscape, thus creating habitats with different characteristics.

Concerning aspect (i), the water's origin determines the water temperature, turbidity and nutrient content, which greatly influence habitat heterogeneity, plant and animal recruitment, and ecosystem productivity. For example, in the Orinoco floodplain, a great heterogeneity develops at smaller scales during the dry season, when the floodplain is fragmented. This time, as can be seen from major ion data, shows a significant but variable contribution of local waters (rain and runoff) (Hamilton & Lewis, 1990). The importance of local tributaries contributions has been demonstrated in several other RFSs (Forsberg et al., 1988; Schemel et al., 2004; Souza-Filho et al., 2004; Thomaz et al., 1997). Data from the Amazon floodplain show that the percentage of local water in a lake, during low waters, depend on the ratio of drainage basin area (BA) to lake area (LA): the greater the BA: LA ratio the more river water was apparently flushed out of a lake and replaced by local water during low waters (Forsberg et al., 1988). The examples from these large basins highlight the importance of lateral tributaries as sources of variation in the floodplain



**Fig. 2** Coefficients of variation estimated for selected limnological variables obtained in ten lakes in the Trombetas River floodplain and in the Trombetas River main channel (Amazon) in the high and low water periods. Each point represents the spatial CV obtained among these ten lakes and Trombetas River in one sampling period. DOC = dissolved organic carbon; DRP = dissolved reactive phosphate; DN = dissolved nitrogen; COND = conductivity; TURB = turbidity; Z = depth; SEC = Secchi disk; TEMP = temperature; PH = pH and SM = suspended matter

habitats and suggest that they may contribute to a large extent to increase the habitat heterogeneity during low waters in RFSs.

In addition to lateral flow, the water of floodplain habitats also originates from seepage. This water has chemical features that are quite different from that of the river channel and it affects the habitats where it emerges upwells. The groundwater is loaded with high concentrations of bio-available solutes in the hyporheic zone and it affects the production and biodiversity of benthos and riparian vegetation. Dynamic convergence of aquifer-riverine components adds physical heterogeneity and functional complexity to floodplain landscapes and the major landscape features of alluvial floodplains include a dynamic mosaic of stream channels and wetlands fed by hyporheic flow (Standford & Ward, 1993). Results obtained in the Val Roseg floodplain (Switzerland) indicate that the concentration of nutrients have small coherence (i.e., low synchrony) in lateral water bodies due to dominance of local hydrological and/or biogeochemical processes, like groundwater upwelling (Tockner et al., 2002). It has been shown that aquatic floodplain habitats that are fed mainly by deep and shallow alluvial groundwater, hyporheic exfiltration, have high variability in terms of both substrate characteristics and faunal compositions (Brunke et al., 2003). Accordingly, Gurnell & Petts (2002) considered a variable mix of river water with surface and groundwater among the causes for vegetation heterogeneity in several European RFSs. It is expected that such increase in habitats heterogeneity, caused by hyporheic water upwelling is more accentuated during low water phases, when flow (and thus, surface connectivity) is lower. In summary, lateral tributaries and hyporheic water largely contribute to the high habitat diversity found in RFSs and such heterogeneity is maintained especially during low water periods.

The importance of wind, in combination with morphometry, (aspect ii above) was shown in the Orinoco floodplain. There an important transition occurred when the mean depth of a lake decreased to such an extent that fine sediments on the bottom of the lake were resuspended by wind-generated currents (Lewis Jr. et al., 2000). In addition to depth, effective fetch, in combination with wind velocity, was important to cause sediment resus-

pension during periods of isolation (Hamilton & Lewis, 1990). These internal loadings, caused by sediment resuspension, are very important in floodplain lagoons during low water periods (Carvalho et al., 2001; Thomaz et al., 1997). In fact, resuspension appears to drive plankton succession during these periods (Train & Rodrigues, 2004).

Resuspension of sediments in shallow lakes may also be caused by other factors, such as animals. In the Upper Paraná floodplain, shallow temporary lagoons (depths < 0.5 m) are subject to cattle and capybaras (the rodent *Hydrochoerus hydrochaeris*) trampling, leading to the highest values of conductivity and nutrient concentrations found among all floodplain water bodies during low water periods (Pagioro et al., 1997; Thomaz et al., 1997; Okada, 1995; Okada et al., 2003).


The morphometry of the water body affects the assemblages of large-bodied organisms. In the Orinoco RFS, the structure of fish assemblages has been largely explained by water transparency that is directly affected by lake area and depth (Rodríguez & Lewis Jr., 1997). In lagoons of the Upper Paraná RFS, depth is an important determinant of fish assemblage structure (Okada, 1995; Okada et al., 2003) and, together with littoral slope, of aquatic macrophyte assemblage structure (Santos, 2004). Floodplain lagoons vary widely in morphometry and wind exposure (fetch) and thus their physical and chemical, as well as biological characteristics are more variable during low water periods.

Another important ecological factor leading to higher dissimilarities in aquatic habitats during low waters is ecological succession (aspect iii). After isolation, each aquatic environment and even specific regions of a same environment follow their own succession whose sequence depends mainly upon the identity of the propagules transported to the water body during the last flood and on the response of these organisms to the localized environmental characteristics that will predominate in the lagoons. This individualistic (Gleasonianan) succession leads to the unique community development in each habitat subsequently creating increased dissimilarity during low waters. Thus, chance (or stochastic events ("who arrives there")) may drive different ecological successions in the habitats.

Even considering the influence of a shared regional species pool, the magnitude of the processes that determine assemblage structures (e.g., competition and predation) will differ from lake to lake, even if we consider a similar start point (Fig. 3). If, during low water levels, we expect a high inter-habitat heterogeneity, we also can infer that, during this period, there is a low level of “spatial predictability” or spatial autocorrelation in the attributes analyzed. Therefore, knowing the attributes of a given lagoon will not help to predict the attributes of another lagoon near to the first. After floods (overflow), we expect a time lag until the effects of homogenization become apparent. This time lag will depend mainly on the intensity of the floods and on the rate of dispersion/migration of the different species that occur in the RFSs. Also, the increase in dissimilarity, after isolation, will be not instantaneous. This time lag will depend, among many other factors, on the intensity of drought and on the species composition that was assembled during the previous high water level period (Fig. 3).

Evidences of the different assemblage trajectories (as seen in Fig. 3) come from several tax-

onomic groups. For example, the rate of succession of vegetation was found to be different in several RFSs across Europe (Jarolimek et al., 2001; Gurnell & Petts, 2002). In the Middle (Neiff, 1979) and Upper Paraná RFS (Santos & Thomaz, in press), the differences in the structure of aquatic macrophyte communities observed among lagoons increased during low waters. This higher dissimilarity depended on the successional course in combination with the reproductive rate of the species and waterbody morphometry (especially depth and wind exposure; Neiff, 1979). Since floods may impede the regrowing and reestablishment of some species of aquatic macrophytes, the interval between floods is also an important determinant of succession of macrophytes in floodplain lagoons (Maltchik et al., 2004). Differences in aquatic plant cover and species composition may lead to different plankton (Cremer et al., 2004; Lansac-Tôha et al., 2004; Izaguirre et al., 2004) and fish assemblages (Pelicice et al., 2005). Thus, the effects of drought increasing the heterogeneity of plankton and fish communities may be also mediated by changes on plant communities.



**Fig. 3** Hypothesized spatial variability in floodplains along a hydrological cycle. Predicting the temporal trajectories and the mean values of the attributes for every lake in the floodplain (numbered squares) is not a simple task during low water periods (scenario a). However, the high level of spatial autocorrelation caused by floods may increase substantially our predictability capacity: knowing the attributes of a community in a given

lake permits the prediction of these attributes in the other lakes (scenario b). Again, after isolation, spatial variability increases (each community may follow a distinct path) due to the local driving forces (e.g., a large number of a predator trapped in one lake but not in the others; scenario c). There is a time lag that contributes to the transitions between scenarios



For fish communities, the similarity between lagoons is low after isolation and a conspicuous difference from the original fauna was found during that period (Rodríguez & Lewis Jr., 1994; Okada et al., 2003). Biotic interactions (e.g., predation by fish and birds and competition) played a major role in these fish assemblages, leading to distinct temporal trajectories in different lagoons of a same RFS (Okada et al., 2003). In two lakes of the Orinoco RFS, the magnitude and direction of the seasonal change of the fish assemblages during the dry season were comparable across years but this temporal change was small relative to differences among lakes (Rodríguez & Lewis Jr., 1994). It shows that in lakes of this RFS there is a strong deterministic component structuring fish assemblages, but it leads to different assemblages in separated lakes. In other words, local processes operating at the habitat scale are the major determinants of the structure of biological communities during the dry season (Rodríguez & Lewis Jr., 1997).

Thus, we may also assume that different paths lead to different communities in water bodies during the low water phases, contributing to increased dissimilarities among habitats (i.e., increased beta diversity).

#### The influence of regional driving force (flooding) during high waters

Increase in river water level enhances the connection between the river and the floodplain aquatic habitats as well as among floodplain habitats. During high water phase, horizontal fluxes are produced from the river course to the floodplain (Neiff, 2001). This increase in connectivity enhances the exchange of water, sediments, minerals and organisms among the different habitats in the floodplain (Agostinho & Zalewski, 1995; Lewis Jr. et al., 2000; Neiff, 2001; Aspetsberger et al., 2002; Zalocar de Domitrovic, 2003; Bonecker et al., 2005). Planktonic organisms may also move freely and passively within and between aquatic habitats. For example, fish larvae have passive dispersion towards the floodplain water bodies during flood pulses (Araujo-Lima & Oliveira, 1998; Nakatani et al., 2004).

Even conspicuous spatial differences within individual lagoons (e.g., comparing areas covered with aquatic macrophytes with bared areas) may be reduced due to the homogenization effects of floods. For example, non-planktonic rotifers have been found in the pelagic zone of floodplain lagoons during high waters as a result of interaction with the littoral during these periods (José de Paggi, 1993; Bonecker & Lansac-Tôha, 1996; Garcia et al., 1998). Similarity between river channel and floodplain lagoons also increases for this group during high waters due to increased connectivity between both habitats (Bonecker et al., 1998). The importance of horizontal exchanges during floods was shown in the Upper Paraná RFS, where the highest species richness of zooplankton was found in connected lagoons, subject to greater water exchanges than isolated lagoons (Aoyagui & Bonecker, 2004).

Large-bodied organisms move passively or actively during high water periods. For example, propagules of aquatic macrophytes disperse passively (e.g., free floating species, that are common in RFSs; Thomaz et al., 2004) but fish disperse actively when the habitats are connected (Rodríguez & Lewis Jr., 1994; Agostinho & Zalewski, 1995). These examples highlight the importance of floods for passive or active exchanges of organisms and propagules among RFS water bodies.

Nevertheless, the increase in similarity between the river channel and floodplain habitats depends on overbank flooding. On the other hand, the ecosystem expansion without overbank flooding ("flow pulses", according to Tockner et al., 2000a), may lead to high similarities in the two first scales considered (within floodplain aquatic habitats and within different sites of a same lagoon) but not between these water bodies and the river channel. These predictions still need to be tested.

#### Possible consequences for biodiversity maintenance

Understanding the consequences of flooding for biology within a RFS, including dispersal, adaptation and ecological succession is of paramount importance to develop efficient conservation and management strategies (Ward & Tockner, 2001). In this sense, our survey in the literature may



bring important insights because they were derived largely from a variety of near natural South-American river-floodplain systems.

It has been proposed that species richness is maximized in RFSs due to the intermediate degree of three inter-related variables: disturbance (Intermediate Disturbance Hypothesis, *sensu* Connell, 1978), ecotone occurrence and hydrological connectivity (Ward & Tockner, 2001). At low connectivity, species diversity should be reduced by the absence of the fluvial dynamics that sustains a diversity of successional stages within RFSs, whereas excessive connectivity will keep all communities in pioneer stages (Ward & Tockner, 2001). Other studies have found that diversity of different groups (fish, macrophytes, macrozoobenthos and amphibians) peaks differently across the gradient of connectivity intensity (Tockner et al., 2000a).

Despite these considerations, we emphasize that even if the highest diversity is reached at intermediate levels of disturbance, intensive floods (which promotes high connectivity) are important for exchange of propagules, nutrients, and organisms among habitats. Such exchanges during high waters (see Lewis Jr. et al., 2000), are mandatory to keep high biodiversity unique of RFSs. This is because the exchanges increase the probability of rare species to disperse and colonize new sites during floods. This statement is even more important if we consider that some species need specific types of habitats to succeed. For example, the successful recruitment of the tree *Calophyllum brasiliense* in the Meander forest (Amazon) requires dispersal to early successional habitats (King, 2003). It is also demonstrated that fish larvae depend on lagoons with appropriate habitat complexity to succeed (Agostinho & Zalewski, 1995; Nakatani et al., 2004).

The short-term exchanges following events of high connectivity cause, at first, a decrease in beta diversity of RFSs (the “homogenization effect”). Nevertheless, they are key to keep long-term high species richness of floodplains (gamma diversity), complementing the role of intermediate disturbances. Consequently, it is expected that artificial flow reductions that lead to reduced connectivity, have serious deleterious effects upon long term RFS biodiversity. The importance of connec-

tivity has lead to its adoption as a natural guideline to design strategies for RFSs rehabilitation (Pedroli et al., 2002). It has been observed, for example, that this process has a key importance in the recover of Lake Batata (Trombetas River – Amazon) from the impact it suffered with bauxite tailings. It is caused basically by organic matter and organism inocula exchanges between the lake and the Trombetas River main channel (RL Bozelli, unpublished data).

It is well known that flow is often a principal river attribute that is changed by different watershed uses (e.g., reservoir construction, hydroways construction, levee alterations and water obstruction; Bunn & Arthington, 2002; Sheldon et al., 2002; Souza-Filho et al., 2004). Great dissimilarity of the fish assemblages among lagoons in the Upper Paraná RFS was recorded after a long period of reduced connectivity, caused by flow reduction associated with reservoir operations (Oliveira et al., 2001). These findings show that the impacts of reservoir operation upon fish assemblages are in accordance with our expectations that in the absence of floods, floodplain habitats are more heterogeneous. Although flow alteration may affect differently the groups of organisms (e.g., long lived aquatic macrophytes versus short lived phytoplankton), interruption of connectivity is deleterious to the whole biodiversity in medium or long term periods due to the reduction in propagules exchanges between habitats. In addition, prolonged low water periods may lead to maximization of competitive exclusion, leading to further local extinctions. It has been shown, for example, that in disconnected forests competitive species may dominate in the absence of floods (van Looy et al., 2004) whereas when floods occur coexistence of several species is possible (Deiller et al., 2001). Accordingly, investigations in a South Brazilian floodplain lagoon showed that the absence of dominance in aquatic macrophyte community was associated with highest number of flood events (Maltchik et al., 2005).

Flow reduction may also contribute to biological invasions by alien species that can cause extinction of native species (Bunn & Arthington, 2002; Agostinho et al., 2005). Large-scale surveys carried out in the Paraná basin showed that

introduced fish succeed in basins with high endemism and/or regulated by dams (Agostinho et al., 2005). Thus, considering that species introductions are considered one of the main causes of extinctions (Clavero & García-Berthou, 2005), the increase in exotic species following flow regulation may be considered a matter of concern for biodiversity maintenance in RFSs.

Complementarity is an important and innovative concept in conservation biology (e.g., Justus & Sarkar, 2002). Basically, it is a measure of how an area possesses species not yet found in other areas. Thus, some areas with low species richness may have high complementarity value due to the presence of rare species. Paradoxically, long droughts may increase the number of areas to be “reserved” (network length) due to the high spatial turnover in species composition during these periods. However, we suspect that the higher spatial turnover during low water periods is associated with low species richness. Thus, this innovative concept should be applied with caution to design networks of reserves in floodplain landscapes. It will be interesting to compare the spatial configuration and the length of the networks that could be generated in high and low water periods.

## Perspectives

Based on several examples collected in different spatial scales and several floodplains, we suggest that the “homogenization of habitats” following floods may be considered a general pattern. The homogenization flood effect is derived from increased connectivity (regional driving force) during high waters, complemented by high isolation (when local driving forces dominate) during low waters. In this way, we suggest as future possible fields for research: (i) analyses of similarity employing multivariate methods in different periods of the flood pulse; (ii) specific studies of organisms movement across river and floodplain habitats during high waters and (iii) studies of succession dynamics and how it affects alpha and beta diversity of the floodplain habitats during low waters. Given the above links between these processes and RFSs’ biodiversity, we believe that these approaches would be helpful in

conserving the unique richness of these fragile and endangered ecosystems.

**Acknowledgments** We acknowledge with appreciation the critical reading and English corrections made by Dr. Michael Lemke (University of Illinois at Springfield) and Dr. Doerthe Mueller-Navarra (University of Hamburg), and the suggestions by Dr. Rosemara Fugì (Nupélia – University of Maringá), MSc. Fernando Mayer Pelicice and MSc. Josilaine Kobayashi (Graduate students at the Program of Ecology in Inland Waters – University of Maringá). The authors acknowledge the Brazilian Council of Research (CNPq) for providing Research Productivity grants and funds for the Long Term Ecological Research Program at the Upper Paraná River Floodplain. RL Bozelli also thanks the colleagues from the Laboratory of Limnology (Federal University of Rio de Janeiro), IBAMA and Mineração Rio do Norte for funding his research at Lake Batata and Trombetas River.

## References

- Agostinho, A. A. & M. Zalewski, 1995. The dependence of fish community structure and dynamics on floodplain riparian ecotone zone in Paraná River, Brazil. *Hydrobiologia* 303: 141–148.
- Agostinho, A. A., S. M. Thomaz & L. C. Gomes, 2005. Conservation of the biodiversity of Brazil’s inland waters. *Conservation Biology* 19: 646–652.
- Agostinho, A. A., S. M. Thomaz, C. V. Mente-Vera & K. O. Winemiller, 2000. Biodiversity in the high Paraná River floodplain. In Gopal B., W. J. Junk & J. A. Davis (eds), *Biodiversity in wetlands: assessment, function and conservation*. Backhuys Publishers, Leiden, 89–118.
- Almeida, J. S., 2000. Influência do regime hidrológico sobre algumas variáveis limnológicas em um lago da planície de inundação do rio Acre (Lago Amapá, Acre, Brasil). Master Thesis. PPG em Ecologia e Manejo de Recursos Naturais, Universidade Federal do Acre.
- Amoros, C. & G. Bornette, 2002. Connectivity and bio-complexity in waterbodies of riverine floodplains. *Freshwater Biology* 47: 761–776.
- Aoyagui, A. S. M. & C. C. Bonecker, 2004. Rotifers in different environments of the Upper Paraná River floodplain (Brazil): richness, abundance and the relationship with connectivity. *Hydrobiologia* 522: 281–290.
- Araujo Lima, C. A. R. M. & E. C. Oliveira, 1998. Transport of larval fish in the Amazon. *Journal of Fish Biology* 53: 297–306.
- Aspetsberger, F., F. Huber, S. Kargl, B. Scharinger, P. Peduzzi & T. Hein, 2002. Particulate organic matter dynamics in a river floodplain system: impact of hydrological connectivity. *Archiv für Hydrobiologie* 156: 23–42.
- Bonecker, C. C. & F. A. Lansac-Tôha, 1996. Community structure of rotifers in two environments of the Upper River Paraná floodplain (MS) – Brazil. *Hydrobiologia* 325: 137–150.

- Bonecker, C. C., C. L. da Costa, L. F. M. Velho, F. A. Lansac-Tôha, 2005. Diversity and abundance of the planktonic rotifers in different environments of the Upper Paraná River floodplain (Paraná State – Mato Grosso do Sul State, Brazil). *Hydrobiologia* 546: 405–414.
- Bonecker, C. C., F. A. Lansac-Tôha & D. Rossa, 1998. Planktonic and non-planktonic rotifers in two environments of the Upper Paraná River floodplain, State of Mato Grosso do Sul, Brazil. *Brazilian Archives of Biology and Technology* 41: 447–456.
- Bozelli, R. L., 1992. Composition of the zooplankton community of Batata and Mussura lakes and the Trombetas River, State of Paraná, Brazil. *Amazoniana-Limnologia et Oecologia Regionalis Systemae Fluminis Amazonas* 12: 239–261.
- Bozelli, R. L., 2000. Zooplâncton. In Bozelli R. L., F. A. Esteves & F. Roland (eds), *Impacto e recuperação de um ecossistema amazônico*. UFRJ/SBL, Rio de Janeiro: 121–138.
- Brunke, M., E. Hoehn & T. Gonser, 2003. Patchiness of river-groundwater interactions within two floodplain landscapes and diversity of aquatic invertebrate communities. *Ecosystems* 6: 707–722.
- Bunn, S. E. & A. H. Arthington, 2002. Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental Management* 30: 492–507.
- Calheiros, D. F. & S. K. Hamilton, 1998. Limnological conditions associated with natural fish kills in the Pantanal wetland of Brazil. *Verhandlungen Internationale Vereinigung für theoretische und angewandte Limnologie* 26: 2189–2193.
- Camargo, A. F. M. & F. A. Esteves, 1995. Influence of water level variation on fertilization of an oxbow lake of Rio Mogi-Guaçu, state of São Paulo, Brazil. *Hydrobiologia* 299: 185–193.
- Carvalho, P., L. M. Bini, S. M. Thomaz, L. G. Oliveira, B. Robertson, W. L. G. Tavechio & A. J. Darwisch, 2001. Comparative limnology of South-American lakes and lagoons. *Acta Scientiarum* 23: 265–273.
- Clavero, M. & E. García-Berthou, 2005. Invasive species are a leading cause of animal extinctions. *Trends in Ecology and Evolution* 20: 110.
- Connel, J. H., 1978. Diversity in tropical rain forest and coral reefs. *Science* 199: 1302–1310.
- Cremer, H., A. D. Buijse, A. F. Lotter, W. Oosterberg & M. Staras, 2004. The palaeolimnological potential of diatom assemblages in floodplain lakes of the Danube Delta, Romania: a pilot study. *Hydrobiologia* 513: 7–26.
- David, G. Q., 2001. Dinâmica de nutrientes em um campo inundável no Pantanal Mato-grossense, Brasil. Master Thesis, PPG em Ecologia, Universidade Federal, Cuiabá.
- Deiller, A. F., J. M. Walter & M. Tremolieres, 2001. Effects of flood interruption on species richness, diversity and floristic composition of woody regeneration in the upper Rhine alluvial hardwood forest Regulated Rivers-Research & Management 17: 393–405.
- Drago, I. E., M. Marchese & K. M. Wantzen, 2004. Benthos of a large neotropical river: spatial patterns and species assemblages in the Lower Paraguay and its floodplain. *Archiv für Hydrobiologie* 160: 347–374.
- Forsberg B. R., A. H. Devol, J. E. Richey, L. A. Martinelli & H. Santos, 1988. Factors controlling nutrient concentrations in Amazon floodplain lakes. *Limnology and Oceanography* 33: 41–56.
- Furch, K. & W. J. Junk, 1985. Dissolved carbon in a floodplain lake of the Amazon and in the river channel. *Mitteilungen des Geologisch-Paläontologischen Institutes der Universität Hamburg* 58: 285–298.
- Garcia, A. P. P., F. A. Lansac-Tôha & C. C. Bonecker, 1998. Species composition and abundance of rotifers in different environments of the floodplain of the Upper Paraná River, Brazil. *Revista Brasileira de Zoologia* 15: 327–343.
- Gopal, B., W. J. Junk & J. A. Davis, 2000. Biodiversity in wetlands: assessment, function and conservation. Backuys Publishers, Leiden.
- Gopal, B., W. J. Junk & J. A. Davis, 2001. Biodiversity in wetlands: assessment, function and conservation. Backuys Publishers, Leiden.
- Guenther, M. & R. Bozelli, 2004. Effects of inorganic turbidity on the phytoplankton of an Amazonian Lake impacted by bauxite talings. *Hydrobiologia* 511: 151–159.
- Gurnell, A. M. & G. E. Petts, 2002. Island-dominated landscapes of large floodplain, a European perspective. *Freshwater Biology* 47: 581–600.
- Hamilton, S. K. & W. M. Lewis, 1990. Basin morphology in relation to chemical and ecological characteristics of lakes on the Orinoco River floodplain, Venezuela. *Archiv für Hydrobiologie* 119: 393–425.
- Huszar, V. L. M., 2000. Fitoplâncton. In Bozelli R. L., F. A. Esteves & F. Roland (eds), *Impacto e recuperação de um ecossistema amazônico*. UFRJ/SBL, Rio de Janeiro, 89–104.
- Izaguirre, I., I. ÓFarrel, F. Unrein, R. Sinistro, M. D. Afonso & G. Tell, 2004. Algal assemblages across a wetland, from a shallow lake to relictual oxbow lakes (Lower Paraná River, South America). *Hydrobiologia* 511: 25–36.
- Jarolimek, I., V. Banasova, H. Otahelova & M. Zaliberova, 2001. Nine year succession of the herbaceous floodplain vegetation in the Morava river. *Ekologia-Bratislava* 20: 92–100.
- José de Paggi, S., 1993. Composition and seasonality of planktonic rotifers in limnetic and littoral regions of a floodplain lake (Paraná River system). *Revue d'Hydrobiologie Tropicale* 26: 53–63.
- Junk, W. J., P. B. Bayley & R. E. Sparks, 1989. The flood pulse concept in river-floodplain systems. *Canadian Special Publication of Fisheries and Aquatic Sciences* 106: 110–127.
- Justus, C. & S. Sarkar, 2002. The principle of complementarity in the design of reserve networks to conserve biodiversity: a preliminary history. *Journal of Biosciences* 27(4): 421–435.

- King, R. T., 2003. Succession and micro-elevation effects on seedling establishment of *Calophyllum brasiliense* camb. (Clusiaceae) in an Amazonian river meander forest. *Biotropica* 35: 462–471.
- Lansac-Tôha F. A., C. C. Bonecker & L. F. M. Velho, 2004. Composition, species richness and abundance of the zooplankton community. In Thomaz S. M., A. A. Agostinho & N. S. Hahn (eds), *The Upper Paraná River and its floodplain: Physical aspects, ecology and conservation*. Backhuys Publishers, Leiden, 145–190.
- Lewis Jr., W. M., S. K. Hamilton, M. A. Lasi, M. Rodríguez, J. F. Saunders III, 2000. Ecological determinism on the Orinoco floodplain. *BioScience* 50: 681–692.
- Maltchik, L., A. S. Rolon & C. Groth, 2004. The effects of flood pulse on the macrophyte community in a shallow lake of Southern Brazil. *Acta Limnologica Brasiliensia* 16: 103–113.
- Maltchik, L., G. R. de Oliveira, A. S. Rolon & C. Stenert, 2005. Diversity and stability of aquatic macrophyte community in three shallow lakes associated to a floodplain system in the south of Brazil. *Interciencia* 30: 166–170.
- Meschiatti, A. J., M. S. Arcifa & N. Fenerich-Verani, 2000. Fish communities associated with macrophytes in Brazilian floodplain lakes. *Environmental Biology of Fishes* 58: 133–143.
- Miranda, L. E., 2005. Fish Assemblages in Oxbow Lakes Relative to Connectivity with the Mississippi River. *Transactions of the American Fisheries Society* 134: 1480–1489.
- Nakatani, K., A. Bialeztki, G. Baumgartner, P. V. Sanches & M. C. Makrakis, 2004. Temporal and spatial dynamics of fish eggs and larvae. In Thomaz S. M., A. A. Agostinho & N. S. Hahn (eds), *The Upper Paraná River and its floodplain: Physical aspects, ecology and conservation*. Backhuys Publishers, Leiden, 293–308.
- Neiff, J. J., 1979. Fluctuaciones de la vegetacion acuatica en ambientes del valle de inundacion del Parana Medio. *Physis* 38: 41–53.
- Neiff, J. J., 1990. Ideas para la interpretacion ecológica del Parana. *Interciencia* 15: 424–441.
- Neiff, J. J., 2001. Diversity in some tropical wetland systems of South América. In Gopal B., W. J. Junk & J. A. Davis (eds), *Biodiversity in wetlands: assessment, function and conservation*. Backhuys Publishers, Leiden, 157–186.
- Okada, E. K., 1995. Diversidade e abundância de peixes em corpos de água sazonalmente isolados na planície alagável do alto rio Paraná e fatores ambientais relacionados. Master Thesis, PPG Ecologia de Ambientes Aquáticos Continentais, Universidade Estadual de Maringá.
- Okada, E. K., A. A. Agostinho, M. Petrere Jr. & T. Penczak, 2003. Factors affecting fish diversity and abundance in drying ponds and lagoons in the Upper Paraná River basin, Brazil. *Ecohydrology & Hydrobiology* 3: 97–110.
- Oliveira, E. F., E. A. Luiz, A. A. Agostinho & E. Benedito-Cecílio, 2001. Fish assemblages in littoral areas of the Upper Paraná River floodplain, Brazil. *Acta Scientiarum* 23: 369–376.
- Pagioro, T. A., M. C. Roberto, F. A. Lansac-Tôha, 1997. Comparative limnological analysis of two lagoons on the floodplain of the Upper Paraná River, Brazil. *International Journal of Environmental Sciences* 23: 229–239.
- Panosso, R. & L. Kubrusly, 2000. Avaliação espacial e temporal das variáveis limnológicas básicas e nutrientes. In Bozelli R. L., F. A. Esteves & F. Roland (eds), *Impacto e recuperação de um ecossistema amazônico*. UFRJ/SBL, Rio de Janeiro, 55–71.
- Pedroli, B., G. de Blust, K. van Looy & S. van Rooij, 2002. Setting targets in strategies for river restoration. *Landscape Ecology* 17: 5–18.
- Pelice, F., A. A. Agostinho & S. M. Thomaz, 2005. Fish assemblages associated with *Egeria* in a tropical reservoir: investigating the effects of plant biomass and diel period. *Acta Oecologica* 27: 9–16.
- Rodríguez, M. A. & W. M. Lewis Jr., 1994. Regulation and stability in fish assemblages of neotropical floodplain lakes. *Oecologia* 99: 166–180.
- Rodríguez, M. A. & W. M. Lewis Jr., 1997. Structure of fish assemblages along environmental gradients in floodplain lakes of the Orinoco River. *Ecological Monographs* 67: 109–128.
- Roland, F., 2000. Produção primária fitoplanctônica. In Bozelli R. L., F. A. Esteves & F. Roland (eds), *Impacto e recuperação de um ecossistema amazônico*. UFRJ/SBL, Rio de Janeiro, 105–117.
- Santos, A. M., S. M. Thomaz, in press. Aquatic macrophytes diversity in lagoons of a tropical floodplain: the role of connectivity and water level. *Austral Ecology*.
- Santos, A. M., 2004. Macrófitas aquáticas da planície de inundação do alto rio Paraná: produtividade primária, decomposição, ciclagem de nutrientes e diversidade. Doctoral Thesis, PPG Ecologia de Ambientes Aquáticos Continentais, Universidade Estadual de Maringá.
- Schemel, L. E., T. R. Somme, A. B. Muller-Solger & W. C. Harrel, 2004. Hydrologic variability, water chemistry and phytoplankton biomass in a large floodplain of the Sacramento River, CA, US. *Hydrobiologia* 513: 129–139.
- Sheldon, F., A. J. Boulton & J. T. Puckridge, 2002. Conservation value of variable connectivity: aquatic invertebrate assemblages of channel and floodplain habitats of a central Australian arid-zone river, Cooper Creek. *Biological Conservation* 103: .
- Souza-Filho, E. E., P. C. Rocha, E. Comunello & J. C. Stevaux, 2004. Effects of Porto Primavera dam on physical environment of the downstream floodplain. In Thomaz S. M., A. A. Agostinho & N. S. Hahn (eds), *The Upper Paraná River and its floodplain: Physical aspects, ecology and conservation*. Backhuys Publishers, Leiden, 55–74.



- Standford, J. A. & J. V. Ward, 1993. An ecosystem perspective of alluvial rivers connectivity and the hyporheic corridor. *Journal of the North American Benthological Society* 12: 48–60.
- Taniguchi, G. M., D. C. Bicudo & P. A. C. Senna, 2004. Abiotic variables in littoral-limnetic gradient of an oxbow lake of Mogi-Guaçu River floodplain, Southeastern, Brazil. *Brazilian Archives of Biology and Technology* 47: 961–971.
- Thomaz, S. M., M. C. Roberto, L. M. Bini, 1997. Caracterização limnológica dos ambientes aquáticos e influência dos níveis fluviométricos. In: Vazzoler A. E. A. M., Agostinho A. A., Hahn N. S., (eds), *Planície de inundação do alto rio Paraná: aspectos físicos, biológicos e socioeconômicos* Eduem Maringá 74–102.
- Thomaz, S. M., L. M. Bini, T. A. Pagioro, K. J. Murphy, A. M. Santos, D. C. Souza, 2004. Aquatic macrophytes: diversity, biomass and decomposition. In Thomaz S. M., A. A. Agostinho & N. S. Hahn (eds), *The Upper Paraná River and its floodplain: Physical aspects, ecology and conservation*. Backhuys Publishers, Leiden, 331–352.
- Tockner, K., D. Pennetzdorfer, N. Reiner, F. Schiemer & J. V. Ward, 1999. Hydrological connectivity, and the exchange of organic matter and nutrients in a dynamic river-floodplain system (Danube, Austria). *Freshwater Biology* 41: 521–535.
- Tockner, K. & J. V. Ward, 1999. Biodiversity along riparian corridors. *Large Rivers* 11: 293–310.
- Tockner, K., F. Malard & J. V. Ward, 2000a. An extension of the flood pulse concept. *Hydrological Processes* 14: 2861–2883.
- Tockner, K., Baumgartner, C., Schiemer, F. & J. V. Ward, 2000b. Biodiversity of a Danubian floodplain: structural, functional and compositional aspects. In Gopal B., W. J. Junk & J. A. Davis (eds), *Biodiversity in wetlands: assessment, function and conservation*. Backhuys Publishers, Leiden, 141–159.
- Tockner, K., F. Malard, U. Uehlinger & J. V. Ward, 2002. Nutrients and organic matter in a glacial river-floodplain system (Val Roseg, Switzerland). *Limnology and Oceanography* 47: 266–277.
- Train, S. & L. C. Rodrigues, 2004. Phytoplanktonic assemblages. In Thomaz S. M., A. A. Agostinho & N. S. Hahn (eds), *The Upper Paraná River and its floodplain: Physical aspects, ecology and conservation*. Backhuys Publishers, Leiden, 103–124.
- van Looy, K., O. Honnay, B. Bossuyt & M. Hermy, 2004. The effects of river embankment and forest fragmentation on the plant species richness and composition of floodplain forests in the Meuse Valley, Belgium. *Belgian Journal of Botany* 136: 97–108.
- Veríssimo, S., 1994. Variações na composição da ictiofauna em três lagoas sazonalmente isoladas, na planície de inundação do alto rio Paraná, ilha de Porto Rico, PT-Brasil. Master Thesis, PPG em Ecologia e Recursos Naturais, Universidade Federal de São Carlos.
- Ward, J. V. & K. Tockner, 2001. Biodiversity: towards a unifying theme for river ecology. *Freshwater Biology* 46: 807–819.

Zalocar de Domitrovic, Y., 2003. Effect of fluctuations of water level on phytoplankton development in three lakes of the Paraná River floodplain (Argentina). *Hydrobiologia* 510: 175–193.



**Sidinei Magela Thomaz** is an aquatic biologist who obtained his PhD degree at the Federal University of São Carlos, Brazil. He is professor and researcher in the Department of Biology and in the Nucleus of Research in Limnology, Ichthyology and Aquaculture (Nupelia) at the Maringá State University (Paraná, Brazil). He also serves as a research scientist for the National Council for Scientific and Technological Development (CNPq). He currently serves as Associate Editor for *Hydrobiologia* since 2003. His main research interests are ecology of aquatic macrophytes (especially patterns of diversity and underlying factors, and interactions of macrophytes with other aquatic organisms). He has studied the Upper Paraná River floodplain and sub-tropical reservoirs for 20 years.



**Luis Mauricio Bini** is an aquatic ecologist who obtained his PhD at the Maringá State University (Paraná, Brazil). He is currently a professor in the Department of General Biology at the Federal University of Goiás and a research scientist for the National Council for Scientific and Technological Development (CNPq). He currently serves as Associate Editor for *Hydrobiologia* since 2004. His research interest is in statistical methods applied to biodiversity analyses and limnology. Current projects involve the analysis of concordance among aquatic assemblages in reservoirs and floodplains.



**Reinaldo Luiz Bozelli** studied Biological Sciences at the Federal University of São Carlos, Brazil where he obtained his PhD in Ecology. In addition he conducted collaborative study at the Max Planck Institut für Limnologie (Germany). Currently he is a professor and research scientist in the Department of Ecology at the Federal University of Rio de Janeiro (UFRJ), and serves as a research scientist with the National Council for Scientific and Technological Development (CNPq). His research emphasis is in Limnology with special interest in the study of the zooplankton ecology relative to evaluation and restoration of anthropogenic degraded coastal lagoons and Amazonian aquatic environments.