

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/258922271>

# Desert dust and human health disorders

Article in *Environment international* · November 2013

DOI: 10.1016/j.envint.2013.10.011 · Source: PubMed

---

CITATIONS

80

---

READS

1,184

1 author:



[Andrew S. Goudie](#)

University of Oxford

378 PUBLICATIONS 7,827 CITATIONS

SEE PROFILE

All content following this page was uploaded by [Andrew S. Goudie](#) on 17 January 2014.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.



## Review

# Desert dust and human health disorders

Andrew S. Goudie

School of Geography and the Environment, University of Oxford, South Parks Road, Oxford OX1 3QY, United Kingdom



## ARTICLE INFO

### Article history:

Received 4 July 2013

Accepted 17 October 2013

Available online xxxx

### Keywords:

Dust storms  
Human health  
Particulates  
Respiratory  
Cardiovascular

## ABSTRACT

Dust storms may originate in many of the world's drylands and have an effect not only on human health in the drylands themselves but also in downwind environments, including some major urban centres, such as Phoenix, Kano, Athens, Madrid, Dubai, Jedda, Tehran, Jaipur, Beijing, Shanghai, Seoul, Taipei, Tokyo, Sydney, Brisbane and Melbourne. In some parts of the world dust storms occur frequently throughout the year. They can transport particulate material, pollutants, and potential allergens over thousands of km from source. The main sources include the Sahara, central and eastern Asia, the Middle East, and parts of the western USA. In some parts of the world, though not all, the frequency of dust storms is changing in response to land use and climatic changes, and in such locations the health implications may become more severe. Data on the PM<sub>10</sub> and P<sub>2.5</sub> loadings of dust events are discussed, as are various pollutants (heavy metals, pesticides, etc.) and biological components (spores, fungi, bacteria, etc.). Particulate loadings can far exceed healthy levels. Among the human health effects of dust storms are respiratory disorders (including asthma, tracheitis, pneumonia, allergic rhinitis and silicosis) cardiovascular disorders (including stroke), conjunctivitis, skin irritations, meningococcal meningitis, valley fever, diseases associated with toxic algal blooms and mortality and injuries related to transport accidents.

© 2013 Elsevier Ltd. All rights reserved.

## Contents

1. Introduction . . . . .	101
2. Desert dust . . . . .	102
2.1. Source areas . . . . .	102
2.2. Long distance transport . . . . .	103
2.3. The changing frequency of dust storms . . . . .	103
2.4. PM <sub>10</sub> and PM <sub>2.5</sub> levels in dust storms . . . . .	104
2.5. The durations of elevated particulate levels . . . . .	105
2.6. Biomaterials . . . . .	106
2.7. Miscellaneous pollutants . . . . .	106
3. Human disorders . . . . .	106
3.1. Respiratory disorders . . . . .	107
3.2. Cardiovascular disorders . . . . .	108
3.3. Miscellaneous other diseases and health issues . . . . .	109
3.3.1. Coccidiomycosis . . . . .	109
3.3.2. Meningococcal meningitis . . . . .	109
3.3.3. Conjunctivitis . . . . .	109
3.3.4. Dermatological disorders . . . . .	109
3.3.5. Algal blooms and red tides . . . . .	109
3.3.6. Deaths and injuries resulting from transport accidents . . . . .	109
4. Conclusions . . . . .	109
References . . . . .	109

## 1. Introduction

The purpose of this paper is firstly to review recent studies on the nature of desert dust storms (e.g. their main sources areas, frequencies,

E-mail address: [Andrew.goudie@stx.ox.ac.uk](mailto:Andrew.goudie@stx.ox.ac.uk).

changing frequencies, durations, and particulate contents), and secondly to consider some of the empirical evidence that is now accumulating to relate these characteristics of dust storms to various human health disorders. It does not seek to review in detail the underlying biological processes that lie behind such disorders.

## 2. Desert dust

Dust storms play an important role in the Earth system (Goudie and Middleton, 2006; Ravi et al., 2011; Shao et al., 2011). They are in most cases the result of turbulent winds, including fronts and convective *haboobs* (Miller et al., 2008), which raise large quantities of dust from desert surfaces and reduce visibility to less than 1 km. Dust reaches concentrations in excess of  $6000 \mu\text{g m}^{-3}$  in severe events (Song et al., 2007). It can be transported over thousands of kilometres and is deposited downwind by wet and dry processes, sometimes in appreciable quantities. Much of this load consists of silt. The silt comes finer with distance from source, and in the case of the Sahara, the coarsest dust (over  $70 \mu\text{m}$ ) occurs close to the Sahara itself, whereas dust that has travelled further tends to be finer silt, with a diameter of 5 to  $30 \mu\text{m}$  or less. Saharan dust collected in the eastern USA may be finer than  $1 \mu\text{m}$  (Perry et al., 1997). Aeolian dust is dominated by  $\text{SiO}_2$  and  $\text{Al}_2\text{O}_3$ , but other significant components are  $\text{Fe}_2\text{O}_3$ ,  $\text{CaO}$  and  $\text{MgO}$ . It may also have a large salt content, an organic content (Zaady et al., 2001), and, crucially from the health point of view can transport pathogens and anthropogenic pollutants.


### 2.1. Source areas

The health impact of dust storms will depend on where human populations are located with respect to the source areas of dust storms and the downwind direction of dust transport from them. Some areas are major generators of dust to the atmosphere. Other areas are, however, much less active. Analysis of Total Ozone Mapping Spectrometer (TOMS) data (Prospero et al., 2002; Varga, 2012; Washington et al., 2003) and other satellite borne sensors such as MODIS (Ginoux et al., 2012) has demonstrated the primacy of the Sahara, and the importance of large basins of internal drainage as dust sources (Bodélé, Taoudenni, Tarim, Seistan, Eyre, Etosha, Mkgadikgadi, Etosha, Uyuni, and the Great Salt Lake) (Engelstädter, 2001). Also, many sources are associated with alluvial deposits (Prospero et al., 2002) or piedmont alluvial fans (Tegen and Schepanski, 2009). Furthermore, TOMS data indicate that many of the world's major dust source areas, with high Aerosol Index (AI) values, are very arid (Table 1). Estimates of the relative strength of dust emissions for different parts of the world are variable but in general they demonstrate the importance, firstly of the Sahara (with over half of the global total), secondly of China and Central Asia (with about 20% of the global total), thirdly of Arabia and fourthly of Australia. Southern Africa and the Americas are relatively minor sources, together accounting for less than about 5% of the total (see, for example, Miller et al. (2004) and Tanaka and Chiba (2006) (Fig. 1)).

**Table 1**

Maximum mean aerosol index (AI) values for major global dust sources determined from TOMS (from Goudie and Middleton, 2006, Table 4.2).

Location	AI value	Average annual rainfall (mm)
Bodélé Depression of south central Sahara	>30	17
West Sahara in Mali and Mauritania	>24	5–100
Arabia (southern Oman/Saudi border)	>21	<100
Eastern Sahara (Libya)	>15	22
Southwest Asia (Makran coast)	>12	98
Taklamakan/Tarim Basin (China)	>11	<25
Etosha Pan (Namibia)	>11	435–530
Lake Eyre Basin (Australia)	>11	150–200
Mkgadikgadi Basin (Botswana)	>8	460
Salar de Uyuni (Bolivia)	>7	178
Great Basin (USA)	>5	400



**Fig. 1.** Estimates of the relative strength of dust emissions for different parts of the world (from data in Miller et al. (2004) (top) and Tanaka and Chiba (2006) (bottom)).

With regard to individual source regions, in the USA the greatest frequency of dust events occurs in the Great Plains and the Great Basin. These areas combine erodible materials with a dry climate and high wind energy (Gillette and Hanson, 1989). With regards to the arid south west, Bach et al. (1996) discuss the spatial and temporal variabilities of dust storms in the Mojave and Colorado Deserts and identify the Coachella Valley as being the dustiest region. Dust blown from the desiccated Owens Valley in California has been a cause of local health concerns. Dust has also been a major health issue in cities such as Phoenix in Arizona (Leathers, 1981), and pneumonia, called 'Haboob Lung Syndrome', has been reported after dust events in Lubbock, Texas (Panikkath et al., 2013).

The main source areas for Saharan dust (Goudie and Middleton, 2001) include: the Bodélé Depression (Bristow et al., 2009; Washington et al., 2006a, 2006b); an area that comprises southern Mauritania, northern Mali and central-southern Algeria (Knippertz and Todd, 2010); southern Morocco and western Algeria; the southern fringes of the Mediterranean Sea in Libya (O'Hara et al., 2006) and Egypt (Koren et al., 2003); and northern Sudan (Brooks et al., 2005). Much dust is moved by the north easterly trades over Nigeria and the Guinea zone to give the Harmattan haze (Breuning-Madsen and Awadzi, 2005) which may play a role in the timing and severity of meningococcal meningitis outbreaks. The Tokar delta of the Sudan, an arid, silty region across which high velocity winds are funnelled by a gap in the Red Sea Hills, is a frequent source of dust over the Red Sea (Hickey and Goudie, 2007) – dust which often moves into Arabia and may lead to air quality deterioration in cities like Jeddah.

Dust storms in the Middle East itself (Middleton, 1986a) are frequent on the alluvial plains of southern Iraq and Kuwait, and concerns have been expressed about their possible health implications (Thalib and Al-Taiar, 2012). In Tehran the great bulk of particulates appear to be derived from the deserts of Iraq and Syria (Givhechi et al., 2013). The Wahiba Sands (Pease et al., 1998) of Oman and the Oman–Saudi Arabia border are other large dust generation areas (Middleton and

Goudie, 2001). Also important is the eastern part of Saudi Arabia to the north of the Rub'Al Khali sand sea. Some dust in the United Arab Emirates may be derived from Iran and central Asia (de Villiers and van Heerden, 2011). Another major region of dust storms occurs in Iraq and Iran (Middleton, 1986b) where much dust is lofted off the Tigris–Euphrates alluvial plain (Hamidi et al., 2013) and at the convergence of the common borders between Iran, Pakistan and Afghanistan. At Zabol, in Iranian Seistan, there are on average 81 dust storm days per year, making it one of Earth's dustiest locations. This is a closed basin fed by the silt-laden Helmand River and through which high velocity winds are channelled by the high mountains that bound it (Hickey and Goudie, 2007). The frequency of dust storms in that region varies greatly from year to year depending on whether or not the Helmand lakes are wet or dry (Miri et al., 2010). The Makran coast is also an area of significant dust storm occurrence, and plumes are often identifiable on satellite images, showing dust being blown off dry river beds.

Middleton (1986c) has mapped dust storm activity in the arid zones of the Indian sub-continent, and has demonstrated that the greatest number of dust storms occur in the Thar Desert, but eastward moving dust clouds move into the heavily populated areas of the North Indian Plains (Gharai et al., 2013).

In the southern former Soviet Union the number of dust storms exceeds 40 per year, and some locations have more than 80, one of the highest occurrences in the world (Groll et al., 2013; Indoitu et al., 2012; Orlovsky et al., 2005). Dust is blown off the desiccated bed of the Aral Sea, with various proposed health implications.

According to Kes and Fedorovich (1976), the Tarim Basin has more dust storms than any other location on Earth, with 100 to 174 per year. Studies of dust loadings and fluxes suggest that there are two main source areas in China: the Taklamakan (Gao and Washington, 2009) and the Badan Jarain (Shao and Wang, 2003). The Gobi of Mongolia is also significant (Natsagdorj et al., 2003). The dust storms can cover immense areas and may transport dust particles to the more humid parts of China (Zhao et al., 2010), Korea, Japan, and beyond. As a consequence some major urban centres, including Beijing, Shanghai, Seoul, Tokyo and Taipei are subjected to what are called 'Asian Dust Storms'.

In southern Africa, satellite images show plumes blowing off the Namib and the Kalahari towards the South Atlantic (Eckardt et al., 2002), often having their sources in ephemeral river channels (Eckardt and Kuring, 2005). However, TOMS analyses indicate that the Etosha Pan in northern Namibia (Bryant, 2003) and the Mkgadikgadi Depression in northern Botswana (Resane et al., 2004) are even bigger sources.

In South America to the west of Buenos Aires in Argentina there are more than 8 dust storms per year (Middleton, 1986b). Dust is also generated from Patagonia (Gassó and Stein, 2007). The presence of extensive areas of closed depressions and of wind fluted topography, combined with the probable importance of salt weathering in the generation of fine material for deflation (Goudie and Wells, 1995), suggest that the dry areas of the Puna and Altiplano should be major source areas for dust storms (Gaiero et al., 2013). TOMS identifies one area where aerosol values are relatively high: this is the Salar de Uyuni, a large closed basin in Bolivia. It is probable that the deflation of fine sediments from its desiccated floor is one of the reasons for the existence of high aerosol values in this region.

Australia, though not as dusty as some regions of the world (McTainsh, 1989), is the largest dust source in the Southern Hemisphere (McTainsh et al., 2010). Within the Lake Eyre basin, one of the main source areas, using MODIS data, Bullard et al. (2008) found that 37% of dust plumes originated in areas of aeolian deposits, 30% from alluvial deposits and flood plains, and 29% from ephemeral lakes. Such lakes make some dust rich in NaCl (Shiga et al., 2011). Dust from this region leaves Australia in two main plumes: one that crosses the Tasman Sea to New Zealand (Marx et al., 2005) and another that heads westwards out into the Indian Ocean (Hesse and McTainsh, 1999). Australian dust may lead to increased respiratory problems in New Zealand (Cowie et al., 2010).

## 2.2. Long distance transport

One of the reasons why dust emissions have impacts on human health at great distances from their source is because of the great lengths over which dust plumes can travel (Zhu et al., 2007). Thus dust from the Lake Eyre basin and other parts of the interior of Australia may accumulate in New Zealand, but it also impacts upon air quality in major Australian cities, including Brisbane, Sydney, Newcastle and Melbourne (Leys et al., 2011). Saharan dust is carried thousands of kilometres to the Americas (Ben-Ami et al., 2010), Europe (Varga et al., 2013) the Near East (Thevenon et al., 2011) and the Arctic (Barkan and Alpert, 2010). Large amounts of Saharan (Harmattan) dust are blown southwards across the West African states into the Gulf of Guinea (Resch et al., 2007) and across the Atlantic to the Americas. The Sahara is also a major source of dust deposition into the Mediterranean Sea and neighbouring countries (Pey et al., 2013; Santese et al., 2007), where its health implications have been explored in such cities as Madrid, Barcelona, Rome and Athens. Dust from the Sahara and Thar reaches Mount Everest and Hong Kong in substantial quantities (Lee et al., 2010). The Ukraine has recently delivered large quantities of dust to central Europe (Birmili et al., 2007). Middle Eastern dust reaches Tehran (Givhechi et al., 2013) and India (Badarinath et al., 2010) and Chinese dust may reach North America and Europe via the Pacific (Uno et al., 2009). Indeed in May 2007 a dust cloud from the Taklamakan made a complete circuit of the globe in just 13 days.

## 2.3. The changing frequency of dust storms

The frequency with which dust storms occur is probably of importance in health terms, and there is an indication that in some areas the incidence of dust storms has increased. Neff et al. (2008), for instance, used analyses of lake cores in the San Juan Mountains of south-western Colorado, USA, to show that dust levels increased by 500% above the late Holocene average following the increased western settlement and livestock grazing during the nineteenth and the early twentieth century. A dust core from the Antarctic Peninsula (McConnell et al., 2007) showed a doubling in dust deposition in the twentieth century, and this is explained by increasing temperatures, decreasing relative humidity, and widespread desertification in the source region – Patagonia and northern Argentina. Finally, analysis of a 3200 year marine core off West Africa shows a marked increase in dust activity at the beginning of the nineteenth century, which was a time that saw the advent of commercial agricultural activity (including the clearing of ground for groundnut production) in the Sahel region (Mulitza et al., 2010).

It remains to be established, however, whether analysis of meteorological data, which has enabled the changing frequency of dust events to be established for the last six decades or so, indicates whether or not increasing dust storm frequencies are the norm. Some areas have indeed shown increasing trends (e.g. the Sahel zone of Africa (Goudie and Middleton, 1992), the eastern Mediterranean (Ganor et al., 2010), the Gobi of Mongolia (Dement'eva et al., 2013), the western USA (Brahney et al., 2013) and Korea (Chun et al., 2001)). However, others have shown declining trends in the late twentieth century (e.g. China (Wang et al., 2004), the Canary Islands (Laken et al., 2013), Turkmenistan (Orlovsky et al., 2005), Central Asia (Indoitu et al., 2012), Pakistan (Hussain et al., 2005) and parts of the USA High Plains (Stout and Lee, 2003), and Utah (Steenburgh et al., 2012). In Australia (McTainsh et al., 2005), the declining trend was followed by a spike of activity in the early years of the present century (e.g. Mitchell et al., 2010). Other areas (e.g. the Kalahari, southwestern Iran and Seistan) have shown marked fluctuations upwards and downwards in response to such factors as lake flooding and desiccation (Bryant et al., 2007; Ghasem et al., 2012; Miri et al., 2010; Rashki et al., 2013b) or climatic fluctuations such as sunspot cycles (Mohammed, 2010).



With respect to China, both natural and anthropogenic factors are implicated in the observed trends (Xu, 2006). There is certainly no clear upward trend in dust storm activity in China when the past 50 years are considered (Yao et al., 2010) though the early years of the present millennium saw some severe events (Kurosaki et al., 2011; Yang et al., 2008). Links have been established between dust emissions from the Tarim basin and the Arctic Oscillation (AO) Index, with dust activity being high during the negative phase of the AO (Gao and Washington, 2010). The general consensus is that natural climatic fluctuations have played a greater role in explaining the observed trends than have changes in human pressures on the land (Wang et al., 2007).

In the Kalahari of southern Africa fluctuations in the emission of dust from the Mkgadikgadi Pan over the period 1980–2000 suggest that dust loadings are intermittently influenced by the extent and frequency of lake inundation, sediment inflows, and surface wind speed variability. The variability of these is in turn influenced by the El Niño–Southern Oscillation and Indian Ocean sea surface temperature anomalies (Bryant et al., 2007).

In the West African Sahel, where drought has been persistent since the mid 1960s, analysis of wind, precipitation and visibility data by Ozer (2003) has shown that there have been remarkable changes in dust emissions since the late 1940s. He indicated that during the pre-drought conditions that existed from the late 1940s to the late 1960s yearly dust production was  $126 \times 10^6$  tons. It rose to  $317 \times 10^6$  tons during the 1970s and has been  $1275 \times 10^6$  tons since 1980, a ten-fold increase. Variability in Sahel dust emissions may be related not only to droughts, but also to an increase in the threshold wind velocity for dust entrainment. It also seems to be related to changes in the North Atlantic Oscillation (NAO) (Engelstaedter et al., 2006), North Atlantic sea surface temperatures (Wong et al., 2008) and the Atlantic Multidecadal Oscillation (Jilbert et al., 2010). The steadily increasing trend of dust storms from Africa in the Eastern Mediterranean over the last five decades in association with changing synoptic conditions has been analysed by Ganor et al. (2010). The slope of the increase is 0.27 days per year. However, African dust incidence in the Mediterranean basin decreased between 2006/2007 and 2011 in parallel with a negative period of the NAO (Pey et al., 2013).

Using a variety of data sources, Mahowald et al. (2010) have tried to estimate the global picture of changes in dust storm activity for the twentieth century. They suggest a doubling of atmospheric desert dust loadings took place over much of the globe.

An important issue from the human health perspective is whether or not dust activity will become even more frequent in coming decades. This will depend on three main factors: anthropogenic modification of desert surfaces (Mahowald and Luo, 2003); natural climatic variability (e.g. in the El Niño Southern Oscillation or the North Atlantic Oscillation); and changes in climate brought about by global warming. With regard to the first of these, increasing human pressures include disturbance of desert surfaces by vehicular traffic, increasing wildfires, removal of vegetation cover for wood supply, grazing and crop production, and desiccation of lakes and soil surfaces by inter-basin water transfers and ground water depletion (see, for example, Pelletier, 2006; Field et al., 2010; Ravi et al., 2011). A good review of the importance of different anthropogenic pressures in different parts of the world is provided by Ginoux et al. (2012). An example of severe change to a major inland sea is that taking place in the Aral Sea. Between 1960 and 1990, largely because of diversions of river flow, it lost more than 40% of its area and about 60% of its volume, and its level fell by more than 14 m. By 2002 its level had fallen another 6 m. By 2008, the area of the Aral Sea was just 15.7% of that in 1961 (Kravtsova and Tarasenko, 2010). This has lowered the artesian water table over a band 80–170 km in width, has exposed extensive expanses of former lake bed to desiccation and deflation. In the USA, water transfers from the Mono and Owens Lake basins have led to exposure of salty flats and the generation of severe dust storms derived from them (see, for example, Oho et al. (2011)).

Some of the highest coarse particulate ( $PM_{10}$ ) values ever recorded have been associated with these two sources (see Table 2).

The impact and occurrence of dust storms will depend a great deal on land management practices. Recent decreases in dust storm activity in North Dakota and the High Plains have resulted from conservation measures (Todhunter and Cihacek, 1999), while in Utah they have been caused by controls on grazing densities (Steenburgh et al., 2012).

Natural climatic variability will doubtless continue as it has in the past, but global warming has the potential to cause major changes in dust emissions. The IPCC (2007) suggests that many dryland areas will suffer from lower rainfall levels and increased amounts of moisture deficits because of higher rates of evapotranspiration. Munson et al. (2011) have argued that with increased drought brought about by reduced precipitation and higher temperatures, there will be a reduction in perennial vegetation cover in the Colorado Plateau and thus an increase in aeolian activity. Modelling studies of southern Africa by Thomas et al. (2005) have suggested that during the course of the twenty-first century most of the currently stable dune systems of the mega-Kalahari will become reactivated and mobile. This could cause substantial winnowing of fines from the weathered dune surfaces and an increase in dust emissions in southern Africa (Bhattachan et al., 2013). In the Bodélé depression of the Central Sahara, in spite of the possibility of higher rainfall amounts predicted by some models, higher wind velocities may increase its dust activity in coming decades (Washington et al., 2009).

In contrast, however, in northern China, there is some evidence that dust storm activity has decreased in recent warming decades, partially in response to changes in the atmospheric circulation and associated wind conditions (Jiang et al., 2009) and might decrease therefore still further in a warming world (Zhu et al., 2008). Indeed, recent research in China has demonstrated that over the last five decades or so there have been changes in wind activity that have been related to warming trends. As warming has occurred wind velocities have fallen (e.g. Wang et al., 2007; Xu, 2006). Various modelling studies have suggested that more generally in low latitudes extreme wind events will become less frequent with global warming, and this has been confirmed for the USA (Breslow and Sailor, 2002). ‘Atmospheric stilling’ has been a widespread feature of recent warming decades in Australia and elsewhere (Vautard et al., 2010).

#### 2.4. $PM_{10}$ and $PM_{2.5}$ levels in dust storms

A very important aspect of dust storms in relationship to health issues, particularly respiratory ones, is the amount of suspended material that they contain. This is normally expressed as  $PM_{10}$  values (aerodynamic diameter less than  $10 \mu m$ ) and  $PM_{2.5}$  (aerodynamic radius less than  $2.5 \mu m$ ). This review concentrates on these two parameters as they are the ones most usually monitored on an international basis. Table 2 presents data on  $PM_{10}$  levels that have been recorded in dust storms, though in many cases these data have not been related to specific health outcomes but are presented to give an indication of the concentrations that have been recorded during recent dust monitoring programmes. Some storms are associated with very high dust concentrations. While this may be the case in proximity to source, it is notable that events at quite large distances downwind can also be elevated, as is evident for some of the data for big cities in East Asia (Wang et al., 2013) and Australia (Leys et al., 2011). Even in southern Europe African dust incursions can produce high particulate levels (Pey et al., 2013), though they show a decline northwards. Table 3 presents data on the concentrations of finer particles ( $PM_{2.5}$ ) and also on the percentage they make of the total coarse and fine particulates. In general the finer particles form a bigger percentage of the total at lower particulate concentrations. In the USA the highest  $PM_{10}$  concentrations are associated with thunderstorm downbursts (Lei and Wang, 2013).

**Table 2**  
Maximum observed PM<sub>10</sub> levels in dust storms.

Source	Location	Date	Concentration ( $\mu\text{g m}^{-3}$ )
<i>(1) Southern Europe</i>			
Masson et al. (2010)	South of France	20/2/2004	150–443
Mallone et al. (2011)	Rome	–	182
Neophytou et al. (2013)	Cyprus	–	>250
Cabello et al. (2012)	S. Spain	October 2008	378
Kaskaoutis et al. (2008)	Crete	17/4/2005	2500
Kaskaoutis et al. (2008)	Athens	17/4/2005	250
Polymenakou et al. (2008)	Crete	February 2006	2800
Birmili et al. (2008)	Central Europe	March 2007	200–1400
<i>(2) Eastern Asia</i>			
Chung et al. (2003)	Korea	21–23/3/2002	3006
Chung et al. (2003)	Korea	7/4/2002	2942
Lee et al. (2013)	Korea	30–31/5/2008	1059
Lee et al. (2013)	Korea	25–26/12/2009	1041
Yuan et al. (2004)	Pescadores (Taiwan)	–	134
Xie et al. (2005)	Beijing	6/4/2000	798
Xie et al. (2005)	Beijing	25/4/2000	849
Fu et al. (2010)	Shanghai	April 2007	648
Wang et al. (2013)	Shanghai	20–21/3/2010	1700
Wang et al. (2013)	Shanghai	26–27/4/2010	236
Dement'eva et al. (2013)	Mongolia	–	1930
<i>(3) Australia</i>			
Merrifield et al. (2013)	Sydney	September 2009	11705
Leys et al. (2011)	Randwick	23/9/09	1734
Leys et al. (2011)	Newcastle	23/9/09	2426
Leys et al. (2011)	Bringelly	23/9/09	15,366
Barnett et al. (2012)	Brisbane	24/9/2009	894
Chan et al. (2005)	Sydney	23/10/2002	266
Chan et al. (2005)	Brisbane	23/10/2002	841
Chan et al. (2005)	Gladstone	23/10/2002	537
Chan et al. (2005)	Mackay	23/10/2002	899
<i>(4) West Africa</i>			
Alastuey et al. (2005)	Canary Islands	July 2002	312
Ozer (2006)	Mauritania	March 2004	2998
Marticorena et al. (2010)	M'Bour (Senegal)	8/3/2006	2500
Marticorena et al. (2010)	Cinzana (Mali)	8/3/2006	3500
Marticorena et al. (2010)	Banizoumbo (Niger)	8/3/2006	5000
<i>(5) North America</i>			
Bennett et al. (2006)	Vancouver (Canada)	April 1998	123
Claiborn et al. (2000)	Spokane (USA)	July 1994	1200
Lei and Wang (2013)	Western USA	–	3543
Cahill et al. (1996)	Owens Lake (USA)	–	40,620
Oho et al. (2011)	Mono Lake (USA)	20/11/2009	65,112
Hahnenberger and Nicoll (2012)	London, Utah (USA)	30/3/2010	424
<i>(6) Middle East</i>			
Rashki et al. (2012)	Iran	–	3094
Shahsavani et al. (2012)	Ahvaz (Iran)	2010	5338
Grishkan et al. (2012)	Haifa (Israel)	23/1/2004	700
Amanollahi et al., 2011	Sanandaj (Iran)	5/7/2009	5619
Krasnov et al. (2013)	Beer-Sheva (Israel)	February 2012	5197

## 2.5. The durations of elevated particulate levels

In general dust storms have a short duration at any one location. In the western USA, [Lei and Wang \(2013\)](#) (Table 2) suggest that most storms last between 2 and 21 h. In Turkmenistan the frequency of dust storms with a duration of 12 h or more is only about 3%, though occasional examples lasting 3 days have been observed there ([Orlovsky and Orlovsky, 2001](#)). In Central Asia more generally, 40–60% of the dust events last on average 1.5–5.4 h ([Indoitu et al., 2012](#)). In the Gobi of Mongolia the average dust storm lasts from 1.6 to 6.0 h ([Natsagdorj et al., 2003](#)), while in China's Taklamakan the most serious dust storm conditions persist for 2 to 4 h ([Yoshino, 1992](#)). More generally in China, [Wang et al. \(2005\)](#) suggest that most dust storms last less than 2 h.

PM<sub>10</sub> values may, however, remain high for extended periods in dusty areas. In the Sistan region of Iran, for example, monthly mean values are high throughout the year, and were found to average

273  $\mu\text{g m}^{-3}$  in the winter, 289 in the spring, 683 in the summer and 484 in the autumn ([Rashki et al., 2012](#)). In Zabol over 30% of days in the year exceeded the US EPA hazardous value of 425  $\mu\text{g m}^{-3}$ . In the Gobi of Mongolia the average annual PM<sub>10</sub> concentration at Zamyn-üüd is 81  $\mu\text{g m}^{-3}$  and the maximum mean monthly concentration in May reaches 139  $\mu\text{g m}^{-3}$  ([Dement'eva et al., 2013](#)). In Ahvaz, Iran, the average PM<sub>10</sub> values for the months of April to September range from 186 to 489  $\mu\text{g m}^{-3}$  ([Shahsavani et al., 2012](#)), while in Zahedan they range from 101 to 182  $\mu\text{g m}^{-3}$  ([Rashki et al., 2013a, 2013b](#)). In 1984, a particularly dry year, the average annual PM<sub>10</sub> concentration in Gouré, Niger, was 245  $\mu\text{g m}^{-3}$  ([de Longueville et al., 2013](#)). In Canyonlands National Park, Utah, [Neff et al. \(2013\)](#) found that in several cases, PM > 10 concentrations exceeded 300  $\mu\text{g m}^{-3}$  with the largest dust-deposition period, in 2010, having a two-week average concentration in excess of 700  $\mu\text{g m}^{-3}$ .

[Engelbrecht et al. \(2009\)](#) presented annual average particulate concentrations for 15 locations in the Middle East (Table 4). This

**Table 3**  
Maximum observed PM<sub>2.5</sub> levels ( $\mu\text{g m}^{-3}$ ) in dust storms.

Source	Location	Date	PM <sub>2.5</sub>	% of PM <sub>10</sub>
<i>(1) Southern Europe</i>				
Mallone et al. (2011)	Rome	–	86	47.3
Jiménez et al. (2010)	Madrid	–	71	47.3
Vanderstraten et al. (2008)	Belgium	March, 2007	43	42.8
<i>(2) Eastern Asia</i>				
Dement'eva et al. (2013)	Mongolia	–	700	36.3
Yuan et al. (2004)	Pescadores (Taiwan)	–	63	47.0
Chung et al. (2003)	Korea	23–24/3/2000	331	11.0
Chung et al. (2003)	Korea	7/4/2002	294	10.0
Jugder et al. (2011)	Mongolia	–	500	60.9
Wang et al. (2013)	Shanghai	20–21/3/2010	469	27.6
Wang et al. (2013)	Shanghai	26–27/4/2010	94	28.8
<i>(3) Others</i>				
Shahsavani et al. (2012)	Ahvaz, Iran	2010	911	17.1
Merrifield et al. (2013)	Sydney	9/2009	1638	14.0
Barnett et al. (2012)	Brisbane	24/9/2009	138	15.4
Chan et al. (2005)	Brisbane	23/10/2002	42	26.1
Claiborn et al. (2000)	Spokane (USA)	07/1994	150	12.5
Grishkan et al. (2012)	Haifa (Israel)	23/1/2004	280	40.0
Hahnenberger and Nicoll (2012)	Lindon, Utah (USA)	30/3/2010	55.7	13.1

demonstrates very powerfully that over large parts of the region, citizens (and military personnel) are exposed to high levels of particulates, with PM<sub>10</sub> values ranging from 72 to 303  $\mu\text{g m}^{-3}$  and PM<sub>2.5</sub> values ranging from 35 to 111  $\mu\text{g m}^{-3}$ . The WHO ([http://www.who.int/phe/health\\_topics/outdoorair/databases/en/](http://www.who.int/phe/health_topics/outdoorair/databases/en/)) (accessed 2nd July, 2013) has provided data on annual average PM<sub>10</sub> values for cities from all over the world, and while some of the cities with high values may be heavily polluted because of industrial and transport emissions, combined with atmospheric inversions that retain particulates in the lower atmosphere, it is noticeable that most of them occur in areas that are also subject to major dust storm incursions (Table 5a). Likewise, World Bank data (Table 5b) (<http://data.worldbank.org/indicator/EN.ATM.PM10.MC.M3>) (accessed 2nd July, 2013) indicates that most countries with high average PM<sub>10</sub> values are in the dusty countries of northern Africa and the Middle East.

## 2.6. Biomaterials

Dust storms pick up and transport biological materials which include bacteria, pollen spores, fungi and viruses (Table 6). These are capable of surviving long-range transport on a global scale. All of these have potential implications for disease incidence, including meningococcal meningitis and coccidiomycosis. This is an area that has been expertly reviewed by Griffin (2007) who notes that although we know that

**Table 4**  
Annual particulate concentrations in Middle East locations (derived from data in Engelbrecht et al., 2009).

Location	PM <sub>10</sub> $\mu\text{g m}^{-3}$	PM <sub>2.5</sub> $\mu\text{g m}^{-3}$	Ratio
Djibouti	72	35	0.49
Bagram (Afghanistan)	108	38	0.35
Khowst (Afghanistan)	127	75	0.59
Qatar	165	67	0.41
UAE	140	52	0.37
Balad (Iraq)	184	56	0.30
Baghdad (Iraq)	250	103	0.41
Tallil (Iraq)	303	65	0.21
Tikrit (Iraq)	298	111	0.37
Taji (Iraq)	213	81	0.38
Al Asda (Iraq)	95	37	0.39
N. Kuwait	211	67	0.32
C. Kuwait	298	87	0.29
Coastal Kuwait	175	60	0.34
Southern Kuwait	199	62	0.31
Mean	189	66	0.35

dust contains pathogens, in most cases the risk associated with them is still largely unknown. Dadvand et al. (2011), working in Barcelona, tested the possibility that infections associated with materials transported from North Africa might have an adverse effect on pregnancies, but found no evidence for a link. Desert dust may contain dangerous cyanotoxins but further research is required to assess their significance (Metcalfe et al., 2012). Nonetheless, in Japan, Watanabe et al. (2011a, 2011b) found that asthma in adults was augmented on days when Asian Dust carried pollen in comparison with dusty days without pollen.

## 2.7. Miscellaneous pollutants

Dust storms can pick and transport anthropogenic material as a result of particulate/pollutant aerosolization and the absorption of such materials as pesticides, herbicides, heavy metals, dioxins, and radioactive isotopes (Table 7). Dust storms that have moved across heavily industrialized and polluted areas, such as northern China may therefore pose a greater risk than those storms that have tracked out of the Sahara to the Atlantic Ocean without passing over heavily developed areas (Almeida-Silva et al., 2013). In addition, dust storms may remove toxic or irritating materials from desiccated lake beds, including various types of salts and arsenic (Morman and Plumlee, 2013). There has been especial concern about the deflation of pesticides from the exposed bed of the shrinking Aral Sea (Ataniyazova et al., 2001; O'Hara et al., 2000). Because they are normally sparsely populated, deserts such as those of China (Lop Nor), Australia (Maralinga), Kazakhstan (Semipalatinsk) USA (Nevada), Algeria (Reggane) and north west India (Pokhran) were the sites of nuclear tests. Radionuclides may be blown from them in dust storms. Dust storms also used to occur in the Chernobyl area and transported cesium towards the Baltic states (Ogorodnikov, 2011). Tailings from uranium mines are another potential source of radionuclides in dust storms (Csavina et al., 2012), with major uranium mines occurring in a number of arid regions, including South Australia, Niger, Utah, Namibia and Kazakhstan.

## 3. Human disorders

There is not space in this paper to consider in detail the specific biological processes by which dust particles of different sizes have an impact on human health. However, as Sandstrom and Forsberg (2008) have pointed out, particle size is a main determinant of where in the respiratory tract the particle will come to rest when inhaled. Because of

**Table 5**

Particulate concentrations in world cities.

a. The world's cities with average annual PM<sub>10</sub> values of 150 and above ( $\mu\text{g m}^{-3}$ ), processed from WHO ([http://www.who.int/phe/health\\_topics/outdoorair/databases/en/](http://www.who.int/phe/health_topics/outdoorair/databases/en/)) (accessed 2nd July, 2013) by author.

Average value	City	Country
372	Ahwaz	Iran
279	Ulanbaator	Mongolia
254	Sanandaj	Iran
251	Ludhiana	India
251	Quetta	Pakistan
229	Kermanshah	Iran
219	Peshawar	Pakistan
216	Gaborone	Botswana
215	Yasouj	Iran
209	Kanpur	India
200	Lahore	Pakistan
198	Delhi	India
193	Karachi	Pakistan
189	Islamabad	Pakistan
185	Rawalpindi	Pakistan
183	Uromiyeh	Iran
176	Qom	Iran
174	Indore	India
168	Khoramabad	Iran
165	Agra	India
158	Al Ain	United Arab Emirates
157	Riyadh	Saudi Arabia
151	Al Hafouf	Saudi Arabia
150	Lanzhou	China

b) World Bank data on countries with average annual PM<sub>10</sub> values greater than  $60 \mu\text{g m}^{-3}$  in 2010 and relating to cities with >100,000 inhabitants (<http://data.worldbank.org/indicator/EN.ATM.PM10.MC.M3>) (accessed 2nd July, 2013).

Average value	Country
137	Sudan
115	Bangladesh
112	Uruguay
111	Mali
97	Trinidad and Tobago
96	Mongolia
96	Niger
96	Saudi Arabia
95	Oman
91	Kuwait
91	Pakistan
89	United Arab Emirates
88	Iraq
83	Chad
78	Egypt
77	Senegal
69	Algeria
68	Mauritania
65	Burkina Faso
65	Libya
65	Sri Lanka
64	Botswana
64	Paraguay
61	Eritrea

their small size, particles on the order of  $\sim 10 \mu\text{m}$  or less (PM<sub>10</sub>) can penetrate the deepest part of the lungs such as the bronchioles or alveoli. Larger particles are generally filtered in the nose and throat via cilia and mucus, but particles smaller than about  $10 \mu\text{m}$ , can settle in the bronchi and lungs and cause health problems. Particles which are smaller than  $2.5 \mu\text{m}$ , tend to penetrate into the gas exchange regions of the lung, and ultra-fine particles ( $<100 \text{ nm}$ ) may pass through the lungs to affect other organs, with possible cardiovascular consequences (Brook et al., 2010; Martinelli et al., 2013).

According to WHO (<http://www.who.int/mediacentre/factsheets/fs313/en/>) (accessed 29th June, 2013) the acceptable annual mean value of PM<sub>10</sub> is  $20 \mu\text{g m}^{-3}$  and the acceptable 24 h mean value is  $50 \mu\text{g m}^{-3}$  (de Longueville et al., 2013). The EU (<http://ec.europa.eu/>

**Table 6**

Examples of biological materials carried in dust storms.

Source	Location	Material
Leski et al. (2011)	Kuwait & Iraq	<i>Mycobacterium</i> , <i>Brucella</i> , <i>Coxiella Burnetii</i> , <i>Clostridium perfringens</i> , <i>Bacillus</i>
Thomson et al. (2009)	West Africa	<i>Neisseria meningitidis</i>
Chen et al. (2010)	Taiwan	Influenza virus
Watanabe et al. (2011a)	Japan	Pollen spores
Polymenakou et al. (2008)	Crete	Bacteria
Jeon et al. (2011)	Korea	Bacteria
Grishkan et al. (2012)	Israel	Fungal communities
Soleimani et al. (2013)	Iran	Fungi: <i>Cladosporium</i> , <i>Alternaria</i> , <i>Aspergillus</i> , <i>Penicillium</i> and <i>Rhizopus</i>
Perfumo and Marchant (2010)	Turkey	Thermophilic bacteria ( <i>Geobacillus</i> )
Schlesinger et al. (2006)	Israel	Bacteria and fungi
Najafi et al. (2013)	Iran	Bacteria and fungi

[environment/air/quality/standards.htm](http://environment.air/quality/standards.htm)) (accessed 29th June 2013) has 1 year mean PM<sub>2.5</sub> values of  $25 \mu\text{g m}^{-3}$ , 1 year PM<sub>10</sub> mean values of 40 and 24 h mean PM<sub>10</sub> values of 50. Such values are regularly exceeded in areas downwind of dust sources (Ozer, 2006). On Dec. 14, 2012 the U.S. Environmental Protection Agency (EPA) strengthened the nation's air quality standards for fine particle pollution to improve public health protection by revising the primary annual PM<sub>2.5</sub> standard to 12 micrograms per cubic meter ( $\mu\text{g m}^{-3}$ ). The EPA also has a 24-h PM<sub>2.5</sub> standard of  $35 \mu\text{g m}^{-3}$  and a 24-h standard of  $150 \mu\text{g m}^{-3}$  for PM<sub>10</sub> (<http://www.epa.gov/airquality/particlepollution/2012/decfsstandards.pdf>) (accessed 29th June, 2013). The EPA regards 24 h PM<sub>2.5</sub> levels between 35.5 and  $55.4 \mu\text{g m}^{-3}$  as being unhealthy for sensitive groups, 55.5–150.4 as being unhealthy, 150.5–250.4 as being very unhealthy, and 250.4–500 as hazardous.

### 3.1. Respiratory disorders

The pathogenic effect of dust inhalation on respiratory tissues can be attributed to the direct physical action of dust particles on the epithelium of the human airways and may be exacerbated by the toxic effects of both trace elements (including arsenic, etc.) and of biologically active compounds (bacteria, fungi, pollen, and viruses) (Leski et al., 2011). It is likely that the population most susceptible to suffering from the short-term effects of suspended particulates are (Jiménez et al., 2010) (a) the elderly, due to their lower immunological capacity and the deterioration in their general health due to the ageing process, (b) subjects affected by chronic cardiopulmonary disorders, and (c) the very young, whose lungs and airways have yet to develop fully (Yu et al., 2012). In Japan, Kanatani et al. (2010) found that Asian dust exacerbated asthma in children and caused increased hospitalisation, while Chien et al. (2012) working in Taipei (Taiwan) found that there was a significant relationship between dust storms and clinic visits for respiratory disease in children.

Questions have been asked in the Americas as to whether or not dust from Asia, notwithstanding the distance over which it has travelled and the relatively low particulate content it displays, has an effect on asthma (Prospero et al., 2008). Gyan et al. (2005) found that there was an association between increased asthma admissions in Trinidad and African dust clouds. On the other hand, Bennett et al. (2006), working in Vancouver (Canada) found that a Gobi desert dust event was not associated with a significant risk to public health.

There are a range of studies in East Asia that have sought to relate Asian Dust events to respiratory problems such as asthma, pneumonia and tracheitis (Yang, 2013). Tao et al. (2012), working in Lanzhou, China, found that dust storms led to increased respiratory hospitalizations, particularly for those aged over 65. Tam et al. (2012), working in Hong Kong, found significant increases in emergency hospital admissions due to respiratory problems, with a 1.05 increase per  $10 \mu\text{g m}^{-3}$ .



**Table 7**

The nature of pollutant materials carried in dust storms.

Source	Location	Nature of substance(s)
Masson et al. (2010)	France	Radioactive cesium ( $^{137}\text{Cs}$ ) from Sahara
Akata et al. (2007) and Fukuyama and Fujiwara (2008)	Japan	Radioactive cesium ( $^{137}\text{Cs}$ ) from China and Mongolia
Kikawada et al. (2012)	Japan	Enriched uranium from Central Asia
Gabrieli et al. (2013)	European Alps	Plutonium from Saharan atomic tests
Lee et al. (2013) and Park and Dam (2010)	Korea	Heavy metals from China
Onishi et al. (2011, 2012)	Japan	Heavy metals from China
Liu et al. (2011), and Li et al., 2013	China	Heavy metals
Li et al. (2012)	China	Sulphates and nitrates
Kwon et al. (2013)	Korea	Polycyclic aromatic hydrocarbons
Hou et al. (2006)	China	Polycyclic aromatic hydrocarbons and fatty acids
O'Hara et al. (2000)	Aral Sea	Phosalone from Aral Sea
Ataniyazova et al. (2001)	Aral Sea	Heavy metals, organochlorine pesticides, Dioxins from Aral Sea
Gill et al. (2002)	Owens Lake, USA	Arsenic
Chi et al. (2008) and Wu et al. (2010)	Taiwan	Dioxins and PCBs from China

Chien et al. (2012), working in Taipei found that compared with weeks before Asian dust events respiratory clinic visits during the weeks after an event increased by 2.54% for children younger than 6, and by 5.03% for children aged 7–14 years. Studies in Seoul, Korea, by Lee and Lee (2013) showed that Asian dust events led to a 22% increase in the rate of asthma treatments with a 6 day lag. Pneumonia is another respiratory complaint that appears to be influenced by Asian dust events. Cheng et al. (2008) and Kang et al. (2012), working in Taipei (Taiwan), found significantly elevated pneumonia admissions. Allergic rhinitis incidence in Taiwan has also been related to Asian dust (Chang et al., 2006).

In Australia, various studies have been undertaken to see the effects of dust storms moving eastwards from the interior. Rutherford et al. (1999), who studied Brisbane, found that dust events were significantly associated with asthma severity. A 15% increase in non-accidental mortality at a lag of 3 days from a dust event was found for Sydney (Johnston et al., 2011), where Merrifield et al. (2013) also found large increases in asthma emergency department visits. In Brisbane, there was a 39% increase in emergency admissions associated with a dust storm in 2009.

The effects of African dust outbreaks on asthma and other respiratory and cardiac afflictions in southern Europe has been the subject of considerable attention (Karanasiou et al., 2012). In Athens, Samoli et al. (2011) found that a  $10 \mu\text{g m}^{-3}$   $\text{PM}_{10}$  level was associated with a 0.71% increase in all deaths, with those over 75 years or older being especially affected. Likewise, in Madrid Jiménez et al. (2010) found a significant statistical association between  $\text{PM}_{10}$  levels and mortality for the elderly on Saharan Dust days. Also in Madrid, Reyes et al. (2013) found that periods with Saharan dust intrusions saw a significant increase in respiratory-caused admissions which could be related to  $\text{PM}_{10}$  levels. In Barcelona, Perez et al. (2008) found that during Saharan Dust days a daily increase of  $10 \mu\text{g m}^{-3}$  of  $\text{PM}_{10-2.5}$  increased daily mortality by 8.4%. In Rome, Mallone et al. (2011) found that such events were associated with an increase in cardiac and respiratory mortality. Similarly in Emilia-Romagna Sajani et al. (2011) found an increased risk of respiratory mortality for people aged 75 or over.

Military personnel stationed in Afghanistan and Iraq have also suffered from dust related pulmonary problems (Furlow, 2013). Other locations where a link between dust events and respiratory problems has been established include Seistan, Iran (Miri et al., 2007). There are also high levels of asthma in the Arabian Gulf region, with, for example, 24% of the population of Saudi Arabia being asthmatic, and dust storms are one possible factor (along with lifestyle, smoking and indoor dust levels) for this (Al-Ghazawy, 2013). On the other hand, a significant association between dust levels and respiratory mortality has not always been found, as demonstrated by the work of Schwartz et al. (1999) in the Spokane area of Washington, USA, by Wiggs et al. (2003) and Bennion et al. (2007) in the Aral Sea region, and by Al-Ta'iar and Thalib (2013) in Kuwait.

Much dust storm material is silt-sized quartz, and it has been argued that this can, if inhaled over a sustained period, cause non-occupational silicosis (also called desert lung syndrome) to develop in human lungs. This appears to be an issue, especially among the elderly, in some areas of High Asia (Derbyshire, 2007) including north west China and Ladakh (India) (Norboo et al., 1991; Saiyed et al., 1991). High incidences of silicosis and pneumoconiosis have been reported in Bedouins in the Negev (Bar-Ziv and Goldberg, 1974). Silicosis may be an important factor in the higher prevalence of tuberculosis in some deserts, including the Thar of India (Mathur and Choudhary, 1997).

Recently, a modelling study has suggested that in the 'dust belt' of North Africa and the Middle East, a substantial proportion of lung cancers may be caused by exposure to desert dust (Giannadaki et al., 2013), especially in countries like Mauritania and Mali.

### 3.2. Cardiovascular disorders

There is an increasing body of epidemiological evidence that high particulate levels in the air cause cardiovascular disease (Brook et al., 2010), including myocardial infarction, stroke, heart failure, arrhythmias, and venous thromboembolism (Martinelli et al., 2013). Schulz et al. (2005) have suggested that the possible biological mechanisms associated with exposure to particulates include '(i) pulmonary and/or systemic inflammatory responses triggering endothelial dysfunction and a pro-coagulatory state with thrombus formation and promotion of atherosclerotic lesions, (ii) dysfunction of the autonomic nervous system in response to direct reflexes from receptors in the lungs and/or to local or systemic inflammatory stimuli, and (iii) cardiac malfunction due to ischemic responses in the myocardium and/or altered ion-channel functions in myocardial cells'.

There is increasing evidence of cardiovascular problems associated with dust events. Yang et al. (2005) and Kang et al. (2013), working in Taiwan, found that Asian Dust Storms were associated with an acute increase in stroke hospitalisation. However, also working in Taiwan, Yang et al. (2009) were unable to find a statistically significant association with congestive heart failure. On the other hand, Meng and Lu (2007), working in Minqin, China, found an association between dust events and hypertension in males. In Nagasaki, Japan, Ueda et al. (2012) reported that heavy Asian dust events caused emergency ambulance dispatches for cardiovascular diseases to rise by 20.8%. Likewise, in Cyprus, Middleton et al. (2008) found that cardiovascular admissions increased after dust episodes, and Neophytou et al. (2013) found there was a 2.43% increase in daily cardiovascular associated with each  $10 \mu\text{g m}^{-3}$  increase in  $\text{PM}_{10}$  concentrations on African Dust days. In Spain, Perez et al. (2012) found an increased cardiovascular mortality associated with  $\text{PM}_{10-2.5}$  levels, as did Alessandrini et al. (2013) in Rome, Italy.

### 3.3. Miscellaneous other diseases and health issues

#### 3.3.1. Coccidiomycosis

Coccidiomycosis is an infection caused by the inhalation of airborne spores of *Coccidioides immitis*, a soil dwelling fungus found in the south west USA and parts of Mexico and central and South America. It affects the lungs, can cause extreme disfigurement, and can be fatal. In the USA it is often known as Valley Fever (Zender and Talamantes, 2006) and around 150,000 cases are reported each year (Anderson, 2013). The association between dust storms and the disease in California and Arizona has been known for a long time (see, for example, Pappagianis and Einstein, 1978; Leathers, 1981), and more recently Comrie (2005) has demonstrated that there is a strong bimodal seasonality of the disease in Pima County, Arizona, with peaks in June–July and August–November, which are the drier and dustier months.

#### 3.3.2. Meningococcal meningitis

Meningococcal meningitis is a serious health problem in West Africa (Thomson et al., 2009) and the incidence of epidemics appears to be related to the timing and severity of Saharan dust intrusions (the Harmattan). Fears have also been expressed about possible transport of the bacteria *Neisseria meningitidis*, to southern Europe (Tobias et al., 2011a, 2011b). During the 1980s there were between 25,000 and 200,000 cases in the Sahel per year (Sultan et al., 2005), and levels continued to be high between 2004 and 2009 (Martigny and Chiapello, 2013), especially in Nigeria, Burkina Faso, the Congo Democratic Republic and Niger. One theory is that dust, combined with the extremely dry air, damages the pharyngeal mucosa, thereby easing bacterial invasion (Agier et al., 2013). There is a clear temporal correlation between times of great atmospheric dust contents and cases of meningitis (i.e. in January to April) (Deroubaix et al., 2013; Martigny and Chiapello, 2013).

#### 3.3.3. Conjunctivitis

Prolonged exposure to dusty air can lead to conjunctivitis. Dust blown by the Irifi wind of the Western (formerly Spanish) Sahara is responsible for the conjunctivitis that is common among the nomads of that country (Morales, 1946). In Taiwan, Yang (2006) found some evidence for an association between Asian Dust Events and clinic visits for conjunctivitis, but the association was not statistically significant. However, in a subsequent study Lien et al. (2013) did find elevated conjunctivitis clinic visits during dust storm periods. An epidemic of conjunctivitis in Lagos, Nigeria has also been related to Saharan dust outbreaks (McMoli et al., 1984).

#### 3.3.4. Dermatological disorders

The presence of heavy metals such as nickel in dust may cause skin irritation (Onishi et al., 2011; Otani et al., 2012). In addition, it has been suggested that Asian dust storm particles can 'exert toxicological effects on human skin through the activation of the cellular detoxification system, the production of pro-inflammatory and immunomodulatory cytokines and changes in the expression of proteins essential in normal epidermal differentiation' (Choi et al., 2011, p. 92).

#### 3.3.5. Algal blooms and red tides

There is increasing evidence that the nutrients carried in dust storms, including iron (Mahowald et al., 2009), can indirectly impact human health by stimulating toxic algal blooms in coastal environments (Griffin and Kellogg, 2004). This is true for the Arabian Gulf (Al-Shehhi et al., 2012), the Arabian Sea (Kayetha et al., 2007), the Yellow Sea off China (Shi et al., in press; Tan et al., 2011), the south east Mediterranean (Herut et al., 1999) but also for Florida, where red tides associated with blooms of *Karenia brevis* and *Trichodesmium* appear to result from inputs of Saharan dust (Lenes et al., 2008; Walsh and Steidinger, 2001; Walsh et al., 2006) and have a range of effects including neurotoxic

shellfish poisoning (NSP) and respiratory complaints (Fleming et al., 2011).

#### 3.3.6. Deaths and injuries resulting from transport accidents

Dust storms can cause death through their role in causing accidents on the road and in the air (Baddock et al., 2013). In November 1991, a series of collisions involving 164 vehicles and causing 17 deaths occurred on Interstate 5 in the San Joaquin Valley in California, USA (Pauley et al., 1996), while in Oregon a dust storm in September 1999 set off a chain reaction of 50 car crashes that killed eight people and injured more than twenty. The loss of visibility may be very sudden when caused by the arrival of a dust wall associated with a dry thunderstorm. Such Haboob dust walls were responsible for 32 multiple accidents between 1968 and 1975 on Interstate 10 in Arizona (Brazel and Hsu, 1981). Between 2001 and 2005 there were 44 deaths in 2323 traffic accidents in New Mexico and 15 deaths in 614 accidents in Arizona ([http://www.azcentral.com/arizonarepublic/news/articles/20120726arizona-researcher-explores-dangers-living-dust.html?nclink\\_check=1](http://www.azcentral.com/arizonarepublic/news/articles/20120726arizona-researcher-explores-dangers-living-dust.html?nclink_check=1)) (accessed 6th September, 2013). In the USA in 2012/3 a survey of newspaper headlines available online indicates that dust storm related multiple traffic incidents occurred in Idaho, California, Texas, Arizona, Nevada and Oklahoma.

Some fatal commercial air crashes have also been attributed to visibility reduction or to the adverse mechanical effects of dust storms (Goudie and Middleton, 2006, p. 51). On 7th May 2002, for example, an EgyptAir aircraft crashed near Tunis killing 18 of 60 people on board (<http://airsafe.com/events/airlines/egyptair.htm>) (accessed 6th September, 2013). On 30th January 2000, a Kenya Airways Airbus crashed in the Ivory Coast with the loss of 179 lives ([http://www.nrlmry.navy.mil/aerosol\\_web/Case\\_studies/20000130\\_ivorycoast/](http://www.nrlmry.navy.mil/aerosol_web/Case_studies/20000130_ivorycoast/)) (accessed 6th September, 2013). On 19th August 2012 31 people died in an air crash at Talodi in Sudan (<http://www.ainonline.com/aviation-news/aviation-international-news/2012-10-05/accidents-october-2012>) (accessed 6th September, 2013) while on 25th May, 2011, 10 people died in a crash in Faridabad, India. (<http://dgca.nic.in/accident/reports/VT-ACF.pdf>) (accessed 6th September, 2013).

## 4. Conclusions

Dust storms are events of considerable extent and of frequent occurrence in deserts and on their margins, and transport large amounts of particulates, pollutants and biological materials for long distances downwind. In some parts of the world, though not all, they appear to be occurring with greater frequency and some forecasts suggest that this will continue in response to increasing land use pressures and lower soil moisture contents resulting from climate change. An increasing corpus of studies, particularly in east Asia, show associations between dust events and a range of human health issues, including respiratory problems, cardiovascular complaints, meningococcal meningitis, conjunctivitis, skin irritation, and deaths and injuries associated with transport accidents. There are, however, some parts of the world, including the Gulf States of the Middle East, and countries in northern and western Africa (De Longueville et al., 2010), where relatively little epidemiological research has been published on the relationship between dust events and health. Nevertheless, the increasing availability of information about dust storm frequencies, trajectories, and particulate contents as a result of the availability of large amounts of data from various satellite-borne sensors, means that there is now a greater potential for their health effects to be explored in greater detail by collaboration between medical and environmental scientists.

## References

- Agier L, Deroubaix A, Martigny N, Yaka P, Djibo A, Broutin H. Seasonality of meningitis in Africa and climate forcing: aerosols stand out. *J R Soc Interface* 2013;10. <http://dx.doi.org/10.1098/rsif.2012.0814>.

- Akita N, Hasegawa H, Kawabata H, Chikuchi Y, Sato T, Ohtsuka Y, et al. Deposition of  $^{137}\text{Cs}$  in Rokkasho, Japan and its relation to Asian dust. *J Environ Radioact* 2007;95:1–9.
- Alastuey A, Querol X, Castillo S, Escudero M, Avila A, Cuevas E, et al. Characterisation of TSP and  $\text{PM}_{2.5}$  at Izaña and Sta Cruz de Tenerife (Canary Islands, Spain) during a Saharan Dust Episode (July 2002). *Atmos Environ* 2005;39:4715–28.
- Alessandrini ER, Stafoggia M, Faustini A, Gobbi GP, Forastiere F. Saharan dust and the association between particulate matter and daily hospitalizations in Rome, Italy. *Occup Environ Med* 2013. <http://dx.doi.org/10.1136/oemed-2012-101182>.
- Al-Ghazawy O. The rising danger of asthma. *Nat Middle East* 2013. <http://dx.doi.org/10.1038/middleeast.2013.79>.
- Almeida-Silva M, Almeida SM, Freitas MC, Pio CA, Nunes T, Cardoso J. Impact of Sahara dust transport on Cape Verde atmospheric element particles. *J Toxicol Environ Health A* 2013;76:240–51.
- Al-Shehhi MR, Gherboudj I, Ghedira H. Temporal-spatial analysis of chlorophyll concentration associated with dust and wind characteristics in the Arabian Gulf. *OCEANS, Yeosu: IEEE*; 2012:1–6.
- Al-Tajer A, Thalib L. Short-term effect of dust storms on the risk of mortality due to respiratory, cardiovascular and all-causes in Kuwait. *Int J Biometeorol* 2013. <http://dx.doi.org/10.1007/s00484-012-0626-7>.
- Amanollahi J, Kabbdvandpour S, Abdullah AM, Ramli MF. Accuracy assessment of moderate resolution image spectroradiometer products for dust storms in semiarid environment. *Int J Environ Sci Technol* 2011;8:373–80.
- Anderson C. Agent-based modelling of coccidioidomycosis. [Doctoral dissertation] USA: University of Pittsburgh; 2013 [<http://d-scholarship.pitt.edu/18000/>]. Accessed 6th September, 2013].
- Ataniyazova OA, Baumann RA, Liem AKD, Mukhopadhyay UA, Vogelaar EF, Boersma ER. Levels of certain metals, organochlorine pesticides and dioxins in cord blood, maternal blood, human milk and some commonly used nutrients in the surroundings of the Aral Sea (Karakalpakstan, Republic of Uzbekistan). *Acta Paediatr* 2001;90:801–8.
- Bach AJ, Brazel AJ, Lancaster N. Temporal and spatial aspects of blowing dust in the Mojave and Colorado Deserts of southern California (1973–1994). *Phys Geogr* 1996;17:329–53.
- Badarinarath KVS, Kharol SK, Kaskaoutis DG, Sharma AR, Ramaswamy V, Kambezidis HD. Long-range transport of dust aerosols over the Arabian Sea and Indian region – a case study using satellite data and ground-based measurements. *Global Planet Change* 2010;72:164–81.
- Baddock MC, Strong CL, Murray PS, McTainsh GH. Aeolian dust as transport hazard. *Atmos Environ* 2013;71:7–14.
- Barkan J, Alpert P. Synoptic analysis of a rare event of Saharan dust reaching the Arctic region. *Weather* 2010;65:208–11.
- Barnett AG, Fraser JF, Munck L. The effects of the 2009 dust storm on emergency admissions to a hospital in Brisbane, Australia. *Int J Meteorol* 2012;56:719–26.
- Bar-Ziv J, Goldberg GM. Simple siliceous pneumoconiosis in Negev Bedouins. *Arch Environ Health* 1974;29:121–6.
- Ben-Ami Y, Koren I, Rudich Y, Artaxo P, Martin ST, Andreae MO. Transport of Saharan dust from the Bodélé Depression to the Amazon Basin: a case study. *Atmos Chem Phys* 2010;10:4345–72.
- Bennett CM, McKendry IG, Kelly S, Denike K, Koch T. Impact of the 1998 Gobi dust event on hospital admissions in the Lower Fraser Valley, British Columbia. *Sci Total Environ* 2006;366:918–25.
- Bennion P, Hubbard R, O'Hara S, Wiggs G, Wegerdt J, Lewis S, et al. The impact of airborne dust on respiratory health in children living in the Aral Sea region. *Int J Epidemiol* 2007;36:1103–10.
- Bhattachan A, D'Odorico P, Okin G, Dintwe K. Potential dust emissions from the southern Kalahari's dunelands. *J Geophys Res Earth Surf* 2013;118:307–14.
- Birmili W, Schepanski K, Ansmann A, Spindler G, Tegen I, Wehner B, et al. An episode of extremely high PM concentrations over central Europe caused by dust emitted over the southern Ukraine. *Atmos Chem Phys* 2007;7:12231–8.
- Birmili W, Schepanski K, Ansmann A, Spindler G, Tegen I, Wehner B, et al. A case of extreme particulate matter concentrations over Central Europe caused by dust emitted over the southern Ukraine. *Atmos Chem Phys* 2008;8:997–1016.
- Brahney J, Ballantyne AP, Sievers C, Neff JC. Increasing  $\text{Ca}^{2+}$  deposition in the western US: the role of mineral aerosols. *Aeol Res* 2013;10:77–87.
- Brazel A, Hsu S. The climatology of hazardous Arizona dust storms. In: Péwé TL, editor. *Desert dust*. 186. *Geol Soc Am Spec Pap*; 1981. p. 293–303.
- Breslow PB, Sailor DJ. Vulnerability of wind power resources to climate change in the continental United States. *Renew Energy* 2002;27:585–98.
- Breuning-Madsen H, Awadzi TW. Harmattan dust deposition and particle size in Ghana. *Catena* 2005;63:23–38.
- Bristow CS, Drake N, Armitage S. Deflation in the dustiest place on Earth: the Bodélé depression Chad. *Geophys J Roy Astron Soc* 2009;105:50–8.
- Brook RD, Rajagopalam S, Arden Pope C, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, et al. Particulate matter air pollution and cardiovascular disease. An update to the scientific statement from the American Heart Association. *Circulation* 2010. <http://dx.doi.org/10.1161/CIR.0b013e3181dbec1>.
- Brooks N, Chiapello I, Di Lernia S, Drake N, Legrand M, Moulin C, et al. The climate–environment–society nexus in the Sahara from prehistoric times to the present day. *J N Afr Stud* 2005;10:253–92.
- Bryant RG. Monitoring hydrological controls on dust emissions: preliminary observations from Etosha Pan, Namibia. *Geophys J Roy Astron Soc* 2003;169:131–41.
- Bryant RG, Bigg GR, Mahowald NM, Eckardt FD, Ross SG. Dust emission response to climate in southern Africa. *J Geophys Res* 2007;112:D09207. <http://dx.doi.org/10.1029/2005JD007025>.
- Bullard J, Baddock M, McTainsh G, Leys J. Sub-basin scale dust source geomorphology detected using MODIS. *Geophys Res Lett* 2008;35:L15404. <http://dx.doi.org/10.1029/2008GL033928>.
- Cabello M, Orza JAG, Barrero MA, Gordo E, Berasaluce A, Canton L, et al. Spatial and temporal variation of the impact of an extreme Saharan dust event. *J Geophys Res* 2012;117:D11204. <http://dx.doi.org/10.1029/2012JD017513>.
- Cahill TA, Gill TE, Reid JS, Gearhart EA, Gillette DA. Saltating particles, playa crusts and dust aerosols at Owens (dry) Lake, California. *Earth Surf Process Landf* 1996;21:621–39.
- Chan YC, McTainsh G, Leys J, McGowan H, Tews K. Influence of the 23 October 2002 dust storm on the air quality of four Australian cities. *Water Air Soil Pollut* 2005;164:329–48.
- Chang CC, Lee IM, Tsai SS, Yang CY. Correlation of Asian dust storm events with daily clinic visits for allergic rhinitis in Taipei, Taiwan. *J Toxicol Environ Health A* 2006;69:229–35.
- Chen PS, Feng TT, Lin CK, Yang CY, Chan CC, Young CY, Lee CH. Ambient influenza and avian influenza virus during dust storm days and background days. *Environ Health Perspect* 2010;118:1211–6.
- Cheng MF, Ho SC, Chiu HF, Wu TN, Chen PS, Yang CY. Consequences of exposure to Asian Dust Events on daily pneumonia hospital admissions in Taipei, Taiwan. *J Toxicol Environ Health A* 2008;71:1295–9.
- Chi HH, Hsu SC, Wang SH, Chang MB. Increases in ambient PCDD/F and PCB concentrations in Northern Taiwan during an Asian dust storm episode. *Sci Total Environ* 2008;401:100–8.
- Chien LC, Yang CH, Yu HL. Estimated effects of Asian Dust Storms on spatiotemporal distributions of clinic visits for respiratory diseases in Taipei children (Taiwan). *Environ Health Perspect* 2012;120:1215–20.
- Choi H, Shin DW, Kim W, Doh SJ, Lee SH, Noh M. Asian dust storm particles induce a broad toxicological transcriptional program in human epidermal keratinocytes. *Toxicol Lett* 2011;200:92–9.
- Chun Y, Boo K-O, Kim Y, Park S-U, Lee M. Synopsis, transport and physical characteristics of Asian dust in Korea. *J Geophys Res* 2001;106(D):18461–9.
- Chung YS, Kim HS, Dulam J, Harris J. On heavy dustfall observed with explosive sandstorms in Chongwon-Chongju, Korea in 2002. *Atmos Environ* 2003;37:3425–33.
- Claiborn CS, Finn D, Larsson TV, Koenig JQ. Windblown dust contributes to high  $\text{PM}_{2.5}$  concentrations. *J Air Waste Manag Assoc* 2000;50:1440–5.
- Comrie AC. Climate factors influencing Coccidioidomycosis seasonality and outbreak. *Environ Health Perspect* 2005;113:688–92.
- Cowie G, Lawson W, Kim N. Australian dust causing respiratory disease admissions in some North Island, New Zealand hospitals. *N Z Med J* 2010;123:87.
- Csavina J, Field J, Taylor MP, Gao S, Landázuri A, Betterton EA, et al. A review on the importance of metals and metalloids in atmospheric dust and aerosol from mining operations. *Sci Total Environ* 2012;433:58–73.
- Dadvand P, Basagana X, Figueras F, Amoly E, Tobias E, de Nazelle A, Querol X, et al. Saharan dust episodes and pregnancy. *J Environ Monit* 2011;13:3222–8.
- De Longueville F, Hountondji Y-C, Henry S, Ozer P. What do we know about effects of desert dust on air quality and human health in West Africa compared to other regions? *Sci Total Environ* 2010;409:1–8.
- De Longueville F, Ozer P, Doumbia S, Henry S. Desert dust impacts on human health: an alarming worldwide reality and a need for studies in West Africa. *Int J Biometeorol* 2013;57:1–19.
- De Villiers M, van Heerden J. Nashi dust storm over the United Arab Emirates. *Weather* 2011;66:79–81.
- Dement'eva AL, Zhamsueva GS, Zayakhanov AS, Tsydyrov VV, Ayurzhanayev AA, Azzayaa D, et al. Mass concentration of  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$  fine-dispersed aerosol fractions in the eastern Gobi Desert. *Russ Meteorol Hydrol* 2013;38:80–7.
- Derbyshire E. Natural minero-genetic dust and human health. *Ambio* 2007;36:73–7.
- Deroubaix A, Martiny N, Chiapello I, Marticoréna B. Suitability of OMI aerosol index to reflect mineral dust surface conditions: preliminary application for studying the link with meningitis epidemics in the Sahel. *Remote Sens Environ* 2013;133:116–27.
- Eckardt F, Kuring N. SeaWiFS identifies dust sources in the Namib Desert. *Int J Remote Sens* 2005;26:4159–67.
- Eckardt F, Washington R, Wilkinson MJ. The origin of dust on the west coast of southern Africa. *Palaeogeogr Afr* 2002;27:207–19.
- Engelbrecht JP, McDonald EV, Gillies JA, Jayanty RKM, Casuccio G, Gertler AW. Characterising mineral dusts and other aerosols from the Middle East – part 1: ambient sampling. *Inhal Technol* 2009;21:297–326.
- Engelstaedter S. Dust storm frequencies and their relationship to land surface conditions. [Diploma thesis] Jena: Friedrich-Schiller-University, Institute for Geoscience; 2001.
- Engelstaedter S, Tegen I, Washington R. North African dust emissions and transport. *Earth Sci Rev* 2006;79:73–100.
- Field JP, Belnap J, Breshears DD, Neff JC, Okin GS, Whicker JJ, et al. The ecology of dust. *Front Ecol Environ* 2010;8:423–30.
- Fleming LE, Kirkpatrick C, Backer L, Walsh CJ, Nierenberg K, Clark J, et al. Review of Florida red tide and human health effects. *Harmful Algae* 2011;10:224–33.
- Fu Q, Zhuang G, Li J, Huang K, Wang Q, Zhang R, Fu J, et al. Source, long-range transport, and characteristics of a heavy dust pollution event in Shanghai. *J Geophys Res* 2010;115:D00K29. <http://dx.doi.org/10.1029/2009JD013208>.
- Fukuyama T, Fujiwara H. Contribution of Asian dust to atmospheric deposition of radioactive cesium ( $^{137}\text{Cs}$ ). *Sci Total Environ* 2008;405:389–95.
- Furlow B. Desert storm: the pulmonary legacies of Afghanistan and Iraq. *Lancet Respir Med* 2013;1:15–6.
- Gabrieli J, Cozzi G, Vailongla P, Schwikowski M, Sigl M, Boutron C, et al. Historical reconstruction on Plutonium contamination in the Swiss–Italian Alps. *E3S Web of Conferences*, 1; 2013. <http://dx.doi.org/10.1051/e3sconf/20130114001>.
- Gaiero DM, Simonella L, Gasso S, Gili S, Stein AF, Soas P, et al. Ground/satellite observations and modeling of dust storms originating in the high Puna–Altiplano desert (South America): implications for the interpretation of paleo-climatic archives. *J Geophys Res Atmos* 2013;118:3817–31.



- Ganor E, Osetinsky I, Stupp A, Alpert P. Increasing trend of African dust, over 49 years, in the eastern Mediterranean. *J Geophys Res* 2010;115:D070201. <http://dx.doi.org/10.1029/2009JD012500>.
- Gao H, Washington R. The spatial and temporal characteristics of TOMS AI over the Tarim Basin, China. *Atmos Environ* 2009;43:1106–15.
- Gao H, Washington R. Arctic oscillation and the interannual variability of dust emissions from the Tarim Basin: a TOMS AI based study. *Climate Dynam* 2010;35:511–22.
- Gassó S, Stein AF. Does dust from Patagonia reach the sub-Antarctic Atlantic Ocean? *Geophys Res Lett* 2007;34:L01801. <http://dx.doi.org/10.1029/2006GL027693>.
- Gharai B, Jose S, Mahalakshmi DV. Monitoring intense dust storms over the Indian region using satellite data—a case study. *Int J Remote Sens* 2013;34:7038–48.
- Ghasem A, Shamsipour A, Miri M, Safarrad T. Synoptic and remote sensing analysis of dust events in southwestern Iran. *Nat Hazards* 2012;64:1625–38.
- Giannadaki D, Pozzer A, Lelieveld J. Modeled global effects of airborne desert dust on air quality and premature mortality. *Atmos Chem Phys Disc* 2013;13:24023–50.
- Gill TE, Gillette DA, Niemeyer T, Winn RT. Elemental geochemistry of wind-erodible playa sediments, Owens Lake, California. *Nucl Instr Meth B* 2002;189:209–13.
- Gillette DA, Hanson KJ. Spatial and temporal variability of dust production caused by wind erosion in the United States. *J Geophys Res* 1989;94(D2):2197–206.
- Ginoux P, Prospero JM, Gill TE, Hsu NC, Zhao M. Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products. *Rev Geophys* 2012;50:RG3005.
- Givethi R, Arhami M, Tajrishy M. Contribution of the Middle Eastern dust source areas to PM<sub>10</sub> levels in urban receptors: case study of Tehran, Iran. *Atmos Environ* 2013;75:287–95.
- Goudie AS, Middleton NJ. The changing frequency of dust storms through time. *Clim Chang* 1992;20:197–225.
- Goudie AS, Middleton NJ. Saharan dust storms: nature and consequences. *Earth Sci Rev* 2001;56:179–204.
- Goudie AS, Middleton NJ. Desert dust in the global system. Heidelberg: Springer Verlag; 2006.
- Goudie AS, Wells GL. The nature, distribution and formation of pans in arid zones. *Earth Sci Rev* 1995;38:1–69.
- Griffin DW. Atmospheric movement of microorganisms in clouds of desert dust and implications for human health. *Clin Microbiol Rev* 2007;20:459–77.
- Griffin DW, Kellogg CA. Dust storms and their impact on ocean and human health: dust in Earth's atmosphere. *Ecohealth* 2004;1:284–95.
- Grishkan I, Schlesinger P, Mamane Y. Influence of dust storms on concentration and content of fungi in the atmosphere of Haifa, Israel. *Aerobiologia*; 2012. <http://dx.doi.org/10.1007/s10453-012-9256-0>.
- Groll M, Opp C, Aslanov I. Spatial and temporal distribution of dust deposition in Central Asia—results from a long term monitoring program. *Aeol Res* 2013;9:49–62.
- Gyan K, Henry W, Lacaille S, Laloo A, Lamsee-Ebanks C, McKay S, et al. African dust clouds are associated with increased paediatric asthma accident and emergency admissions on the Caribbean island of Trinidad. *Int J Biometeorol* 2005;49:371–6.
- Hahnenberger M, Nicoll K. Meteorological characteristics of dust storm events in the eastern Great Basin of Utah, U.S.A. *Atmos Environ* 2012;60:601–12.
- Hamidi M, Kavianpour MR, Shao Y. Synoptic analysis of dust storms in the Middle East. *Asia Pac J Atmos Sci* 2013;49:279–86.
- Herut B, Krom MD, Pan G, Mortimer R. Atmospheric input of nitrogen and phosphorus to the Southeast Mediterranean: sources, fluxes, and possible impact. *Limnol Oceanogr* 1999;44:1683–92.
- Hesse PP, McTainsh GH. Last glacial maximum to early Holocene wind strength in the mid-latitudes of the southern hemisphere from aeolian dust in the Tasman Sea. *Quat Res* 1999;52:343–9.
- Hickey B, Goudie AS. The use of TOMS and MODIS to identify dust storm source areas: the Tokar delta (Sudan) and the Seistan basin (south west Asia). In: Goudie AS, Kalvoda J, editors. *Geomorphological Variations*, P3K, Prague; 2007. p. 37–57.
- Hou X, Zhuang G, Sun Y, An Z. Characteristics and sources of polycyclic aromatic hydrocarbons and fatty acids in PM<sub>2.5</sub> aerosols in dust season in China. *Atmos Environ* 2006;40:3251–62.
- Hussain A, Mir H, Afzal M. Analysis of dust storms frequency over Pakistan during (1961–2000). *Pak J Meteorol* 2005;2:49–68.
- Indoito R, Orlovsky L, Orlovsky N. Dust storms in Central Asia: spatial and temporal variations. *J Arid Environ* 2012;85:62–70.
- IPCC. Climate change: the physical science basis. In: Solomon S, et al, editors. *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge and New York: Cambridge University Press; 2007.
- Jeon EM, Kim HJ, Jung K, Kim JH, Kim MY, Kim YP, et al. Impact of Asian dust events on airborne bacterial community assessed by molecular analyses. *Atmos Environ* 2011;45:4313–21.
- Jiang Y, Luo Y, Zhao Z, Tao S. Changes in wind speed over China during 1956–2004. *Theor Appl Climatol* 2009. <http://dx.doi.org/10.1007/s00704-009-0152-7>.
- Jilbert T, Reichard G-J, Aeschlimann B, Günther D, Boer W, de Lange G. Climate-controlled multidecadal variability in North African dust transport to the Mediterranean. *Geology* 2010;38:19–22.
- Jiménez E, Linares C, Martínez D, Díaz J. Role of Saharan dust in the relationship between particulate matter and short-term daily mortality among the elderly in Madrid (Spain). *Sci Total Environ* 2010;408:5729–36.
- Johnston F, Hanigan I, Henderson S, Morgan G, Bowman D. Extreme air pollution events from bushfires and dust storms and their association with mortality in Sydney, Australia 1994–2007. *Environ Res* 2011;111:811–6.
- Jugder D, Shinoda M, Sugimoto N, Matsui I, Nishikawa M, Park SU, et al. Spatial and temporal variations of dust concentrations in the Gobi Desert of Mongolia. *Global Planet Change* 2011;78:14–22.
- Kanatani KT, et al. Desert dust exposure is associated with increased risk of asthma hospitalization in children. *Am J Respir Crit Care Med* 2010;182:1475–81.
- Kang JH, Keller JI, Chen SC, Lin HC. Asian dust storm events are associated with an acute increase in pneumonia hospitalization. *Ann Epidemiol* 2012;22:257–63.
- Kang JH, Liu TC, Keller J, Lin HC. Asian storm events are associated with an acute increase in stroke hospitalization. *J Epidemiol Community Health* 2013;67:125–31.
- Karanasiou A, Moreno N, Moreno T, Viana M, de Leeuw F, Querol X. Health effects from Sahara dust episodes in Europe: literature review and research gaps. *Environ Int* 2012;47:107–14.
- Kaskaoutis DG, Kambezidis HD, Nastos PT, Kosmopoulos PG. Study on an intense dust storm over Greece. *Atmos Environ* 2008;42:6884–96.
- Kayetha VK, Senthilkumar J, Prasad AK, Cervone G, Singh RP. Effect of dust storm on ocean color and snow parameters. *J Ind Soc Remote Sens* 2007;35:1–9.
- Kes AS, Fedorovich BA. Process of forming of aeolian dust in space and time. 23rd Int. Geogr. Cong., Section 1; 1976. p. 174–7.
- Kikawada Y, Oda K, Nomura M, Honda T, Oi T, Hirose K, et al. Origin of enriched uranium contained in Japanese atmospheric deposits. *Nat Sci* 2012;4(Special Issue):936–42.
- Knippertz P, Todd MC. The central west Saharan dust hot spot and its relation to African easterly waves and extratropical disturbances. *J Geophys Res* 2010;115:D12117. <http://dx.doi.org/10.1029/2009JD012819>.
- Koren I, Joseph JH, Israelevich P. Detection of dust plumes and their sources in northeastern Libya. *Can J Remote Sens* 2003;29:792–6.
- Krasnov H, Katra I, Koutrakis P, Friger D. Contribution of desert dust storms to PM<sub>10</sub> levels in an urban arid environment, Beer-Sheva, Negev, Israel. *J Air Waste Manag Assoc* 2013. <http://dx.doi.org/10.1080/10962247.2013.841598>.
- Kravtsova VI, Tarasenko TV. Space monitoring of Aral Sea degradation. *Water Res* 2010;37:285–96.
- Kurosaki Y, Shinoda M, Mikami M. What caused a recent increase in dust outbreaks over East Asia? *Geophys Res Lett* 2011;38:L11702. <http://dx.doi.org/10.1029/2011GL047494>.
- Kwon SH, Kim YP, Lee JY. Impact of dust storms on the organic composition in the ambient aerosol. *Aerosol Air Qual Res* 2013;13:97–106.
- Laken BA, Parviainen H, Pallé E, Shabaz T. Saharan mineral dust outbreaks observed over the North Atlantic island of La Palma in summertime between 1984 and 2012. *Q J Roy Meteorol Soc* 2013. <http://dx.doi.org/10.1002/qj.2170>.
- Leathers CR. Plant components of desert dust in Arizona and their significance for man. *Geol Soc Am Spec Pap* 1981;186:191–206.
- Lee JW, Lee KK. Effects of Asian dust events on daily asthma patients in Seoul, Korea. *Meteorol Appl* 2013. <http://dx.doi.org/10.1002/met.1351>.
- Lee YC, Yang X, Wenig M. Transport of dusts from east Asian and non-East Asian sources to Hong Kong during dust storm related events 1996–2007. *Atmos Environ* 2010;44:3728–38.
- Lee PK, Yoon SJ, Jo HY. Heavy metal concentrations and contamination levels from Asian dust and identification of sources: a case-study. *Chemosphere* 2013;91:1018–25.
- Lei H, Wang JXL. Characteristics of dust storm events over the western United States. *Atmos Chem Phys Disc* 2013;13:14195–220.
- Lenes JM, Darrow BA, Walsh JJ, Prospero JM, He R, Weisberg RH, et al. Saharan dust and phosphatic fidelity: a three-dimensional biogeochemical model of *Trichodesmium* as a nutrient source for red tides on the West Florida Shelf. *Cont Shelf Res* 2008;28:1091–115.
- Leski TA, Malanoski AP, Gregory MJ, Lin B, Stenger DA. Application of a broad-range resequencing array for detection of pathogens in desert dust samples from Kuwait and Iraq. *Appl Environ Microbiol* 2011;77:4285–92.
- Leys JF, Heidenreich SK, Strong CL, McTainsh GH, Quigley S. PM<sub>10</sub> concentrations and mass transport during “red Dawn”—Sydney 23 September 2009. *Aeolian Res* 2011;3:327–42.
- Li J, Wang Z, Zhuang G, Luo G, Sun Y, Wang Q. Mixing of Asian mineral dust with anthropogenic pollutants over East Asia: a model case study of a super-duststorm in March 2010. *Atmos Chem Phys* 2012;12:7591–607.
- Li X, Feng L, Huang C, Yan X, Zhang X. Potential hazardous elements (PHEs) in atmospheric particulate matter (APM) in the south of Xi'an during the dust episodes of 2001–2012 (NW China): chemical fractionation, ecological and health risk assessment. *Environ Earth Sci* 2013. <http://dx.doi.org/10.1007/s12665-013-2800-6>.
- Lien YJ, Chien LC, Yang CH, Yu HL. Health impact from climatic extremes: a case study of Asian dust storms in Taiwan. *Geophys Res Abstr* 2013;15:EGU2013–5817.
- Liu D, Abuduwalili J, Lei J, Wu G. Deposition rate and chemical composition of the aeolian dust from a bare saline playa, Ebinur Lake, Xinjiang, China. *Water Air Soil Pollut* 2011;218:175–84.
- Mahowald NM, Luo C. A less dusty future? *Geophys Res Lett* 2003;30. <http://dx.doi.org/10.1029/2003GL017880>.
- Mahowald NM, Engelstaedter S, Luo C, Sealy A, Artaxo P, Benitez-Nelson C, et al. Atmospheric iron deposition: global distribution, variability, and human perturbations. *Ann Rev Mar Sci* 2009;1:245–78.
- Mahowald NM, Kloster S, Engelstaedter S, Moore JK, Mukhopadhyay S, McConnell JR, et al. Observed 20th century desert dust variability: impact on climate and biogeochemistry. *Atmos Chem Phys* 2010;10:10875–93.
- Mallone S, Staffagia M, Faustini A, Gobbi GP, Marconi A, Forastiere F. Saharan dust and associations between particulate matter and daily mortality in Rome, Italy. *Environ Health Perspect* 2011;119:1409–14.
- Marticorena B, Chatenet B, Rajot JL, Traore S, Coulibaly M, Diallo A, et al. Temporal variability of mineral dust concentrations over West Africa: analyses of a pluriannual monitoring from the AMMA Sahelian Dust Transect. *Atmos Chem Phys* 2010;10:8899–915.
- Martigny N, Chiappello I. Assessments for the impact of mineral dust on the meningitis incidence in West Africa. *Atmos Environ* 2013;70:245–53.



- Martinelli N, Olivieri O, Girelli D. Air particulate matter and cardiovascular disease: a narrative review. *Eur J Intern Med* 2013;24:295–302.
- Marx SK, Kamber BS, McGowan HA. Estimates of Australian dust flux into New Zealand: quantifying the eastern Australian dust plume pathway using trace element calibrated  $^{210}\text{Pb}$  as a monitor. *Earth Planet Sci Lett* 2005;239:336–51.
- Masson O, Piga D, Gurriaran R, D'Amico D. Impact of an exceptional Saharan dust outbreak in France:  $\text{PM}_{10}$  and artificial radionuclides in air and in dust deposit. *Atmos Environ* 2010;44:2478–86.
- Mathur ML, Choudhary RC. Desert lung syndrome in rural dwellers of the Thar Desert, India. *J Arid Environ* 1997;35:559–62.
- McConnell JR, Aristarain AJ, Banta JR, Edwards PR, Simões JC. 20th-century doubling in dust archived in an Antarctic Peninsula ice core parallels climate change and desertification in South America. *Proc Natl Acad Sci U S A* 2007;104:5743–8.
- McMoli TE, Bordon AN, Munube GMK, Bell EJ. Epidemic acute haemorrhagic conjunctivitis in Lagos, Nigeria. *Br J Ophthalmol* 1984;68:401–4.
- McTainsh GH. Quaternary aeolian dust processes and sediments in the Australian region. *Quat Sci Rev* 1989;8:235–53.
- McTainsh GH, Chan Y-C, McGowan H, Leys J, Tews K. The 23rd October 2002 dust storm in eastern Australia: characteristics and meteorological conditions. *Atmos Environ* 2005;39:1227–36.
- McTainsh GH, Leys J, Baker M, Tews K, Barton K. Wind erosion histories, model input data and community DustWatch. Brisbane: Griffith University; 2010.
- Meng Z, Lu B. Dust events as a risk factor for daily hospitalization for respiratory and cardiovascular diseases in Mingin, China. *Atmos Environ* 2007;41:7048–58.
- Merrifield A, Schindler S, Jalaludin B, Smith W. Health effects of the September 2009 dust storm in Sydney, Australia: did emergency department visits and hospital admissions increase? *Environ Health* 2013. <http://dx.doi.org/10.1186/1476-069X-12-32>.
- Metcalfe JS, Richer R, Cox PA, Godd GA. Cyanotoxins in desert environments may present a risk to human health. *Sci Total Environ* 2012;421–422:118–23.
- Middleton NJ. A geography of dust storms in south-west Asia. *J Climatol* 1986a;6:183–96.
- Middleton NJ. The Geography of Dust Storms. Unpublished DPhil Thesis. University of Oxford, 1986b.
- Middleton NJ. Dust storms in the Middle East. *J Arid Environ* 1986c;10:83–96.
- Middleton NJ, Goudie AS. Saharan dust: sources and trajectories. *Trans Inst Brit Geogr* 2001;NS 26:165–81.
- Middleton N, Yiallouris P, Kleanthous S, Kolokotroni O, Schwartz J, Dockery DW, et al. A 10-year time-series analysis of respiratory and cardiovascular morbidity in Nicosia, Cyprus: the effect of short-term changes in air pollution and dust storms. *Environ Health* 2008. <http://dx.doi.org/10.1186/1476-069X-7-39>.
- Miller RL, Tegen I, Perlwitz J. Surface radiative forcing by soil dust aerosols and the hydrological cycle. *J Geophys Res* 2004;109(D). <http://dx.doi.org/10.1029/2003JD004085>.
- Miller LD, Kuciauskas AP, Liu M, Ji Q, Reid JS, Breed DW, et al. Haboob dust storms of the southern Arabian Peninsula. *J Geophys Res* 2008;113:D01202. <http://dx.doi.org/10.1029/2007JD008550>.
- Miri A, Ahmadi H, Ghanbari A, Moghaddamnia A. Dust storms impacts on air pollution and public health under hot and dry climate. *Int J Energy Environ* 2007;1:101–5.
- Miri A, Moghaddamnia A, Pahlavanrabi A, Panjehkek N. Dust storm frequency after the 1999 drought in the Sistan region, Iran. *Clim Res* 2010;41:83–90.
- Mitchell RM, Campbell SK, Qin Y. Recent increase in aerosol loading over the Australian arid zone. *Atmos Chem Phys* 2010;10:1689–99.
- Mohammed BAR. Correlation between solar cycles and dust activity pver Iraq. *J Al Narain Univ* 2010;13:82–7.
- Morales AF. El Sahara Español. Alta Comisaria de España en Marruecos, Madrid; 1946.
- Morman SA, Plumlee GS. The role of airborne dusts in human disease. *Aeolian Res* 2013;9:203–12.
- Mulitza S, Heslop D, Pittauerova D, Fischer HW, Meyuer I, Stuut JB, et al. Increase in African dust flux at the onset of commercial agriculture in the Sahel region. *Nature* 2010;466:226–8.
- Munson S, Belnap J, Okin GS. Responses of wind erosion to climate-induced vegetation changes on the Colorado Plateau. *Proc Natl Acad Sci U S A* 2011;108:3854–9.
- Najafi MS, Khoshaklagh F, Zamanzadeh SM, Shiraz MH, Samadi M, Hajikhani S. Characteristics of TSP loads during the Middle East Springtime Dust Storm (MESDS) in western Iran. *Arab J Geosci* 2013. <http://dx.doi.org/10.1007/S12517-013-1086-z>.
- Natsagdorj DL, Jugder D, Chung YS. Analysis of dust storms observed in Mongolia during 1937–1999. *Atmos Environ* 2003;37:1401–11.
- Neff JC, Ballantyne AP, Farmer GL, Mahowald NM, Conroy JL, Landry CC, et al. Increasing eolian dust deposition in the western United States linked to human activity. *Nat Geosci* 2008;1:189–95.
- Neff JC, Reynolds RL, Munson SM, Fernandez D, Belnap J. The role of dust storms in total atmosphere particle concentrations at two sites in the western US. *J Geophys Res* 2013. <http://dx.doi.org/10.1002/jgrd.50855>.
- Neophytou A, Yiallouris P, Coull BA, Kleanthous S, Pavlou P, Pashiardis S, et al. Particulate matter concentrations during desert dust outbreaks and daily mortality in Nicosia, Cyprus. *J Expo Sci Environ Epidemiol* 2013;23:275–80.
- Norboo T, Angchuk PT, Yahya M, Kamat SR, Pooley FD, Corrin B, et al. Silicosis in a Himalayan village population: role of environmental dust. *Thorax* 1991;46:341–3.
- Ogorodnikov BI. A dust storm over the Ukraine and Belarus territory contaminated by radionuclides after the Chernobyl accident. *Russ Meteorol Hydrol* 2011;36:613–23.
- O'Hara SL, Wiggs G, Mamedov B, Davidson G, Hubbard RB. Exposure to airborne dust contaminated with pesticide in the Aral Sea region. *Lancet* 2000;355:627–8.
- O'Hara SL, Clarke ML, Elatrash MS. Field measurements of desert dust deposition in Libya. *Atmos Environ* 2006;40:3881–97.
- Oho D, Kiddoo P, Howard C, David G. Application of a combined measurement and modelling method to quantify windblown dust emissions from the exposed playa at Mono Lake, California; 2011 [[http://gubapcd.org/outgoing/dono/Mono%20Lake%20Dust%20ID%202011%20\(3-3-2011\).pdf](http://gubapcd.org/outgoing/dono/Mono%20Lake%20Dust%20ID%202011%20(3-3-2011).pdf)] (accessed 3/7/2003)].
- Onishi K, Otani S, Yoshida A, Mu H, Kurozawa Y. Adverse health effects of Asian dust particles and heavy metals in Japan. *Asia Pac J Public Health* 2011. <http://dx.doi.org/10.1177/1010539511428667>.
- Onishi K, Kurosaki Y, Otani S, Yoshida A, Sugimoto N, Kurozawa Y. Atmospheric transport route determines components of Asian dust and health effects in Japan. *Atmos Environ* 2012;49:94–102.
- Orlovsky N, Orlovsky L. White sandstorms in Central Asia. In: Yang Y, Squires V, Lu Q, editors. Global alarm: dust and sandstorms from the world's drylands. Bangkok: United Nations; 2001. p. 169–201.
- Orlovsky L, Orlovsky N, Durdiev A. Dust storms in Turkmenistan. *J Arid Environ* 2005;60:83–97.
- Otani S, Onishi K, Mu H, Yokoyama Y, Hosoda T, Okamoto M, et al. The relationship between skin symptoms and allergic reactions to Asian dust. *Int J Environ Res Public Health* 2012;9:4606–14.
- Ozer P. Fifty years of African mineral dust production. *Bull Sci Ac Roy Sci* 2003;49:371–96. [d'Outre-Mer].
- Ozer P. Dust in the wind and public health: example from Mauritania. International Conference on Desertification: Migration, Health, remediation and Local Governance, Brussels, 22 September, 2006; 2006. p. 55–74.
- Panikkath R, Jumper CA, Mulkey Z. Multilobar lung infiltrates after exposure to dust storm: the Haboob Lung Syndrome. *Am J Med* 2013;126:5–7.
- Pappagianis D, Einstein H. Tempest from tehachap takes toll or Coccilioides conveyed aloft and afar. *West J Med* 1978;129:527–30.
- Park K, Dam HD. Characterization of metal aerosols in  $\text{PM}_{10}$  from urban, industrial and Asian dust sources. *Environ Monit Assess* 2010;160:289–300.
- Pauley PM, Baker NL, Barker EH. An observational study of the "Interstate 5" dust storm case. *Bull Am Meteorol Soc* 1996;77:693–720.
- Pease PP, Tchakerian VP, Tindale NW. Aerosols over the Arabian Sea: geochemistry and source areas for aeolian desert dust. *J Arid Environ* 1998;39:477–96.
- Pelletier JD. Sensitivity of playa wind-blown dust emissions to climatic and anthropogenic change. *J Arid Environ* 2006;66:62–75.
- Perez L, Tobias A, Querol X, Kunzli N, Pey J, Alastuey A, et al. Coarse particles from Saharan dust and daily mortality. *Epidemiology* 2008;19:800–7.
- Perez L, Tobias A, Querol X, Pey J, Alastuey A, Diaz J, et al. Saharan dust, particulate matter and cause-specific mortality: a case-crossover study in Barcelona (Spain). *Environ Int* 2012;48:150–5.
- Perfumo A, Marchant R. Global transport of thermophilic bacteria in atmospheric dust. *Environ Microbiol Rep* 2010;2:333–9.
- Perry KD, Cahill TA, Eldred RA, Dutcher DD. Long-range transport of North African dust to the eastern United States. *J Geophys Res* 1997;102:11225–38.
- Pey J, Querol X, Alastuey A, Forastiere F, Stafoggia M. African dust outbreaks over the Mediterranean Basin during 2001–2011:  $\text{PM}_{10}$  concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology. *Atmos Chem Phys* 2013;13:1395–410.
- Polymenakou PN, Mandalakis M, Stephanou EG, Tselepidis A. Particle size distribution of airborne microorganisms and pathogens during an intensive African dust event in the eastern Mediterranean. *Environ Health Perspect* 2008;116:292–6.
- Prospero JM, Ginoux P, Torres O, Nicholson SE, Gill TE. Environmental characterisation of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product. *Rev Geophys* 2002;40:2–31.
- Prospero JM, Blades E, Naidu R, Mathison G, Thani H, Lavoie MC. Relationship between African dust carried in the Atlantic trade winds and surges in pediatric asthma attendances in the Caribbean. *Int J Biometeorol* 2008;52:823–32.
- Rashki A, Kaskoutis DG, Rautenbach CJ deW, Eriksson PG, Qiang M, Gupta P. Dust storms and their horizontal dust loading in the Sistan region, Iran. *Aeolian Res* 2012;5:51–62.
- Rashki A, Rautenbach CJ deW, Eriksson PG, Kaskoutis DG, Gupta P. Temporal changes of particulate concentration in the ambient air over the city of Zahedan, Iran. *Air Qual Atmos Health* 2013a;6:123–35.
- Rashki AR, Kaskoutis DG, Goudie AS, Kahn RA. Dryness of ephemeral lakes and consequences for dust activity: the case of the Hamoun drainage basin, southeastern Iran. *Sci Total Environ* 2013b;463–464:552–64.
- Ravi S, D'Odorico P, Breshears DD, Field JP, Goudie AS, Kahn RA, et al. Aeolian processes and the biosphere. *Rev Geophys* 2011;49:RG3001. <http://dx.doi.org/10.1029/2010RG000328>.
- Resane T, Annegarn H, Freiman T. The day of the white rain: origin of unusual dust deposition in Johannesburg, South Africa. *S Afr J Sci* 2004;100:483–7.
- Resch F, Sunnu A, Afeti G. Saharan dust flux and deposition rate near the Gulf of Guinea. *Tellus* 2007. <http://dx.doi.org/10.1111/j.1600-0889.2007.00286.x>.
- Reyes M, Díaz J, Tobias A, Montero JC, Linares C. Impact of Saharan dust particles on hospital admissions in Madrid (Spain). *Int J Environ Health Res* 2013. <http://dx.doi.org/10.1080/09603123.782604>.
- Rutherford S, Clark E, McTainsh GH, Simpson R, Mitchell C. Characteristics of rural dust events shown to impact on asthma severity in Brisbane, Australia. *Int J Biometeorol* 1999;42:217–25.
- Saiyed HN, Sharma YK, Sadhu HG, Norboo T, Patel PD, Venkaiah K, et al. Non-occupational pneumoconiosis at high altitude villages in central Ladakh. *Br J Ind Med* 1991;48:825–9.
- Sajani SZ, Miglio R, Bonasoni P, Cristofanelli P, Marinoni A, Sartini C, et al. Saharan dust and daily mortality in Emilia-Romagna (Italy). *Occup Environ Med* 2011;68:446–51.
- Samoli E, Kougea E, Kassomenos P, Analitis A, Katsouyanni K. Does the presence of desert dust modify the effect of  $\text{PM}_{10}$  on mortality in Athens, Greece? *Sci Total Environ* 2011;409:2049–54.

- Sandstrom T, Forsberg B. Desert dust. An unrecognized source of dangerous pollution? *Epidemiology* 2008;19:808–9.
- Santese M, De Tomasi F, Perrone ME. Moderate Resolution Imaging Spectroradiometer (MODIS) and Aerosol Robotic Network (AERONET) retrievals during dust outbreaks over the Mediterranean. *J Geophys Res* 2007;112:D18201. <http://dx.doi.org/10.1029/2007JD008482>.
- Schlesinger P, Mamane Y, Grishkan I. Transport of microorganisms to Israel during Saharan dust events. *Aerobiologia* 2006;22:259–73.
- Schulz H, Harder V, Ibalid-Mulli A, Khandoga A, Koenig W, Krombach F, et al. Cardiovascular effects of fine and ultrafine particles. *J Aerosol Med* 2005;18:1–22.
- Schwartz J, Norris G, Larson T, Sheppard L, Claiborne C, Koenig J. Episodes of high coarse particle concentrations are not associated with increased mortality. *Environ Health Perspect* 1999;107:339–42.
- Shahsavani A, Naddafi K, Haghighifard NJ, Mesdaghinia A, Yunesian M, Nabizadeh R, et al. The evaluation of PM<sub>10</sub>, PM<sub>2.5</sub> and PM<sub>1</sub> concentrations during the Middle Eastern Dust (MED) events in Ahvaz, Iran, from April through September 2010. *J Arid Environ* 2012;77:72–83.
- Shao Y, Wang JJ. A climatology of Northeast Asian dust events. *Meteorol Z* 2003;12:175–83.
- Shao Y, Wyrwoll KH, Chappell A, Huang J, Lin Z, McTainsh GH, et al. Dust cycle: an emerging core theme in Earth system science. *Aeolian Res* 2011;2:181–204.
- Shi JH, Zhang J, Gao HW, Tan SC, Yao XH, Ren JL. Concentration, solubility and deposition flux of atmospheric particulate nutrients over the Yellow Sea. *Deep-Sea Res* 2013. <http://dx.doi.org/10.1016/j.dsr.2013.05.004>. [in press].
- Shiga Y, Greene RSB, Scott KM, Stelcer E. Recognising terrestrially-derived salt (NaCl) in ESE Australian dust. *Aerobiologia* 2011;27:215–20.
- Soleimani Z, Goudarzi G, Naddafi K, Sadeghinejad B, Lattifi SM, Parhizgari N, et al. Determination of culturable indoor airborne fungi during normal and dust event days in Ahvaz, Iran. *Aerobiologia* 2013;29:279–90.
- Song Z, Wang J, Wang S. Quantitative classification of northeast Asian dust events. *J Geophys Res* 2007;112:D04211. <http://dx.doi.org/10.1029/2006JD007048>.
- Steenburgh WJ, Massey JD, Painter TH. Episodic dust events of Utah's Wasatch front and adjoining region. *J Appl Meteorol Climatol* 2012;51:1654–69.
- Stout JE, Lee JA. Indirect evidence of wind erosion trends on the southern High Plains of North America. *J Arid Environ* 2003;55:43–61.
- Sultan B, Labadi K, Guégan J-F, Janicot S. Climate drives the Meningitis epidemics onset in West Africa. *PLoS Med* 2005;2:44–9.
- Tam VVWS, Wong TW, Wong AHS, Hui DSC. Effect of dust storm events on daily emergency admissions for respiratory diseases. *Respirology* 2012;17:143–8.
- Tan SC, Shi GY, Shi JH, Gao HW, Yao X. Correlation of Asian dust with chlorophyll and primary productivity in the coastal seas of China during the period from 1998 to 2008. *J Geophys Res* 2011;116:G2029. <http://dx.doi.org/10.1029/2010JG001456>.
- Tanaka TY, Chiba M. A numerical study of the contributions of dust source regions to the global dust budget. *Global Planet Change* 2006;52:88–104.
- Tao Y, An Z, Sun Z, Hou Q, Wang Y. Association between dust weather and number of admissions for patients with respiratory diseases in spring in Lanzhou. *Sci Total Environ* 2012;423:8–21.
- Tegen I, Schepanski K. The global distribution of mineral dust. *IOP Conf Ser Earth Environ Sci* 2009;7. <http://dx.doi.org/10.1088/1755-1307/7/1/012001>.
- Thalib L, Al-Tajer A. Dust storms and the risk of asthma admissions to hospitals in Kuwait. *Sci Total Environ* 2012;433:347–51.
- Thevenon F, Chiaradia M, Adatte T, Hueglin C, Poté J. Ancient versus modern mineral dust transported to high-altitude alpine glaciers evidences Saharan sources and atmospheric circulation changes. *Atmos Chem Phys* 2011;11:859–84.
- Thomas DSG, Knight M, Wiggs GFS. Remobilization of southern African desert dune systems by twenty-first century global warming. *Nature* 2005;435:1218–21.
- Thomson MC, Jeanne I, Djingarey M. Dust and epidemic meningitis in the Sahel: a public health and operational research perspective. *IOP Conf Ser Earth Environ Sci* 2009;7. <http://dx.doi.org/10.1088/1755-1307/7/1/012017>.
- Tobias A, Cayla JA, Pey J, Alastuey A, Querol X. Are Saharan dust emissions increasing the risk of meningococcal meningitis? *Int J Infect Dis* 2011a;15. <http://dx.doi.org/10.1016/j.ijid.2011.03.008>.
- Tobias A, Perez L, Diaz J, Linares C, Pey J, Alastuey A, et al. Short-term effects of particulate matter on total mortality during Saharan dust outbreaks: a case-crossover analysis in Madrid (Spain). *Sci Total Environ* 2011b;412:413–386–9.
- Todhunter PE, Cihacek LJ. Historical reduction of airborne dust in the Red River Valley of the North. *J Soil Water Conserv* 1999;54:543–51.
- Ueda K, Shimizu A, Nitta H, Inoue K. Long-range transported Asian Dust and emergency ambulance dispatches. *Inhal Toxicol* 2012;24:858–67.
- Uno I, Eguchi K, Yumimoto K, Takemura T, Shimizu A, Uematsu M, et al. Asian dust transport one full circuit around the globe. *Nat Geosci* 2009;2:557–60.
- Vanderstraten P, Lenelle Y, Meurrens A, Carati D, Brenig I, Declou A, et al. Dust storm originate from Sahara covering Western Europe: a case study. *Atmos Environ* 2008;42:5489–93.
- Varga G. Spatio-temporal distribution of dust storms—a global coverage using NASA TOMS aerosol measurements. *Hung Geogr Bull* 2012;61:275–98.
- Varga G, Kovács J, Ujvári G. Analysis of Saharan dust intrusions into the Carpathian Basin (Central Europe) over the period 1979–2011. *Global Planet Change* 2013;100:333–42.
- Vautard R, Cattiaux J, Yiou P, Thépaut J-N, Ciais P. Northern Hemisphere atmospheric stilling partly attributed to an increase in surface roughness. *Nat Geosci* 2010;3:756–61.
- Walsh JJ, Steidinger KA. Saharan dust and Florida red tides: the cyanophyte connection. *J Geophys Res* 2001;106(C6):11597–612.
- Walsh JJ, Jolliff JK, Darrow BP, Lenos JM, Milroy SP, Remsen A, et al. Red tides in the Gulf of Mexico: where, when and why? *J Geophys Res C* 2006;111:1–46.
- Wang X, Dong Z, Zhang J, Liu L. Modern dust storms in China: an overview. *J Arid Environ* 2004;55:59–74.
- Wang Z, Dong Z, Yan P, Zhang J, Qian G. Wind energy environments and dune field activity in the Chinese deserts. *Geophys J Roy Astron Soc* 2005;65:33–48.
- Wang X, Eerdun H, Zhou Z, Liu X. Significance of variations in the wind energy environment over the past 50 years with respect to dune activity and desertification in arid and semiarid northern China. *Geophys J Roy Astron Soc* 2007;86:252–66.
- Wang L, et al. Consecutive transport of anthropogenic air masses and dust storm plume: two case events at Shanghai, China. *Atmos Res* 2013;127:22–33.
- Washington R, Todd M, Middleton N, Goudie AS. Global dust storm source areas determined by the Total Ozone Monitoring Spectrometer and ground observations. *Ann Assoc Am Geogr* 2003;93:297–313.
- Washington R, Todd MC, Engelstaedter S, Mbainayel S, Mitchell F. Dust and the low-level circulation over the Bodélé depression, Chad: observations from BoDEx 2005. *J Geophys Res* 2006a;111:D03201. <http://dx.doi.org/10.1029/2005JD006502>.
- Washington R, Todd MC, Lizcano G, Tegen I, Flamant C, Koren I, et al. Links between topography, wind, deflation, lakes and dust: the case of the Bodélé depression, Chad. *Geophys Res Lett* 2006b;33:L09401. <http://dx.doi.org/10.1029/2006GL025827>.
- Washington R, Bouet C, Cautenet G, Mackenzie E, Ashpole I, Engelstaedter S, et al. Dust as a tipping element: the Bodélé Depression, Chad. *Proc Natl Acad Sci U S A* 2009;106:20564–71.
- Watanabe M, Igishi T, Burioka N, Yamasaki A, Kurai J, Takeuchi H, et al. Pollen augments the influence of desert dust on symptoms of adult asthma patients. *Allergol Int* 2011a;60:517–24.
- Watanabe M, Yamasaki A, Burioka N, Kurai J, Yoneda K, Yoshida A, et al. Correlation between Asian dust storms and worsening asthma in western Japan. *Allergol Int* 2011b;60:267–75.
- Wiggs GFS, O'Hara S, Wegerdt J, Van Der Meer J, Small I, Hubbard R. The dynamics and characteristics of aeolian dust in dryland Central Asia: possible impacts on human exposure and respiratory health in the Aral Sea basin. *Geophys J Roy Astron Soc* 2003;169:142–57.
- Wong S, Dessler AE, Mahowald N, Colarco PR, da Silva A. Long-term variability in Saharan dust transport and its link to North Atlantic sea surface temperature. *Geophys Res Lett* 2008;35:L07812. <http://dx.doi.org/10.1029/2007GL032297>.
- Wu TY, Lin YC, Chen YM, Chang-Chien GP. Impact of dust storms on PCDD/Fs and regulated harmful matter in a basin area. *Atmos Res* 2010;97:279–93.
- Xie S, Yu T, Zhang Y, Zeng L, Qi L, Tang X. Characteristics of PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>x</sub> and O<sub>3</sub> in ambient air during the dust storm period in Beijing. *Sci Total Environ* 2005;345:153–64.
- Xu J. Sand-dust storms in and around the Ordos Plateau of China as influenced by land use change and desertification. *Catena* 2006;65:279–84.
- Yang CY. Effects of Asian dust storm events on daily clinical visits for conjunctivitis in Taipei, Taiwan. *J Toxicol Environ Health A* 2006;69:1673–80.
- Yang ZH. Association of inhalable particles in dust events with daily outpatient visits for Tracheitis. *J Environ Occup Med* 2013;30:88–92.
- Yang CY, Chen YS, Chiu HF, Goggins WB. Effects of Asian dust storm events on daily stroke admissions in Taipei, Taiwan. *Environ Res* 2005;99:79–84.
- Yang YQ, Hou Q, Zhou CH, Liu HL, Wang YQ, Niu T. Sand/dust storm processes in Northeast Asia and associated large-scale circulations. *Atmos Chem Phys* 2008;8:25–33.
- Yang CY, Cheng MH, Chen CC. Effects of Asian dust storm events on hospital admissions for congestive heart failure in Taipei, Taiwan. *J Toxicol Environ Health A* 2009;72:324–8.
- Yao Z, Xiao J, Li C, Zhu K. Regional characteristics of dust storms observed in the Alxa Plateau of China from 1961 to 2005. *Environ Earth Sci* 2010. <http://dx.doi.org/10.1007/s12665-010-0845-3>.
- Yoshino M. Wind and rain in the desert region of Xinjiang, northwest China. *Erdkunde* 1992;46:203–16.
- Yu HL, Chien LC, Yang CH. Asian dust storm elevates children's respiratory health risks: a spatiotemporal analysis of children's clinic visits across Taipei (Taiwan). *PLoS ONE* 2012;7(7):e41317. <http://dx.doi.org/10.1371/journal.pone.0041317>.
- Yuan CS, Sau CC, Chen MC, Huang MH, Chang SW, Lin YC, et al. Mass concentration and size-resolved chemical composition of atmospheric aerosols sampled at the Pescadores Islands during Asian dust storm periods in the years of 2001 and 2002. *Terr Atmos Ocean Sci* 2004;15:857–79.
- Zaady E, Offer ZY, Shachak M. The content and contribution of deposited aeolian organic matter in a dry land ecosystem of the Negev Desert, Israel. *Atmos Environ* 2001;35:769–76.
- Zender CS, Talamantes J. Climate controls on valley fever incidence in Kern County, California. *Int J Biometeorol* 2006;50:174–82.
- Zhao Q, He K, Rahn KA, Ma Y, Yang F, Duna F, et al. Dust storms come to Central and Southwestern China, too: implications from a major dust event in Chongqing. *Atmos Chem Phys* 2010;10:2615–30.
- Zhu A, Ramanathan V, Li F, Kim D. Dust plumes over the Pacific, Indian, and Atlantic Oceans: climatology and radiative impact. *J Geophys Res* 2007;112:D16208. <http://dx.doi.org/10.1029/2007JD008427>.
- Zhu C, Wang B, Qian W. Why do dust storms decrease in northern China concurrently with the recent global warming? *Geophys Res Lett* 2008;35:L18702. <http://dx.doi.org/10.1029/2008GL034886>.