

A 10-day heatwave at flowering superimposed on climate change conditions strongly affects production of 22 barley accessions

Ingvordsen, Cathrine Heinz; Lyngkjær, Michael F.; Peltonen-Sainio, Pirjo; Mikkelsen, Teis Nørgaard; Stockmarr, Anders; Bagger Jørgensen, Rikke

Publication date:
2015

[Link to publication](#)

Citation (APA):

Ingvordsen, C. H., Lyngkjær, M. F., Peltonen-Sainio, P., Mikkelsen, T. N., Stockmarr, A., & Bagger Jørgensen, R. (2015). A 10-day heatwave at flowering superimposed on climate change conditions strongly affects production of 22 barley accessions. Poster session presented at Agriculture and Climate Change: Adapting Crops to Increased Uncertainty, Amsterdam, Netherlands.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A 10-days heatwave at flowering superimposed on climate change conditions strongly affects production of 22 barley accessions

CH Ingvordsen¹, MF Lyngkjær², P Peltonen-Sainio³, TN. Mikkelsen¹, A Stockmarr and RB Jørgensen¹

¹Technical University of Denmark, Denmark. ²University of Copenhagen, Denmark. ³MTT Agrifood Research Finland, Finland

INTRODUCTION Extreme climate events are projected to be among the future most challenging constraints to plant development. Heatwaves as well as floods and droughts cause acute changes in the growth environment determining our primary production (Collins et al., 2013). Europe experienced extreme heatwaves in 2003 and 2006. In 2003, a 21 % decrease in the French wheat production was found from temperatures up to 6 °C above long-term means and precipitation being less than 50 % of the average (Ciais et al., 2005). One strategy to mitigate the this decrease from heatwaves is to identify resilient cultivars and incorporate them in breeding programs.

EXPERIMENT Four basic treatments with projected levels of temperature and [CO₂] as might be expected at the end of this century (~RCP8.5; IPCC, 2013) was applied as single factors or in combination throughout the growing period (Fig. 1). At the time of flowering a 10 day-heatwave of 33/28 °C (day/night) was superimposed on the treatments (Fig. 1).

At maturity plants were harvested and threshed individually and grain, vegetative biomass, harvest index (HI) and grain yield per ear were determined.

RESULTS The applied heatwave decreased grain yield with the strongest decrease in the treatment with elevated [CO₂] (43 %; Fig. 2). However, vast variation existed between accessions (Fig. 3). In the treatment of elevated temperature and [CO₂] combined, the superimposed heatwave decreased grain yield of accessions 2-80 %. The heatwave further changed allocation by increasing aboveground vegetative biomass. The most likely future climate scenario of elevated temperature and [CO₂] in combination and with the superimposed heatwave was found to decrease overall grain yield by 52 % compared to present day conditions.

CONCLUSION The identified variation in the set of barley accessions in response to multifactor climate treatments was high as such, however, the ability to cope with heatwaves and avoid severe yield losses of less than 29 % was not identified in the accessions included. In fact, these results emphasize the threat that temperature extremes exert to crop production systems.

Acknowledgements

COBRA (Core Organic II) and NordForsk ('Sustainable primary production in a changing climate')

References

Ciais P et al. 2005. Nature 437, 529–33.
Collins M et al. 2013. WG1, AR5, IPCC.
Bencze S et al. 2004. Cereal Res Commun 32, 75–82.

Contact

Rikke B. Jørgensen
rijq@kt.dtu.dk
Senior Scientist, ECO,
Technical University of Denmark

