# Malaria risk and temperature: Influences from global climate change and local land use practices

## Jonathan A. Patz\* and Sarah H. Olson


Center for Sustainability and the Global Environment (SAGE), The Nelson Institute and Department of Population Health Sciences, University of Wisconsin, 1710 University Avenue, Madison, WI 53726

n estimated 700,000 to 2.7 million people die of malaria each year, and 75% of those are African children (www. cdc.gov/malaria). Recent resurgence in the East African highlands involves multiple factors, from climate and land use change to drug resistance, variable disease control efforts, and other sociodemographic factors (1). But malaria is an extremely climate-sensitive tropical disease, making the assessment of potential change in risk due to past and projected warming trends one of the most important climate change/health questions to resolve. Pascual *et al.* (2) now provide important new insights toward answering this malaria/climate question in their article in this issue of PNAS.

#### **Warming Trends and Malaria**

Pascual's research team used a detrended time series of temperature and documented a warming trend in the East African highlands from 1950 to 2002, concomitant with increases in malaria incidence. Moreover, their findings confirm the importance of the well recognized nonlinear and threshold responses of malaria (a biological system) to the effect of regional temperature change (Fig. 1). For example, showing that the biological response of mosquito populations to warming can be more than an order of magnitude larger than the measured change in temperature represents a stunning finding, critical in advancing risk assessment of climate change impacts. Those who argue that we need not worry about small shifts in temperature should pause after considering the findings of Pascual et al. (2) that a mere half-degree centigrade increase in temperature trend can translate into a 30-100% increase in mosquito abundance, in other words "biological amplification" of temperature effects. In the African highlands, where mosquito populations are relatively low compared with lowland areas (3), such biological responses may be especially significant to determining the risk of malaria.

As a general rule, temperatures decrease on average by 6°C for every 1,000 m of elevation gained. Referring to Fig. 1, minimum temperature for parasite



**Fig. 1.** Relationship between temperature and malaria parasite development time inside the mosquito ("extrinsic incubation period" or EIP). EIP shortens at higher temperatures, so mosquitoes become infectious sooner. Note the nonlinear response to temperature, as well as relative threshold limits for malaria parasite development (≈18°C and 15°C for *P. falciparum*, and *P. vivax*, respectively) (adapted from ref. 16).

development of Plasmodium falciparum and Plasmodium vivax approximates 18°C and 15°C, respectively, limiting the spread of malaria at higher altitudes. There is also a relationship between increasing altitude and decreasing mosquito abundance in African highlands (4). Projecting into the future, Ebi et al. (5) have compared climate suitability maps for malaria in the topographically diverse country of Zimbabwe and found that the projected warming from global climate models would make the country's entire highland area climatologically more favorable to support malaria by year 2050 (Fig. 2).

Large epidemics of malaria elsewhere have been associated with climate and temperature anomalies, such as in the Indian subcontinent (6), Colombia (7), and Uganda (8). Thomson *et al.* (9) recently showed that, in Botswana, indices of El Niño-related climate variability can serve as the basis of malaria risk prediction and early warning.

# Malaria and Local Effects on Climate from Land Use Change

Changing landscapes can significantly affect local weather more acutely than long-term climate change. Land cover change can influence microclimatic conditions, including temperature, evapotranspiration, and surface runoff (10,

11), all key to determining mosquito abundance and survivorship. In Kenya, Yan and colleagues (12) have found that open, treeless habitats experience warmer midday temperatures than forested habitats and also affect indoor hut temperatures. As a result, the gonotrophic cycle of female Anopheles gambiae was found to be shortened by 2.6 days (52%) and 2.9 days (21%) during the dry and rainy seasons, respectively, compared with forested sites. Similar findings have been documented in Uganda where higher temperatures have been measured in communities bordering cultivated fields compared with those adjacent to natural wetlands, and the number of A. gambiae s.l. per house increased along with minimum temperatures after adjustment for potential confounding variables (13).

Higher maximum and mean temperatures of aquatic breeding sites found in farmlands also hasten larval development and pupation rates (14). Increased canopy cover in western Kenya is negatively associated with the presence of A.

Conflict of interest statement: No conflicts declared. See companion article on page 5829.

<sup>\*</sup>To whom correspondence should be addressed. E-mail: patz@wisc.edu.

<sup>© 2006</sup> by The National Academy of Sciences of the USA




Fig. 2. Climate suitability for stable malaria transmission across the diverse topography of Zimbabwe, determined by fuzzy logic analysis and based on United  $Kingdom\ Meteorological\ Office\ (UKMO)\ global\ climate\ scenarios\ for\ year\ 2000\ (\textit{Left})\ and\ with\ warming\ to\ the\ year\ 2050\ (\textit{Right}).\ Orange-red\ colors\ indicate\ more$ suitable climate conditions for malaria transmission. [Adapted from Ebi et al. (5) and Hartman et al. (17); maps provided by K. Ebi.]

gambiae complex and Anopheles funestus larvae in natural aquatic habitats (3). In artificial pools, survivorship of A. gambiae larvae in sunlit open areas was 50 times the survivorship in forested areas and also related to assemblages of predatory species (15). In short, deforestation and cultivation of natural swamps in the African highlands create conditions favorable for the survival of A. gambiae larvae, making analysis of land use change on local climate, habitat, and biodiversity central to malaria risk assessments.

### Conclusions

Studies of the association of malaria and past climate in the African highlands

- 1. Patz, J. A., Hulme, M., Rosenzweig, C., Mitchell, T. D., Goldberg, R. A., Githeko, A. K., Lele, S., McMichael, A. J. & Le Sueur, D. (2002) Nature 420, 627-628.
- 2. Pascual, M., Ahumada, J. A., Chaves, L. F., Rodó, X. & Bouma, M. (2006) Proc. Natl. Acad. Sci. USA 103, 5829-5834.
- 3. Minakawa, N., Sonye, G., Mogi, M., Githeko, A. & Yan, G. (2002) J. Med. Entomol. 39, 833-841.
- 4. Bodker, R., Akida, J., Shayo, D., Kisinza, W., Msangeni, H. A., Pedersen, E. M. & Lindsay, S. W. (2003) J. Med. Entomol. 40, 706-717.
- 5. Ebi, K. L., Hartman, J., Chan, N., McConnell, K. J., Schlesinger, M. & Weyant, J. (2005) Clim. Change 73, 375-393.
- 6. Bouma, M. J. & van der Kaay, H. J. (1996) Trop. Med. Int. Health 1, 86-96.

have been controversial in part due to the varying quality of long-term disease data across sites in Africa, as well as the difficulty in adequately controlling for sociodemographic and biological (drug resistance) data. Not only have Pascual and her team (2) improved upon temperature trend analysis in the region, but, by inclusion of an entomological model, they have enhanced the predictability of how changing temperature trends may alter malaria risk. Accurate risk prediction, of course, still requires improved monitoring and modeling of mosquito populations and human malaria epidemiology, as well as effects on the parasite and disease itself.

The relentless expansion of malaria in the highlands is exacerbated by rapid

- 7. Poveda, G., Rojas, W., Quinones, M. L., Velez, I. D., Mantilla, R. I., Ruiz, D., Zuluaga, J. S. & Rua, G. L. (2001) Environ. Health Perspect. 109, 489-493
- 8. Lindblade, K. A., Walker, E. D., Onapa, A. W., Katungu, J. & Wilson, M. L. (1999) Trans. R. Soc. Trop. Med. Hyg. 93, 480-487.
- 9. Thomson, M. C., Doblas-Reyes, F. J., Mason, S. J., Hagedorn, R., Connor, S. J., Phindela, T., Morse, A. P. & Palmer, T. N. (2006) Nature 439, 576-579
- 10. Foley, J. A., Defries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T., Daily, G. C., Gibbs, H. K., et al. (2005) Science 309, 570-574.
- 11. Patz, J. A., Campbell-Lendrum, D., Holloway, T. & Foley, J. A. (2005) Nature 438, 310-317.

population growth and massive land use changes (such as deforestation) that can favor mosquito breeding. Combined with poor access to effective health care and inefficient vector control measures, these challenges demand collaboration across ministries of health, environment, and finance/development at the local, national, and regional levels to best address malaria in the African highlands. At the international level, industrialized nations must confront the deleterious health effects that their greenhouse gas emissions (causing global warming) are having around the world.

We thank Dr. Kris Ebi (Exponent Health Sciences Group, Washington, DC) for providing Fig. 2.

- 12. Afrane, Y. A., Lawson, B. W., Githeko, A. K. & Yan, G. (2005) J. Med. Entomol. 42, 974-
- 13. Lindblade, K. A., Walker, E. D., Onapa, A. W., Katungu, J. & Wilson, M. L. (2000) Trop. Med. Int. Health 5, 263-274.
- 14. Munga, S., Minakawa, N., Zhou, G., Mushinzimana, E., Barrack, O. O., Githeko, A. K. & Yan, G. (2006) Am. J. Trop. Med. Hyg. 74,
- 15. Tuno, N., Okeka, W., Minakawa, N., Takagi, M. & Yan, G. (2005) J. Med. Entomol. 42, 270-277.
- 16. Gilles, H. M. (1999) Epidemiology (Oxford Univ. Press, Oxford).
- Hartman, J., Ebi, K., McConnell, K. J., Chan, N. & Weyant, J. (2002) Glob. Change Hum. Health 3, 42-54.