

# A general non-Markovian master equation for time-dependent Hamiltonians with coupling that is weak, strong, or anything in between

Nike Dattani\*

Harvard-Smithsonian Center for Astrophysics

Camilo Chaparro Sogamoso<sup>†</sup>

National University of Colombia

(Dated: 19th September 2018)

We start with a time-dependent Hamiltonian of the form:

$$H(t) = H_S(t) + H_I + H_B, \quad (1)$$

$$H_S(t) = \varepsilon_0(t) |0\rangle\langle 0| + \varepsilon_1(t) |1\rangle\langle 1| + V_{10}(t) |1\rangle\langle 0| + V_{01}(t) |0\rangle\langle 1|, \quad (2)$$

$$H_I = |0\rangle\langle 0| \sum_{\mathbf{k}} \left( g_{0\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{0\mathbf{k}}^* b_{\mathbf{k}} \right) + |1\rangle\langle 1| \sum_{\mathbf{k}} \left( g_{1\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{1\mathbf{k}}^* b_{\mathbf{k}} \right), \quad (3)$$

$$H_B = \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}. \quad (4)$$

For the states  $|0\rangle, |1\rangle$  we have the orthonormal condition:

$$\langle i|j\rangle = \delta_{ij}. \quad (5)$$

## I. UNITARY TRANSFORMATION INTO THE VARIATIONALLY OPTIMIZABLE FRAME

We will apply to  $H(t)$ , the unitary transformation defined by  $e^{\pm V(t)}$  where is the variationally optimizable anti-Hermitian operator:

$$V(t) \equiv |0\rangle\langle 0| \sum_{\mathbf{k}} \left( \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right) + |1\rangle\langle 1| \sum_{\mathbf{k}} \left( \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right). \quad (6)$$

in terms of the variational scalar parameters  $v_{i\mathbf{k}}(t)$  defined as:

$$v_{i\mathbf{k}}(t) = \omega_{\mathbf{k}} \alpha_{i\mathbf{k}}(t). \quad (7)$$

which will soon be optimized in order to give the most accurate possible master equation for the system's dynamics in the presence of this bath. We define the following notation for the function (6):

$$\hat{\varphi}_i(t) \equiv \sum_{\mathbf{k}} \left( \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right), \quad (8)$$

$$V(t) = |0\rangle\langle 0| \hat{\varphi}_0(t) + |1\rangle\langle 1| \hat{\varphi}_1(t). \quad (9)$$

Here  $*$  denotes the complex conjugate. Expanding  $e^{\pm V(t)}$  using the notation (6) will give us the following result:

$$e^{\pm V(t)} = e^{\pm(|0\rangle\langle 0|\hat{\varphi}_0(t) + |1\rangle\langle 1|\hat{\varphi}_1(t))} \quad (10)$$

$$= \mathbb{I} \pm (|0\rangle\langle 0|\hat{\varphi}_0(t) + |1\rangle\langle 1|\hat{\varphi}_1(t)) + \frac{(\pm(|0\rangle\langle 0|\hat{\varphi}_0(t) + |1\rangle\langle 1|\hat{\varphi}_1(t)))^2}{2!} + \dots \quad (11)$$

$$= |0\rangle\langle 0| + |1\rangle\langle 1| \pm (|0\rangle\langle 0|\hat{\varphi}_0(t) + |1\rangle\langle 1|\hat{\varphi}_1(t)) + \frac{|0\rangle\langle 0|\hat{\varphi}_0^2(t)}{2!} + \frac{|1\rangle\langle 1|\hat{\varphi}_1^2(t)}{2!} + \dots \quad (12)$$

$$= |0\rangle\langle 0| \left( \mathbb{I} \pm \hat{\varphi}_0(t) + \frac{\hat{\varphi}_0^2(t)}{2!} \pm \dots \right) + |1\rangle\langle 1| \left( \mathbb{I} \pm \hat{\varphi}_1(t) + \frac{\hat{\varphi}_1^2(t)}{2!} \pm \dots \right) \quad (13)$$

$$= |0\rangle\langle 0| e^{\pm \hat{\varphi}_0(t)} + |1\rangle\langle 1| e^{\pm \hat{\varphi}_1(t)} \quad (14)$$

$$= |0\rangle\langle 0| e^{\pm \sum_{\mathbf{k}} (\alpha_{0\mathbf{k}}(t) b_{\mathbf{k}}^\dagger - \alpha_{0\mathbf{k}}^*(t) b_{\mathbf{k}})} + |1\rangle\langle 1| e^{\pm \sum_{\mathbf{k}} (\alpha_{1\mathbf{k}}(t) b_{\mathbf{k}}^\dagger - \alpha_{1\mathbf{k}}^*(t) b_{\mathbf{k}})} \quad (15)$$

$$= |0\rangle\langle 0| B_0^\pm(t) + |1\rangle\langle 1| B_1^\pm(t), \quad (16)$$

$$B_i^\pm(t) \equiv e^{\pm \sum_{\mathbf{k}} \left( \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right)}. \quad (17)$$

Let's recall the Zassenhaus formula:

$$e^{r(X+Y)} = e^{rX} e^{rY} e^{-\frac{r^2}{2}[X,Y]} e^{\frac{r^3}{6}(2[Y,[X,Y]] + [X,[X,Y]])} e^{-\frac{r^4}{24}([[[X,Y],X],X] + 3[[[X,Y],X],Y] + 3[[[X,Y],Y],Y])} \dots \quad (18)$$

Since  $\left[ \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}, \frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}^\dagger - \frac{v_{j\mathbf{k}'}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'} \right] = 0$  for all  $\mathbf{k}', \mathbf{k}$  and  $i, j$  we can show making  $r = 1$  in (18) the following result:

$$e^{\left( \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right) + \left( \frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}^\dagger - \frac{v_{j\mathbf{k}'}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'} \right)} = e^{\frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}} e^{\frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}^\dagger - \frac{v_{j\mathbf{k}'}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}} e^{-\frac{1}{2} \left[ \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}, \frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}^\dagger - \frac{v_{j\mathbf{k}'}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'} \right]} \dots \quad (19)$$

$$= e^{\frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}} e^{\frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}^\dagger - \frac{v_{j\mathbf{k}'}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}} e^{-\frac{1}{2} 0} \dots \quad (20)$$

$$= e^{\frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}} e^{\frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}^\dagger - \frac{v_{j\mathbf{k}'}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}'}}. \quad (21)$$

By induction of this result we can write an expression of  $B_i^\pm(t)$  (shown in equation (17)) as a product of exponentials, which we will call “displacement” operators  $D(\pm v_{i\mathbf{k}}(t))$ :

$$D(\pm v_{i\mathbf{k}}(t)) \equiv e^{\pm \left( \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right)}, \quad (22)$$

$$B_i^\pm(t) = \prod_{\mathbf{k}} D\left(\pm \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right). \quad (23)$$

this will help us to write operators  $O(t)$  transformed in the variational frame as:

$$\overline{O(t)} \equiv e^{V(t)} O(t) e^{-V(t)}. \quad (24)$$

We will use the following identities:

$$\overline{|0\rangle\langle 0|}(t) = e^{V(t)}|0\rangle\langle 0|e^{-V(t)} \quad (25)$$

$$= (|0\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|B_1^+(t))|0\rangle\langle 0|(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (26)$$

$$= (|0\rangle\langle 0|0\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|0\rangle\langle 0|B_1^+(t))(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (27)$$

$$= |0\rangle\langle 0|B_0^+(t)(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (28)$$

$$= |0\rangle\langle 0|0\rangle\langle 0|B_0^+(t)B_0^-(t) + |0\rangle\langle 0|1\rangle\langle 1|B_0^+(t)B_1^-(t) \quad (29)$$

$$= |0\rangle\langle 0|, \quad (30)$$

$$\overline{|1\rangle\langle 1|}(t) = (|0\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|B_1^+(t))|1\rangle\langle 1|(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (31)$$

$$= (|0\rangle\langle 0|1\rangle\langle 1|B_0^+(t) + |1\rangle\langle 1|1\rangle\langle 1|B_1^+(t))(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (32)$$

$$= |1\rangle\langle 1|B_1^+(t)(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (33)$$

$$= |1\rangle\langle 1|0\rangle\langle 0|B_1^+(t)B_0^-(t) + B_1^+(t)|1\rangle\langle 1|1\rangle\langle 1|B_1^-(t) \quad (34)$$

$$= B_1^+(t)|1\rangle\langle 1|1\rangle\langle 1|B_1^-(t) \quad (35)$$

$$= |1\rangle\langle 1|, \quad (36)$$

$$\overline{|0\rangle\langle 1|}(t) = e^{V(t)}|0\rangle\langle 1|e^{-V(t)} \quad (37)$$

$$= (|0\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|B_1^+(t))|0\rangle\langle 1|(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (38)$$

$$= (|0\rangle\langle 0|0\rangle\langle 1|B_0^+(t) + |1\rangle\langle 1|B_1^+(t)|0\rangle\langle 1|)(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (39)$$

$$= (|0\rangle\langle 0|0\rangle\langle 1|B_0^+(t) + |1\rangle\langle 1|0\rangle\langle 1|B_1^+(t))(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (40)$$

$$= |0\rangle\langle 1|B_0^+(t)(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (41)$$

$$= |0\rangle\langle 1|0\rangle\langle 0|B_0^+(t)B_0^-(t) + |0\rangle\langle 1|1\rangle\langle 1|B_0^+(t)B_1^-(t) \quad (42)$$

$$= |0\rangle\langle 1|B_0^+(t)B_1^-(t), \quad (43)$$

$$\overline{|1\rangle\langle 0|}(t) = e^{V(t)}|1\rangle\langle 0|e^{-V(t)} \quad (44)$$

$$= (|0\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|B_1^+(t))|1\rangle\langle 0|(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (45)$$

$$= (|0\rangle\langle 0|1\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|B_1^+(t)|1\rangle\langle 0|)(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (46)$$

$$= (|0\rangle\langle 0|1\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|1\rangle\langle 0|B_1^+(t))(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (47)$$

$$= |1\rangle\langle 0|0\rangle\langle 0|B_1^+(t)B_0^-(t) + |1\rangle\langle 0|1\rangle\langle 1|B_1^+(t)B_1^-(t) \quad (48)$$

$$= |1\rangle\langle 0|B_1^+(t)B_0^-(t), \quad (49)$$

$$\overline{b_{\mathbf{k}}}(t) = e^{V(t)}b_{\mathbf{k}}e^{-V(t)} \quad (50)$$

$$= (|0\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|B_1^+(t))b_{\mathbf{k}}(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (51)$$

$$= |0\rangle\langle 0|B_0^+(t)b_{\mathbf{k}}B_0^-(t)|0\rangle\langle 0| + |0\rangle\langle 0|B_0^+(t)b_{\mathbf{k}}|1\rangle\langle 1|B_1^-(t) + |1\rangle\langle 1|B_1^+(t)b_{\mathbf{k}}|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^+(t)b_{\mathbf{k}}B_1^-(t)|1\rangle\langle 1| \quad (52)$$

$$= |0\rangle\langle 0|0\rangle\langle 0|B_0^+(t)b_{\mathbf{k}}B_0^-(t) + |0\rangle\langle 0|1\rangle\langle 1|B_0^+(t)b_{\mathbf{k}}B_1^-(t) + |1\rangle\langle 1|0\rangle\langle 0|B_1^+(t)b_{\mathbf{k}}B_0^-(t) + |1\rangle\langle 1|B_1^+(t)b_{\mathbf{k}}B_1^-(t) \quad (53)$$

$$= |0\rangle\langle 0|\left(b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}}{\omega_{\mathbf{k}}}\right) + |1\rangle\langle 1|\left(b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}}{\omega_{\mathbf{k}}}\right) \quad (54)$$

$$= (|0\rangle\langle 0| + |1\rangle\langle 1|)b_{\mathbf{k}} - |1\rangle\langle 1|\frac{v_{1\mathbf{k}}}{\omega_{\mathbf{k}}} - |0\rangle\langle 0|\frac{v_{0\mathbf{k}}}{\omega_{\mathbf{k}}} \quad (55)$$

$$= b_{\mathbf{k}} - |1\rangle\langle 1|\frac{v_{1\mathbf{k}}}{\omega_{\mathbf{k}}} - |0\rangle\langle 0|\frac{v_{0\mathbf{k}}}{\omega_{\mathbf{k}}}, \quad (56)$$

$$\overline{b_{\mathbf{k}}(t)}^\dagger = e^{V(t)}b_{\mathbf{k}}^\dagger e^{-V(t)} \quad (57)$$

$$= (|0\rangle\langle 0|B_0^+(t) + |1\rangle\langle 1|B_1^+(t))b_{\mathbf{k}}^\dagger(|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^-(t)) \quad (58)$$

$$= |0\rangle\langle 0|B_0^+(t)b_{\mathbf{k}}^\dagger B_0^-(t)|0\rangle\langle 0| + |0\rangle\langle 0|B_0^+(t)b_{\mathbf{k}}^\dagger|1\rangle\langle 1|B_1^-(t) + |1\rangle\langle 1|B_1^+(t)b_{\mathbf{k}}^\dagger|0\rangle\langle 0|B_0^-(t) + |1\rangle\langle 1|B_1^+(t)b_{\mathbf{k}}^\dagger B_1^-(t)|1\rangle\langle 1| \quad (59)$$

$$= |0\rangle\langle 0|0\rangle\langle 0|B_0^+(t)b_{\mathbf{k}}^\dagger B_0^-(t) + |0\rangle\langle 0|1\rangle\langle 1|B_0^+(t)b_{\mathbf{k}}^\dagger B_1^-(t) + |1\rangle\langle 1|0\rangle\langle 0|B_1^+(t)b_{\mathbf{k}}^\dagger B_0^-(t) + |1\rangle\langle 1|1\rangle\langle 1|B_1^+(t)b_{\mathbf{k}}^\dagger B_1^-(t) \quad (60)$$

$$= |0\rangle\langle 0|\left(b_{\mathbf{k}}^\dagger - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}}\right) + |1\rangle\langle 1|\left(b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}}\right) \quad (61)$$

$$= b_{\mathbf{k}}^\dagger - |1\rangle\langle 1|\frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} - |0\rangle\langle 0|\frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}}. \quad (62)$$

We have used the following results as well to obtain the transformed  $b_{\mathbf{k}}$  and  $b_{\mathbf{k}}^\dagger$ :

$$B_i^+(t) b_{\mathbf{k}} B_i^-(t) = b_{\mathbf{k}} - \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}}, \quad (63)$$

$$B_i^+(t) b_{\mathbf{k}}^\dagger B_i^-(t) = b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}}. \quad (64)$$

We therefore have the following relationships:

$$\overline{\varepsilon_0(t) |0\rangle\langle 0| (t)} = \varepsilon_0(t) |0\rangle\langle 0|, \quad (65)$$

$$\overline{\varepsilon_1(t) |1\rangle\langle 1| (t)} = \varepsilon_1(t) |1\rangle\langle 1|, \quad (66)$$

$$\overline{V_{10}(t) |1\rangle\langle 0| (t)} = V_{10}(t) |1\rangle\langle 0| B_1^+(t) B_0^-(t), \quad (67)$$

$$\overline{V_{01}(t) |0\rangle\langle 1| (t)} = V_{01}(t) |0\rangle\langle 1| B_0^+(t) B_1^-(t), \quad (68)$$

$$\overline{(g_{i\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{i\mathbf{k}}^* b_{\mathbf{k}})(t)} = g_{i\mathbf{k}} \left( |0\rangle\langle 0| \left( b_{\mathbf{k}}^\dagger - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) + |1\rangle\langle 1| \left( b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) \right) + g_{i\mathbf{k}}^* \left( |0\rangle\langle 0| \left( b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) + |1\rangle\langle 1| \left( b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right) \quad (69)$$

$$= g_{i\mathbf{k}} \left( (|0\rangle\langle 0| + |1\rangle\langle 1|) b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} |1\rangle\langle 1| - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} |0\rangle\langle 0| \right) + g_{i\mathbf{k}}^* \left( (|0\rangle\langle 0| + |1\rangle\langle 1|) b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} |1\rangle\langle 1| - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} |0\rangle\langle 0| \right) \quad (70)$$

$$= g_{i\mathbf{k}} \left( (|0\rangle\langle 0| + |1\rangle\langle 1|) b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} |1\rangle\langle 1| - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} |0\rangle\langle 0| \right) + g_{i\mathbf{k}}^* \left( (|0\rangle\langle 0| + |1\rangle\langle 1|) b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} |1\rangle\langle 1| - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} |0\rangle\langle 0| \right) \quad (71)$$

$$= g_{i\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{i\mathbf{k}}^* b_{\mathbf{k}} - g_{i\mathbf{k}} \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} |0\rangle\langle 0| - g_{i\mathbf{k}}^* \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} |0\rangle\langle 0| - g_{i\mathbf{k}} \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} |1\rangle\langle 1| - g_{i\mathbf{k}}^* \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} |1\rangle\langle 1| \quad (72)$$

$$= g_{i\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{i\mathbf{k}}^* b_{\mathbf{k}} - \left( g_{i\mathbf{k}} \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} + g_{i\mathbf{k}}^* \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) |0\rangle\langle 0| - \left( g_{i\mathbf{k}} \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} + g_{i\mathbf{k}}^* \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) |1\rangle\langle 1|, \quad (73)$$

$$\overline{|0\rangle\langle 0| (g_{0\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{0\mathbf{k}}^* b_{\mathbf{k}})(t)} = (|0\rangle\langle 0| B_0^+(t) + |1\rangle\langle 1| B_1^+(t)) |0\rangle\langle 0| (g_{0\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{0\mathbf{k}}^* b_{\mathbf{k}}) (|0\rangle\langle 0| B_0^-(t) + |1\rangle\langle 1| B_1^-(t)) \quad (74)$$

$$= |0\rangle\langle 0| B_0^+(t) |0\rangle\langle 0| (g_{0\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{0\mathbf{k}}^* b_{\mathbf{k}}) |0\rangle\langle 0| B_0^-(t) \quad (75)$$

$$= |0\rangle\langle 0| B_0^+(t) (g_{0\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{0\mathbf{k}}^* b_{\mathbf{k}}) B_0^-(t) \quad (76)$$

$$= |0\rangle\langle 0| \left( g_{0\mathbf{k}} \left( b_{\mathbf{k}}^\dagger - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) + g_{0\mathbf{k}}^* \left( b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right), \quad (77)$$

$$\overline{|1\rangle\langle 1| (g_{1\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{1\mathbf{k}}^* b_{\mathbf{k}})(t)} = (|0\rangle\langle 0| B_0^+(t) + |1\rangle\langle 1| B_1^+(t)) |1\rangle\langle 1| (g_{1\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{1\mathbf{k}}^* b_{\mathbf{k}}) (|0\rangle\langle 0| B_0^-(t) + |1\rangle\langle 1| B_1^-(t)) \quad (78)$$

$$= |1\rangle\langle 1| B_1^+(t) |1\rangle\langle 1| (g_{1\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{1\mathbf{k}}^* b_{\mathbf{k}}) |1\rangle\langle 1| B_1^-(t) \quad (79)$$

$$= |1\rangle\langle 1| B_1^+(t) (g_{1\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{1\mathbf{k}}^* b_{\mathbf{k}}) B_1^-(t) \quad (80)$$

$$= |1\rangle\langle 1| \left( g_{1\mathbf{k}} \left( b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) + g_{1\mathbf{k}}^* \left( b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right), \quad (81)$$

$$\overline{\omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}(t)} = \omega_{\mathbf{k}} (|0\rangle\langle 0| B_0^+(t) + |1\rangle\langle 1| B_1^+(t)) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} (|0\rangle\langle 0| B_0^-(t) + |1\rangle\langle 1| B_1^-(t)) \quad (82)$$

$$= \omega_{\mathbf{k}} (|0\rangle\langle 0| \Pi_{\mathbf{k}'} D\left(\frac{v_{0\mathbf{k}'}(t)}{\omega_{\mathbf{k}'}}\right) + |1\rangle\langle 1| \Pi_{\mathbf{k}'} D\left(\frac{v_{1\mathbf{k}'}(t)}{\omega_{\mathbf{k}'}}\right)) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} (|0\rangle\langle 0| \Pi_{\mathbf{k}'} D\left(-\frac{v_{0\mathbf{k}'}(t)}{\omega_{\mathbf{k}'}}\right) + |1\rangle\langle 1| \Pi_{\mathbf{k}'} D\left(-\frac{v_{1\mathbf{k}'}(t)}{\omega_{\mathbf{k}'}}\right)) \quad (83)$$

$$= \omega_{\mathbf{k}} (|0\rangle\langle 0| B_0^+(t) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} B_0^-(t) + |1\rangle\langle 1| B_1^+(t) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} B_1^-(t)) \quad (84)$$

$$= \omega_{\mathbf{k}} \sum_j |j\rangle\langle j| D\left(\frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} D\left(-\frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) \prod_{\mathbf{k}' \neq \mathbf{k}} \left( D\left(\frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}'}}\right) D\left(-\frac{v_{j\mathbf{k}'}(t)}{\omega_{\mathbf{k}'}}\right) \right) \quad (85)$$

$$= \omega_{\mathbf{k}} \left( |0\rangle\langle 0| D\left(\frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} D\left(-\frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) \mathbb{I} + |1\rangle\langle 1| D\left(\frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} D\left(-\frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) \mathbb{I} \right) \quad (86)$$

$$= \omega_{\mathbf{k}} \left( |0\rangle\langle 0| \left( b_{\mathbf{k}}^\dagger - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) \left( b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) + |1\rangle\langle 1| \left( b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) \left( b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right) \quad (87)$$

$$= \omega_{\mathbf{k}} \left( |0\rangle\langle 0| \left( b_{\mathbf{k}}^\dagger b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger + \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \right) + |1\rangle\langle 1| \left( b_{\mathbf{k}}^\dagger b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger + \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \right) \right) \quad (88)$$

$$= \omega_{\mathbf{k}} \left( |0\rangle\langle 0| b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |1\rangle\langle 1| b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |1\rangle\langle 1| \left( \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger \right) + |0\rangle\langle 0| \left( \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger \right) \right) \quad (89)$$

$$= \omega_{\mathbf{k}} \left( b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |1\rangle\langle 1| \left( \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger \right) + |0\rangle\langle 0| \left( \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger \right) \right) \quad (90)$$

$$= \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \omega_{\mathbf{k}} \left( |1\rangle\langle 1| \left( \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger \right) + |0\rangle\langle 0| \left( \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger \right) \right) \quad (91)$$

$$= \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |1\rangle\langle 1| \left( \frac{|v_{1\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - v_{1\mathbf{k}}^*(t) b_{\mathbf{k}} - v_{1\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) + |0\rangle\langle 0| \left( \frac{|v_{0\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - v_{0\mathbf{k}}^*(t) b_{\mathbf{k}} - v_{0\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) \quad (92)$$

$$= \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |1\rangle\langle 1| \left( \frac{|v_{1\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( v_{1\mathbf{k}}^*(t) b_{\mathbf{k}} + v_{1\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) \right) + |0\rangle\langle 0| \left( \frac{|v_{0\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( v_{0\mathbf{k}}^*(t) b_{\mathbf{k}} + v_{0\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) \right). \quad (93)$$

So all parts of  $H(t)$  can be written in the variationally optimizable frame now:

$$\overline{H_S(t)} = \overline{\varepsilon_0(t)} |0\rangle\langle 0| + \overline{\varepsilon_1(t)} |1\rangle\langle 1| + \overline{V_{10}(t)} |1\rangle\langle 0| + \overline{V_{01}(t)} |0\rangle\langle 1| \quad (94)$$

$$= \varepsilon_0(t) |0\rangle\langle 0| + \varepsilon_1(t) |1\rangle\langle 1| + V_{10}(t) |1\rangle\langle 0| B_1^+(t) B_0^-(t) + V_{01}(t) |0\rangle\langle 1| B_0^+(t) B_1^-(t), \quad (95)$$

$$\overline{H_I} = \sum_{\mathbf{k}} |0\rangle\langle 0| \left( g_{0\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{0\mathbf{k}}^* b_{\mathbf{k}} \right) + \sum_{\mathbf{k}} |1\rangle\langle 1| \left( g_{1\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{1\mathbf{k}}^* b_{\mathbf{k}} \right) \quad (96)$$

$$= \sum_{\mathbf{k}} |0\rangle\langle 0| \left( g_{0\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{0\mathbf{k}}^* b_{\mathbf{k}} \right) + \sum_{\mathbf{k}} |1\rangle\langle 1| \left( g_{1\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{1\mathbf{k}}^* b_{\mathbf{k}} \right) \quad (97)$$

$$= \sum_{\mathbf{k}} |0\rangle\langle 0| \left( g_{0\mathbf{k}} \left( b_{\mathbf{k}}^\dagger - \frac{v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) + g_{0\mathbf{k}}^* \left( b_{\mathbf{k}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right) + \sum_{\mathbf{k}} |1\rangle\langle 1| \left( g_{1\mathbf{k}} \left( b_{\mathbf{k}}^\dagger - \frac{v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} \right) + g_{1\mathbf{k}}^* \left( b_{\mathbf{k}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right) \quad (98)$$

$$= \sum_{\mathbf{k}, i} |i\rangle\langle i| \left( g_{i\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{i\mathbf{k}}^* b_{\mathbf{k}} - \left( g_{i\mathbf{k}} \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} + g_{i\mathbf{k}}^* \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right), \quad (99)$$

$$\overline{H_B} = \sum_{\mathbf{k}} \overline{\omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}} \quad (100)$$

$$= \sum_{\mathbf{k}} \left( \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |1\rangle\langle 1| \left( \frac{|v_{1\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( v_{1\mathbf{k}}^*(t) b_{\mathbf{k}} + v_{1\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) \right) + |0\rangle\langle 0| \left( \frac{|v_{0\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( v_{0\mathbf{k}}^*(t) b_{\mathbf{k}} + v_{0\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) \right) \right) \quad (101)$$

$$= \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{\mathbf{k}} \left( |1\rangle\langle 1| \left( \frac{|v_{1\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( v_{1\mathbf{k}}^*(t) b_{\mathbf{k}} + v_{1\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) \right) + |0\rangle\langle 0| \left( \frac{|v_{0\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( v_{0\mathbf{k}}^*(t) b_{\mathbf{k}} + v_{0\mathbf{k}}(t) b_{\mathbf{k}}^\dagger \right) \right) \right). \quad (102)$$

Finally merging these expressions gives the transformed Hamiltonian:

$$\overline{H(t)} = \sum_j \varepsilon_j(t) |j\rangle\langle j| + \sum_{j \neq j'} V_{jj'}(t) |j\rangle\langle j'| B_j^+(t) B_{j'}^-(t) + \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{j\mathbf{k}} |j\rangle\langle j| \left( \left( g_{j\mathbf{k}} - v_{j\mathbf{k}}(t) \right) b_{\mathbf{k}}^\dagger + \left( g_{j\mathbf{k}} - v_{j\mathbf{k}}(t) \right)^* b_{\mathbf{k}} + \frac{|v_{j\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( g_{j\mathbf{k}} \frac{v_{j\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} + g_{j\mathbf{k}}^* \frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right). \quad (103)$$

Also we may write this transformed Hamiltonian as a sum of the form:

$$\overline{H(t)} = \overline{H_S}(t) + \overline{H_I}(t) + \overline{H_B}. \quad (104)$$

Let's define:

$$R_i(t) \equiv \sum_{\mathbf{k}} \left( \frac{|v_{i\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( g_{i\mathbf{k}} \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} + g_{i\mathbf{k}}^* \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right), \quad (105)$$

$$B_{iz}(t) \equiv \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger + (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* b_{\mathbf{k}} \right), \quad (106)$$

$$\chi_{ij}(t) \equiv \sum_{\mathbf{k}} \frac{1}{2} \left( \frac{v_{i\mathbf{k}}^*(t) v_{j\mathbf{k}}(t) - v_{i\mathbf{k}}(t) v_{j\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right). \quad (107)$$

$\chi_{ij}(t)$  is an imaginary number so  $e^{\chi_{ij}(t)}$  is the phase associated to  $B_{ij}(t)$  as we will show. We can summarize these definitions with other that we will proof later and use from now in the following matrix:

$$\begin{pmatrix} B_{iz}(t) & B_i^\pm(t) \\ B_x(t) & B_i(t) \\ B_y(t) & B_{ij}(t) \end{pmatrix} \equiv \begin{pmatrix} \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger + (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* b_{\mathbf{k}} \right) & e^{\pm \sum_{\mathbf{k}} \left( \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right)} \\ \frac{B_1^+(t)B_0^-(t) + B_0^+(t)B_1^-(t) - B_{10}(t) - B_{01}(t)}{2} & e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \\ \frac{B_0^+(t)B_1^-(t) - B_1^+(t)B_0^-(t) + B_{10}(t) - B_{01}(t)}{2i} & e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{i\mathbf{k}}(t) - v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} e^{\chi_{ij}(t)} \end{pmatrix}, \quad (108)$$

$$(\cdot)^{\Re} \equiv \Re(\cdot), \quad (109)$$

$$(\cdot)^{\Im} \equiv \Im(\cdot). \quad (110)$$

We reduced the lenght of the expression for the real and imaginary part as shown before. We assume that the bath is at equilibrium with inverse temperature  $\beta = \frac{1}{k_B T}$ , considering the stationary bath state as reference written in the following way:

$$\rho_B = \frac{e^{-\beta H_B}}{\text{Tr}(e^{-\beta H_B})}. \quad (111)$$

We can show using the coherence representation of the creation and annihilation operators that:

$$b^\dagger = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & \dots \\ \sqrt{1} & 0 & 0 & \dots & 0 & \dots \\ 0 & \sqrt{2} & 0 & \dots & 0 & \dots \\ 0 & 0 & \sqrt{3} & \dots & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots & \vdots & \dots \\ 0 & 0 & 0 & \dots & \sqrt{n} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad (112)$$

$$b = \begin{pmatrix} 0 & \sqrt{1} & 0 & 0 & \dots & 0 & \dots \\ 0 & 0 & \sqrt{2} & 0 & \dots & 0 & \dots \\ 0 & 0 & 0 & \sqrt{3} & \dots & 0 & \dots \\ 0 & 0 & 0 & 0 & \ddots & \vdots & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots & \sqrt{n} & \dots \\ 0 & 0 & 0 & 0 & \dots & 0 & \ddots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}. \quad (113)$$

So the product of the matrix representation of  $b^\dagger$  and  $b$  with  $-\beta$  is:

$$-\beta\omega b^\dagger b = -\beta\omega \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & \dots \\ 0 & 1 & 0 & \dots & 0 & \dots \\ 0 & 0 & 2 & \dots & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots & \vdots & \dots \\ 0 & 0 & 0 & \dots & n & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad (114)$$

$$= \sum_{j=0}^{\infty} -j\beta\omega |j\rangle\langle j|, \quad (115)$$

So the density matrix  $\rho_B$  written in the coherence representation can be obtained using the Zassenhaus formula and the fact that  $[|j\rangle\langle j|, |i\rangle\langle i|] = 0$  for all  $i, j$ .

$$e^{-\beta\omega_{\mathbf{k}}b_{\mathbf{k}}^{\dagger}b_{\mathbf{k}}} = \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}}\beta\omega_{\mathbf{k}}}|j_{\mathbf{k}}\rangle\langle j_{\mathbf{k}}|, \quad (116)$$

$$e^{-\beta\sum_{\mathbf{k}}\omega_{\mathbf{k}}b_{\mathbf{k}}^{\dagger}b_{\mathbf{k}}} = \prod_{\mathbf{k}} \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}}\beta\omega_{\mathbf{k}}}|j_{\mathbf{k}}\rangle\langle j_{\mathbf{k}}|. \quad (117)$$

The value of  $\text{Tr} \left( e^{-\beta\sum_{\mathbf{k}}\omega_{\mathbf{k}}b_{\mathbf{k}}^{\dagger}b_{\mathbf{k}}} \right)$  is:

$$\text{Tr} \left( e^{-\beta\omega_{\mathbf{k}}b_{\mathbf{k}}^{\dagger}b_{\mathbf{k}}} \right) = \text{Tr} \left( \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}}\beta\omega_{\mathbf{k}}}|j_{\mathbf{k}}\rangle\langle j_{\mathbf{k}}| \right) \quad (118)$$

$$= \sum_{j_{\mathbf{k}}} (e^{-\beta\omega_{\mathbf{k}}})^{j_{\mathbf{k}}} \quad (119)$$

$$= \frac{1}{1 - e^{-\beta\omega_{\mathbf{k}}}} \quad (\text{by geometric series}) \quad (120)$$

$$\equiv f_{\text{Bose-Einstein}}(-\beta\omega_{\mathbf{k}}), \quad (121)$$

$$\text{Tr} \left( e^{-\beta\sum_{\mathbf{k}}\omega_{\mathbf{k}}b_{\mathbf{k}}^{\dagger}b_{\mathbf{k}}} \right) = \text{Tr} \left( \prod_{\mathbf{k}} \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}}\beta\omega_{\mathbf{k}}}|j_{\mathbf{k}}\rangle\langle j_{\mathbf{k}}| \right) \quad (122)$$

$$= \prod_{\mathbf{k}} \text{Tr} \left( \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}}\beta\omega_{\mathbf{k}}}|j_{\mathbf{k}}\rangle\langle j_{\mathbf{k}}| \right) \quad (123)$$

$$= \prod_{\mathbf{k}} f_{\text{Bose-Einstein}}(-\beta\omega_{\mathbf{k}}). \quad (124)$$

So the density matrix of the bath is:

$$\rho_B = \frac{e^{-\beta H_B}}{\text{Tr}(e^{-\beta H_B})} \quad (125)$$

$$= \frac{\prod_{\mathbf{k}} \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}}\beta\omega_{\mathbf{k}}}|j_{\mathbf{k}}\rangle\langle j_{\mathbf{k}}|}{\prod_{\mathbf{k}} f_{\text{Bose-Einstein}}(-\beta\omega_{\mathbf{k}})} \quad (126)$$

$$= \prod_{\mathbf{k}} \frac{\sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}}\beta\omega_{\mathbf{k}}}|j_{\mathbf{k}}\rangle\langle j_{\mathbf{k}}|}{f_{\text{Bose-Einstein}}(-\beta\omega_{\mathbf{k}})}. \quad (127)$$

Now, given that creation and annihilation satisfy:

$$b_{\mathbf{k}} |j_{\mathbf{k}}\rangle = \sqrt{j_{\mathbf{k}}}|j_{\mathbf{k}}-1\rangle, \quad (128)$$

$$b_{\mathbf{k}}^{\dagger} |j_{\mathbf{k}}\rangle = \sqrt{j_{\mathbf{k}}+1}|j_{\mathbf{k}}+1\rangle. \quad (129)$$

Then we can prove that  $\langle B_{iz} \rangle_{\overline{H_B}} = 0$  using the following property based on (128)-(129):

$$\langle B_{iz}(t) \rangle_{\overline{H_B}} = \text{Tr} (B_{iz}(t) \rho_B) \quad (130)$$

$$= \text{Tr} \left( \left( \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger + (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* b_{\mathbf{k}} \right) \right) \rho_B \right) \quad (131)$$

$$= \sum_{\mathbf{k}} \text{Tr} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger \rho_B \right) + \sum_{\mathbf{k}} \text{Tr} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* b_{\mathbf{k}} \rho_B \right) \quad (132)$$

$$= \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \text{Tr} \left( b_{\mathbf{k}}^\dagger \rho_B \right) + \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* \text{Tr} (b_{\mathbf{k}} \rho_B) \quad (133)$$

$$= \sum_{\mathbf{k}} \text{Tr} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger \prod_{\mathbf{k}} \frac{\sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}|}{f_{\text{Bose-Einstein}}(-\beta \omega_{\mathbf{k}})} \right) + \sum_{\mathbf{k}} \text{Tr} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* b_{\mathbf{k}} \prod_{\mathbf{k}} \frac{\sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}|}{f_{\text{Bose-Einstein}}(-\beta \omega_{\mathbf{k}})} \right) \quad (134)$$

$$= \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \text{Tr} \left( b_{\mathbf{k}}^\dagger \prod_{\mathbf{k}} \frac{\sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}|}{f_{\text{Bose-Einstein}}(-\beta \omega_{\mathbf{k}})} \right) + \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* \text{Tr} \left( b_{\mathbf{k}} \prod_{\mathbf{k}} \frac{\sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}|}{f_{\text{Bose-Einstein}}(-\beta \omega_{\mathbf{k}})} \right), \quad (135)$$

$$\text{Tr} \left( b_{\mathbf{k}}^\dagger \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}| \right) = \text{Tr} \left( \left( \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} \right) b_{\mathbf{k}}^\dagger |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}| \right) \quad (\text{by cyclic permutivity of trace, move } b_{\mathbf{k}}^\dagger) \quad (136)$$

$$= \text{Tr} \left( \left( \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} \right) \sqrt{j_{\mathbf{k}} + 1} |j_{\mathbf{k}} + 1 \rangle \langle j_{\mathbf{k}}| \right) \quad (137)$$

$$= 0, \quad (138)$$

$$\text{Tr} \left( b_{\mathbf{k}} \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}| \right) = \text{Tr} \left( \left( \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} \right) b_{\mathbf{k}} |j_{\mathbf{k}} \rangle \langle j_{\mathbf{k}}| \right) \quad (\text{by cyclic permutivity of trace, move } b_{\mathbf{k}}) \quad (139)$$

$$= \text{Tr} \left( \left( \sum_{j_{\mathbf{k}}} e^{-j_{\mathbf{k}} \beta \omega_{\mathbf{k}}} \right) \sqrt{j_{\mathbf{k}}} |j_{\mathbf{k}} - 1 \rangle \langle j_{\mathbf{k}}| \right) \quad (140)$$

$$= 0. \quad (141)$$

we therefore find that:

$$\langle B_{iz}(t) \rangle_{\overline{H_B}} = 0. \quad (142)$$

Another important expected value is  $B(t) = \langle B^\pm(t) \rangle_{\overline{H_B}}$ , where  $B^\pm(t) = e^{\pm \sum_{\mathbf{k}} \left( \frac{v_{\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right)}$  is given by:

$$\langle B^\pm(t) \rangle_{H_B} = \text{Tr} (\rho_B B^\pm(t)) = \text{Tr} (B^\pm(t) \rho_B) \quad (143)$$

$$= \text{Tr} \left( e^{\pm \sum_{\mathbf{k}} \left( \frac{v_{\mathbf{k}}(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}}^\dagger - \frac{v_{\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} b_{\mathbf{k}} \right)} \rho_B \right) \quad (144)$$

$$= \prod_{\mathbf{k}} \text{Tr} (D(\pm \alpha_{\mathbf{k}}(t)) \rho_B) \quad (145)$$

$$= \prod_{\mathbf{k}} \langle D(\pm \alpha_{\mathbf{k}}(t)) \rangle. \quad (146)$$

Given that we can write a density operator as:

$$\rho = \int P(\alpha) |\alpha \rangle \langle \alpha| d^2 \alpha. \quad (147)$$

where  $P(\alpha)$  satisfies  $\int P(\alpha) d^2 \alpha = 1$  and describes the state. It follows that the expectation value of an operator  $A$  with respect to the density operator described by  $P(\alpha)$  is given by:

$$\langle A \rangle = \text{Tr} (A \rho) \quad (148)$$

$$= \int P(\alpha) \langle \alpha | A | \alpha \rangle d^2 \alpha. \quad (149)$$



We are typically interested in thermal state density operators, for which it can be shown that  $P(\alpha) = \frac{1}{\pi N} e^{-\frac{|\alpha|^2}{N}}$  where  $N = (e^{\beta\omega} - 1)^{-1}$  is the average number of excitations in an oscillator of frequency  $\omega$  at inverse temperature  $\beta = \frac{1}{k_B T}$ .

Using the integral representation (149) we could obtain that the expected value for the displacement operator  $D(h)$  with  $h \in \mathbb{C}$  is equal to:

$$\langle D(h) \rangle = \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} \langle \alpha | D(h) | \alpha \rangle d^2 \alpha \quad (150)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} \langle 0 | D(-\alpha) D(h) D(\alpha) | 0 \rangle d^2 \alpha, \quad (151)$$

$$D(h) D(\alpha) = D(h + \alpha) e^{\frac{1}{2}(h\alpha^* - h^* \alpha)}, \quad (152)$$

$$D(-\alpha) (D(h) D(\alpha)) = D(-\alpha) D(h + \alpha) e^{\frac{1}{2}(h\alpha^* - h^* \alpha)} \quad (153)$$

$$= D(h) e^{\frac{1}{2}(-\alpha(h+\alpha)^* + \alpha^*(h+\alpha))} e^{\frac{1}{2}(h\alpha^* - h^* \alpha)} \quad (154)$$

$$= D(\alpha) e^{\frac{1}{2}(-\alpha h^* - |\alpha|^2 + \alpha^* h + |\alpha|^2)} e^{\frac{1}{2}(h\alpha^* - h^* \alpha)} \quad (155)$$

$$= D(\alpha) e^{(h\alpha^* - h^* \alpha)}, \quad (156)$$

$$\langle D(h) \rangle = \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} \langle 0 | D(h) e^{(h\alpha^* - h^* \alpha)} | 0 \rangle d^2 \alpha \quad (157)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} e^{(h\alpha^* - h^* \alpha)} \langle 0 | D(h) | 0 \rangle d^2 \alpha \quad (158)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} e^{(h\alpha^* - h^* \alpha)} \langle 0 | h \rangle d^2 \alpha, \quad (159)$$

$$|\alpha\rangle = e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle, \quad (160)$$

$$\langle D(h) \rangle = \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} e^{(h\alpha^* - h^* \alpha)} \langle 0 | e^{-\frac{|h|^2}{2}} \sum_{n=0}^{\infty} \frac{h^n}{\sqrt{n!}} |n\rangle d^2 \alpha \quad (161)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} e^{(h\alpha^* - h^* \alpha)} e^{-\frac{|h|^2}{2}} d^2 \alpha \quad (162)$$

$$= \frac{e^{-\frac{|h|^2}{2}}}{\pi N} \int e^{-\frac{|\alpha|^2}{N} + h\alpha^* - h^* \alpha} d^2 \alpha, \quad (163)$$

$$\alpha = x + iy, \quad (164)$$

$$\langle D(h) \rangle = \frac{e^{-\frac{|h|^2}{2}}}{\pi N} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-\frac{x^2+y^2}{N} + h(x-iy) - h^*(x+iy)} dx dy \quad (165)$$

$$= \frac{e^{-\frac{|h|^2}{2}}}{\pi N} \int_{-\infty}^{\infty} e^{-\frac{x^2}{N} + hx - h^* x} dx \int_{-\infty}^{\infty} e^{-\frac{y^2}{N} + hy - h^* y} dy, \quad (166)$$

$$-\frac{x^2}{N} + hx - h^* x = -\frac{1}{N} (x^2 - Nhx + Nh^* x) \quad (167)$$

$$= -\frac{1}{N} \left( x + \frac{(Nh^* - Nh)}{2} \right)^2 + \frac{N(h^* - h)^2}{4}, \quad (168)$$

$$-\frac{y^2}{N} - ihy - ih^* y = -\frac{1}{N} (y^2 + iNhy + iNh^* y) \quad (169)$$

$$= -\frac{1}{N} \left( y^2 + \frac{iN(h + h^*)}{2} \right) - \frac{N(h + h^*)^2}{4}, \quad (170)$$

$$\sqrt{2\pi}\sigma = \int_{-\infty}^{\infty} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx, \quad (171)$$

$$\langle D(h) \rangle = \frac{e^{-\frac{|h|^2}{2} + \frac{N(h^* - h)^2}{4} - \frac{N(h + h^*)^2}{4}}}{\pi N} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-\frac{1}{N} \left( x + \frac{(Nh^* - Nh)}{2} \right)^2 - \frac{1}{N} \left( y^2 + \frac{iN(h + h^*)}{2} \right)} dx dy, \quad (172)$$

$$\langle D(h) \rangle = \frac{e^{-\frac{|h|^2}{2} + \frac{N(h^* - h)^2}{4} - \frac{N(h + h^*)^2}{4}}}{\pi N} \int_{-\infty}^{\infty} e^{-\frac{\left( x + \frac{(Nh^* - Nh)}{2} \right)^2}{2 \left( \sqrt{\frac{N}{2}} \right)^2}} dx \int_{-\infty}^{\infty} e^{-\frac{\left( y^2 + \frac{iN(h + h^*)}{2} \right)}{2 \left( \sqrt{\frac{N}{2}} \right)^2}} dy \quad (173)$$

$$= \frac{e^{-\frac{|h|^2}{2} + \frac{N(h^* - h)^2}{4} - \frac{N(h + h^*)^2}{4}}}{\pi N} \left( \sqrt{2\pi} \sqrt{\frac{N}{2}} \right)^2 \quad (174)$$

$$= e^{-\frac{|h|^2}{2} + \frac{N(h^* - h)^2}{4} - \frac{N(h + h^*)^2}{4}} \quad (175)$$

$$= e^{-\frac{|h|^2}{2} + \frac{N(h^{*2} - 2hh^* + h^2)}{4} - \frac{N(h^2 + 2hh^* + h^{*2})}{4}} \quad (176)$$

$$= e^{-|h|^2 \left( N + \frac{1}{2} \right)} \quad (177)$$

$$= e^{-|h|^2 \left( \frac{1}{e^{\beta\omega} - 1} + \frac{1}{2} \right)} \quad (178)$$

$$= e^{-\frac{|h|^2}{2} \left( \frac{e^{\beta\omega} + 1}{e^{\beta\omega} - 1} \right)} \quad (179)$$

$$= e^{-\frac{|h|^2}{2} \coth\left(\frac{\beta\omega}{2}\right)}. \quad (180)$$

In the last line we used  $\frac{e^{\beta\omega} + 1}{e^{\beta\omega} - 1} = \coth\left(\frac{\beta\omega}{2}\right)$ . So the value of (145) using (??) is given by:

$$B = e^{-\sum_{\mathbf{k}} \frac{|\alpha_{\mathbf{k}}|^2}{2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}. \quad (181)$$

We will now force  $\langle \overline{H_I}(t) \rangle_{\overline{H_B}} = 0$ . We will also introduce the bath renormalizing driving in  $\overline{H_S}(t)$  to treat it non-perturbatively in the subsequent formalism, we associate the terms related with  $B_i^+(t) \sigma^+$  and  $B_i^-(t) \sigma^-$  with the interaction part of the Hamiltonian  $\overline{H_I}(t)$  and we subtract their expected value in order to satisfy  $\langle \overline{H_I}(t) \rangle_{\overline{H_B}} = 0$ .

A final form of the terms of the Hamiltonian  $\overline{H}(t)$  is:

$$\overline{H(t)} = \sum_j \varepsilon_j(t) |j\rangle \langle j| + \sum_{j \neq j'} V_{jj'}(t) |j\rangle \langle j'| B_j^+(t) B_{j'}^-(t) + \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{j\mathbf{k}} |j\rangle \langle j| \left( (g_{j\mathbf{k}} - v_{j\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger + (g_{j\mathbf{k}} - v_{j\mathbf{k}}(t))^* b_{\mathbf{k}} + \frac{|v_{j\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( g_{j\mathbf{k}} \frac{v_{j\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} + g_{j\mathbf{k}}^* \frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right) \quad (182)$$

$$= \sum_j \varepsilon_j(t) |j\rangle \langle j| + \sum_{j \neq j'} V_{jj'}(t) |j\rangle \langle j'| B_{jj'}(t) + \sum_j |j\rangle \langle j| B_{jz}(t) + \sum_{j \neq j'} V_{jj'}(t) |j\rangle \langle j'| (B_j^+(t) B_{j'}^-(t) - B_{jj'}(t)) + \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \quad (183)$$

$$\equiv \overline{H_S}(t) + \overline{H_I}(t) + \overline{H_B}. \quad (184)$$

The parts of the Hamiltonian splitted are obtained using the following expected value:

$$\langle B_i^+(t) B_j^-(t) \rangle = B_{ij}(t) \quad (185)$$

$$= \left\langle \prod_{\mathbf{k}} D\left(\frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) \prod_{\mathbf{k}} D\left(-\frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) \right\rangle \quad (186)$$

$$= \left\langle \prod_{\mathbf{k}} \left( D\left(\frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) D\left(-\frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}}\right) \right) \right\rangle \quad (187)$$

$$= \left\langle \prod_{\mathbf{k}} \left( D \left( \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) e^{\frac{1}{2} \left( \frac{v_{i\mathbf{k}}^*(t) v_{j\mathbf{k}}(t) - v_{i\mathbf{k}}(t) v_{j\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right) \right\rangle \quad (188)$$

$$= \prod_{\mathbf{k}} \left\langle D \left( \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right\rangle e^{\frac{1}{2} \left( \frac{v_{i\mathbf{k}}^*(t) v_{j\mathbf{k}}(t) - v_{i\mathbf{k}}(t) v_{j\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \quad (189)$$

$$= \prod_{\mathbf{k}} e^{-\frac{1}{2} \left| \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} e^{\frac{1}{2} \left( \frac{v_{i\mathbf{k}}^*(t) v_{j\mathbf{k}}(t) - v_{i\mathbf{k}}(t) v_{j\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \quad (190)$$

$$= e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{j\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{i\mathbf{k}}^*(t) v_{j\mathbf{k}}(t) - v_{i\mathbf{k}}(t) v_{j\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)}. \quad (191)$$

From the definition  $B_{01}(t) = \langle B_0^+(t) B_1^-(t) \rangle$  using the displacement operator we have:

$$\langle B_0^+(t) B_1^-(t) \rangle = B_{01}(t) \quad (192)$$

$$= \left\langle \prod_{\mathbf{k}} D \left( \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \prod_{\mathbf{k}} D \left( -\frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right\rangle \quad (193)$$

$$= \left\langle \prod_{\mathbf{k}} \left( D \left( \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) D \left( -\frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right) \right\rangle \quad (194)$$

$$= \left\langle \prod_{\mathbf{k}} \left( D \left( \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) e^{\frac{1}{2} \left( \frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right) \right\rangle \quad (195)$$

$$= \prod_{\mathbf{k}} \left( \left\langle D \left( \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right\rangle e^{\frac{1}{2} \left( \frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right) \quad (196)$$

$$= \prod_{\mathbf{k}} \left( e^{-\frac{1}{2} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} e^{\frac{1}{2} \left( \frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right) \quad (197)$$

$$= e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)}. \quad (198)$$

We can check:

$$\langle B_0^+(t) B_1^-(t) \rangle = B_{01}(t) \quad (199)$$

$$= e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \quad (200)$$

$$= e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)}^* \quad (201)$$

$$= \langle B_1^+(t) B_0^-(t) \rangle^* \quad (202)$$

$$= B_{10}^*(t). \quad (203)$$

The parts of the splitted Hamiltonian are:

$$\overline{H_{\bar{S}}(t)} \equiv (\varepsilon_0(t) + R_0(t)) |0\rangle\langle 0| + (\varepsilon_1(t) + R_1(t)) |1\rangle\langle 1| + V_{10}(t) B_{10} \sigma^+ + V_{01}(t) B_{01} \sigma^-, \quad (204)$$

$$\overline{H_{\bar{I}}(t)} \equiv V_{10}(t) (B_1^+(t) B_0^-(t) - B_{10}(t)) \sigma^+ + V_{01}(t) (B_0^+(t) B_1^-(t) - B_{01}(t)) \sigma^- + |0\rangle\langle 0| B_{0z}(t) + |1\rangle\langle 1| B_{1z}(t), \quad (205)$$

$$\overline{H_B} \equiv \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \quad (206)$$

$$= H_B. \quad (207)$$

Note that  $\overline{H_B}$ , which is the bath acting on the effective “system”  $\tilde{S}$  in the variational frame, is just the original bath,  $H_B$ , before transforming to the variational frame.

For the Hamiltonian (205) we can verify the condition  $\langle \overline{H_I} \rangle_{\overline{H_B}} = 0$  in the following way:

$$\langle \overline{H_I} \rangle_{\overline{H_B}} = \left\langle \sum_{n\mathbf{k}} \left( (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger + (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t))^* b_{\mathbf{k}} \right) |n\rangle\langle n| + \sum_{j \neq j'} V_{jj'}(t) |j\rangle\langle j'| \left( B_j^+(t) B_{j'}^-(t) - B_{jj'}(t) \right) \right\rangle_{\overline{H_B}} \quad (208)$$

$$= \left\langle \sum_{n\mathbf{k}} \left( (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger + (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t))^* b_{\mathbf{k}} \right) |n\rangle\langle n| \right\rangle_{\overline{H_B}} + \left\langle \sum_{j \neq j'} V_{jj'}(t) |j\rangle\langle j'| \left( B_j^+(t) B_{j'}^-(t) - B_{jj'}(t) \right) \right\rangle_{\overline{H_B}} \quad (209)$$

$$= \sum_{n\mathbf{k}} \left( \left\langle (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger \right\rangle_{\overline{H_B}} + \left\langle (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t))^* b_{\mathbf{k}} \right\rangle_{\overline{H_B}} \right) |n\rangle\langle n| + \sum_{j \neq j'} |j\rangle\langle j'| \left( \left\langle V_{jj'}(t) B_j^+(t) B_{j'}^-(t) \right\rangle_{\overline{H_B}} - \left\langle V_{jj'}(t) B_{jj'}(t) \right\rangle_{\overline{H_B}} \right) \quad (210)$$

$$= \sum_{n\mathbf{k}} \left( (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t)) \left\langle b_{\mathbf{k}}^\dagger \right\rangle_{\overline{H_B}} + (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t))^* \left\langle b_{\mathbf{k}} \right\rangle_{\overline{H_B}} \right) |n\rangle\langle n| + \sum_{j \neq j'} |j\rangle\langle j'| V_{jj'}(t) \left( \left\langle B_j^+(t) B_{j'}^-(t) \right\rangle_{\overline{H_B}} - \left\langle B_{jj'}(t) \right\rangle_{\overline{H_B}} \right) \quad (211)$$

$$= \sum_{n\mathbf{k}} \left( (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t)) \left\langle b_{\mathbf{k}}^\dagger \right\rangle_{\overline{H_B}} + (g_{n\mathbf{k}} - v_{n\mathbf{k}}(t))^* \left\langle b_{\mathbf{k}} \right\rangle_{\overline{H_B}} \right) |n\rangle\langle n| + \sum_{j \neq j'} |j\rangle\langle j'| V_{jj'}(t) (B_{jj'}(t) - B_{jj'}(t)) \quad (212)$$

$$= 0. \quad (213)$$

We used (142) and (??) to evaluate the expression. Let's consider the following Hermitian combinations:

$$B_x(t) = B_x^\dagger(t) \quad (214)$$

$$= \frac{B_1^+(t) B_0^-(t) + B_0^+(t) B_1^-(t) - B_{10}(t) - B_{01}(t)}{2}, \quad (215)$$

$$B_y(t) = B_y^\dagger(t) \quad (216)$$

$$= \frac{B_0^+(t) B_1^-(t) - B_1^+(t) B_0^-(t) + B_{10}(t) - B_{01}(t)}{2i}, \quad (217)$$

$$B_{iz}(t) = B_{iz}^\dagger(t) \quad (218)$$

$$= \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) b_{\mathbf{k}}^\dagger + (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* b_{\mathbf{k}} \right). \quad (219)$$

Writing the equations (204) and (205) using the previous combinations we obtain that:

$$\overline{H_S}(t) = \sum_{j \in \{0,1\}} (\varepsilon_j(t) + R_j(t)) |j\rangle\langle j| + V_{10}(t) B_{10}(t) \sigma^+ + V_{01}(t) B_{01}(t) \sigma^- \quad (220)$$

$$= \sum_{j \in \{0,1\}} (\varepsilon_j(t) + R_j(t)) |j\rangle\langle j| + V_{10}(t) B_{10}(t) \frac{\sigma_x + i\sigma_y}{2} + V_{01}(t) B_{01}(t) \frac{\sigma_x - i\sigma_y}{2} \quad (221)$$

$$= \sum_{j \in \{0,1\}} (\varepsilon_j(t) + R_j(t)) |j\rangle\langle j| + V_{10}(t) \left( B_{10}^{\Re}(t) + iB_{10}^{\Im}(t) \right) \frac{\sigma_x + i\sigma_y}{2} + V_{01}(t) \left( B_{10}^{\Re}(t) - iB_{10}^{\Im}(t) \right) \frac{\sigma_x - i\sigma_y}{2} \quad (222)$$

$$= \sum_{j \in \{0,1\}} (\varepsilon_j(t) + R_j(t)) |j\rangle\langle j| + B_{10}^{\Re}(t) \left( V_{10}(t) \frac{\sigma_x + i\sigma_y}{2} + V_{01}(t) \frac{\sigma_x - i\sigma_y}{2} \right) + iB_{10}^{\Im}(t) \left( V_{10}(t) \frac{\sigma_x + i\sigma_y}{2} - V_{01}(t) \frac{\sigma_x - i\sigma_y}{2} \right) \quad (223)$$

$$= \sum_{j \in \{0,1\}} (\varepsilon_j(t) + R_j(t)) |j\rangle\langle j| + B_{10}^{\Re}(t) \left( \sigma_x \frac{V_{10}(t) + V_{01}(t)}{2} + i\sigma_y \frac{V_{10}(t) - V_{01}(t)}{2} \right) + iB_{10}^{\Im}(t) \left( \sigma_x \frac{V_{10}(t) - V_{01}(t)}{2} + i\sigma_y \frac{V_{10}(t) + V_{01}(t)}{2} \right) \quad (224)$$

$$= \sum_{j \in \{0,1\}} (\varepsilon_j(t) + R_j(t)) |j\rangle\langle j| + B_{10}^{\Re}(t) \left( \sigma_x V_{10}^{\Re}(t) - \sigma_y V_{10}^{\Im}(t) \right) + iB_{10}^{\Im}(t) \left( i\sigma_x V_{10}^{\Im}(t) + i\sigma_y V_{10}^{\Re}(t) \right) \quad (225)$$

$$= (\varepsilon_0(t) + R_0(t)) |0\rangle\langle 0| + (\varepsilon_1(t) + R_1(t)) |1\rangle\langle 1| + B_{10}^{\Re}(t) \left( \sigma_x V_{10}^{\Re}(t) - \sigma_y V_{10}^{\Im}(t) \right) + iB_{10}^{\Im}(t) \left( i\sigma_x V_{10}^{\Im}(t) + i\sigma_y V_{10}^{\Re}(t) \right) \quad (226)$$

$$= (\varepsilon_0(t) + R_0(t)) |0\rangle\langle 0| + (\varepsilon_1(t) + R_1(t)) |1\rangle\langle 1| + \left( \sigma_x B_{10}^{\Re}(t) V_{10}^{\Re}(t) - \sigma_y B_{10}^{\Re}(t) V_{10}^{\Im}(t) \right) - \left( \sigma_x B_{10}^{\Im}(t) V_{10}^{\Im}(t) + \sigma_y B_{10}^{\Im}(t) V_{10}^{\Re}(t) \right) \quad (227)$$

$$= (\varepsilon_0(t) + R_0(t)) |0\rangle\langle 0| + (\varepsilon_1(t) + R_1(t)) |1\rangle\langle 1| + \sigma_x \left( B_{10}^{\Re}(t) V_{10}^{\Re}(t) - B_{10}^{\Im}(t) V_{10}^{\Im}(t) \right) - \sigma_y \left( B_{10}^{\Re}(t) V_{10}^{\Im}(t) + B_{10}^{\Im}(t) V_{10}^{\Re}(t) \right) \quad (228)$$

$$= (\epsilon_0(t) + R_0(t)) |0\rangle\langle 0| + (\epsilon_1(t) + R_1(t)) |1\rangle\langle 1| + \sigma_x (B_{10}^{\Re}(t) V_{10}^{\Re}(t) - B_{10}^{\Im}(t) V_{10}^{\Im}(t)) - \sigma_y (B_{10}^{\Re}(t) V_{10}^{\Im}(t) + B_{10}^{\Im}(t) V_{10}^{\Re}(t)), \quad (229)$$

$$\overline{H_I} = V_{10}(t) (\sigma^+ B_1^+(t) B_0^-(t) - \sigma^+ B_{10}(t)) + V_{01}(t) (\sigma^- B_0^+(t) B_1^-(t) - \sigma^- B_{01}(t)) + |0\rangle\langle 0| B_{0z}(t) + |1\rangle\langle 1| B_{1z}(t) \quad (230)$$

$$= |0\rangle\langle 0| B_{0z}(t) + |1\rangle\langle 1| B_{1z}(t) + (V_{10}^{\Re}(t) + i V_{10}^{\Im}(t)) (\sigma^+ B_1^+(t) B_0^-(t) - \sigma^+ B_{10}(t)) + (V_{10}^{\Re}(t) - i V_{10}^{\Im}(t)) (\sigma^- B_0^+(t) B_1^-(t) - \sigma^- B_{01}(t)) \quad (231)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) (\sigma^+ B_1^+(t) B_0^-(t) - \sigma^+ B_{10}(t) + \sigma^- B_0^+(t) B_1^-(t) - \sigma^- B_{01}(t)) + i V_{10}^{\Im}(t) (\sigma^+ B_1^+(t) B_0^-(t) - \sigma^+ B_{10}(t) - \sigma^- B_0^+(t) B_1^-(t) + \sigma^- B_{01}(t)) \quad (232)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) \left( \frac{\sigma_x + i\sigma_y}{2} B_1^+(t) B_0^-(t) - \frac{\sigma_x + i\sigma_y}{2} B_{10}(t) + \frac{\sigma_x - i\sigma_y}{2} B_0^+(t) B_1^-(t) - \frac{\sigma_x - i\sigma_y}{2} B_{01}(t) \right) \quad (233)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) \left( \frac{\sigma_x + i\sigma_y}{2} B_1^+(t) B_0^-(t) - \frac{\sigma_x + i\sigma_y}{2} B_{10}(t) + \frac{\sigma_x - i\sigma_y}{2} B_0^+(t) B_1^-(t) - \frac{\sigma_x - i\sigma_y}{2} B_{01}(t) \right) \quad (234)$$

$$+ i V_{10}^{\Im}(t) \left( \frac{\sigma_x + i\sigma_y}{2} B_1^+(t) B_0^-(t) - \frac{\sigma_x + i\sigma_y}{2} B_{10}(t) - \frac{\sigma_x - i\sigma_y}{2} B_0^+(t) B_1^-(t) + \frac{\sigma_x - i\sigma_y}{2} B_{01}(t) \right) \quad (235)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) \left( \sigma_x \frac{B_1^+(t) B_0^-(t) + B_0^+(t) B_1^-(t) - B_{10}(t) - B_{01}(t)}{2} + i\sigma_y \frac{B_1^+(t) B_0^-(t) - B_0^+(t) B_1^-(t) - B_{10}(t) + B_{01}(t)}{2} \right) \quad (236)$$

$$+ i V_{10}^{\Im}(t) \left( \sigma_x \frac{B_1^+(t) B_0^-(t) - B_0^+(t) B_1^-(t) - B_{10}(t) + B_{01}(t)}{2} + i\sigma_y \frac{B_1^+(t) B_0^-(t) + B_0^+(t) B_1^-(t) - B_{10}(t) - B_{01}(t)}{2} \right) \quad (237)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) (\sigma_x B_x(t) + \sigma_y B_y(t)) + i V_{10}^{\Im}(t) \left( i\sigma_x \frac{B_1^+(t) B_0^-(t) - B_0^+(t) B_1^-(t) - B_{10}(t) + B_{01}(t)}{2} - \sigma_y \frac{B_1^+(t) B_0^-(t) + B_0^+(t) B_1^-(t) - B_{10}(t) - B_{01}(t)}{2} \right) \quad (238)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) (\sigma_x B_x(t) + \sigma_y B_y(t)) + i V_{10}^{\Im}(t) \left( i^2 \sigma_x \frac{B_1^+(t) B_0^-(t) - B_0^+(t) B_1^-(t) - B_{10}(t) + B_{01}(t)}{2i} - \sigma_y \frac{B_1^+(t) B_0^-(t) + B_0^+(t) B_1^-(t) - B_{10}(t) - B_{01}(t)}{2} \right) \quad (239)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) (\sigma_x B_x(t) + \sigma_y B_y(t)) + i V_{10}^{\Im}(t) \left( i^2 \sigma_x \frac{B_1^+(t) B_0^-(t) - B_0^+(t) B_1^-(t) - B_{10}(t) + B_{01}(t)}{2i} - \sigma_y \frac{B_1^+(t) B_0^-(t) + B_0^+(t) B_1^-(t) - B_{10}(t) - B_{01}(t)}{2} \right) \quad (240)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) (\sigma_x B_x(t) + \sigma_y B_y(t)) + V_{10}^{\Im}(t) (i^2 \sigma_x (-B_y(t)) - \sigma_y B_x(t)) \quad (241)$$

$$= \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) (\sigma_x B_x(t) + \sigma_y B_y(t)) + V_{10}^{\Im}(t) (\sigma_x B_y(t) - \sigma_y B_x(t)). \quad (242)$$

## II. FREE-ENERGY MINIMIZATION

The true free energy  $E_{\text{Free}}(t)$  is bounded by the Bogoliubov inequality:

$$E_{\text{Free}}(t) \leq E_{\text{Free,B}}(t) \equiv -\frac{1}{\beta} \ln \left( \text{Tr} \left( e^{-\beta \overline{H_S}(t) + \overline{H_B}} \right) \right) + \langle \overline{H_I}(t) \rangle_{\overline{H_S}(t) + \overline{H_B}} + O \left( \langle \overline{H_I}^2(t) \rangle_{\overline{H_S}(t) + \overline{H_B}} \right). \quad (243)$$

We will optimize the set of variational parameters  $\{v_{\mathbf{k}}(t)\}$  in order to minimize  $E_{\text{Free,B}}(t)$  (i.e. to make it as close to the true free energy  $E_{\text{Free}}(t)$  as possible). Neglecting the higher order terms and using  $\langle \overline{H_I}(t) \rangle_{\overline{H_S}(t) + \overline{H_B}} = 0$  we can obtain the following condition to obtain the set  $\{v_{\mathbf{k}}(t)\}$ :

$$\frac{\partial E_{\text{Free,B}}(t)}{\partial v_{i\mathbf{k}}(t)} = 0. \quad (244)$$

Using this condition and given that  $[\overline{H_S}(t), \overline{H_B}] = 0$ , we have:

$$e^{-\beta(\overline{H_S}(t) + \overline{H_B})} = e^{-\beta \overline{H_S}(t)} e^{-\beta \overline{H_B}}. \quad (245)$$

Then using the fact that  $\overline{H_S}(t)$  and  $\overline{H_B}$  relate to different Hilbert spaces, we obtain:

$$\text{Tr} \left( e^{-\beta \overline{H_S}(t)} e^{-\beta \overline{H_B}} \right) = \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right) \text{Tr} \left( e^{-\beta \overline{H_B}} \right). \quad (246)$$

So Eq. (244) becomes:

$$\frac{\partial E_{\text{Free,B}}(t)}{\partial v_{i\mathbf{k}}(t)} = -\frac{1}{\beta} \frac{\partial \ln \left( \text{Tr} \left( e^{-\beta(\overline{H_S}(t) + \overline{H_B})} \right) \right)}{\partial v_{i\mathbf{k}}(t)} \quad (247)$$

$$= -\frac{1}{\beta} \frac{\partial \ln \left( \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right) \text{Tr} \left( e^{-\beta \overline{H_B}} \right) \right)}{\partial v_{i\mathbf{k}}(t)} \quad (248)$$

$$= -\frac{1}{\beta} \frac{\partial \left( \ln \left( \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right) \right) + \ln \left( \text{Tr} \left( e^{-\beta \overline{H_B}} \right) \right) \right)}{\partial v_{i\mathbf{k}}(t)} \quad (249)$$

$$= -\frac{1}{\beta} \frac{\partial \ln \left( \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right) \right)}{\partial v_{i\mathbf{k}}(t)} - \frac{1}{\beta} \frac{\partial \ln \left( \text{Tr} \left( e^{-\beta \overline{H_B}} \right) \right)}{\partial v_{i\mathbf{k}}(t)} \quad (250)$$

$$= 0 \quad (\text{by Eq. (244)}). \quad (251)$$

But since  $\overline{H_B} = H_B$  which doesn't contain any  $v_{i\mathbf{k}}(t)$ , a derivative of any function of  $H_B$  that does not introduce new  $v_{i\mathbf{k}}(t)$  will be zero. We therefore require the following:

$$\frac{\partial \ln \left( \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right) \right)}{\partial v_{i\mathbf{k}}(t)} = \frac{1}{e^{-\beta \overline{H_S}(t)}} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right)}{\partial v_{i\mathbf{k}}(t)} \quad (252)$$

$$= 0. \quad (253)$$

This means we need to impose:

$$\frac{\partial \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right)}{\partial v_{i\mathbf{k}}(t)} = 0. \quad (254)$$

First we look at:

$$-\beta \overline{H_S}(t) = -\beta \left( (\varepsilon_0(t) + R_0(t)) |0\rangle\langle 0| + (\varepsilon_1(t) + R_1(t)) |1\rangle\langle 1| + V_{10}(t) B_{10}(t) \sigma^+ + V_{01}(t) B_{01}(t) \sigma^- \right). \quad (255)$$

Then the eigenvalues of  $-\beta \overline{H_S}(t)$  satisfy the following relationship deduced from the Caley-Hamilton theorem:

$$\lambda^2 - \text{Tr} \left( -\beta \overline{H_S}(t) \right) + \text{Det} \left( -\beta \overline{H_S}(t) \right) = 0. \quad (256)$$

Let's define:

$$\varepsilon(t) \equiv \text{Tr} \left( \overline{H_S}(t) \right), \quad (257)$$

$$\eta \equiv \sqrt{\left( \text{Tr} \left( \overline{H_S}(t) \right) \right)^2 - 4 \text{Det} \left( \overline{H_S}(t) \right)}. \quad (258)$$

The solutions of the equation (256) are:

$$\lambda = \beta \frac{-\text{Tr} \left( \overline{H_S}(t) \right) \pm \sqrt{\left( \text{Tr} \left( \overline{H_S}(t) \right) \right)^2 - 4 \text{Det} \left( \overline{H_S}(t) \right)}}{2} \quad (259)$$

$$= \beta \frac{-\varepsilon(t) \pm \eta(t)}{2} \quad (260)$$

$$= -\beta \frac{\varepsilon(t) \mp \eta(t)}{2}. \quad (261)$$

The value of  $\text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right)$  can be written in terms of this eigenvalues as (since there's only 2 eigenvalues of a  $2 \times 2$  matrix):

$$\text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right) = e^{-\frac{\varepsilon(t)\beta}{2}} e^{\frac{\eta(t)\beta}{2}} + e^{-\frac{\varepsilon(t)\beta}{2}} e^{-\frac{\eta(t)\beta}{2}} \quad (262)$$

$$= 2e^{-\frac{\varepsilon(t)\beta}{2}} \cosh \left( \frac{\eta(t)\beta}{2} \right). \quad (263)$$

Given that  $v_{i\mathbf{k}}(t)$  is a complex number then we will optimize in the real and complex parts of this element, this can be seen in the following reasoning.

Using the chain rule on the function  $\text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right) = A(\varepsilon(t), \eta(t))$  to calculate  $\frac{\partial \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right)}{\partial v_{i\mathbf{k}}^{\Re}(t)}$  can lead to:

$$\frac{\partial \text{Tr} \left( e^{-\beta \overline{H_S}(t)} \right)}{\partial v_{i\mathbf{k}}^{\Re}(t)} = \frac{\partial \left( 2e^{-\frac{\varepsilon(t)\beta}{2}} \cosh \left( \frac{\eta(t)\beta}{2} \right) \right)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \quad (264)$$

$$= 2 \left( -\frac{\beta}{2} \frac{\partial \varepsilon(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \right) e^{-\frac{\varepsilon(t)\beta}{2}} \cosh \left( \frac{\eta(t)\beta}{2} \right) + 2 \left( \frac{\beta}{2} \frac{\partial \eta(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \right) e^{-\frac{\varepsilon(t)\beta}{2}} \sinh \left( \frac{\eta(t)\beta}{2} \right) \quad (265)$$

$$= -\beta e^{-\frac{\varepsilon(t)\beta}{2}} \left( \frac{\partial \varepsilon(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \cosh \left( \frac{\eta(t)\beta}{2} \right) - \frac{\partial \eta(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \sinh \left( \frac{\eta(t)\beta}{2} \right) \right). \quad (266)$$

Making the derivate equal to zero make us suitable to write:

$$\frac{\partial \varepsilon(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \cosh \left( \frac{\eta(t)\beta}{2} \right) - \frac{\partial \eta(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \sinh \left( \frac{\eta(t)\beta}{2} \right) = 0. \quad (267)$$

The derivates included in the expression given are related to:

$$\langle B_0^+(t) B_1^-(t) \rangle = \left( \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right) e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth \left( \frac{\beta \omega_{\mathbf{k}}}{2} \right)} \quad (268)$$

$$= \left( \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right)^* e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth \left( \frac{\beta \omega_{\mathbf{k}}}{2} \right)} \quad (269)$$

$$= \langle B_1^+(t) B_0^-(t) \rangle^*, \quad (270)$$

$$R_i(t) = \sum_{\mathbf{k}} \left( \frac{|v_{i\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - \left( g_{i\mathbf{k}} \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} + g_{i\mathbf{k}}^* \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) \right) \quad (271)$$

$$= \sum_{\mathbf{k}} \left( \frac{|v_{i\mathbf{k}}(t)|^2}{\omega_{\mathbf{k}}} - g_{i\mathbf{k}} \frac{v_{i\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}} - g_{i\mathbf{k}}^* \frac{v_{i\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right), \quad (272)$$

$$\langle B_0^+(t) B_1^-(t) \rangle = \left( \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right) e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth \left( \frac{\beta \omega_{\mathbf{k}}}{2} \right)} \quad (273)$$

$$= \left( \prod_{\mathbf{k}} e^{\frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{2\omega_{\mathbf{k}}^2}} \right) e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth \left( \frac{\beta \omega_{\mathbf{k}}}{2} \right)}, \quad (274)$$

$$v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t) = (v_{0\mathbf{k}}^{\Re}(t) - i v_{0\mathbf{k}}^{\Im}(t)) (v_{1\mathbf{k}}^{\Re}(t) + i v_{1\mathbf{k}}^{\Im}(t)) - (v_{0\mathbf{k}}^{\Re}(t) + i v_{0\mathbf{k}}^{\Im}(t)) (v_{1\mathbf{k}}^{\Re}(t) - i v_{1\mathbf{k}}^{\Im}(t)) \quad (275)$$

$$= (v_{0\mathbf{k}}^{\Re}(t) v_{1\mathbf{k}}^{\Re}(t) + i v_{0\mathbf{k}}^{\Re}(t) v_{1\mathbf{k}}^{\Im}(t) - i v_{0\mathbf{k}}^{\Im}(t) v_{1\mathbf{k}}^{\Re}(t) + v_{0\mathbf{k}}^{\Im}(t) v_{1\mathbf{k}}^{\Im}(t)) \quad (276)$$

$$- (v_{0\mathbf{k}}^{\Re}(t) v_{1\mathbf{k}}^{\Re}(t) - i v_{0\mathbf{k}}^{\Re}(t) v_{1\mathbf{k}}^{\Im}(t) + i v_{0\mathbf{k}}^{\Im}(t) v_{1\mathbf{k}}^{\Re}(t) + v_{0\mathbf{k}}^{\Im}(t) v_{1\mathbf{k}}^{\Im}(t)) \quad (277)$$

$$= 2i (v_{0\mathbf{k}}^{\Re}(t) v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t) v_{1\mathbf{k}}^{\Re}(t)), \quad (278)$$

$$|v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)|^2 = (v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) (v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))^* \quad (279)$$

$$= |v_{1\mathbf{k}}(t)|^2 + |v_{0\mathbf{k}}(t)|^2 - (v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t) + v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t)) \quad (280)$$

$$= (v_{1\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t))^2 + (v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{0\mathbf{k}}^{\Im}(t))^2 - 2(v_{1\mathbf{k}}^{\Re}(t) v_{0\mathbf{k}}^{\Re}(t) + v_{1\mathbf{k}}^{\Im}(t) v_{0\mathbf{k}}^{\Im}(t)) \quad (281)$$

$$= (v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))^2. \quad (282)$$

Rewriting in terms of real and imaginary parts.

$$R_i(t) = \sum_{\mathbf{k}} \left( \frac{(v_{i\mathbf{k}}^{\Re}(t))^2 + (v_{i\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}} - \left( g_{i\mathbf{k}} \frac{v_{i\mathbf{k}}^{\Re}(t) - i v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} + g_{i\mathbf{k}}^* \frac{v_{i\mathbf{k}}^{\Re}(t) + i v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} \right) \right) \quad (283)$$

$$= \sum_{\mathbf{k}} \left( \frac{(v_{i\mathbf{k}}^{\Re}(t))^2 + (v_{i\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}} - v_{i\mathbf{k}}^{\Re}(t) \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} - i v_{i\mathbf{k}}^{\Im}(t) \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right), \quad (284)$$

$$\langle B_0^+(t) B_1^-(t) \rangle = \left( \prod_{\mathbf{k}} e^{\frac{v_{0\mathbf{k}}^*(t) v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) v_{1\mathbf{k}}^*(t)}{2\omega_{\mathbf{k}}^2}} \right) \left( e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (285)$$

$$= \left( \prod_{\mathbf{k}} e^{\frac{2i(v_{0\mathbf{k}}^{\Re}(t) v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t) v_{1\mathbf{k}}^{\Re}(t))}{2\omega_{\mathbf{k}}^2}} \right) \left( e^{-\frac{1}{2} \sum_{\mathbf{k}} \frac{(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (286)$$

$$= \left( \prod_{\mathbf{k}} e^{\frac{i(v_{0\mathbf{k}}^{\Re}(t) v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t) v_{1\mathbf{k}}^{\Re}(t))}{\omega_{\mathbf{k}}^2}} \right) \left( e^{-\frac{1}{2} \sum_{\mathbf{k}} \frac{(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right), \quad (287)$$

Calculating the derivates respect to  $\alpha_{i\mathbf{k}}^{\Re}$  and  $\alpha_{i\mathbf{k}}^{\Im}$  we have:

$$\frac{\partial \varepsilon(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} = \frac{\partial (\varepsilon_1(t) + R_1 + \varepsilon_0(t) + R_0(t))}{\partial v_{i\mathbf{k}}^{\Re}(t)} \quad (288)$$

$$= \frac{\partial \left( \left( \frac{(v_{i\mathbf{k}}^{\Re}(t))^2 + (v_{i\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}} - v_{i\mathbf{k}}^{\Re}(t) \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} - i v_{i\mathbf{k}}^{\Im}(t) \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right) \right)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \quad (289)$$

$$= \frac{2v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} - \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}}, \quad (290)$$

$$\frac{\partial |B_{10}(t)|^2}{\partial v_{i\mathbf{k}}^{\Re}(t)} = \frac{\partial \left( e^{-\sum_{\mathbf{k}} \frac{(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)}{\partial v_{i\mathbf{k}}^{\Re}(t)} \quad (291)$$

$$= -\frac{2(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))}{\omega_{\mathbf{k}}^2} \frac{\partial (v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))}{\partial v_{i\mathbf{k}}^{\Re}(t)} e^{-\sum_{\mathbf{k}} \frac{(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (292)$$

$$= -\frac{2(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))}{\omega_{\mathbf{k}}^2} \frac{\partial (v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))}{\partial v_{i\mathbf{k}}^{\Re}(t)} |B_{10}(t)|^2, \quad (293)$$

$$\frac{\partial \eta(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} = \frac{\partial \sqrt{(\text{Tr}(\overline{H_{\bar{S}}(t)}))^2 - 4\text{Det}(\overline{H_{\bar{S}}(t)})}}{\partial v_{i\mathbf{k}}^{\Re}(t)} \quad (294)$$

$$= \frac{2\text{Tr}(\overline{H_{\bar{S}}(t)}) \frac{\partial \text{Tr}(\overline{H_{\bar{S}}(t)})}{\partial v_{i\mathbf{k}}^{\Re}(t)} - 4 \frac{\partial \text{Det}(\overline{H_{\bar{S}}(t)})}{\partial v_{i\mathbf{k}}^{\Re}(t)}}{2\sqrt{(\text{Tr}(\overline{H_{\bar{S}}(t)}))^2 - 4\text{Det}(\overline{H_{\bar{S}}(t)})}} \quad (295)$$

$$= \frac{\varepsilon(t) \left( \frac{2v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} - \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} \right) - 2 \frac{\partial ((\varepsilon_1(t) + R_1(t))(\varepsilon_0(t) + R_0(t)) - |V_{10}(t)|^2 |B_{10}(t)|^2)}{\partial v_{i\mathbf{k}}^{\Re}(t)}}{\eta(t)} \quad (296)$$



$$= \frac{\varepsilon(t) \left( \frac{2v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} - \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} \right) - 2 \left( (\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \left( \frac{2v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} - \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} \right) + \frac{2(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))}{\omega_{\mathbf{k}}^2} \frac{\partial(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))}{\partial v_{i\mathbf{k}}^{\Re}(t)} |B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) \right)}{\eta(t)} \quad (297)$$

$$= \frac{\varepsilon(t) \left( \frac{2v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} - \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} \right) - 2 \left( (\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \left( \frac{2v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} - \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} \right) + \frac{2(v_{i\mathbf{k}}^{\Re}(t) - v_{i'\mathbf{k}}^{\Re}(t))}{\omega_{\mathbf{k}}^2} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) \right)}{\eta(t)} \quad (298)$$

$$= \frac{v_{i\mathbf{k}}^{\Re}(t) \left( 2\varepsilon(t) - 4(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{4}{\omega_{\mathbf{k}}} |B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) \right)}{\eta(t)} + \frac{1}{\eta(t)} \left( -\frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} \varepsilon(t) + 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} \right) \quad (299)$$

$$+ 4 \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}^2} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) \quad (300)$$

From the equation (267) and replacing the derivates obtained we have:

$$\tanh\left(\frac{\beta\eta(t)}{2}\right) = \frac{\frac{\partial\varepsilon(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)}}{\frac{\partial\eta(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)}} \quad (301)$$

$$= \frac{\frac{2v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} - \frac{2g_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}}}{\frac{v_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} \left( 2\varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2}{\omega_{\mathbf{k}}} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) \right) + 2 \frac{(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \frac{g_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} + 2 \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) - \frac{g_{i\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} \varepsilon(t)}{\eta(t)}} \quad (302)$$

Rearranging this equation will lead to:

$$\tanh\left(\frac{\beta\eta(t)}{2}\right) = \frac{(2v_{i\mathbf{k}}^{\Re}(t) - g_{i\mathbf{k}}^* \omega_{\mathbf{k}}) \eta(t)}{v_{i\mathbf{k}}^{\Re}(t) \left( 2\varepsilon(t) - 4(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{4|V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right) - (g_{i\mathbf{k}} + g_{i\mathbf{k}}^*) (\varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t))) + 4 \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} |V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (303)$$

$$= \frac{(2v_{i\mathbf{k}}^{\Re}(t) - 2g_{i\mathbf{k}}^{\Re}(t)) \eta(t)}{v_{i\mathbf{k}}^{\Re}(t) \left( 2\varepsilon(t) - 4(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{4|V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right) - 2g_{i\mathbf{k}}^{\Re}(t) (\varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t))) + 4 \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} |V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (304)$$

$$= \frac{(2v_{i\mathbf{k}}^{\Re}(t) - 2g_{i\mathbf{k}}^{\Re}(t)) \eta(t)}{v_{i\mathbf{k}}^{\Re}(t) \left( 2\varepsilon(t) - 4(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{4|V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right) - 2g_{i\mathbf{k}}^{\Re}(t) (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t)) + 4 \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} |V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (305)$$

$$= \frac{(v_{i\mathbf{k}}^{\Re}(t) - g_{i\mathbf{k}}^{\Re}(t)) \eta(t)}{v_{i\mathbf{k}}^{\Re}(t) \left( \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right) - g_{i\mathbf{k}}^{\Re}(t) (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t)) + 2 \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} |V_{10}(t)B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (306)$$

Separating (305) such that the terms with  $v_{i\mathbf{k}}$  are located at one side of the equation permit us to write:

$$\frac{(v_{i\mathbf{k}}^{\Re}(t) - g_{i\mathbf{k}}^{\Re}(t)) \eta(t)}{\tanh\left(\frac{\beta\eta(t)}{2}\right)} = v_{i\mathbf{k}}^{\Re}(t) \left( \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|B_{10}(t)V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right) - g_{i\mathbf{k}}^{\Re}(t) (2(\varepsilon_i(t) + R_i(t)) - \varepsilon(t)) + 2 \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} |B_{10}(t)V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right), \quad (307)$$

$$v_{i\mathbf{k}}^{\Re}(t) - g_{i\mathbf{k}}^{\Re}(t) = v_{i\mathbf{k}}^{\Re}(t) \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left( \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|V_{10}(t)|^2 |B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right) - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} g_{i\mathbf{k}}^{\Re}(t) (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t)) \quad (308)$$

$$+ 2 \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \frac{v_{i'\mathbf{k}}^{\Re}(t)}{\omega_{\mathbf{k}}} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right), \quad (309)$$

$$v_{i\mathbf{k}}^{\Re}(t) = \frac{g_{i\mathbf{k}}^{\Re}(t) \left( 1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t)) + 2 \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)\omega_{\mathbf{k}}} \frac{v_{i'\mathbf{k}}^{\Re}(t)}{g_{i\mathbf{k}}^{\Re}(t)} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) \right)}{1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left( \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|V_{10}(t)|^2 |B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right)}, \quad (310)$$

$$v_{i\mathbf{k}}^{\Re}(t) = \frac{g_{i\mathbf{k}}^{\Re}(t) \left( 1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t)) + 2 \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)\omega_{\mathbf{k}}} \frac{v_{i'\mathbf{k}}^{\Re}(t)}{g_{i\mathbf{k}}^{\Re}(t)} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right) \right)}{1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left( \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|V_{10}(t)|^2 |B_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right)}. \quad (311)$$

The imaginary part can be found in the following way:

$$\frac{\partial \varepsilon(t)}{\partial v_{i\mathbf{k}}^{\Im}(t)} = \frac{\partial (\varepsilon_1(t) + R_1(t) + \varepsilon_0(t) + R_0(t))}{\partial v_{i\mathbf{k}}^{\Im}(t)} \quad (312)$$

$$= \frac{\partial \left( \left( \frac{(v_{i\mathbf{k}}^{\Re}(t))^2 + (v_{i\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}} - v_{i\mathbf{k}}^{\Re}(t) \frac{g_{i\mathbf{k}} + g_{i\mathbf{k}}^*}{\omega_{\mathbf{k}}} - i v_{i\mathbf{k}}^{\Im}(t) \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right) \right)}{\partial v_{i\mathbf{k}}^{\Im}(t)} \quad (313)$$

$$= 2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} - i \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}}, \quad (314)$$

$$\frac{\partial |B_{10}(t)|^2}{\partial v_{i\mathbf{k}}^{\Im}(t)} = \frac{\partial \left( e^{-\sum_{\mathbf{k}} \frac{(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right)} \right)}{\partial v_{i\mathbf{k}}^{\Im}(t)} \quad (315)$$

$$= - \frac{2(v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))}{\omega_{\mathbf{k}}^2} \frac{\partial (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))}{\partial v_{i\mathbf{k}}^{\Im}(t)} e^{-\sum_{\mathbf{k}} \frac{(v_{1\mathbf{k}}^{\Re}(t) - v_{0\mathbf{k}}^{\Re}(t))^2 + (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))^2}{\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right)} \quad (316)$$

$$= - \frac{2(v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))}{\omega_{\mathbf{k}}^2} \frac{\partial (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))}{\partial v_{i\mathbf{k}}^{\Im}(t)} |B_{10}(t)|^2, \quad (317)$$

$$\frac{\partial \eta(t)}{\partial v_{i\mathbf{k}}^{\Re}(t)} = \frac{\partial \sqrt{\left( \text{Tr} \left( \overline{H_{\bar{S}}(t)} \right) \right)^2 - 4 \text{Det} \left( \overline{H_{\bar{S}}(t)} \right)}}{\partial v_{i\mathbf{k}}^{\Re}(t)} \quad (318)$$

$$= \frac{2 \text{Tr} \left( \overline{H_{\bar{S}}(t)} \right) \frac{\partial \text{Tr} \left( \overline{H_{\bar{S}}(t)} \right)}{\partial v_{i\mathbf{k}}^{\Re}(t)} - 4 \frac{\partial \text{Det} \left( \overline{H_{\bar{S}}(t)} \right)}{\partial v_{i\mathbf{k}}^{\Re}(t)}}{2 \sqrt{\left( \text{Tr} \left( \overline{H_{\bar{S}}(t)} \right) \right)^2 - 4 \text{Det} \left( \overline{H_{\bar{S}}(t)} \right)}} \quad (319)$$

$$= \frac{\varepsilon(t) \left( 2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} - i \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right) - 2 \frac{\partial ((\varepsilon_1(t) + R_1(t))(\varepsilon_0(t) + R_0(t)) - |B_{10}(t) V_{10}(t)|^2)}{\partial v_{i\mathbf{k}}^{\Re}(t)}}{\eta(t)} \quad (320)$$

$$= \frac{\varepsilon(t) \left( 2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} - i \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right) - 2 \left( (\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \left( 2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} - i \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right) + \frac{2(v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))}{\omega_{\mathbf{k}}^2} \frac{\partial (v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))}{\partial v_{i\mathbf{k}}^{\Re}(t)} |B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right) \right)}{\eta(t)} \quad (321)$$

$$= \frac{\varepsilon(t) \left( 2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} - i \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right) - 2 \left( (\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \left( 2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} - i \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}} \right) + \frac{2(v_{1\mathbf{k}}^{\Im}(t) - v_{0\mathbf{k}}^{\Im}(t))}{\omega_{\mathbf{k}}^2} |B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right) \right)}{\eta(t)} \quad (322)$$

$$= \frac{v_{i\mathbf{k}}^{\Im}(t) 4(\varepsilon_i(t) + R_i(t)) - 2\varepsilon(t) - \frac{4|B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}}}{\eta(t)} + \frac{1}{\eta(t)} \left( 2 \frac{g_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} \varepsilon(t) - 4(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \frac{g_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} + 4 \frac{v_{i\mathbf{k}}^{\Im}(t) |B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}^2} \right). \quad (323)$$

From the equation (267) and replacing the derivatives obtained we have:

$$\frac{\frac{\partial \varepsilon(t)}{\partial v_{i\mathbf{k}}^{\Im}(t)}}{\frac{\partial \eta(t)}{\partial v_{i\mathbf{k}}^{\Im}(t)}} \tanh\left(\frac{\beta \eta(t)}{2}\right) \quad (324)$$

$$= \frac{2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} - i \frac{g_{i\mathbf{k}}^* - g_{i\mathbf{k}}}{\omega_{\mathbf{k}}}}{v_{i\mathbf{k}}^{\Im}(t) \left( \frac{2\varepsilon(t) - 4(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{4}{\omega_{\mathbf{k}}} |B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right)}{\eta(t) \omega_{\mathbf{k}}} \right) + \frac{2}{\eta(t)} \left( \frac{g_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) \frac{g_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}} + 2 \frac{v_{i\mathbf{k}}^{\Im}(t)}{\omega_{\mathbf{k}}^2} |B_{10}(t) V_{10}(t)|^2 \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right) \right)}. \quad (325)$$

Rearranging this equation will lead to:



$$a_i(\omega_{\mathbf{k}}, t) = \frac{\left(1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t))\right)}{1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left(\varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|B_{10}(t)|^2|V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}}\right)}, \quad (339)$$

$$b_i(\omega_{\mathbf{k}}, t) = \frac{2\frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \frac{1}{\omega_{\mathbf{k}}} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left(\varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|B_{10}(t)|^2|V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}}\right)}. \quad (340)$$

So the equation (335) written in explicit form is:

$$v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t) = g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) + v_{1\mathbf{k}}(\omega_{\mathbf{k}}, t) b_0(\omega_{\mathbf{k}}, t), \quad (341)$$

$$v_{1\mathbf{k}}(\omega_{\mathbf{k}}, t) = g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) + v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t). \quad (342)$$

This system of equations has the following solutions:

$$v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t) = g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) + v_{1\mathbf{k}}(\omega_{\mathbf{k}}, t) b_0(\omega_{\mathbf{k}}, t) \quad (343)$$

$$= g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) + (g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) + v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)) b_0(\omega_{\mathbf{k}}, t) \quad (344)$$

$$= g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) + g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) b_0(\omega_{\mathbf{k}}, t) + v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t) b_0(\omega_{\mathbf{k}}, t) \quad (345)$$

$$v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t)(1 - b_1(\omega_{\mathbf{k}}, t)b_0(\omega_{\mathbf{k}}, t)) = g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) + g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) b_0(\omega_{\mathbf{k}}, t) \quad (346)$$

$$v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t) = \frac{g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) + g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) b_0(\omega_{\mathbf{k}}, t)}{1 - b_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)}, \quad (347)$$

$$v_{1\mathbf{k}}(\omega_{\mathbf{k}}, t) = g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) + \frac{g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) + g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) b_0(\omega_{\mathbf{k}}, t)}{1 - b_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)} b_1(\omega_{\mathbf{k}}, t) \quad (348)$$

$$= \frac{g_1(\omega_{\mathbf{k}}) a_1(\omega_{\mathbf{k}}, t) + g_0(\omega_{\mathbf{k}}) a_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)}{1 - b_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)}. \quad (349)$$

For a shorter representation let's define:

$$r_i(\omega_{\mathbf{k}}, t) = \frac{a_i(\omega_{\mathbf{k}}, t)}{1 - b_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)}, \quad (350)$$

$$s_i(\omega_{\mathbf{k}}, t) = \frac{a_{(i+1) \bmod 2}(\omega_{\mathbf{k}}, t) b_{i \bmod 2}(\omega_{\mathbf{k}}, t)}{1 - b_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)}. \quad (351)$$

So the variational parameters are given by:

$$\begin{pmatrix} v_{0\mathbf{k}}(\omega_{\mathbf{k}}, t) \\ v_{1\mathbf{k}}(\omega_{\mathbf{k}}, t) \end{pmatrix} \equiv \begin{pmatrix} r_0(\omega_{\mathbf{k}}, t) & s_0(\omega_{\mathbf{k}}, t) \\ r_1(\omega_{\mathbf{k}}, t) & s_1(\omega_{\mathbf{k}}, t) \end{pmatrix} \begin{pmatrix} g_0(\omega_{\mathbf{k}}) \\ g_1(\omega_{\mathbf{k}}) \end{pmatrix}. \quad (352)$$

Given that  $v_{i\mathbf{k}}(\omega_{\mathbf{k}}, t) \equiv g_i(\omega_{\mathbf{k}}) F_i(\omega_{\mathbf{k}}, t)$  then we can write:

$$F_0(\omega_{\mathbf{k}}, t) = r_0(\omega_{\mathbf{k}}, t) + \frac{g_1(\omega_{\mathbf{k}})}{g_0(\omega_{\mathbf{k}})} s_0(\omega_{\mathbf{k}}, t) \quad (353)$$

$$F_1(\omega_{\mathbf{k}}, t) = \frac{g_0(\omega_{\mathbf{k}})}{g_1(\omega_{\mathbf{k}})} r_1(\omega_{\mathbf{k}}, t) + s_1(\omega_{\mathbf{k}}, t) \quad (354)$$

### III. MASTER EQUATION

In order to describe the dynamics of the QD under the influence of the phonon environment, we use the time-convolutionless projection operator technique. The initial density operator is  $\rho_T(0) = \rho_S(0) \otimes \rho_B^{\text{Thermal}}$ , where  $\rho_B^{\text{Thermal}} \equiv \rho_B(0) \equiv \rho_B$ , so the transformed density operator is equal to:

$$\overline{\rho_T(0)} \equiv e^{V(0)} \rho_T(0) e^{-V(0)} \quad (355)$$

$$= (|0\rangle\langle 0|B_0^+(0) + |1\rangle\langle 1|B_1^+(0)) (\rho_S(0) \otimes \rho_B) (|0\rangle\langle 0|B_0^-(0) + |1\rangle\langle 1|B_1^-(0)), \quad (356)$$

$$\text{for } \rho_S(0) = |0\rangle\langle 0|: \quad |0\rangle\langle 0|B_0^+(0)\langle 0|\rho_B|0\rangle\langle 0|B_0^-(0) \quad (357)$$

$$= |0\rangle\langle 0|B_0^+(0)\langle 0|\rho_B|0\rangle\langle 0|B_0^-(0) \quad (358)$$

$$= |0\rangle\langle 0| \otimes B_0^+(0) \rho_B B_0^-(0), \quad (359)$$

$$\text{for } \rho_S(0) = |1\rangle\langle 1|: \quad |1\rangle\langle 1|B_1^+(0) |1\rangle\langle 1|\rho_B |1\rangle\langle 1|B_1^-(0) \quad (360)$$

$$= |1\rangle\langle 1|B_1^+(0) \rho_B B_1^-(0) \quad (361)$$

$$= |1\rangle\langle 1| \otimes B_1^+(0) \rho_B B_1^-(0), \quad (362)$$

$$\text{for } \rho_S(0) = |0\rangle\langle 1|: \quad |0\rangle\langle 0|B_0^+(0) |0\rangle\langle 1|\rho_B |1\rangle\langle 1|B_1^-(0) \quad (363)$$

$$= |0\rangle\langle 1|B_0^+(0) \rho_B |1\rangle\langle 1|B_1^-(0) \quad (364)$$

$$= |0\rangle\langle 1|1\rangle\langle 1|B_0^+(0) \rho_B B_1^-(0) \quad (365)$$

$$= |0\rangle\langle 1| \otimes B_0^+(0) \rho_B B_1^-(0), \quad (366)$$

$$\text{for } \rho_S(0) = |1\rangle\langle 0|: \quad |1\rangle\langle 1|B_1^+(0) |1\rangle\langle 0|\rho_B |0\rangle\langle 0|B_0^-(0) \quad (367)$$

$$= |1\rangle\langle 0| \otimes B_1^+(0) \rho_B B_0^-(0). \quad (368)$$

We transform any operator  $O$  into the interaction picture in the following way:

$$\tilde{O}(t) \equiv U^\dagger(t) O(t) U(t), \quad (369)$$

$$U(t) \equiv \mathcal{T} \exp \left( -i \int_0^t dt' \overline{H_S}(t') \right). \quad (370)$$

Here  $\mathcal{T}$  denotes a time ordering operator. Therefore:

$$\widetilde{\overline{\rho_S}}(t) = U^\dagger(t) \overline{\rho_S}(t) U(t), \text{ where} \quad (371)$$

$$\overline{\rho_S}(t) = \text{Tr}_B(\overline{\rho_T}(t)). \quad (372)$$

In order to separate the Hamiltonian we define the matrix  $\Lambda(t)$  such that  $\Lambda_{1i}(t) = A_i$ ,  $\Lambda_{2i}(t) = B_i$  and  $\Lambda_{3i}(t) = C_i(t)$  written as:

$$\begin{pmatrix} A(t) \\ B(t) \\ C(t) \end{pmatrix} = \begin{pmatrix} \sigma_x & \sigma_y & \frac{I-\sigma_z}{2} & \sigma_x & \sigma_y & \frac{I+\sigma_z}{2} \\ B_x(t) & B_y(t) & B_{1z}(t) & B_y(t) & B_x(t) & B_{0z}(t) \\ V_{10}^{\Re}(t) & V_{10}^{\Re}(t) & 1 & V_{10}^{\Im}(t) & -V_{10}^{\Im}(t) & 1 \end{pmatrix}. \quad (373)$$

In this case  $|1\rangle\langle 1| = \frac{I-\sigma_z}{2}$  and  $|0\rangle\langle 0| = \frac{I+\sigma_z}{2}$  with  $\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = |0\rangle\langle 0| - |1\rangle\langle 1|$ .

The previous notation allows us to write the interaction Hamiltonian  $\overline{H_I}(t)$  as pointed in the equation (??):

$$\overline{H_I}(t) = \sum_i B_{iz}(t) |i\rangle\langle i| + V_{10}^{\Re}(t) (\sigma_x B_x(t) + \sigma_y B_y(t)) + V_{10}^{\Im}(t) (\sigma_x B_y(t) - \sigma_y B_x(t)) \quad (374)$$

$$= B_{0z}(t) |0\rangle\langle 0| + B_{1z}(t) |1\rangle\langle 1| + V_{10}^{\Re}(t) \sigma_x B_x(t) + V_{10}^{\Re}(t) \sigma_y B_y(t) + V_{10}^{\Im}(t) \sigma_x B_y(t) - V_{10}^{\Im}(t) \sigma_y B_x(t) \quad (375)$$

$$= \sum_i C_i(t) (A_i \otimes B_i(t)). \quad (376)$$

As the combined system and environment is closed, within the interaction picture the system-environment density operator evolves according to:

$$\frac{d\widetilde{\overline{\rho_T}}(t)}{dt} = -i[\widetilde{\overline{H_I}}(t), \widetilde{\overline{\rho_T}}(t)]. \quad (377)$$

This equation has the formal solution

$$\widetilde{\overline{\rho_T}}(t) = \overline{\rho_T}(0) - i \int_0^t [\widetilde{\overline{H_I}}(t'), \widetilde{\overline{\rho_T}}(t')] dt'. \quad (378)$$

Replacing the equation (378) in the equation (377) gives us:

$$\frac{d\widetilde{\overline{\rho_T}}(t)}{dt} = -i[\widetilde{\overline{H_I}}(t), \overline{\rho_T}(0)] - \int_0^t [\widetilde{\overline{H_I}}(t), [\widetilde{\overline{H_I}}(t'), \widetilde{\overline{\rho_T}}(t')]] dt'. \quad (379)$$

This equation allow us to iterate and write in terms of a series expansion with  $\overline{\rho_T}(0)$  the solution as:

$$\widetilde{\overline{\rho_T}}(t) = \overline{\rho_T}(0) + \sum_{n=0}^{\infty} (-i)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} dt_n [\widetilde{\overline{H_I}}(t_1), [\widetilde{\overline{H_I}}(t_2), \dots [\widetilde{\overline{H_I}}(t_n), \overline{\rho_T}(0)]] \dots]. \quad (380)$$

Taking the trace over the environmental degrees of freedom, we find

$$\widetilde{\overline{\rho_S}}(t) = \overline{\rho_S}(0) + \sum_{n=1}^{\infty} (-i)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} dt_n \text{Tr}_B [\widetilde{\overline{H_I}}(t_1), [\widetilde{\overline{H_I}}(t_2), \dots [\widetilde{\overline{H_I}}(t_n), \overline{\rho_S}(0) \rho_B]] \dots]. \quad (381)$$

here we have assumed that  $\overline{\rho_T}(0) = \overline{\rho_S}(0) \otimes \rho_B$ . Consider the following notation:

$$\widetilde{\overline{\rho_S}}(t) = (1 + W_1(t) + W_2(t) + \dots) \overline{\rho_S}(0) \quad (382)$$

$$= W(t) \overline{\rho_S}(0). \quad (383)$$

in this case

$$W_n(t) = (-i)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} dt_n \text{Tr}_B [\widetilde{\overline{H_I}}(t_1), [\widetilde{\overline{H_I}}(t_2), \dots [\widetilde{\overline{H_I}}(t_n), (\cdot) \rho_B]] \dots]. \quad (384)$$

are superoperators acting on the initial system density operator. Differentiating with respect to time, we have:

$$\frac{d\widetilde{\overline{\rho_S}}(t)}{dt} = (\dot{W}_1(t) + \dot{W}_2(t) + \dots) \overline{\rho_S}(0) \quad (385)$$

$$= (\dot{W}_1(t) + \dot{W}_2(t) + \dots) W(t)^{-1} W(t) \overline{\rho_S}(0) \quad (386)$$

$$= (\dot{W}_1(t) + \dot{W}_2(t) + \dots) W(t)^{-1} \widetilde{\overline{\rho_S}}(t). \quad (387)$$

where we assumed that  $W(t)$  is invertible. Usually, it is convenient (and possible) to define the interaction Hamiltonian such that  $\text{Tr}_B [\widetilde{\overline{H_I}}(t) \rho_B] = 0$  so  $W_1(t) = 0$ . Thus, to second order and approximating  $W(t) \approx \mathbb{I}$  then the equation (385) becomes:

$$\frac{d\widetilde{\overline{\rho_S}}(t)}{dt} = \dot{W}_2(t) \widetilde{\overline{\rho_S}}(t) \quad (388)$$

$$= - \int_0^t dt_1 \text{Tr}_B [\widetilde{\overline{H_I}}(t), [\widetilde{\overline{H_I}}(t_1), \widetilde{\overline{\rho_S}}(t) \rho_B]]. \quad (389)$$

Replacing  $t_1 \rightarrow t - \tau$

$$\frac{d\overline{\rho_S}(t)}{dt} = -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \int_0^t d\tau \text{Tr}_B [\overline{H_I}(t), [\widetilde{\overline{H_I}}(-\tau), \overline{\rho_S}(t) \rho_B]] . \quad (390)$$

From the interaction picture applied on  $\overline{H_I}(t)$  we find:

$$\widetilde{\overline{H_I}}(t) = U^\dagger(t) e^{iH_B t} \overline{H_I}(t) e^{-iH_B t} U(t) . \quad (391)$$

we use the time-ordering operator  $\mathcal{T}$  because in general  $\overline{H_S}(t)$  doesn't commute with itself at two different times. We write the interaction Hamiltonian as:

$$\widetilde{\overline{H_I}}(t) = \sum_i C_i(t) (\widetilde{A_i}(t) \otimes \widetilde{B_i}(t)) , \quad (392)$$

$$\widetilde{A_i}(t) = U^\dagger(t) e^{iH_B t} A_i e^{-iH_B t} U(t) \quad (393)$$

$$= U^\dagger(t) A_i U(t) e^{iH_B t} e^{-iH_B t} \quad (394)$$

$$= U^\dagger(t) A_i U(t) \mathbb{I} \quad (395)$$

$$= U^\dagger(t) A_i U(t) , \quad (396)$$

$$\widetilde{B_i}(t) = U^\dagger(t) e^{iH_B t} B_i(t) e^{-iH_B t} U(t) \quad (397)$$

$$= U^\dagger(t) U(t) e^{iH_B t} B_i(t) e^{-iH_B t} \quad (398)$$

$$= \mathbb{I} e^{iH_B t} B_i(t) e^{-iH_B t} \quad (399)$$

$$= e^{iH_B t} B_i(t) e^{-iH_B t} . \quad (400)$$

Here we have used the fact that  $[\overline{H_S}(t), H_B] = 0$  because these operators belong to different Hilbert spaces, so  $[U(t), e^{iH_B t}] = 0$ .

Using the expression (392) to replace it in the equation (389)

$$\frac{d\widetilde{\overline{\rho_S}}(t)}{dt} = - \int_0^t \text{Tr}_B [\widetilde{\overline{H_I}}(t), [\widetilde{\overline{H_I}}(t'), \widetilde{\overline{\rho_S}}(t) \rho_B]] dt' \quad (401)$$

$$= - \int_0^t \text{Tr}_B \left[ \sum_j C_j(t) (\widetilde{A_j}(t) \otimes \widetilde{B_j}(t)) , \left[ \sum_i C_i(t') (\widetilde{A_i}(t') \otimes \widetilde{B_i}(t')) , \widetilde{\overline{\rho_S}}(t) \rho_B \right] \right] dt' \quad (402)$$

$$= - \int_0^t \text{Tr}_B [\sum_j C_j(t) (\widetilde{A_j}(t) \otimes \widetilde{B_j}(t)) , \sum_i C_i(t') (\widetilde{A_i}(t') \otimes \widetilde{B_i}(t')) \widetilde{\overline{\rho_S}}(t) \rho_B - \widetilde{\overline{\rho_S}}(t) \rho_B \sum_i C_i(t') (\widetilde{A_i}(t') \otimes \widetilde{B_i}(t'))] dt' \quad (403)$$

$$= - \int_0^t \text{Tr}_B (\sum_j C_j(t) (\widetilde{A_j}(t) \otimes \widetilde{B_j}(t)) \sum_i C_i(t') (\widetilde{A_i}(t') \otimes \widetilde{B_i}(t')) \widetilde{\overline{\rho_S}}(t) \rho_B - \sum_j C_j(t) (\widetilde{A_j}(t) \otimes \widetilde{B_j}(t)) \widetilde{\overline{\rho_S}}(t) \rho_B \sum_i C_i(t') (\widetilde{A_i}(t') \otimes \widetilde{B_i}(t')) \quad (404)$$

$$- \sum_i C_i(t') (\widetilde{A_i}(t') \otimes \widetilde{B_i}(t')) \widetilde{\overline{\rho_S}}(t) \rho_B \sum_j C_j(t) (\widetilde{A_j}(t) \otimes \widetilde{B_j}(t)) + \widetilde{\overline{\rho_S}}(t) \rho_B \sum_i C_i(t') (\widetilde{A_i}(t') \otimes \widetilde{B_i}(t')) \sum_j C_j(t) (\widetilde{A_j}(t) \otimes \widetilde{B_j}(t))) dt' . \quad (405)$$

In order to calculate the correlation functions we define:

$$\mathcal{B}_{ij}(t, t') = \text{Tr}_B (\widetilde{B_i}(t) \widetilde{B_j}(t') \rho_B) . \quad (406)$$

An useful property is

$$\mathcal{B}_{ji}^*(t, t') = \text{Tr}_B (\widetilde{B_j}(t) \widetilde{B_i}(t') \rho_B)^\dagger \quad (407)$$

$$= \text{Tr}_B (\rho_B^\dagger \widetilde{B_i}^\dagger(t') \widetilde{B_j}^\dagger(t)) \quad (408)$$

$$= \text{Tr}_B (\rho_B \widetilde{B_i}(t') \widetilde{B_j}(t)) \quad (409)$$

$$= \text{Tr}_B (\widetilde{B_i}(t') \widetilde{B_j}(t) \rho_B) \quad (410)$$

$$= \mathcal{B}_{ij}(t', t) . \quad (411)$$

The correlation functions relevant that appear in the equation (405) are:

$$\text{Tr}_B \left( \widetilde{B}_j(t) \widetilde{B}_i(t') \rho_B \right) = \left\langle \widetilde{B}_j(t) \widetilde{B}_i(t') \right\rangle_B \quad (412)$$

$$= \mathcal{B}_{ji}(t, t') \quad (413)$$

$$= \mathcal{B}_{ij}^*(t', t) \quad (414)$$

$$\text{Tr}_B \left( \widetilde{B}_j(t) \rho_B \widetilde{B}_i(t') \right) = \text{Tr}_B \left( \widetilde{B}_i(t') \widetilde{B}_j(t) \rho_B \right) \quad (415)$$

$$= \mathcal{B}_{ij}(t', t) \quad (416)$$

$$\text{Tr}_B \left( \widetilde{B}_i(t') \rho_B \widetilde{B}_j(t) \right) = \text{Tr}_B \left( \widetilde{B}_j(t) \widetilde{B}_i(t') \rho_B \right) \quad (417)$$

$$= \mathcal{B}_{ij}^*(t', t) \quad (418)$$

$$\text{Tr}_B \left( \rho_B \widetilde{B}_i(t') \widetilde{B}_j(t) \right) = \text{Tr}_B \left( \widetilde{B}_i(t') \widetilde{B}_j(t) \rho_B \right) \quad (419)$$

$$= \mathcal{B}_{ij}(t', t) \quad (420)$$

The cyclic property of the trace was use widely in the development of equations (412) and (420). Replacing in (405)

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \text{Tr}_B \left( \sum_j C_j(t) (\widetilde{A}_j(t) \otimes \widetilde{B}_j(t)) \sum_i C_i(t') (\widetilde{A}_i(t') \otimes \widetilde{B}_i(t')) \widetilde{\rho_S}(t) \rho_B - \sum_j C_j(t) (\widetilde{A}_j(t) \otimes \widetilde{B}_j(t)) \widetilde{\rho_S}(t) \rho_B \sum_i C_i(t') (\widetilde{A}_i(t') \otimes \widetilde{B}_i(t')) \right) \quad (421)$$

$$- \sum_i C_i(t') (\widetilde{A}_i(t') \otimes \widetilde{B}_i(t')) \widetilde{\rho_S}(t) \rho_B \sum_j C_j(t) (\widetilde{A}_j(t) \otimes \widetilde{B}_j(t)) + \widetilde{\rho_S}(t) \rho_B \sum_i C_i(t') (\widetilde{A}_i(t') \otimes \widetilde{B}_i(t')) \sum_j C_j(t) (\widetilde{A}_j(t) \otimes \widetilde{B}_j(t)) dt'. \quad (422)$$

$$= - \int_0^t \text{Tr}_B \left( \sum_{ji} C_j(t) C_i(t') (\widetilde{A}_j(t) \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{B}_j(t) \widetilde{B}_i(t') \rho_B - \widetilde{A}_j(t) \widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{B}_j(t) \rho_B \widetilde{B}_i(t') \right) \quad (423)$$

$$+ \sum_{ij} C_i(t') C_j(t) (\widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{A}_j(t) \rho_B \widetilde{B}_i(t') \widetilde{B}_j(t) - \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{A}_j(t) \widetilde{B}_i(t') \rho_B \widetilde{B}_j(t)) dt' \quad (424)$$

$$= - \int_0^t \text{Tr}_B \left( \sum_{ji} C_j(t) C_i(t') (\widetilde{A}_j(t) \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{B}_j(t) \widetilde{B}_i(t') \rho_B - \widetilde{A}_j(t) \widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{B}_j(t) \rho_B \widetilde{B}_i(t') \right) \quad (425)$$

$$+ \sum_{ij} C_i(t') C_j(t) (\widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{A}_j(t) \rho_B \widetilde{B}_i(t') \widetilde{B}_j(t) - \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{A}_j(t) \widetilde{B}_i(t') \rho_B \widetilde{B}_j(t)) dt' \quad (426)$$

$$= - \int_0^t \text{Tr}_B \left( \sum_{ij} C_j(t) C_i(t') (\widetilde{A}_j(t) \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{B}_j(t) \widetilde{B}_i(t') \rho_B - \widetilde{A}_j(t) \widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{B}_j(t) \rho_B \widetilde{B}_i(t') \right) \text{ (by permuting i and j because } ij \in J) \quad (427)$$

$$+ \sum_{ij} C_i(t') C_j(t) (\widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{A}_j(t) \rho_B \widetilde{B}_i(t') \widetilde{B}_j(t) - \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{A}_j(t) \widetilde{B}_i(t') \rho_B \widetilde{B}_j(t)) dt' \quad (428)$$

$$= - \int_0^t \text{Tr}_B \left( \sum_{ij} C_j(t) C_i(t') (\widetilde{A}_j(t) \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{B}_j(t) \widetilde{B}_i(t') \rho_B - \widetilde{A}_j(t) \widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{B}_j(t) \rho_B \widetilde{B}_i(t') \right) \quad (429)$$

$$+ \widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{A}_j(t) \rho_B \widetilde{B}_i(t') \widetilde{B}_j(t) - \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{A}_j(t) \widetilde{B}_i(t') \rho_B \widetilde{B}_j(t)) dt' \quad (430)$$

$$= - \int_0^t \left( \sum_{ij} C_j(t) C_i(t') (\widetilde{A}_j(t) \widetilde{A}_i(t') \widetilde{\rho_S}(t) \mathcal{B}_{ji}(t, t') - \widetilde{A}_j(t) \widetilde{\rho_S}(t) \widetilde{A}_i(t') \mathcal{B}_{ij}(t', t) \right) \quad (431)$$

$$+ \widetilde{\rho_S}(t) \widetilde{A}_i(t') \widetilde{A}_j(t) \mathcal{B}_{ij}(t, t) - \widetilde{A}_i(t') \widetilde{\rho_S}(t) \widetilde{A}_j(t) \mathcal{B}_{ji}(t, t)) dt' \quad (432)$$

$$= - \int_0^t \left( \sum_{ij} C_j(t) C_i(t') \left( \mathcal{B}_{ji}(t, t') \left[ \widetilde{A}_j(t), \widetilde{A}_i(t') \widetilde{\rho_S}(t) \right] + \mathcal{B}_{ij}(t', t) \left[ \widetilde{\rho_S}(t) \widetilde{A}_i(t'), \widetilde{A}_j(t) \right] \right) \right) dt' \quad (433)$$

$$= - \int_0^t \left( \sum_{ij} C_i(t) C_j(t') \left( \mathcal{B}_{ij}(t, t') \left[ \widetilde{A}_i(t), \widetilde{A}_j(t') \widetilde{\rho_S}(t) \right] + \mathcal{B}_{ji}(t', t) \left[ \widetilde{\rho_S}(t) \widetilde{A}_j(t'), \widetilde{A}_i(t) \right] \right) \right) dt' \text{ (exchanging i and j)} \quad (434)$$

$$= - \int_0^t \left( \sum_{ij} C_i(t) C_j(t') \left( \mathcal{B}_{ij}(t, t') \left[ \widetilde{A}_i(t), \widetilde{A}_j(t') \widetilde{\rho_S}(t) \right] + \mathcal{B}_{ij}^*(t, t') \left[ \widetilde{\rho_S}(t) \widetilde{A}_j(t'), \widetilde{A}_i(t) \right] \right) \right) dt' \quad (435)$$

$$= - \int_0^t \left( \sum_{ij} C_i(t) C_j(t') \left( \mathcal{B}_{ij}(t, t') \left[ \widetilde{A}_i(t), \widetilde{A}_j(t') \widetilde{\rho_S}(t) \right] - \mathcal{B}_{ij}^*(t, t') \left[ \widetilde{A}_i(t), \widetilde{\rho_S}(t) \widetilde{A}_j(t') \right] \right) \right) dt' \quad (436)$$

We could identify the following commutators in the equation deduced:

$$\mathcal{B}_{ij}(t, t') \widetilde{A}_i(t) \widetilde{A}_j(t') \widetilde{\rho_S}(t) - \mathcal{B}_{ij}(t, t') \widetilde{A}_j(t') \widetilde{\rho_S}(t) \widetilde{A}_i(t) = \mathcal{B}_{ij}(t, t') \left[ \widetilde{A}_i(t), \widetilde{A}_j(t') \widetilde{\rho_S}(t) \right], \quad (437)$$

$$\mathcal{B}_{ij}^*(t, t') \widetilde{\rho_S}(t) \widetilde{A}_j(t') \widetilde{A}_i(t) - \mathcal{B}_{ij}^*(t, t') \widetilde{A}_i(t) \widetilde{\rho_S}(t) \widetilde{A}_j(t') = \mathcal{B}_{ij}^*(t, t') \left[ \widetilde{\rho_S}(t) \widetilde{A}_j(t'), \widetilde{A}_i(t) \right]. \quad (438)$$

Returning to the Schroedinger picture we have:



$$U(t) \widetilde{A}_i(t) \widetilde{A}_j(t') \widetilde{\rho_S}(t) U^\dagger(t) = U(t) \widetilde{A}_i(t) U^\dagger(t) U(t) \widetilde{A}_j(t') U^\dagger(t) U(t) \widetilde{\rho_S}(t) U^\dagger(t), \quad (439)$$

$$= \left( U(t) \widetilde{A}_i(t) U^\dagger(t) \right) \left( U(t) \widetilde{A}_j(t') U^\dagger(t) \right) \left( U(t) \widetilde{\rho_S}(t) U^\dagger(t) \right), \quad (440)$$

$$= A_i(t) \widetilde{A}_j(t', t) \overline{\rho_S}(t). \quad (441)$$

This procedure applying to the relevant commutators give us:

$$U(t) \left[ \widetilde{A}_i(t), \widetilde{A}_j(t') \widetilde{\rho_S}(t) \right] U^\dagger(t) = \left( U(t) \widetilde{A}_i(t) \widetilde{A}_j(t') \widetilde{\rho_S}(t) U^\dagger(t) - U(t) \widetilde{A}_j(t') \widetilde{\rho_S}(t) \widetilde{A}_i(t) U^\dagger(t) \right) \quad (442)$$

$$= A_i(t) \widetilde{A}_j(t', t) \overline{\rho_S}(t) - \widetilde{A}_j(t', t) \overline{\rho_S}(t) A_i \quad (443)$$

$$= \left[ A_i(t), \widetilde{A}_j(t', t) \overline{\rho_S}(t) \right]. \quad (444)$$

Introducing this transformed commutators in the equation (436) allow us to obtain the master equation of the system written as an integro-differential equation with the correlation functions  $\mathcal{B}_{ij}(\tau)$  as defined before, this equations has the following form:

$$\frac{d\overline{\rho_S}(t)}{dt} = -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ij} \int_0^t ds C_i(t) C_j(t') \left( \mathcal{B}_{ij}(t, t') \left[ A_i(t), \widetilde{A}_j(t', t) \overline{\rho_S}(t) \right] + \mathcal{B}_{ij}^*(t, t') \left[ \overline{\rho_S}(t) \widetilde{A}_j(t', t), A_i \right] \right), \quad (445)$$

$$t' = t - \tau \text{ (Change of variables in the integration process),} \quad (446)$$

$$\frac{d\overline{\rho_S}(t)}{dt} = -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ij} \int_0^t d\tau C_i(t) C_j(t') \left( \mathcal{B}_{ij}(t, t') \left[ A_i(t), \widetilde{A}_j(t', t) \overline{\rho_S}(t) \right] + \mathcal{B}_{ij}^*(t, t') \left[ \overline{\rho_S}(t) \widetilde{A}_j(t', t), A_i(t) \right] \right). \quad (447)$$

where  $i, j \in \{1, 2, 3, 4, 5, 6\}$  and  $t' = t - \tau$ .

Here  $\widetilde{A}_j(t - \tau, t) = U(t) U^\dagger(t - \tau) A_j(t) U(t - \tau) U^\dagger(t)$  where  $U(t)$  is given by (370). The equation obtained is a non-Markovian master equation which describes the QD exciton dynamics in the variational frame with a general time-dependent Hamiltonian, and valid at second order in  $H_I(t)$ . In order to write in a simplified way we define the following notation:

$$\mathcal{B}_{ij}(t, t') = \text{Tr}_B \left( \widetilde{B}_i(t) \widetilde{B}_j(t') \rho_B \right) \quad (448)$$

$$= \text{Tr}_B \left( e^{iH_B t} B_i(t) e^{-iH_B t} e^{iH_B t'} B_j(t') e^{-iH_B t'} \rho_B \right) \quad (449)$$

$$e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!} \quad (450)$$

$$e^{-iH_B t'} e^{-\beta H_B} = \sum_{m=0}^{\infty} \frac{(-iH_B t')^m}{m!} \sum_{n=0}^{\infty} \frac{(-\beta H_B)^n}{n!} \quad (451)$$

$$= \sum_{m,n} \frac{(-iH_B t')^m}{m!} \frac{(-\beta H_B)^n}{n!} \quad (452)$$

$$= \sum_{m,n} \frac{(-it')^m}{m!} \frac{(-\beta)^n}{n!} H_B^m H_B^n \quad (453)$$

$$= \sum_{m,n} \frac{(-it')^m}{m!} \frac{(-\beta)^n}{n!} H_B^n H_B^m \text{ (because the powers of a matrix commute)} \quad (454)$$

$$= \sum_{m,n} \frac{(-\beta)^n}{n!} H_B^n \frac{(-it')^m}{m!} H_B^m \quad (455)$$

$$= \sum_{m,n} \frac{(-\beta H_B)^n}{n!} \frac{(-it' H_B)^m}{m!} \quad (456)$$

$$= \sum_{n=0}^{\infty} \frac{(-\beta H_B)^n}{n!} \sum_{m=0}^{\infty} \frac{(-i H_B t')^m}{m!} \quad (457)$$

$$= e^{-\beta H_B} e^{-i H_B t'} \quad (458)$$

$$0 = e^{-i H_B t'} e^{-\beta H_B} - e^{-\beta H_B} e^{-i H_B t'} \text{ (then } e^{-i H_B t'} \text{ and } \rho_B \text{ commute)} \quad (459)$$

$$\mathcal{B}_{ij}(t, t') = \text{Tr}_B \left( e^{i H_B t} B_i(t) e^{-i H_B t} e^{i H_B t'} B_j(t') \rho_B e^{-i H_B t'} \right) \text{ (by permuting } e^{-i H_B t'} \text{ and } \rho_B) \quad (460)$$

$$= \text{Tr}_B \left( \left( e^{i H_B t} B_i(t) e^{-i H_B t} e^{i H_B t'} B_j(t') \right) \rho_B e^{-i H_B t'} \right) \text{ (by associative property)} \quad (461)$$

$$= \text{Tr}_B \left( e^{-i H_B t'} \left( e^{i H_B t} B_i(t) e^{-i H_B t} e^{i H_B t'} B_j(t') \right) \rho_B \right) \text{ (by cyclic property of the trace)} \quad (462)$$

$$= \text{Tr}_B \left( \left( e^{-i H_B t'} e^{i H_B t} \right) B_i(t) \left( e^{-i H_B t} e^{i H_B t'} \right) B_j(t') \rho_B \right) \text{ (by associative property)} \quad (463)$$

$$[i H_B t, -i H_B t'] = i H_B t (-i H_B t') - (-i H_B t') i H_B t \quad (464)$$

$$= t t' H_B^2 - t t' H_B^2 \quad (465)$$

$$= 0 \text{ (so } i H_B t \text{ and } -i H_B t' \text{ commute)} \quad (466)$$

$$e^{-i H_B t'} e^{i H_B t} = e^{i H_B t - i H_B t'} \text{ (by the Zassenhaus formula because } i H_B t \text{ and } -i H_B t' \text{ commute)} \quad (467)$$

$$= e^{i H_B (t - t')} \quad (468)$$

$$= e^{i H_B \tau} \quad (469)$$

$$e^{i H_B t'} e^{-i H_B t} = e^{-i H_B t + i H_B t'} \text{ (by the Zassenhaus formula because } -i H_B t \text{ and } i H_B t' \text{ commute)} \quad (470)$$

$$= e^{i H_B (-t + t')} \quad (471)$$

$$= e^{-i H_B \tau} \quad (472)$$

$$\mathcal{B}_{ij}(t, t') = \text{Tr}_B \left( e^{i H_B \tau} B_i(t) e^{-i H_B \tau} B_j(t') \rho_B \right) \quad (473)$$

$$B_i(t, \tau) \equiv e^{i H_B \tau} B_i(t) e^{-i H_B \tau} \quad (474)$$

$$\mathcal{B}_{ij}(t, t') = \text{Tr}_B \left( e^{i H_B (t - t')} B_i(t) e^{-i H_B (t - t')} B_j(t') \rho_B \right) \quad (475)$$

$$t' = t - \tau \quad (476)$$

$$\mathcal{B}_{ij}(t, t') = \text{Tr}_B \left( e^{i H_B \tau} B_i(t) e^{-i H_B \tau} B_j(t') \rho_B \right) \quad (477)$$

$$= \text{Tr}_B \left( B_i(t, \tau) B_j(t', 0) \rho_B \right) \quad (478)$$

For the following results  $i, j \in \{3, 6\}$ , calculating the correlation functions allow us to obtain:

$$\left\langle \widetilde{B_{j \bmod 2z}(t)} \widetilde{B_{j \bmod 2z}(t')} \right\rangle_B = \text{Tr}_B \left( B_{j \bmod 2z}(t, \tau) B_{j \bmod 2z}(t', 0) \rho_B \right) \quad (479)$$

$$= \int d^2 \alpha P(\alpha) \langle \alpha | B_{j \bmod 2z}(t, \tau) B_{j \bmod 2z}(t', 0) | \alpha \rangle \quad (480)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} \langle \alpha | B_{j \bmod 2z}(t, \tau) B_{j \bmod 2z}(t', 0) | \alpha \rangle d^2 \alpha, \quad (481)$$

$$q_{j\mathbf{k}}(t) = g_{j \bmod 2\mathbf{k}} - v_{j \bmod 2\mathbf{k}}(t) \quad (482)$$

$$B_{j \bmod 2z}(t, \tau) = \sum_{\mathbf{k}} \left( q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau} \right), \quad (483)$$

$$B_{j \bmod 2z}(t', 0) = \sum_{\mathbf{k}'} \left( q_{j\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{j\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right), \quad (484)$$

$$\left\langle \widetilde{B_{j \bmod 2z}(t)} \widetilde{B_{j \bmod 2z}(t')} \right\rangle_B = \text{Tr}_B \left( B_{j \bmod 2z}(t, \tau) B_{j \bmod 2z}(t', 0) \rho_B \right) \quad (485)$$

$$= \text{Tr}_B \left( \sum_{\mathbf{k}} \left( q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau} \right) \sum_{\mathbf{k}'} \left( q_{j\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{j\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right) \rho_B \right) \quad (486)$$

$$= \text{Tr}_B \left( \sum_{\mathbf{k}} (q_{j\mathbf{k}}(t) b_{\mathbf{k}}^{\dagger} e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau}) \sum_{\mathbf{k}'} (q_{j\mathbf{k}'}(t') b_{\mathbf{k}'}^{\dagger} + q_{j\mathbf{k}'}^*(t') b_{\mathbf{k}'}) \rho_B \right) \quad (487)$$

$$= \text{Tr}_B \left( \sum_{\mathbf{k} \neq \mathbf{k}'} \left( q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau} \right) \left( q_{j\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{j\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right) \rho_B \right) \quad (488)$$

$$+\mathrm{Tr}_B\left(\sum_{\mathbf{k}}\left(q_{j\mathbf{k}}(t)b_{\mathbf{k}}^{\dagger}e^{i\omega_{\mathbf{k}}\tau}+q_{j\mathbf{k}}^{*}(t)b_{\mathbf{k}}e^{-i\omega_{\mathbf{k}}\tau}\right)\left(q_{j\mathbf{k}}(t')b_{\mathbf{k}}^{\dagger}+q_{j\mathbf{k}}^{*}(t')b_{\mathbf{k}}\right)\rho_B\right), \quad (489)$$

$$\langle \widetilde{B_{\text{mod}2z}(t)} \widetilde{B_{\text{mod}2z}(t')} \rangle_B = \text{Tr}_B \left( \sum_{\mathbf{k} \neq \mathbf{k}'} \left( q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau} \right) \left( q_{j\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{j\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right) \rho_B \right) \quad (490)$$

$$+ \text{Tr}_B \left( \sum_{\mathbf{k}} \left( q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau} \right) \left( q_{j\mathbf{k}}(t') b_{\mathbf{k}}^\dagger + q_{j\mathbf{k}}^*(t') b_{\mathbf{k}} \right) \rho_B \right) \quad (491)$$

$$0 = \text{Tr}_B \left( \sum_{\mathbf{k} \neq \mathbf{k}'} \left( q_{j\mathbf{k}}(t) b_{\mathbf{k}}^{\dagger} e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau} \right) \left( q_{j\mathbf{k}'}(t') b_{\mathbf{k}'}^{\dagger} + q_{j\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right) \rho_B \right) \quad (492)$$

$$\left\langle \widetilde{B_{j\text{mod}2z}(t)} \widetilde{B_{j\text{mod}2z}(t')} \right\rangle_B = 0 + \text{Tr}_B \left( \sum_{\mathbf{k}} \left( q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau} \right) \left( q_{j\mathbf{k}}(t') b_{\mathbf{k}}^\dagger + q_{j\mathbf{k}}^*(t') b_{\mathbf{k}} \right) \rho_B \right) \quad (493)$$

$$= \text{Tr}_B \sum_{\mathbf{k}} \left( q_{j\mathbf{k}}(t) q_{j\mathbf{k}}(t') \left( b_{\mathbf{k}}^\dagger \right)^2 e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') b_{\mathbf{k}}^\dagger b_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') b_{\mathbf{k}} b_{\mathbf{k}}^\dagger e^{-i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}^*(t') b_{\mathbf{k}}^2 e^{-i\omega_{\mathbf{k}}\tau} \right) \rho_B \quad (494)$$

$$= \text{Tr}_B \left( \sum_{\mathbf{k}} q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') b_{\mathbf{k}}^\dagger b_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \rho_B \right) + \text{Tr}_B \left( \sum_{\mathbf{k}} q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') b_{\mathbf{k}} b_{\mathbf{k}}^\dagger e^{-i\omega_{\mathbf{k}}\tau} \rho_B \right) \quad (495)$$

$$= \sum_{\mathbf{k}} q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} \text{Tr}_B \left( b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \rho_B \right) + \sum_{\mathbf{k}} q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \text{Tr}_B \left( b_{\mathbf{k}} b_{\mathbf{k}}^\dagger \rho_B \right) \quad (496)$$

$$= \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\left|\frac{\alpha_{\mathbf{k}}}{N}\right|^2} \left( q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} \left\langle \alpha_{\mathbf{k}} \left| b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \right| \alpha_{\mathbf{k}} \right\rangle + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \left\langle \alpha_{\mathbf{k}} \left| b_{\mathbf{k}} b_{\mathbf{k}}^\dagger \right| \alpha_{\mathbf{k}} \right\rangle \right) d^2 \alpha_{\mathbf{k}} \quad (497)$$

$$= \sum_{\mathbf{k}} q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') \left( e^{i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \left\langle 0 \left| D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}}^\dagger b_{\mathbf{k}} D(\alpha_{\mathbf{k}}) \right| 0 \right\rangle d^2\alpha_{\mathbf{k}} \right) \quad (498)$$

$$+ \sum_{\mathbf{k}} q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') \left( e^{-i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}} b_{\mathbf{k}}^\dagger D(\alpha_{\mathbf{k}}) | 0 \rangle d^2 \alpha_{\mathbf{k}} \right) \quad (499)$$

$$= \sum_{\mathbf{k}} q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') \left( e^{i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}}^\dagger D(\alpha_{\mathbf{k}}) D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}} D(\alpha_{\mathbf{k}}) | 0 \rangle d^2\alpha_{\mathbf{k}} \right) \quad (500)$$

$$+ \sum_{\mathbf{k}} q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') \left( e^{-i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \left\langle 0 \left| D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}} D(\alpha_{\mathbf{k}}) D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}}^\dagger D(\alpha_{\mathbf{k}}) \right| 0 \right\rangle d^2\alpha_{\mathbf{k}} \right) \quad (501)$$

$$= \sum_{\mathbf{k}} q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') \left( e^{i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | (b_{\mathbf{k}}^\dagger + \alpha_{\mathbf{k}}^*) (b_{\mathbf{k}} + \alpha_{\mathbf{k}}) | 0 \rangle d^2\alpha_{\mathbf{k}} \right) \quad (502)$$

$$+ \sum_{\mathbf{k}} q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') \left( e^{-i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | (b_{\mathbf{k}} + \alpha_{\mathbf{k}}) (b_{\mathbf{k}}^\dagger + \alpha_{\mathbf{k}}^*) | 0 \rangle d^2 \alpha_{\mathbf{k}} \right), \quad (503)$$

$$= \sum_{\mathbf{k}} \left( q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} \frac{\tau}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \left\langle 0 \left| b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + b_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} + b_{\mathbf{k}} \alpha_{\mathbf{k}}^* + |\alpha_{\mathbf{k}}|^2 \right| 0 \right\rangle d^2 \alpha_{\mathbf{k}} \right) \quad (504)$$

$$+q_{j\mathbf{k}}^*(t)q_{j\mathbf{k}}(t')e^{-i\omega_{\mathbf{k}}\tau}\frac{1}{\pi N}\int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}}\langle 0|b_{\mathbf{k}}b_{\mathbf{k}}^\dagger+b_{\mathbf{k}}^\dagger\alpha_{\mathbf{k}}+b_{\mathbf{k}}\alpha_{\mathbf{k}}^*+|\alpha_{\mathbf{k}}|^2|0\rangle d^2\alpha_{\mathbf{k}}\rangle \quad (505)$$

$$= \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \left( q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} \langle 0 | b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |\alpha_{\mathbf{k}}|^2 | 0 \rangle + q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} \langle 0 | b_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} + b_{\mathbf{k}} \alpha_{\mathbf{k}}^* | 0 \rangle \right) \quad (506)$$

$$+ q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \langle 0 | b_{\mathbf{k}} b_{\mathbf{k}}^\dagger + |\alpha_{\mathbf{k}}|^2 | 0 \rangle + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \langle 0 | b_{\mathbf{k}}^\dagger \alpha_{\mathbf{k}} + b_{\mathbf{k}} \alpha_{\mathbf{k}}^* | 0 \rangle \rangle d^2 \alpha_{\mathbf{k}} \quad (507)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \sum_{\mathbf{k}} \left( \langle 0 | q_{\mathbf{k}}(t) q_{\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} (b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |\alpha_{\mathbf{k}}|^2) + q_{\mathbf{k}}^*(t) q_{\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} (b_{\mathbf{k}} b_{\mathbf{k}}^\dagger + |\alpha_{\mathbf{k}}|^2) | 0 \rangle \right) d^2 \alpha_{\mathbf{k}} \quad (508)$$

$$= \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} \left( \langle 0 | |\alpha_{\mathbf{k}}|^2 | 0 \rangle + \langle 0 | b_{\mathbf{k}}^\dagger b_{\mathbf{k}} | 0 \rangle \right) d^2 \alpha_{\mathbf{k}} \quad (509)$$

$$+ \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \left( \langle 0 | b_{\mathbf{k}} b_{\mathbf{k}}^\dagger | 0 \rangle + \langle 0 | |\alpha_{\mathbf{k}}|^2 | 0 \rangle \right) d^2\alpha_{\mathbf{k}}, \quad (510)$$

$$1 = \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} d^2 \alpha_{\mathbf{k}}, \quad (511)$$

$$b_{\mathbf{k}}^\dagger b_{\mathbf{k}} |0\rangle = 0, \quad (512)$$

$$b_{\mathbf{k}} b_{\mathbf{k}}^{\dagger} |0\rangle = |0\rangle, \quad (513)$$

$$\langle \widetilde{B_{j\text{mod}2z}}(t) \widetilde{B_{j\text{mod}2z}}(t') \rangle_B = \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \left( (q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau}) \langle 0 | |\alpha_{\mathbf{k}}|^2 | 0 \rangle \right. \quad (514)$$

$$\left. + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') \left( e^{-i\omega_{\mathbf{k}}\tau} \langle 0 | b_{\mathbf{k}} b_{\mathbf{k}}^\dagger | 0 \rangle \right) \right) d^2 \alpha_{\mathbf{k}} \quad (515)$$

$$= \frac{1}{\pi N} \int \sum_{\mathbf{k}} e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \left( (q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau}) |\alpha_{\mathbf{k}}|^2 + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \right) d^2 \alpha_{\mathbf{k}}, \quad (516)$$

$$\int_0^{2\pi} \int_0^{+\infty} r^2 e^{-\frac{r^2}{N}} r dr d\theta = \int |\alpha_{\mathbf{k}}|^2 e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} d^2 \alpha_{\mathbf{k}} \quad (517)$$

$$= \pi N^2 \quad (518)$$

$$\langle \widetilde{B_{j\text{mod}2z}}(t) \widetilde{B_{j\text{mod}2z}}(t') \rangle_B = \sum_{\mathbf{k}} \left( (q_{j\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau}) N + q_{j\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \right) \quad (519)$$

$$\langle \widetilde{B_{j\text{mod}2z}}(t) \widetilde{B_{j'\text{mod}2z}}(t') \rangle_B = \text{Tr}_B (B_{j\text{mod}2z}(t, \tau) B_{j'\text{mod}2z}(t', 0) \rho_B) \quad (520)$$

$$= \int d^2 \alpha P(\alpha) \langle \alpha | B_{j\text{mod}2z}(t, \tau) B_{j'\text{mod}2z}(t', 0) | \alpha \rangle \quad (521)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha|^2}{N}} \langle \alpha | B_{j\text{mod}2z}(t, \tau) B_{j'\text{mod}2z}(t', 0) | \alpha \rangle d^2 \alpha \quad (522)$$

$$= \langle \alpha | \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} (q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau}) \sum_{\mathbf{k}'} (q_{j'\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{j'\mathbf{k}'}^*(t') b_{\mathbf{k}'}) | \alpha \rangle d^2 \alpha_{\mathbf{k}} \quad (523)$$

$$= \langle \alpha | \sum_{\mathbf{k} \neq \mathbf{k}'} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} (q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau}) (q_{j'\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{j'\mathbf{k}'}^*(t') b_{\mathbf{k}'}) | \alpha \rangle d^2 \alpha_{\mathbf{k}} \quad (524)$$

$$+ \langle \alpha | \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} (q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau}) (q_{j'\mathbf{k}'}(t') b_{\mathbf{k}}^\dagger + q_{j'\mathbf{k}}^*(t') b_{\mathbf{k}}) | \alpha \rangle d^2 \alpha_{\mathbf{k}} \quad (525)$$

$$= \langle \alpha | \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} (q_{j\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}}\tau} + q_{j\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}}\tau}) (q_{j'\mathbf{k}}(t') b_{\mathbf{k}}^\dagger + q_{j'\mathbf{k}}^*(t') b_{\mathbf{k}}) | \alpha \rangle d^2 \alpha_{\mathbf{k}} \quad (526)$$

$$= \left\langle \alpha \left| \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} q_{j\mathbf{k}}(t) q_{j'\mathbf{k}}^*(t') b_{\mathbf{k}}^\dagger b_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \right| \alpha \right\rangle d^2 \alpha_{\mathbf{k}} \quad (527)$$

$$+ \left\langle \alpha \left| \sum_{\mathbf{k}} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} q_{j\mathbf{k}}^*(t) q_{j'\mathbf{k}}(t') b_{\mathbf{k}} b_{\mathbf{k}}^\dagger e^{-i\omega_{\mathbf{k}}\tau} \right| \alpha \right\rangle d^2 \alpha_{\mathbf{k}} \quad (528)$$

$$= \sum_{\mathbf{k}} q_{j\mathbf{k}}(t) q_{j'\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle \alpha_{\mathbf{k}} | b_{\mathbf{k}}^\dagger b_{\mathbf{k}} | \alpha_{\mathbf{k}} \rangle d^2 \alpha_{\mathbf{k}} \quad (529)$$

$$+ \sum_{\mathbf{k}} q_{j\mathbf{k}}^*(t) q_{j'\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle \alpha_{\mathbf{k}} | b_{\mathbf{k}} b_{\mathbf{k}}^\dagger | \alpha_{\mathbf{k}} \rangle d^2 \alpha_{\mathbf{k}}, \quad (530)$$

$$\langle b_{\mathbf{k}} b_{\mathbf{k}}^\dagger \rangle_B = \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}} D(\alpha_{\mathbf{k}}) D(-\alpha_{\mathbf{k}}) b_{\mathbf{k}}^\dagger D(\alpha_{\mathbf{k}}) | 0 \rangle d^2 \alpha_{\mathbf{k}} \quad (531)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | (b_{\mathbf{k}} + \alpha_{\mathbf{k}}) (b_{\mathbf{k}}^\dagger + \alpha_{\mathbf{k}}^*) | 0 \rangle d^2 \alpha_{\mathbf{k}} \quad (532)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | b_{\mathbf{k}} b_{\mathbf{k}}^\dagger + \alpha_{\mathbf{k}} b_{\mathbf{k}}^\dagger + b_{\mathbf{k}} \alpha_{\mathbf{k}}^* + |\alpha_{\mathbf{k}}|^2 | 0 \rangle d^2 \alpha_{\mathbf{k}} \quad (533)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | b_{\mathbf{k}} b_{\mathbf{k}}^\dagger + |\alpha_{\mathbf{k}}|^2 | 0 \rangle d^2 \alpha_{\mathbf{k}} \quad (534)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | |\alpha_{\mathbf{k}}|^2 | 0 \rangle d^2 \alpha_{\mathbf{k}} + \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | b_{\mathbf{k}} b_{\mathbf{k}}^\dagger | 0 \rangle d^2 \alpha_{\mathbf{k}} \quad (535)$$

$$= N + 1, \quad (536)$$

$$\langle b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \rangle_B = \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} \langle 0 | (b_{\mathbf{k}}^\dagger + \alpha_{\mathbf{k}}^*) (b_{\mathbf{k}} + \alpha_{\mathbf{k}}) | 0 \rangle d^2 \alpha_{\mathbf{k}} \quad (537)$$

$$= \frac{1}{\pi N} \int e^{-\frac{|\alpha_{\mathbf{k}}|^2}{N}} |\alpha_{\mathbf{k}}|^2 d^2 \alpha_{\mathbf{k}} \quad (538)$$

$$= N, \quad (539)$$

$$\left\langle \widetilde{B_{j \bmod 2z}}(t) \widetilde{B_{j' \bmod 2z}}(t') \right\rangle_B = \sum_{\mathbf{k}} (q_{j\mathbf{k}}(t) q_{j'\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau} N + q_{j\mathbf{k}}^*(t) q_{j'\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} (N+1)) \quad (540)$$

$$= \sum_{\mathbf{k}} 2N (q_{j\mathbf{k}}(t) q_{j'\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}}\tau})^{\Re} + \sum_{\mathbf{k}} q_{j\mathbf{k}}^*(t) q_{j'\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}}\tau} \quad (541)$$

$$D(h') D(h) = e^{\frac{1}{2}(h'h^* - h'^*h)} D(h' + h), \quad (542)$$

$$\langle D(h') D(h) \rangle_B = \text{Tr}_B \left( e^{\frac{1}{2}(h'h^* - h'^*h)} D(h' + h) \rho_B \right) \quad (543)$$

$$= e^{\frac{1}{2}(h'h^* - h'^*h)} \text{Tr}_B (D(h' + h) \rho_B) \quad (544)$$

$$= e^{\frac{1}{2}(h'h^* - h'^*h)} \frac{1}{\pi N} \int d^2\alpha P(\alpha) \langle \alpha | D(h' + h) | \alpha \rangle \quad (545)$$

$$= e^{\frac{1}{2}(h'h^* - h'^*h)} e^{-\frac{|h+h'|^2}{2} \coth\left(\frac{\beta\omega}{2}\right)}, \quad (546)$$

$$h' = h e^{i\omega\tau}, \quad (547)$$

$$\langle D(h e^{i\omega\tau}) D(h) \rangle_B = e^{\frac{1}{2}(h h^* e^{i\omega\tau} - h^* h e^{-i\omega\tau})} e^{-\frac{|h+h e^{i\omega\tau}|^2}{2} \coth\left(\frac{\beta\omega}{2}\right)}, \quad (548)$$

$$\frac{1}{2} |h|^2 (e^{i\omega\tau} - e^{-i\omega\tau}) = \frac{1}{2} (h h^* e^{i\omega\tau} - h^* h e^{-i\omega\tau}) \quad (549)$$

$$= \frac{1}{2} |h|^2 (\cos(\omega\tau) + i \sin(\omega\tau) - \cos(\omega\tau) + i \sin(\omega\tau)) \quad (550)$$

$$= \frac{1}{2} |h|^2 (2i \sin(\omega\tau)) \quad (551)$$

$$= i |h|^2 \sin(\omega\tau), \quad (552)$$

$$-\frac{|h + h e^{i\omega\tau}|^2}{2} = -|h|^2 \frac{|1 + e^{i\omega\tau}|^2}{2} \quad (553)$$

$$= -|h|^2 \frac{(1 + 2 \cos(\omega\tau) + \cos^2(\omega\tau)) + \sin^2(\omega\tau)}{2} \quad (554)$$

$$= -|h|^2 \frac{2 + 2 \cos(\omega\tau)}{2} \quad (555)$$

$$= -|h|^2 (1 + \cos(\omega\tau)), \quad (556)$$

$$\langle D(h e^{i\omega\tau}) D(h) \rangle_B = e^{i|h|^2 \sin(\omega\tau)} e^{-|h|^2 (1 + \cos(\omega\tau)) \coth\left(\frac{\beta\omega}{2}\right)} \quad (557)$$

$$= e^{i|h|^2 \sin(\omega\tau) - |h|^2 (1 + \cos(\omega\tau)) \coth\left(\frac{\beta\omega}{2}\right)} \quad (558)$$

$$= e^{-|h|^2 (-i \sin(\omega\tau) + \cos(\omega\tau) \coth\left(\frac{\beta\omega}{2}\right))} e^{-|h|^2 \coth\left(\frac{\beta\omega}{2}\right)} \quad (559)$$

$$= \langle D(h) \rangle_B e^{-\phi(\tau)}, \quad (560)$$

$$e^{-\phi(\tau)} = e^{-|h|^2 (\cos(\omega\tau) \coth\left(\frac{\beta\omega}{2}\right) - i \sin(\omega\tau))}, \quad (561)$$

$$\phi(\tau) = |h|^2 \left( \cos(\omega\tau) \coth\left(\frac{\beta\omega}{2}\right) - i \sin(\omega\tau) \right), \quad (562)$$

$$\langle D(h') D(h) \rangle_B = e^{\frac{1}{2}(h'h^* - h'^*h)} e^{-\frac{|h+h'|^2}{2} \coth\left(\frac{\beta\omega}{2}\right)}, \quad (563)$$

$$h' = v e^{i\omega\tau}, \quad (564)$$

$$m_{\mathbf{k}}(t) = \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}}, \quad (565)$$

$$\Gamma_{\mathbf{k}}(t) = \frac{v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \quad (566)$$

$$\left\langle \widetilde{B_1^+ B_0^-}(t) \widetilde{B_1^+ B_0^-}(t') \right\rangle_B = \langle B_1^+ B_0^-(t, \tau) B_1^+ B_0^-(t', 0) \rangle_B \quad (567)$$

$$= \langle B_{10}(t, \tau) B_{10}(t', 0) \rangle_B \quad (568)$$

$$= \text{Tr}_B (B_{10}(t, \tau) B_{10}(t', 0) \rho_B) \quad (569)$$

$$= e^{\chi_{10}(t) + \chi_{10}(t')} \text{Tr}_B \left( \prod_{\mathbf{k}} (D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau})) \prod_{\mathbf{k}} (D(m_{\mathbf{k}}(t'))) \rho_B \right) \quad (570)$$

$$= e^{\chi_{10}(t) + \chi_{10}(t')} \text{Tr}_B \left( \prod_{\mathbf{k}} (D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau}) D(m_{\mathbf{k}}(t'))) \rho_B \right) \quad (571)$$

$$= e^{\chi_{10}(t) + \chi_{10}(t')} \prod_{\mathbf{k}} e^{\frac{1}{2} (m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau} m_{\mathbf{k}}^*(t') - (m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau})^* m_{\mathbf{k}}(t')) - \frac{|m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau} + m_{\mathbf{k}}(t')|^2}{2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (572)$$

$$= e^{\chi_{10}(t) + \chi_{10}(t')} \prod_{\mathbf{k}} e^{i(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau} m_{\mathbf{k}}^*(t'))^{\Im} - \frac{|v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (573)$$

$$= e^{\chi_{10}(t) + \chi_{10}(t')} \prod_{\mathbf{k}} e^{i(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau} m_{\mathbf{k}}^*(t'))^{\Im} - \frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (574)$$

$$\left\langle \widetilde{B_0^+ B_1^-}(t) \widetilde{B_0^+ B_1^-}(t') \right\rangle_B = e^{\chi_{10}(t) + \chi_{10}(t')} \prod_{\mathbf{k}} \left( e^{i(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau} m_{\mathbf{k}}^*(t'))^{\Im} - \frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (575)$$

$$\langle D(h) b \rangle_B = \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle \alpha | D(h) b | \alpha \rangle \quad (576)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle \alpha | D(-\alpha) D(h) b D(\alpha) | \alpha \rangle \quad (577)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle 0 | D(-\alpha) D(h) b D(\alpha) | 0 \rangle \quad (578)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle 0 | D(-\alpha) D(h) D(\alpha) D(-\alpha) b D(\alpha) | 0 \rangle \quad (579)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle 0 | D(-\alpha) D(h) D(\alpha) (b + \alpha) | 0 \rangle \quad (580)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) (b + \alpha) | 0 \rangle \quad (581)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) b | 0 \rangle + \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) \alpha | 0 \rangle \quad (582)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) \alpha | 0 \rangle \quad (583)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} e^{-\frac{|h|^2}{2}} d^2\alpha \quad (584)$$

$$= hN \langle D(h) \rangle_B, \quad (585)$$

$$\langle D(h) b^\dagger \rangle_B = \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle \alpha | D(h) b^\dagger | \alpha \rangle \quad (586)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle 0 | D(-\alpha) D(h) b^\dagger D(\alpha) | 0 \rangle \quad (587)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle 0 | D(-\alpha) D(h) b^\dagger D(\alpha) | 0 \rangle \quad (588)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle 0 | D(-\alpha) D(h) D(\alpha) D(-\alpha) b^\dagger D(\alpha) | 0 \rangle \quad (589)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} \langle 0 | D(-\alpha) D(h) D(\alpha) (b^\dagger + \alpha^*) | 0 \rangle \quad (590)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) (b^\dagger + \alpha^*) | 0 \rangle \quad (591)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) b^\dagger | 0 \rangle + \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) \alpha^* | 0 \rangle \quad (592)$$

$$= \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \langle 0 | D(h) | 1 \rangle + \frac{1}{\pi N} \int d^2\alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^*\alpha} \alpha^* \langle 0 | D(h) | 0 \rangle \quad (593)$$

$$= \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} \langle -h|1 \rangle + \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} \alpha^* \langle 0|D(h)|0 \rangle, \quad (594)$$

$$\langle -h| = e^{-\frac{|-h^*|^2}{2}} \sum_n \frac{(-h^*)^n}{\sqrt{n!}} \langle n| \quad (595)$$

$$\langle -h|1 \rangle = e^{-\frac{|-h^*|^2}{2}} \sum_n \frac{(-h^*)^n}{\sqrt{n!}} \langle n|1 \rangle \quad (596)$$

$$\langle -h|1 \rangle = e^{-\frac{|-h^*|^2}{2}} (-h^*), \quad (597)$$

$$\langle D(h)b^\dagger \rangle_B = \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} e^{-\frac{|-h^*|^2}{2}} (-h^*) + \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} \alpha^* e^{-\frac{|-h^*|^2}{2}} \quad (598)$$

$$\langle D(h)b^\dagger \rangle_B = \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} e^{-\frac{|-h^*|^2}{2}} (-h^*) + \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} \alpha^* e^{-\frac{|-h^*|^2}{2}} \quad (599)$$

$$= -h^* \langle D(h) \rangle_B (N+1), \quad (600)$$

$$\langle bD(h) \rangle_B = \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} \langle \alpha | bD(h) | \alpha \rangle \quad (601)$$

$$= \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} e^{-\frac{|h|^2}{2}} h + \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} e^{-\frac{|h|^2}{2}} \quad (602)$$

$$= h \langle D(h) \rangle_B (N+1), \quad (603)$$

$$\langle b^\dagger D(h) \rangle_B = \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} \langle \alpha | b^\dagger D(h) | \alpha \rangle \quad (604)$$

$$= \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} e^{-\frac{|h|^2}{2}} h + \frac{1}{\pi N} \int d^2 \alpha e^{-\frac{|\alpha|^2}{2}} e^{h\alpha^* - h^* \alpha} e^{-\frac{|h|^2}{2}} \quad (605)$$

$$= -h^* \langle D(h) \rangle_B N. \quad (606)$$

The correlation functions can be found readily as:

$$B_1^+ B_0^- (t, \tau) = \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} e^{\frac{\Gamma_{\mathbf{k}}(t)}{2}} \right), \quad (607)$$

$$B_0^+ B_1^- (t, \tau) = \prod_{\mathbf{k}} \left( D(-m_{\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} e^{\frac{\Gamma_{\mathbf{k}}(t)}{2}} \right), \quad (608)$$

$$B_{10}(t) = e^{\chi_{10}(t)} \left( e^{-\frac{1}{2} \sum_{\mathbf{k}} |m_{\mathbf{k}}(t)|^2 \coth\left(\frac{\beta \omega_{\mathbf{k}}}{2}\right)} \right), \quad (609)$$

$$B_x(t, \tau) = \frac{B_1^+ B_0^- (t, \tau) + B_0^+ B_1^- (t, \tau) - B_{10}(t) - B_{01}(t)}{2}, \quad (610)$$

$$B_y(t, \tau) = \frac{B_0^+ B_1^- (t, \tau) - B_1^+ B_0^- (t, \tau) + B_{10}(t) - B_{01}(t)}{2i}, \quad (611)$$

$$B_{i \bmod 2z}(t, \tau) = \sum_{\mathbf{k}} \left( q_{i\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}} \tau} + q_{i\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}} \tau} \right), \quad (612)$$

$$\left\langle \widetilde{B_{i \bmod 2z}}(t) \widetilde{B_{j \bmod 2z}}(t') \right\rangle_B = \langle B_{i \bmod 2z}(t, \tau) B_{j \bmod 2z}(t', 0) \rangle_B \quad (613)$$

$$= \left\langle \sum_{\mathbf{k}} \left( q_{i\mathbf{k}}(t) b_{\mathbf{k}}^\dagger e^{i\omega_{\mathbf{k}} \tau} + q_{i\mathbf{k}}^*(t) b_{\mathbf{k}} e^{-i\omega_{\mathbf{k}} \tau} \right) \sum_{\mathbf{k}} \left( q_{j\mathbf{k}}(t') b_{\mathbf{k}}^\dagger + q_{j\mathbf{k}}^*(t') b_{\mathbf{k}} \right) \right\rangle_B \quad (614)$$

$$= \sum_{\mathbf{k}} q_{i\mathbf{k}}(t) q_{j\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}} \tau} N_{\mathbf{k}} + \sum_{\mathbf{k}} q_{i\mathbf{k}}^*(t) q_{j\mathbf{k}}(t') e^{-i\omega_{\mathbf{k}} \tau} (N_{\mathbf{k}} + 1), \quad (615)$$

$$\left\langle \widetilde{B_x}(t) \widetilde{B_x}(t') \right\rangle_B = \langle B_x(t, \tau) B_x(t', 0) \rangle_B \quad (616)$$

$$= \left\langle \left( \frac{B_1^+ B_0^- (t, \tau) + B_0^+ B_1^- (t, \tau) - B_{10}(t) - B_{01}(t)}{2} \right) \left( \frac{B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t', 0) - B_{10}(t') - B_{01}(t')}{2} \right) \right\rangle_B \quad (617)$$

$$= \frac{1}{4} \left\langle \left( B_1^+ B_0^- (t, \tau) + B_0^+ B_1^- (t, \tau) - B_{10}(t) - B_{01}(t) \right) \left( B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t', 0) - B_{10}(t') - B_{01}(t') \right) \right\rangle_B \quad (618)$$

$$= \frac{1}{4} \left\langle B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) + B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) - B_1^+ B_0^- (t, \tau) B_{10}(t') - B_1^+ B_0^- (t, \tau) B_{01}(t') \right. \quad (619)$$

$$\left. + B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) - B_0^+ B_1^- (t, \tau) B_{10}(t') - B_0^+ B_1^- (t, \tau) B_{01}(t') \right. \quad (620)$$

$$\left. - B_{10}(t) B_1^+ B_0^- (t', 0) - B_{10}(t) B_0^+ B_1^- (t', 0) + B_{10}(t) B_{10}(t') + B_{10}(t) B_{01}(t') - B_{01}(t) B_1^+ B_0^- (t', 0) \right. \quad (621)$$

$$\left. - B_{01}(t) B_0^+ B_1^- (t', 0) + B_{01}(t) B_{10}(t') + B_{01}(t) B_{01}(t') \right\rangle \quad (622)$$

$$= \frac{1}{4} \left\langle B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) + B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) + B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) \right. \quad (623)$$

$$\left. + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) \right\rangle - \frac{(B_{01}(t) + B_{10}(t))(B_{01}(t') + B_{10}(t'))}{4}, \quad (624)$$

$$U_{10}(t, t') = \prod_{\mathbf{k}} e^{i(m_{\mathbf{k}}(t) m_{\mathbf{k}}^*(t') e^{i\omega_{\mathbf{k}} \tau})^3}, \quad (625)$$

$$\langle B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) \rangle_B = \left\langle \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}} \tau}) e^{\frac{\Gamma_{\mathbf{k}}(t)}{2}} \right) \prod_{\mathbf{k}} \left( D(-m_{\mathbf{k}}(t')) e^{-\frac{\Gamma_{\mathbf{k}}(t')}{2}} \right) \right\rangle_B \quad (626)$$

$$= e^{\chi_{10}(t) + \chi_{01}(t')} \left\langle \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}} \tau}) \right) \prod_{\mathbf{k}} \left( D(-m_{\mathbf{k}}(t')) \right) \right\rangle_B \quad (627)$$

$$= e^{\chi_{10}(t) + \chi_{01}(t')} \prod_{\mathbf{k}} \left\langle \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}} \tau}) D(-m_{\mathbf{k}}(t')) \right) \right\rangle_B \quad (628)$$

$$= e^{\chi_{10}(t) + \chi_{01}(t')} U_{10}^*(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (629)$$

$$\langle B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) \rangle_B = \left\langle \prod_{\mathbf{k}} \left( D(-m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}} \tau}) e^{-\frac{\Gamma_{\mathbf{k}}(t)}{2}} \right) \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t')) e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \right) \right\rangle_B \quad (630)$$

$$= \prod_{\mathbf{k}} e^{-\frac{\Gamma_{\mathbf{k}}(t)}{2}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \left\langle D(-m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}} \tau}) D(m_{\mathbf{k}}(t')) \right\rangle_B \quad (631)$$

$$= e^{\chi_{01}(t) + \chi_{10}(t')} \prod_{\mathbf{k}} \left\langle D(-m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}} \tau}) D(m_{\mathbf{k}}(t')) \right\rangle_B \quad (632)$$

$$= e^{\chi_{01}(t) + \chi_{10}(t')} U_{10}^*(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} + (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (633)$$

$$\langle B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) \rangle_B = e^{\chi_{10}(t) + \chi_{10}(t')} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (634)$$

$$\langle B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) \rangle_B = e^{\chi_{01}(t) + \chi_{01}(t')} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (635)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_x(t') \rangle_B = \frac{1}{4} \left\langle B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) + B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) + B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) \right. \quad (636)$$

$$\left. + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) \right\rangle - \frac{(B_{01}(t) + B_{10}(t))(B_{01}(t') + B_{10}(t'))}{4}, \quad (637)$$

$$= \frac{1}{4} \left( 2U_{10}(t, t') \left( e^{\chi_{10}(t) + \chi_{10}(t')} \right)^{\Re} \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (638)$$

$$\left. + 2U_{10}^*(t, t') \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (639)$$

$$- \left( e^{\chi_{10}(t)} e^{-\frac{1}{2} \sum_{\mathbf{k}} |m_{\mathbf{k}}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Re} \left( e^{\chi_{01}(t')} e^{-\frac{1}{2} \sum_{\mathbf{k}} |m_{\mathbf{k}}(t')|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Re} \quad (640)$$

$$= \frac{1}{2} \left( \left( e^{\chi_{10}(t) + \chi_{10}(t')} \right)^{\Re} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}} \tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (641)$$



$$+ \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (642)$$

$$- \left( e^{\chi_{10}(t)} e^{-\frac{1}{2} \sum_{\mathbf{k}} |m_{\mathbf{k}}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Re} \left( e^{\chi_{01}(t')} e^{-\frac{1}{2} \sum_{\mathbf{k}} (|m_{\mathbf{k}}(t')|^2) \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Re} \quad (643)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_y(t') \rangle_B = \langle B_y(t, \tau) B_y(t', 0) \rangle_B \quad (644)$$

$$= \frac{1}{2} \left( \left( e^{\chi_{10}(t) + \chi_{10}(t')} \right)^{\Re} U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \right. \quad (645)$$

$$\left. + \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (646)$$

$$= \left\langle \left( \frac{B_0^+ B_1^-(t, \tau) - B_1^+ B_0^-(t, \tau) + B_{10}(t) - B_{01}(t)}{2i} \right) \left( \frac{B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t', 0) + B_{10}(t') - B_{01}(t')}{2i} \right) \right\rangle_B \quad (647)$$

$$= -\frac{1}{4} \left\langle \left( B_0^+ B_1^-(t, \tau) - B_1^+ B_0^-(t, \tau) + B_{10}(t) - B_{01}(t) \right) \left( B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t', 0) + B_{10}(t') - B_{01}(t') \right) \right\rangle_B \quad (648)$$

$$= -\frac{1}{4} \left\langle B_0^+ B_1^-(t, \tau) B_0^+ B_1^-(t', 0) - B_0^+ B_1^-(t, \tau) B_1^+ B_0^-(t', 0) + B_0^+ B_1^-(t, \tau) B_{10}(t') - B_0^+ B_1^-(t, \tau) B_{01}(t') - B_1^+ B_0^-(t, \tau) B_0^+ B_1^-(t', 0) \right. \quad (649)$$

$$\left. + B_1^+ B_0^-(t, \tau) B_1^+ B_0^-(t', 0) - B_1^+ B_0^-(t, \tau) B_{10}(t') + B_1^+ B_0^-(t, \tau) B_{01}(t') + B_{10}(t) B_0^+ B_1^-(t', 0) - B_{10}(t) B_1^+ B_0^-(t', 0) \right. \quad (650)$$

$$\left. + B_{10}(t) B_{10}(t') - B_{10}(t) B_{01}(t') - B_{01}(t) B_0^+ B_1^-(t', 0) + B_{01}(t) B_1^+ B_0^-(t', 0) - B_{01}(t) B_{10}(t') + B_{01}(t) B_{01}(t') \right\rangle \quad (651)$$

$$= -\frac{1}{4} \left\langle B_0^+ B_1^-(t, \tau) B_0^+ B_1^-(t', 0) - B_0^+ B_1^-(t, \tau) B_1^+ B_0^-(t', 0) - B_1^+ B_0^-(t, \tau) B_0^+ B_1^-(t', 0) + B_1^+ B_0^-(t, \tau) B_1^+ B_0^-(t', 0) \right. \quad (652)$$

$$\left. + (B_{01}(t))^{\Im} (B_{10}(t'))^{\Im} \right\rangle \quad (653)$$

$$= -\frac{1}{4} \left( 2 \left( e^{\chi_{10}(t) + \chi_{10}(t')} \right)^{\Re} U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (654)$$

$$\left. - 2 \left( e^{\chi_{01}(t) + \chi_{10}(t')} \right)^{\Re} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (655)$$

$$+ \left( e^{\chi_{01}(t)} e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Im} \left( e^{\chi_{10}(t')} e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t')}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Im} \quad (656)$$

$$= -\frac{1}{2} \left( \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Re} U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (657)$$

$$\left. - \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (658)$$

$$+ \left( e^{\chi_{01}(t)} e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Im} \left( e^{\chi_{10}(t')} e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t')}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Im} \quad (659)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_y(t') \rangle_B = \langle B_x(t, \tau) B_y(t', 0) \rangle_B \quad (660)$$

$$= \left\langle \left( \frac{B_1^+ B_0^-(t, \tau) + B_0^+ B_1^-(t, \tau) - B_{10}(t) - B_{01}(t)}{2} \right) \left( \frac{B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t', 0) + B_{10}(t') - B_{01}(t')}{2i} \right) \right\rangle_B \quad (661)$$

$$= \frac{1}{4i} \left\langle B_1^+ B_0^-(t, \tau) B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t, \tau) B_1^+ B_0^-(t', 0) + B_1^+ B_0^-(t, \tau) B_{10}(t') - B_1^+ B_0^-(t, \tau) B_{01}(t') + B_{01}(t) B_1^+ B_0^-(t', 0) \right. \quad (662)$$

$$\left. + B_0^+ B_1^-(t, \tau) B_0^+ B_1^-(t', 0) - B_0^+ B_1^-(t, \tau) B_1^+ B_0^-(t', 0) + B_0^+ B_1^-(t, \tau) B_{10}(t') - B_0^+ B_1^-(t, \tau) B_{01}(t') + B_{01}(t) B_{01}(t') \right. \quad (663)$$

$$\left. - B_{10}(t) B_0^+ B_1^-(t', 0) + B_{10}(t) B_1^+ B_0^-(t', 0) - B_{10}(t) B_{10}(t') + B_{10}(t) B_{01}(t') - B_{01}(t) B_0^+ B_1^-(t', 0) - B_{01}(t) B_{10}(t') \right\rangle_B \quad (664)$$

$$= \frac{1}{4i} \left\langle B_1^+ B_0^-(t, \tau) B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t, \tau) B_1^+ B_0^-(t', 0) + B_1^+ B_0^-(t, \tau) B_{10}(t') - B_1^+ B_0^-(t, \tau) B_{01}(t') \right. \quad (665)$$

$$\left. + B_0^+ B_1^-(t, \tau) B_0^+ B_1^-(t', 0) - B_0^+ B_1^-(t, \tau) B_1^+ B_0^-(t', 0) + B_0^+ B_1^-(t, \tau) B_{10}(t') - B_0^+ B_1^-(t, \tau) B_{01}(t') \right. \quad (666)$$

$$\left. - B_{10}(t) B_0^+ B_1^-(t', 0) + B_{10}(t) B_1^+ B_0^-(t', 0) - B_{10}(t) B_{10}(t') + B_{10}(t) B_{01}(t') - B_{01}(t) B_0^+ B_1^-(t', 0) \right. \quad (667)$$

$$\left. + B_{01}(t) B_1^+ B_0^-(t', 0) - B_{01}(t) B_{10}(t') + B_{01}(t) B_{01}(t') \right\rangle_B \quad (668)$$

$$= \frac{1}{4i} \left\langle B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) - B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) - B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) \right\rangle \quad (669)$$

$$+ \frac{1}{4i} (B_{10} (t) + B_{01} (t)) (B_{10} (t') - B_{01} (t')) \quad (670)$$

$$= \frac{1}{4i} \left\langle B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) - B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) \right. \quad (671)$$

$$\left. - B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) \right\rangle + \frac{1}{4i} (B_{10} (t) + B_{01} (t)) (B_{10} (t') - B_{01} (t')) \quad (672)$$

$$= \frac{1}{4i} \left\langle B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) - B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) \right. \quad (673)$$

$$\left. - B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) \right\rangle + (B_{10} (t))^{\Re} (B_{10} (t'))^{\Im} \quad (674)$$

$$= \frac{1}{4i} \left( \left( e^{\chi_{10}(t) + \chi_{01}(t')} - e^{\chi_{01}(t) + \chi_{10}(t')} \right) U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (675)$$

$$\left. + \left( e^{\chi_{01}(t) + \chi_{01}(t')} - e^{\chi_{10}(t) + \chi_{10}(t')} \right) U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (676)$$

$$+ (B_{10} (t))^{\Re} (B_{10} (t'))^{\Im} \quad (677)$$

$$= \frac{1}{2} \left( \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (678)$$

$$\left. + \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Im} U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) + (B_{10}(t))^{\Re} (B_{10}(t'))^{\Im} \quad (679)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_x(t') \rangle_B = \left\langle \left( \frac{B_0^+ B_1^- (t, \tau) - B_1^+ B_0^- (t, \tau) + B_{10} (t) - B_{01} (t)}{2i} \right) \left( \frac{B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t', 0) - B_{10} (t') - B_{01} (t')}{2} \right) \right\rangle_B \quad (680)$$

$$= \frac{1}{4i} \left\langle \left( B_0^+ B_1^- (t, \tau) - B_1^+ B_0^- (t, \tau) + B_{10} (t) - B_{01} (t) \right) \left( B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t', 0) - B_{10} (t') - B_{01} (t') \right) \right\rangle_B \quad (681)$$

$$= \frac{1}{4i} \left\langle B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) - B_0^+ B_1^- (t, \tau) B_{10} (t') - B_0^+ B_1^- (t, \tau) B_{01} (t') + B_{01} (t) B_{01} (t') \right. \quad (682)$$

$$\left. - B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) - B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) + B_1^+ B_0^- (t, \tau) B_{10} (t') + B_1^+ B_0^- (t, \tau) B_{01} (t') \right. \quad (683)$$

$$\left. + B_{10} (t) B_0^+ B_1^- (t', 0) - B_{10} (t) B_{10} (t') - B_{10} (t) B_{01} (t') - B_{01} (t) B_1^+ B_0^- (t', 0) - B_{01} (t) B_0^+ B_1^- (t', 0) \right. \quad (684)$$

$$\left. + B_{01} (t) B_{10} (t') + B_{10} (t) B_1^+ B_0^- (t', 0) \right\rangle \quad (685)$$

$$= \frac{1}{4i} \left\langle B_0^+ B_1^- (t, \tau) B_1^+ B_0^- (t', 0) + B_0^+ B_1^- (t, \tau) B_0^+ B_1^- (t', 0) - B_1^+ B_0^- (t, \tau) B_1^+ B_0^- (t', 0) - B_1^+ B_0^- (t, \tau) B_0^+ B_1^- (t', 0) \right\rangle \quad (686)$$

$$+ (B_{10} (t))^{\Im} (B_{10} (t'))^{\Re} \quad (687)$$

$$= \frac{1}{4i} \left( e^{\chi_{01}(t) + \chi_{10}(t')} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (688)$$

$$\left. + e^{\chi_{01}(t) + \chi_{01}(t')} U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (689)$$

$$- e^{\chi_{10}(t) + \chi_{10}(t')} U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (690)$$

$$\left. - e^{\chi_{10}(t) + \chi_{01}(t')} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) + (B_{10} (t))^{\Im} (B_{10} (t'))^{\Re} \quad (691)$$

$$= \frac{1}{4i} \left( 2i \left( e^{\chi_{01}(t) + \chi_{10}(t')} \right)^{\Im} U_{10}^* (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (692)$$

$$\left. + 2i \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Im} U_{10} (t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) + (B_{10}(t))^{\Im} (B_{10}(t'))^{\Re} \quad (693)$$

$$= (B_{10}(t))^{\Im} (B_{10}(t'))^{\Re} + \frac{1}{2} \left( \left( e^{\chi_{01}(t) + \chi_{10}(t')} \right)^{\Im} U_{10}^*(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (694)$$

$$\left. + \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (695)$$

$$\langle b^\dagger D(h) \rangle_B = -h^* \langle D(h) \rangle_B N \quad (696)$$

$$\langle b D(h) \rangle_B = h \langle D(h) \rangle_B (N+1) \quad (697)$$

$$\langle D(h) b^\dagger \rangle_B = -h^* \langle D(h) \rangle_B (N+1) \quad (698)$$

$$\langle D(h) b \rangle_B = h \langle D(h) \rangle_B N \quad (699)$$

$$g_{j\mathbf{k}} - v_{j\mathbf{k}}(t) = q_{j\mathbf{k}}(t) \quad (700)$$

$$\langle B_1^+ B_0^-(t, \tau) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger \rangle_B = \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \left\langle \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau}) \right) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger \right\rangle_B \quad (701)$$

$$= e^{\chi_{10}(t)} \left\langle D(m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau}) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger \right\rangle_B \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau}) \right) \right\rangle_B \quad (702)$$

$$= e^{\chi_{10}(t)} q_{i\mathbf{k}'}(t') \left\langle D(m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau}) b_{\mathbf{k}'}^\dagger \right\rangle_B \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau}) \right) \right\rangle_B \quad (703)$$

$$= e^{\chi_{10}(t)} q_{i\mathbf{k}'}(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) \left\langle \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau}) \right) \right\rangle_B \quad (704)$$

$$= q_{i\mathbf{k}'}(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) e^{\chi_{10}(t)} \left\langle \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau}) \right) \right\rangle_B \quad (705)$$

$$= -q_{i\mathbf{k}'}(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{10}(t) \quad (706)$$

$$\langle B_1^+ B_0^-(t, \tau) q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \rangle_B = q_{i\mathbf{k}'}^*(t') \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} \left\langle \prod_{\mathbf{k}} D(m_{\mathbf{k}}(t) e^{i\omega_{\mathbf{k}}\tau}) \right\rangle \quad (707)$$

$$= q_{i\mathbf{k}'}^*(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} B_{10}(t), \quad (708)$$

$$\langle B_0^+ B_1^-(t, \tau) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger \rangle_B = -q_{i\mathbf{k}'}(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{01}(t), \quad (709)$$

$$\langle B_0^+ B_1^-(t, \tau) q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \rangle_B = q_{i\mathbf{k}'}^*(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} B_{01}(t), \quad (710)$$

$$q_{i\mathbf{k}'}(0) = g_{i\mathbf{k}'} - v_{i\mathbf{k}'} \quad (711)$$

$$\langle B_x(t, \tau) B_{\text{imod}2z}(t', 0) \rangle_B = \left\langle \left( \frac{B_{1+} B_{0-}(\tau) + B_{0+} B_{1-}(\tau) - B_{10} - B_{10}^*}{2} \right) \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(0) b_{\mathbf{k}'}^\dagger + q_{i\mathbf{k}'}^*(0) b_{\mathbf{k}'} \right) \right\rangle_B \quad (712)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left\langle (B_{1+} B_{0-}(\tau) + B_{0+} B_{1-}(\tau) - B_{10} - B_{10}^*) \left( q_{i\mathbf{k}'}(0) b_{\mathbf{k}'}^\dagger + q_{i\mathbf{k}'}^*(0) b_{\mathbf{k}'} \right) \right\rangle_B \quad (713)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left\langle (B_{1+} B_{0-}(\tau) + B_{0+} B_{1-}(\tau)) \left( q_{i\mathbf{k}'}(0) b_{\mathbf{k}'}^\dagger + q_{i\mathbf{k}'}^*(0) b_{\mathbf{k}'} \right) \right\rangle_B \quad (714)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left\langle B_{1+} B_{0-}(\tau) q_{i\mathbf{k}'}(0) b_{\mathbf{k}'}^\dagger + B_{0+} B_{1-}(\tau) q_{i\mathbf{k}'}(0) b_{\mathbf{k}'}^\dagger + B_{1+} B_{0-}(\tau) q_{i\mathbf{k}'}^*(0) b_{\mathbf{k}'} \right. \quad (715)$$

$$\left. + B_{0+} B_{1-}(\tau) q_{i\mathbf{k}'}^*(0) b_{\mathbf{k}'} \right\rangle_B \quad (716)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left( -q_{i\mathbf{k}'}(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{10}(t) + q_{i\mathbf{k}'}(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{01}(t) \right. \quad (717)$$

$$\left. + q_{i\mathbf{k}'}^*(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} B_{10}(t) + q_{i\mathbf{k}'}^*(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} B_{01}(t) \right) \quad (718)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left( -q_{i\mathbf{k}'}(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{10}(t) + q_{i\mathbf{k}'}(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{01}(t) \right. \quad (719)$$

$$\left. + q_{i\mathbf{k}'}^*(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} B_{10}(t) + q_{i\mathbf{k}'}^*(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} B_{01}(t) \right) \quad (720)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left( -q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \left( \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* B_{10}(t) + \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* B_{01}(t) \right) \right. \quad (721)$$

$$\left. + q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} B_{10}(t) - m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} B_{01}(t) \right) \right) \quad (722)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left( -q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \left( \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* B_{10}(t) - \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* B_{01}(t) \right) \right. \quad (723)$$

$$\left. + q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} B_{10}(t) - m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} B_{01}(t) \right) \right) \quad (724)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left( -q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (B_{10}(t) - B_{01}(t)) + q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} (B_{10}(t) \right. \quad (725)$$

$$-B_{01}(t))) \quad (726)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} 2iB_{10}^{\mathbb{S}}(t) \left( q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) - q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* \right) \quad (727)$$

$$= i \sum_{\mathbf{k}'} B_{10}^{\mathbb{S}}(t) \left( q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} - q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* \right) \quad (728)$$

$$= i \sum_{\mathbf{k}'} B_{10}^{\mathbb{S}}(t) \left( q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} - q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) (m_{\mathbf{k}'}(t))^* e^{-i\omega_{\mathbf{k}'}\tau} \right), \quad (729)$$

$$= i \sum_{\mathbf{k}'} B_{10}^{\mathbb{S}}(t) \left( q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} - q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) (m_{\mathbf{k}'}(t))^* e^{-i\omega_{\mathbf{k}'}\tau} \right) \quad (730)$$

$$\langle B_{\text{imod}2z}(t, \tau) B_x(t', 0) \rangle_B = \left\langle \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} \right) \left( \frac{B_1^+ B_0^-(t', 0) + B_0^+ B_1^-(t', 0) - B_{10}(t') - B_{01}(t')}{2} \right) \right\rangle_B \quad (731)$$

$$= \sum_{\mathbf{k}'} \left\langle \left( q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} \right) \left( \frac{B_1^+ B_0^-(t', 0) + B_0^+ B_1^-(t', 0) - B_{10}(t') - B_{01}(t')}{2} \right) \right\rangle_B \quad (732)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left\langle \left( q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} \right) \left( B_1^+ B_0^-(t', 0) + B_0^+ B_1^-(t', 0) - B_{10}(t') - B_{01}(t') \right) \right\rangle_B \quad (733)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left\langle \left( q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} \right) \left( B_1^+ B_0^-(t', 0) + B_0^+ B_1^-(t', 0) \right) \right\rangle_B \quad (734)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left\langle q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} B_1^+ B_0^-(t', 0) + q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} B_0^+ B_1^-(t', 0) + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} B_1^+ B_0^-(t', 0) \right. \quad (735)$$

$$\left. + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} B_0^+ B_1^-(t', 0) \right\rangle, \quad (736)$$

$$\langle q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} B_1^+ B_0^-(t', 0) \rangle_B = q_{i\mathbf{k}'}(t) \langle b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} B_1^+ B_0^-(t', 0) \rangle_B \quad (737)$$

$$= q_{i\mathbf{k}'}(t) \left\langle b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t')) e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \right) \right\rangle_B \quad (738)$$

$$= q_{i\mathbf{k}'}(t) \left\langle b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} \left( D(m_{\mathbf{k}'}(t')) e^{\frac{\Gamma_{\mathbf{k}'}(t')}{2}} \right) \right\rangle_B \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} \left( D(m_{\mathbf{k}}(t')) e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \right) \right\rangle_B \quad (739)$$

$$= q_{i\mathbf{k}'}(t) \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} D(m_{\mathbf{k}}(t')) \right\rangle_B \langle b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} D(m_{\mathbf{k}'}(t')) \rangle_B \quad (740)$$

$$= q_{i\mathbf{k}'}(t) \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} D(m_{\mathbf{k}}(t')) \right\rangle_B \langle b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} D(m_{\mathbf{k}'}(t')) \rangle_B \quad (741)$$

$$= q_{i\mathbf{k}'}(t) \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} D(m_{\mathbf{k}}(t')) \right\rangle_B \langle b_{\mathbf{k}'}^\dagger D(m_{\mathbf{k}'}(t')) \rangle_B e^{i\omega_{\mathbf{k}'}\tau} \quad (742)$$

$$= q_{i\mathbf{k}'}(t) \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} D(m_{\mathbf{k}}(t')) \right\rangle_B (-m_{\mathbf{k}'}^*(t') \langle D(m_{\mathbf{k}'}(t')) \rangle_B N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} \quad (743)$$

$$= -m_{\mathbf{k}'}^*(t') q_{i\mathbf{k}'}(t) \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \left\langle \prod_{\mathbf{k}} D(m_{\mathbf{k}}(t')) \right\rangle_B N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} \quad (744)$$

$$= -m_{\mathbf{k}'}^*(t') q_{i\mathbf{k}'}(t) B_{10}(t') N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau}, \quad (745)$$

$$\langle q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} B_0^+ B_1^-(t', 0) \rangle_B = m_{\mathbf{k}'}^*(t') q_{i\mathbf{k}'}(t) B_{01}(t') N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau}, \quad (746)$$

$$\langle q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} B_1^+ B_0^-(t', 0) \rangle_B = q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} \langle b_{\mathbf{k}'} B_1^+ B_0^-(t', 0) \rangle_B \quad (747)$$

$$= q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} \left\langle b_{\mathbf{k}'} \prod_{\mathbf{k}} \left( D(m_{\mathbf{k}}(t')) e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \right) \right\rangle_B \quad (748)$$

$$= q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} \langle b_{\mathbf{k}'} D(m_{\mathbf{k}'}(t')) \rangle_B \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} (D(m_{\mathbf{k}}(t')))) \right\rangle_B \quad (749)$$

$$= q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \langle D(m_{\mathbf{k}'}(t')) \rangle_B \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} (D(m_{\mathbf{k}}(t')))) \right\rangle_B \quad (750)$$

$$= q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} \prod_{\mathbf{k}} e^{\frac{\Gamma_{\mathbf{k}}(t')}{2}} m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \langle D(m_{\mathbf{k}'}(t')) \rangle_B \left\langle \prod_{\mathbf{k} \neq \mathbf{k}'} (D(m_{\mathbf{k}}(t')))) \right\rangle_B \quad (751)$$

$$= q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{10}(t'), \quad (752)$$

$$\langle (q_{i\mathbf{k}'}(t))^* b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} B_{0+} B_1^-(t', 0) \rangle_B = q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} (-m_{\mathbf{k}'}(t')) (N_{\mathbf{k}'} + 1) B_{01}(t'), \quad (753)$$

$$\langle B_{i\text{mod}2z}(t, \tau) B_x(t', 0) \rangle_B = \frac{1}{2} \sum_{\mathbf{k}'} \left( -m_{\mathbf{k}'}^*(t') q_{i\mathbf{k}'}(t) B_{10}(t') N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} - (-m_{\mathbf{k}'}^*(t')) q_{i\mathbf{k}'}(t) B_{01}(t') N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} \right. \quad (754)$$

$$\left. + q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{10}(t') + q_{i\mathbf{k}'}^*(t) e^{-i\omega_{\mathbf{k}'}\tau} (-m_{\mathbf{k}'}(t')) (N_{\mathbf{k}'} + 1) B_{01}(t') \right) \quad (755)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t) N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}^*(t') (B_{01}(t') - B_{10}(t')) + q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') e^{-i\omega_{\mathbf{k}'}\tau} (N_{\mathbf{k}'} + 1) (B_{10}(t') - B_{01}(t')) \right) \quad (756)$$

$$+ 1) (B_{10}(t') - B_{01}(t')) \quad (757)$$

$$= \frac{1}{2} \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t) N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}^*(t') (B_{01}(t') - B_{10}(t')) - q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') e^{-i\omega_{\mathbf{k}'}\tau} (N_{\mathbf{k}'} + 1) (B_{01}(t') - B_{10}(t')) \right) \quad (758)$$

$$+ 1) (B_{01}(t') - B_{10}(t')) \quad (759)$$

$$= i \sum_{\mathbf{k}'} B_{01}^{\mathfrak{S}}(t') \left( q_{i\mathbf{k}'}(t) N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}^*(t') - q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') e^{-i\omega_{\mathbf{k}'}\tau} (N_{\mathbf{k}'} + 1) \right), \quad (760)$$

$$\langle B_y(t, \tau) B_{i\text{mod}2z}(t', 0) \rangle_B = \left\langle \left( \frac{B_0^+ B_1^-(t, \tau) - B_1^+ B_0^-(t, \tau) + B_{10}(t) - B_{01}(t)}{2i} \right) \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right) \right\rangle_B \quad (761)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left\langle (B_0^+ B_1^-(t, \tau) - B_1^+ B_0^-(t, \tau) + B_{10}(t) - B_{01}(t)) \left( q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right) \right\rangle_B \quad (762)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left\langle (B_0^+ B_1^-(t, \tau) - B_1^+ B_0^-(t, \tau)) \left( q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right) \right\rangle_B \quad (763)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left\langle B_0^+ B_1^-(t, \tau) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger - B_1^+ B_0^-(t, \tau) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger + B_0^+ B_1^-(t, \tau) q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right. \quad (764)$$

$$\left. - B_1^+ B_0^-(t, \tau) q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \right\rangle, \quad (765)$$

$$\langle B_0^+ B_1^-(t, \tau) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger \rangle_B = -q_{i\mathbf{k}'}(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{01}(t), \quad (766)$$

$$\langle B_0^+ B_1^-(t, \tau) q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \rangle_B = q_{i\mathbf{k}'}^*(t') \left( -m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right) N_{\mathbf{k}'} B_{01}(t), \quad (767)$$

$$\langle B_1^+ B_0^-(t, \tau) q_{i\mathbf{k}'}(t') b_{\mathbf{k}'}^\dagger \rangle_B = -q_{i\mathbf{k}'}(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) B_{10}(t), \quad (768)$$

$$\langle B_1^+ B_0^-(t, \tau) q_{i\mathbf{k}'}^*(t') b_{\mathbf{k}'} \rangle_B = q_{i\mathbf{k}'}^*(t') m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} N_{\mathbf{k}'} B_{10}(t), \quad (769)$$

$$\langle B_y(t, \tau) B_{i\text{mod}2z}(t', 0) \rangle_B = \frac{1}{2i} \sum_{\mathbf{k}'} (B_{01}(t) + B_{10}(t)) \left( q_{i\mathbf{k}'}(t') \left( m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} \right)^* (N_{\mathbf{k}'} + 1) - q_{i\mathbf{k}'}^*(t') m_{\mathbf{k}'}(t) e^{i\omega_{\mathbf{k}'}\tau} N_{\mathbf{k}'} \right) \quad (770)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}^*(t') (N_{\mathbf{k}'} + 1) e^{-i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}^*(t) - q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}(t) \right) (B_{10}(t) + B_{01}(t)), \quad (771)$$

$$= \frac{2}{2i} \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) e^{-i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}^*(t) (B_{10}(t))^{\mathfrak{R}} - q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}(t) (B_{10}(t))^{\mathfrak{R}} \right), \quad (772)$$

$$= \frac{(B_{10}(t))^{\mathfrak{R}}}{i} \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) e^{-i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}^*(t) - q_{i\mathbf{k}'}^*(t') N_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} m_{\mathbf{k}'}(t) \right), \quad (773)$$

$$\langle B_{i\text{mod}2z}(t, \tau) B_y(t', 0) \rangle_B = \left\langle \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} \right) \left( \frac{B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t', 0) + B_{10}(t') - B_{01}(t')}{2i} \right) \right\rangle_B \quad (774)$$

$$= \frac{1}{2i} \left\langle \sum_{\mathbf{k}'} \left( q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} \right) (B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t', 0) + B_{10}(t') \right. \quad (775)$$

$$\left. - B_{01}(t') \right) \rangle_B \quad (776)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left\langle \left( q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} + q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} \right) (B_0^+ B_1^-(t', 0) - B_1^+ B_0^-(t', 0)) \right\rangle_B \quad (777)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left\langle q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} B_0^+ B_1^-(t', 0) - q_{i\mathbf{k}'}(t) b_{\mathbf{k}'}^\dagger e^{i\omega_{\mathbf{k}'}\tau} B_1^+ B_0^-(t', 0) \right. \quad (778)$$

$$+q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} B_0^+ B_1^- (t', 0) - q_{i\mathbf{k}'}^*(t) b_{\mathbf{k}'} e^{-i\omega_{\mathbf{k}'}\tau} B_1^+ B_0^- (t', 0) \rangle \quad (779)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left\langle e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) \langle b_{\mathbf{k}'}^\dagger B_0^+ B_1^- (t', 0) \rangle - e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) \langle b_{\mathbf{k}'}^\dagger B_1^+ B_0^- (t', 0) \rangle \right. \quad (780)$$

$$\left. + e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) \langle b_{\mathbf{k}'} B_0^+ B_1^- (t', 0) \rangle - e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) \langle b_{\mathbf{k}'} B_1^+ B_0^- (t', 0) \rangle \right\rangle \quad (781)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) \langle b_{\mathbf{k}'}^\dagger B_0^+ B_1^- (t', 0) \rangle - e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) \langle b_{\mathbf{k}'}^\dagger B_1^+ B_0^- (t', 0) \rangle + e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) \langle b_{\mathbf{k}'} B_0^+ B_1^- (t', 0) \rangle \quad (782)$$

$$- e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) \langle b_{\mathbf{k}'} B_1^+ B_0^- (t', 0) \rangle \quad (783)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left( e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) \langle b_{\mathbf{k}'}^\dagger B_0^+ B_1^- (t', 0) \rangle - e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) \langle b_{\mathbf{k}'}^\dagger B_1^+ B_0^- (t', 0) \rangle + e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) \langle b_{\mathbf{k}'} B_0^+ B_1^- (t', 0) \rangle \right. \quad (784)$$

$$\left. - e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) \langle b_{\mathbf{k}'} B_1^+ B_0^- (t', 0) \rangle \right) \quad (785)$$

$$\langle b_{\mathbf{k}'}^\dagger B_1^+ B_0^- (t', 0) \rangle_B = -m_{\mathbf{k}'}^*(t') B_{10}(t') N_{\mathbf{k}'}, \quad (786)$$

$$\langle b_{\mathbf{k}'}^\dagger B_0^+ B_1^- (t', 0) \rangle_B = m_{\mathbf{k}'}^*(t') B_{01}(t') N_{\mathbf{k}'}, \quad (787)$$

$$\langle b_{\mathbf{k}'} B_1^+ B_0^- (t', 0) \rangle_B = m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{10}(t'), \quad (788)$$

$$\langle b_{\mathbf{k}'} B_0^+ B_1^- (t', 0) \rangle_B = -m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{01}(t'), \quad (789)$$

$$\langle B_{i\text{mod}2z}(t, \tau) B_y(t', 0) \rangle_B = \frac{1}{2i} \sum_{\mathbf{k}'} \left( e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) (-(-m_{\mathbf{k}'}^*(t')) B_{01}(t') N_{\mathbf{k}'}) - e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) (-m_{\mathbf{k}'}^*(t') B_{10}(t') N_{\mathbf{k}'}) \right. \quad (790)$$

$$\left. + e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) (-m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{01}(t')) - e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{10}(t') \right) \quad (791)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} \left( e^{i\omega_{\mathbf{k}'}\tau} (-q_{i\mathbf{k}'}(t) (-m_{\mathbf{k}'}^*(t')) B_{01}(t') N_{\mathbf{k}'} + q_{i\mathbf{k}'}(t) m_{\mathbf{k}'}^*(t') B_{10}(t') N_{\mathbf{k}'}) \right. \quad (792)$$

$$\left. + e^{-i\omega_{\mathbf{k}'}\tau} (q_{i\mathbf{k}'}^*(t) (-m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{01}(t') - q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) B_{10}(t')) \right) \quad (793)$$

$$= \frac{1}{2i} \sum_{\mathbf{k}'} (B_{10}(t') + B_{01}(t')) \left( e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) m_{\mathbf{k}'}^*(t') N_{\mathbf{k}'} - e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) \right) \quad (794)$$

$$= \frac{1}{i} \sum_{\mathbf{k}'} \left( e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) m_{\mathbf{k}'}^*(t') B_{10}^{\Re}(t') N_{\mathbf{k}'} - e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') B_{10}^{\Re}(t') (N_{\mathbf{k}'} + 1) \right) \quad (795)$$

$$= i \sum_{\mathbf{k}'} \left( e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') B_{10}^{\Re}(t') (N_{\mathbf{k}'} + 1) - e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) m_{\mathbf{k}'}^*(t') B_{10}^{\Re}(t') N_{\mathbf{k}'} \right) \quad (796)$$

$$= i \sum_{\mathbf{k}'} \left( e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') B_{10}^{\Re}(t') (N_{\mathbf{k}'} + 1) - e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) m_{\mathbf{k}'}^*(t') B_{10}^{\Re}(t') N_{\mathbf{k}'} \right) \quad (797)$$

$$= i B_{10}^{\Re}(t') \sum_{\mathbf{k}'} \left( e^{-i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}^*(t) m_{\mathbf{k}'}(t') (N_{\mathbf{k}'} + 1) - e^{i\omega_{\mathbf{k}'}\tau} q_{i\mathbf{k}'}(t) m_{\mathbf{k}'}^*(t') N_{\mathbf{k}'} \right). \quad (798)$$

The correlation functions are equal to:

$$\langle \widetilde{B_{i\text{mod}2z}}(t) \widetilde{B_{j\text{mod}2z}}(t') \rangle_B = \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) (g_{j\mathbf{k}} - v_{j\mathbf{k}}(t'))^* e^{i\omega_{\mathbf{k}}\tau} N_{\mathbf{k}} + \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* (g_{j\mathbf{k}} - v_{j\mathbf{k}}(t')) e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1), \quad (799)$$

$$\langle \widetilde{B_x}(t) \widetilde{B_x}(t') \rangle_B = \frac{1}{2} \left( \left( e^{\chi_{10}(t) + \chi_{10}(t')} \right)^{\Re} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (800)$$

$$\left. + \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}^*(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (801)$$

$$- \left( e^{\chi_{10}(t)} e^{-\frac{1}{2} \sum_{\mathbf{k}} \left( \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \right) \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Re} \left( e^{\chi_{01}(t')} e^{-\frac{1}{2} \sum_{\mathbf{k}} \left( \left| \frac{v_{1\mathbf{k}}(t')}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right|^2 \right) \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right)^{\Re} \quad (802)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_y(t') \rangle_B = -\frac{1}{2} \left( \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (803)$$

$$\left. - \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}^*(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \quad (804)$$

$$+ \left( e^{\chi_{01}(t)} \left( e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \right)^{\Im} \left( e^{\chi_{10}(t')} \left( e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t')}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) \right)^{\Im} \quad (805)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_y(t') \rangle_B = \frac{1}{2} \left( \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}^*(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (806)$$

$$+ \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) + (B_{10}(t))^{\Re} (B_{10}(t'))^{\Im} \quad (807)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_x(t') \rangle_B = \frac{1}{2} \left( \left( e^{\chi_{01}(t) + \chi_{10}(t')} \right)^{\Im} U_{10}^*(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right. \quad (808)$$

$$+ \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}(t, t') \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right) + (B_{10}(t))^{\Im} (B_{10}(t'))^{\Re} \quad (809)$$

$$\langle \widetilde{B}_{i\text{mod}2z}(t) \widetilde{B}_x(t') \rangle_B = i \sum_{\mathbf{k}} B_{01}^{\Im}(t') \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) N_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \left( \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right)^* - (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1) \right), \quad (810)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_{i\text{mod}2z}(t') \rangle_B = i B_{10}^{\Im}(t) \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t'))^* N_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) - (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t')) \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right)^* e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1) \right), \quad (811)$$

$$\langle \widetilde{B}_{i\text{mod}2z}(t) \widetilde{B}_y(t') \rangle_B = i B_{10}^{\Re}(t') \sum_{\mathbf{k}} \left( e^{-i\omega_{\mathbf{k}}\tau} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \left( \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right) (N_{\mathbf{k}} + 1) - e^{i\omega_{\mathbf{k}}\tau} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \left( \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right)^* N_{\mathbf{k}} \right), \quad (812)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_{i\text{mod}2z}(t') \rangle_B = i B_{10}^{\Re}(t) \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t'))^* N_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) - (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t')) (N_{\mathbf{k}} + 1) e^{-i\omega_{\mathbf{k}}\tau} \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right)^* \right). \quad (813)$$

Let's consider the following expression related to the sum of coupling constants for a bath over all the frequencies:

$$L_i(\omega) \equiv \sum_{\mathbf{k}} g_{i\mathbf{k}} \sqrt{\delta(\omega - \omega_{\mathbf{k}})}. \quad (814)$$

Under this definition we have the following expression for a function  $f(\omega) \in L^2$ :

$$\int_0^\infty f(\omega) L_i(\omega) L_j^*(\omega) d\omega \approx \int_0^\infty f(\omega) \sum_{\mathbf{k}} g_i(\omega_{\mathbf{k}}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} \sum_{\mathbf{k}'} g_j(\omega_{\mathbf{k}'}) \sqrt{\delta(\omega - \omega_{\mathbf{k}'})} d\omega \quad (815)$$

$$= \int_0^\infty f(\omega) \sum_{\mathbf{k}, \mathbf{k}'} g_i(\omega_{\mathbf{k}}) g_j(\omega_{\mathbf{k}'}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} \sqrt{\delta(\omega - \omega_{\mathbf{k}'})} d\omega, \quad (816)$$

$$\int_0^\infty f(\omega) \sum_{\mathbf{k}} g_i(\omega_{\mathbf{k}}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} d\omega = \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} d\omega. \quad (817)$$

Now we will approach to the function  $\sqrt{\delta(\omega - \omega_{\mathbf{k}})}$  using the normal distribution, so:

$$\delta(\omega - \omega_{\mathbf{k}}) = \lim_{\sigma \rightarrow 0^+} \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(\omega - \omega_{\mathbf{k}})^2}{2\sigma^2}} \quad (818)$$

$$\sqrt{\delta(\omega - \omega_{\mathbf{k}})} = \lim_{\sigma \rightarrow 0^+} \sqrt{\frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(\omega - \omega_{\mathbf{k}})^2}{2\sigma^2}}} \quad (819)$$

$$= \lim_{\sigma \rightarrow 0^+} \sqrt{\sqrt{2\pi}\sigma} \frac{\sqrt{2}}{\sqrt{2\pi}\sqrt{2\sigma}} e^{-\frac{(\omega - \omega_{\mathbf{k}})^2}{4\sigma^2}} \quad (820)$$

$$= \lim_{\sigma \rightarrow 0^+} \sqrt{\sqrt{2\pi}\sigma} \frac{\sqrt{2}}{\sqrt{2\pi}\sqrt{2\sigma}} e^{-\frac{(\omega - \omega_{\mathbf{k}})^2}{2(\sqrt{2}\sigma)^2}} \quad (821)$$

$$= \lim_{\sigma \rightarrow 0^+} \sqrt{2} \sqrt{\sqrt{2\pi}\sigma} N(x; \omega_{\mathbf{k}}, \sqrt{2}\sigma). \quad (822)$$

So we can obtain that:

$$\sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} d\omega = \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} d\omega \quad (823)$$

$$= \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) \left( \lim_{\sigma \rightarrow 0^+} \sqrt{2} \sqrt{\sqrt{2\pi}\sigma} N(x; \omega_{\mathbf{k}}, \sqrt{2}\sigma) \right) d\omega \quad (824)$$

$$= \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) \left( \lim_{\sigma \rightarrow 0^+} \sqrt{2} \sqrt{\sqrt{2\pi}\sigma} N(x; \omega_{\mathbf{k}}, \sqrt{2}\sigma) \right) d\omega \quad (825)$$

$$= \lim_{\sigma \rightarrow 0^+} \sqrt{2} \sqrt{\sqrt{2\pi}\sigma} \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) N(x; \omega_{\mathbf{k}}, \sqrt{2}\sigma) d\omega \quad (826)$$

$$= \sum_{\mathbf{k}} \left( \lim_{\sigma \rightarrow 0^+} \sqrt{2} \sqrt{\sqrt{2\pi}\sigma} \right) \left( \lim_{\sigma \rightarrow 0^+} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) N(x; \omega_{\mathbf{k}}, \sqrt{2}\sigma) d\omega \right) \quad (827)$$

$$= \sum_{\mathbf{k}} \left( \lim_{\sigma \rightarrow 0^+} \sqrt{2} \sqrt{\sqrt{2\pi}\sigma} \right) f(\omega_{\mathbf{k}}) g_i(\omega_{\mathbf{k}}) \text{ (with } f(\omega), g_i(\omega) \in L^2 \text{)} \quad (828)$$

$$= \lim_{\sigma \rightarrow 0^+} \sqrt{2} \sqrt{\sqrt{2\pi}\sigma} \sum_{\mathbf{k}} f(\omega_{\mathbf{k}}) g_i(\omega_{\mathbf{k}}) \text{ (with } f(\omega), g_i(\omega) \in L^2 \text{)} \quad (829)$$

$$= 0 \text{ (because the sum } \sum_{\mathbf{k}} f(\omega_{\mathbf{k}}) g_i(\omega_{\mathbf{k}}) \text{ is finite).} \quad (830)$$

Then we can proof the following:

$$\int_0^\infty f(\omega) L_i(\omega) L_j^*(\omega) d\omega \approx \int_0^\infty f(\omega) \sum_{\mathbf{k}} g_i(\omega_{\mathbf{k}}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} \sum_{\mathbf{k}'} g_j^*(\omega_{\mathbf{k}'}) \sqrt{\delta(\omega - \omega_{\mathbf{k}'})} d\omega \quad (831)$$

$$= \int_0^\infty f(\omega) \sum_{\mathbf{k}, \mathbf{k}'} g_i(\omega_{\mathbf{k}}) g_j^*(\omega_{\mathbf{k}'}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} \sqrt{\delta(\omega - \omega_{\mathbf{k}'})} d\omega \quad (832)$$

$$= \sum_{\mathbf{k} \neq \mathbf{k}'} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) g_j^*(\omega_{\mathbf{k}'}) \sqrt{\delta(\omega - \omega_{\mathbf{k}})} \sqrt{\delta(\omega - \omega_{\mathbf{k}'})} d\omega + \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) g_j^*(\omega_{\mathbf{k}}) \delta(\omega - \omega_{\mathbf{k}}) d\omega \quad (833)$$

$$= 0 + \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) g_j^*(\omega_{\mathbf{k}}) \delta(\omega - \omega_{\mathbf{k}}) d\omega \quad (834)$$

$$= \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) g_j^*(\omega_{\mathbf{k}}) \delta(\omega - \omega_{\mathbf{k}}) d\omega \quad (835)$$

$$= \sum_{\mathbf{k}} f(\omega_{\mathbf{k}}) g_i(\omega_{\mathbf{k}}) g_j^*(\omega_{\mathbf{k}}) \quad (836)$$

if  $i = j$  we recover the spectral density defined in the usual way when we integrate for a function  $f(\omega)$  that belongs to the set  $L^2$ :



$$\sum_{\mathbf{k}} f(\omega_{\mathbf{k}}) g_i(\omega_{\mathbf{k}}) g_i^*(\omega_{\mathbf{k}}) = \sum_{\mathbf{k}} \int_0^\infty f(\omega) g_i(\omega_{\mathbf{k}}) g_i^*(\omega_{\mathbf{k}}) \delta(\omega - \omega_{\mathbf{k}}) d\omega \quad (837)$$

$$= \int_0^\infty f(\omega) J_{ii}(\omega) d\omega \quad (838)$$

$$= \int_0^\infty f(\omega) |L_i(\omega)|^2 d\omega. \quad (839)$$

where

$$J_{ii}(\omega) = \sum_{\mathbf{k}} |g_{i\mathbf{k}}|^2 \delta(\omega - \omega_{\mathbf{k}}), \quad (840)$$

$$v_{i\mathbf{k}}(\omega_{\mathbf{k}}, t) = g_{i\mathbf{k}} F_i(\omega_{\mathbf{k}}, t). \quad (841)$$

In this case  $g_i(\omega)$  and  $v_i(\omega, t)$  are the continuous version of  $g_i(\omega_{\mathbf{k}})$  and  $v_{i\mathbf{k}}(\omega_{\mathbf{k}}, t)$  respectively.

The integral version of the correlation functions can be obtained as follows:

$$\langle \widetilde{B_{iz}}(t) \widetilde{B_{j \bmod 2z}}(t') \rangle_B = \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) (g_{j\mathbf{k}} - v_{j\mathbf{k}}(t'))^* e^{i\omega_{\mathbf{k}}\tau} N_{\mathbf{k}} + \sum_{\mathbf{k}} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* (g_{j\mathbf{k}} - v_{j\mathbf{k}}(t')) e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1), \quad (842)$$

$$= \sum_{\mathbf{k}} \left( g_{i\mathbf{k}} (1 - F_i(\omega_{\mathbf{k}}, t)) g_{j\mathbf{k}}^* (1 - F_j(\omega_{\mathbf{k}}, t'))^* e^{i\omega_{\mathbf{k}}\tau} N_{\mathbf{k}} + g_{i\mathbf{k}}^* (1 - F_i(\omega_{\mathbf{k}}, t))^* g_{j\mathbf{k}} (1 - F_j(\omega_{\mathbf{k}}, t')) e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1) \right) \quad (843)$$

$$\approx \int_0^\infty \left( L_i(\omega) L_j^*(\omega) (1 - F_i(\omega, t)) (1 - F_j^*(\omega, t')) e^{i\omega\tau} N(\omega) + L_i^*(\omega) L_j(\omega) (1 - F_i^*(\omega, t)) (1 - F_j(\omega, t')) e^{-i\omega\tau} (N(\omega) + 1) \right) d\omega, \quad (844)$$

$$\chi_{10}(t) = \sum_{\mathbf{k}} \frac{1}{2} \left( \frac{v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right) \quad (845)$$

$$= \sum_{\mathbf{k}} \frac{1}{2} \left( \frac{g_{1\mathbf{k}}^* F_1^*(\omega_{\mathbf{k}}, t) g_{0\mathbf{k}} F_0(\omega_{\mathbf{k}}, t) - g_{1\mathbf{k}} F_1(\omega_{\mathbf{k}}, t) g_{0\mathbf{k}}^* F_0^*(\omega_{\mathbf{k}}, t)}{\omega_{\mathbf{k}}^2} \right) \quad (846)$$

$$= \sum_{\mathbf{k}} \frac{1}{2} \left( \frac{g_{1\mathbf{k}}^* g_{0\mathbf{k}} F_1^*(\omega_{\mathbf{k}}, t) F_0(\omega_{\mathbf{k}}, t) - g_{1\mathbf{k}} g_{0\mathbf{k}}^* F_1(\omega_{\mathbf{k}}, t) F_0^*(\omega_{\mathbf{k}}, t)}{\omega_{\mathbf{k}}^2} \right) \quad (847)$$

$$\approx \int_0^\infty \frac{L_0(\omega) L_1^*(\omega) F_1^*(\omega, t) F_0(\omega, t) - L_1(\omega) L_0^*(\omega) F_1(\omega, t) F_0^*(\omega, t)}{2\omega^2} d\omega, \quad (848)$$

$$U_{10}(t, t') = \prod_{\mathbf{k}} e^{i \left( \frac{(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))(v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))^* e^{i\omega_{\mathbf{k}}\tau}}{\omega_{\mathbf{k}}^2} \right)} \quad (849)$$

$$= e^{i \sum_{\mathbf{k}} \left( \frac{(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))(v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))^* e^{i\omega_{\mathbf{k}}\tau}}{\omega_{\mathbf{k}}^2} \right)} \quad (850)$$

$$= e^{i \left( \sum_{\mathbf{k}} \frac{(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))(v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))^* e^{i\omega_{\mathbf{k}}\tau}}{\omega_{\mathbf{k}}^2} \right)} \quad (851)$$

$$= e^{i \left( \sum_{\mathbf{k}} \frac{(g_{1\mathbf{k}} F_1(\omega_{\mathbf{k}}, t) - g_{0\mathbf{k}} F_0(\omega_{\mathbf{k}}, t))(g_{1\mathbf{k}} F_1(\omega_{\mathbf{k}}, t') - g_{0\mathbf{k}} F_0(\omega_{\mathbf{k}}, t'))^* e^{i\omega_{\mathbf{k}}\tau}}{\omega_{\mathbf{k}}^2} \right)} \quad (852)$$

$$\approx e^{i \left( \int_0^\infty \frac{(L_1(\omega) F_1(\omega, t) - L_0(\omega) F_0(\omega, t))(L_1(\omega) F_1(\omega, t') - L_0(\omega) F_0(\omega, t'))^* e^{i\omega\tau}}{\omega^2} d\omega \right)} \quad (853)$$

$$B_{10}(t) = \left( \prod_{\mathbf{k}} e^{\frac{1}{2} \left( \frac{v_{1\mathbf{k}}^*(t) v_{0\mathbf{k}}(t) - v_{1\mathbf{k}}(t) v_{0\mathbf{k}}^*(t)}{\omega_{\mathbf{k}}^2} \right)} \right) \left( e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{v_{1\mathbf{k}}(t)}{\omega_{\mathbf{k}}} - \frac{v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \right), \quad (854)$$

$$= e^{\chi_{10}(t)} e^{-\frac{1}{2} \sum_{\mathbf{k}} \left| \frac{g_{1\mathbf{k}} F_1(\omega_{\mathbf{k}}, t) - g_{0\mathbf{k}} F_0(\omega_{\mathbf{k}}, t)}{\omega_{\mathbf{k}}} \right|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (855)$$

$$\approx e^{\chi_{10}(t)} e^{-\frac{1}{2} \int_0^\infty \left| \frac{L_1(\omega) F_1(\omega, t) - L_0(\omega) F_0(\omega, t)}{\omega} \right|^2 \coth\left(\frac{\beta\omega}{2}\right) d\omega} \quad (856)$$

$$\xi^+(t, t') = \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (857)$$

$$= e^{-\sum_{\mathbf{k}} \frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (858)$$

$$= e^{-\sum_{\mathbf{k}} \frac{|(g_{1\mathbf{k}}F_1(\omega_{\mathbf{k}}, t) - g_{0\mathbf{k}}F_0(\omega_{\mathbf{k}}, t))e^{i\omega_{\mathbf{k}}\tau} + g_{1\mathbf{k}}F_1(\omega_{\mathbf{k}}, t') - g_{0\mathbf{k}}F_0(\omega_{\mathbf{k}}, t')|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (859)$$

$$\approx e^{-\int_0^\infty \frac{|(L_1(\omega)F_1(\omega, t) - L_0(\omega)F_0(\omega, t))e^{i\omega\tau} + L_1(\omega)F_1(\omega, t') - L_0(\omega)F_0(\omega, t')|^2}{2\omega^2} \coth\left(\frac{\beta\omega}{2}\right) d\omega} \quad (860)$$

$$\xi^-(t, t') = \prod_{\mathbf{k}} e^{-\frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (861)$$

$$= e^{-\sum_{\mathbf{k}} \frac{|(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t))e^{i\omega_{\mathbf{k}}\tau} - (v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (862)$$

$$= e^{-\sum_{\mathbf{k}} \frac{|(g_{1\mathbf{k}}F_1(\omega_{\mathbf{k}}, t) - g_{0\mathbf{k}}F_0(\omega_{\mathbf{k}}, t))e^{i\omega_{\mathbf{k}}\tau} - (g_{1\mathbf{k}}F_1(\omega_{\mathbf{k}}, t') - g_{0\mathbf{k}}F_0(\omega_{\mathbf{k}}, t'))|^2}{2\omega_{\mathbf{k}}^2} \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)} \quad (863)$$

$$\approx e^{-\int_0^\infty \frac{|(L_1(\omega)F_1(\omega, t) - L_0(\omega)F_0(\omega, t))e^{i\omega\tau} - (L_1(\omega)F_1(\omega, t') - L_0(\omega)F_0(\omega, t'))|^2}{2\omega^2} \coth\left(\frac{\beta\omega}{2}\right) d\omega} \quad (864)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_x(t') \rangle_B = \frac{1}{2} \left( \left( e^{\chi_{10}(t) + \chi_{10}(t')} \right)^{\Re} U_{10}(t, t') \xi^+(t, t') + \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}^*(t, t') \xi^-(t, t') \right) - (B_{10}(t))^{\Re} (B_{01}(t'))^{\Re} \quad (865)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_y(t') \rangle_B = -\frac{1}{2} \left( \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}(t, t') \xi^+(t, t') - \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Re} U_{10}^*(t, t') \xi^-(t, t') \right) + (B_{01}(t))^{\Im} (B_{10}(t'))^{\Im} \quad (866)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_y(t') \rangle_B = \frac{1}{2} \left( \left( e^{\chi_{10}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}^*(t, t') \xi^-(t, t') + \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}(t, t') \xi^+(t, t') \right) + (B_{10}(t))^{\Re} (B_{10}(t'))^{\Im} \quad (867)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_x(t') \rangle_B = \frac{1}{2} \left( \left( e^{\chi_{01}(t) + \chi_{10}(t')} \right)^{\Im} U_{10}^*(t, t') \xi^-(t, t') + \left( e^{\chi_{01}(t) + \chi_{01}(t')} \right)^{\Im} U_{10}(t, t') \xi^+(t, t') \right) + (B_{10}(t))^{\Im} (B_{10}(t'))^{\Re} \quad (868)$$

$$\langle \widetilde{B}_{iz}(t) \widetilde{B}_x(t') \rangle_B = iB_{01}^{\Im}(t') \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \left( \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right)^* N_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} - (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1) \right), \quad (869)$$

$$= iB_{01}^{\Im}(t') \sum_{\mathbf{k}} \left( -g_{i\mathbf{k}}^* (1 - F_i(\omega_{\mathbf{k}}, t))^* \frac{g_{1\mathbf{k}}F_1(\omega_{\mathbf{k}}, t') - g_{0\mathbf{k}}F_0(\omega_{\mathbf{k}}, t')}{\omega_{\mathbf{k}}} e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1) \right. \quad (870)$$

$$\left. + g_{i\mathbf{k}} (1 - F_i(\omega_{\mathbf{k}}, t)) \left( \frac{g_{1\mathbf{k}}F_1(\omega_{\mathbf{k}}, t') - g_{0\mathbf{k}}F_0(\omega_{\mathbf{k}}, t')}{\omega_{\mathbf{k}}} \right)^* N_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \right), \quad (871)$$

$$Q(\omega, t) = \frac{L_1(\omega)F_1(\omega, t) - L_0(\omega)F_0(\omega, t)}{\omega} \quad (872)$$

$$\langle \widetilde{B}_{iz}(t) \widetilde{B}_x(t') \rangle_B \approx iB_{01}^{\Im}(t') \int_0^\infty \left( L_i(\omega) (1 - F_i(\omega, t)) Q^*(\omega, t') N(\omega) e^{i\omega\tau} - L_i^*(\omega) (1 - F_i^*(\omega, t)) Q(\omega, t') e^{-i\omega\tau} (N(\omega) + 1) \right) d\omega, \quad (873)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_{iz}(t') \rangle_B = iB_{10}^{\Im}(t) \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t'))^* N_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) - (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right)^* e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1) \right), \quad (874)$$

$$\approx iB_{01}^{\Im}(t) \int_0^\infty \left( L_i^*(\omega) (1 - F_i^*(\omega, t')) Q(\omega, t) N(\omega) e^{i\omega\tau} - L_i(\omega) (1 - F_i(\omega, t')) Q^*(\omega, t) e^{-i\omega\tau} (N(\omega) + 1) \right) d\omega \quad (875)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_y(t') \rangle_B = iB_{10}^{\Re}(t') \sum_{\mathbf{k}} \left( e^{-i\omega_{\mathbf{k}}\tau} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t))^* \left( \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right) (N_{\mathbf{k}} + 1) - e^{i\omega_{\mathbf{k}}\tau} (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t)) \left( \frac{v_{1\mathbf{k}}(t') - v_{0\mathbf{k}}(t')}{\omega_{\mathbf{k}}} \right)^* N_{\mathbf{k}} \right), \quad (876)$$

$$\approx iB_{10}^{\Re}(t') \int_0^\infty \left( L_i^*(\omega) (1 - F_i^*(\omega, t')) Q(\omega, t') (N(\omega) + 1) e^{-i\omega\tau} - L_i(\omega) (1 - F_i(\omega, t')) Q^*(\omega, t') e^{i\omega\tau} N(\omega) \right) d\omega \quad (877)$$

$$\langle \widetilde{B}_y(t) \widetilde{B}_{iz}(t') \rangle_B = iB_{10}^{\Re}(t) \sum_{\mathbf{k}} \left( (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t'))^* N_{\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right) - (g_{i\mathbf{k}} - v_{i\mathbf{k}}(t')) (N_{\mathbf{k}} + 1) e^{-i\omega_{\mathbf{k}}\tau} \left( \frac{v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)}{\omega_{\mathbf{k}}} \right)^* \right) \quad (878)$$

$$\approx iB_{10}^{\Re}(t) \int_0^\infty \left( L_i^*(\omega) (1 - F_i^*(\omega, t')) Q(\omega, t) N(\omega) e^{i\omega\tau} - L_i(\omega) (1 - F_i(\omega, t')) Q^*(\omega, t) e^{-i\omega\tau} (N(\omega) + 1) \right) d\omega. \quad (879)$$

The integral version of  $F_0(\omega, t)$  and  $F_1(\omega, t)$  are:

$$a_i(\omega_{\mathbf{k}}, t) = \frac{\left( 1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t)) \right)}{1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left( \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|B_{10}(t)|^2|V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right)}, \quad (880)$$

$$b_i(\omega_{\mathbf{k}}, t) = \frac{2 \frac{\tanh(\frac{\beta\eta(t)}{2})}{\eta(t)} \frac{1}{\omega_{\mathbf{k}}} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{1 - \frac{\tanh(\frac{\beta\eta(t)}{2})}{\eta(t)} \left( \varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega_{\mathbf{k}}}{2}\right)}{\omega_{\mathbf{k}}} \right)}, \quad (881)$$

$$r_i(\omega_{\mathbf{k}}, t) = \frac{a_i(\omega_{\mathbf{k}}, t)}{1 - b_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)}, \quad (882)$$

$$s_i(\omega_{\mathbf{k}}, t) = \frac{a_{(i+1) \bmod 2}(\omega_{\mathbf{k}}, t) b_{i \bmod 2}(\omega_{\mathbf{k}}, t)}{1 - b_0(\omega_{\mathbf{k}}, t) b_1(\omega_{\mathbf{k}}, t)}. \quad (883)$$

$$F_0(\omega, t) = r_0(\omega_{\mathbf{k}}, t) + \frac{g_1(\omega_{\mathbf{k}})}{g_0(\omega_{\mathbf{k}})} s_0(\omega_{\mathbf{k}}, t) \quad (884)$$

$$\approx r_0(\omega, t) + \frac{g_1(\omega)}{g_0(\omega)} s_0(\omega, t) \quad (885)$$

$$= r_0(\omega, t) + \frac{L_1(\omega)}{L_0(\omega)} s_0(\omega, t), \quad (886)$$

$$F_1(\omega, t) = \frac{g_0(\omega_{\mathbf{k}})}{g_1(\omega_{\mathbf{k}})} r_1(\omega_{\mathbf{k}}, t) + s_1(\omega_{\mathbf{k}}, t) \quad (887)$$

$$\approx \frac{g_0(\omega)}{g_1(\omega)} r_1(\omega, t) + s_1(\omega, t) \quad (888)$$

$$= \frac{L_0(\omega)}{L_1(\omega)} r_1(\omega, t) + s_1(\omega, t). \quad (889)$$

The expression showed are well defined because the relevant products present in the correlations functions are of the form:

$$\int_0^\infty f(\omega) L_j(\omega) F_j(\omega, t) L_i^*(\omega) F_i^*(\omega, t) d\omega = \int_0^\infty f(\omega) L_j(\omega) \left( r_j(\omega, t) + \frac{L_j(\omega)}{L_j(\omega)} s_j(\omega, t) \right) L_i^*(\omega) \left( r_i^*(\omega, t) + \frac{L_j^*(\omega)}{L_i^*(\omega)} s_i^*(\omega, t) \right) d\omega \quad (890)$$

$$= \int_0^\infty f(\omega) (L_j(\omega) r_j(\omega, t) + L_i(\omega) s_j(\omega, t)) (L_i^*(\omega) r_i^*(\omega, t) + L_j^*(\omega) s_i^*(\omega, t)) d\omega \quad (891)$$

$$= \int_0^\infty f(\omega) (L_j(\omega) L_i^*(\omega) r_j(\omega, t) r_i^*(\omega, t) + |L_j(\omega)|^2 r_j(\omega, t) s_i^*(\omega, t) \quad (892)$$

$$+ |L_i(\omega)|^2 s_j(\omega, t) r_i^*(\omega, t) + L_i(\omega) L_j^*(\omega) s_j(\omega, t) s_i^*(\omega, t)) d\omega \quad (893)$$

here  $f(\omega) \in L^2$ . As we could proof these integral are convergent.

So the integral version of the correlation functions  $\mathcal{B}_{ij}(t, t')$  is can be written in a neater form as:

$$\mathcal{B}(t, t') = \begin{pmatrix} \mathcal{B}_{11}(t, t') & \mathcal{B}_{12}(t, t') & \mathcal{B}_{13}(t, t') & \mathcal{B}_{12}(t, t') & \mathcal{B}_{11}(t, t') & \mathcal{B}_{16}(t, t') \\ \mathcal{B}_{21}(t, t') & \mathcal{B}_{22}(t, t') & \mathcal{B}_{23}(t, t') & \mathcal{B}_{22}(t, t') & \mathcal{B}_{21}(t, t') & \mathcal{B}_{26}(t, t') \\ \mathcal{B}_{31}(t, t') & \mathcal{B}_{32}(t, t') & \mathcal{B}_{33}(t, t') & \mathcal{B}_{32}(t, t') & \mathcal{B}_{31}(t, t') & \mathcal{B}_{36}(t, t') \\ \mathcal{B}_{21}(t, t') & \mathcal{B}_{22}(t, t') & \mathcal{B}_{23}(t, t') & \mathcal{B}_{22}(t, t') & \mathcal{B}_{21}(t, t') & \mathcal{B}_{26}(t, t') \\ \mathcal{B}_{11}(t, t') & \mathcal{B}_{12}(t, t') & \mathcal{B}_{13}(t, t') & \mathcal{B}_{12}(t, t') & \mathcal{B}_{11}(t, t') & \mathcal{B}_{16}(t, t') \\ \mathcal{B}_{61}(t, t') & \mathcal{B}_{62}(t, t') & \mathcal{B}_{63}(t, t') & \mathcal{B}_{62}(t, t') & \mathcal{B}_{61}(t, t') & \mathcal{B}_{66}(t, t') \end{pmatrix}, \quad (894)$$

$$\mathcal{B}_{11}(t, t') = \frac{1}{2} \left( \Re \left( e^{X_{10}(t) + X_{10}(t')} \right) \zeta_{10}(t, t') \xi_{10}^+(t, t') + \Re \left( e^{X_{10}(t) + X_{01}(t')} \right) \zeta_{10}^*(t, t') \xi_{10}^-(t, t') \right) - B_{10}^{\Re}(t) B_{01}^{\Re}(t'), \quad (895)$$

$$\mathcal{B}_{22}(t, t') = -\frac{1}{2} \left( \Re \left( e^{X_{01}(t) + X_{01}(t')} \right) \zeta_{10}(t, t') \xi_{10}^+(t, t') - \Re \left( e^{X_{10}(t) + X_{01}(t')} \right) \zeta_{10}^*(t, t') \xi_{10}^-(t, t') \right) + B_{01}^{\Im}(t) B_{10}^{\Im}(t'), \quad (896)$$

$$\mathcal{B}_{12}(t, t') = \frac{1}{2} \left( \Im \left( e^{X_{10}(t) + X_{01}(t')} \right) \zeta_{10}^*(t, t') \xi_{10}^-(t, t') + \Im \left( e^{X_{01}(t) + X_{01}(t')} \right) \zeta_{10}(t, t') \xi_{10}^+(t, t') \right) + B_{10}^{\Re}(t) B_{10}^{\Im}(t'), \quad (897)$$

$$\mathcal{B}_{21}(t, t') = \frac{1}{2} \left( \Im \left( e^{X_{01}(t) + X_{10}(t')} \right) \zeta_{10}^*(t, t') \xi_{10}^-(t, t') + \Im \left( e^{X_{01}(t) + X_{01}(t')} \right) \zeta_{10}(t, t') \xi_{10}^+(t, t') \right) + B_{10}^{\Im}(t) B_{10}^{\Re}(t'), \quad (898)$$

$$\begin{aligned}
\mathcal{B}_{ij}(t, t') &= \int_0^\infty (P_i(\omega, t) P_j^*(\omega, t') e^{i\omega\tau} N(\omega) + P_i^*(\omega, t) P_j(\omega, t') e^{-i\omega\tau} (N(\omega) + 1)) d\omega, i, j \in \{3, 6\}, \\
\mathcal{B}_{i1}(t, t') &= iB_{01}^{\mathfrak{S}}(t') \int_0^\infty (P_i(\omega, t) Q_{10}^*(\omega, t') N(\omega) e^{i\omega\tau} - P_i^*(\omega, t) Q_{10}(\omega, t') e^{-i\omega\tau} (N(\omega) + 1)) d\omega, i \in \{3, 6\}, \\
\mathcal{B}_{1i}(t, t') &= iB_{01}^{\mathfrak{S}}(t) \int_0^\infty (P_i^*(\omega, t') Q_{10}(\omega, t) N(\omega) e^{i\omega\tau} - P_i(\omega, t') Q_{10}^*(\omega, t) e^{-i\omega\tau} (N(\omega) + 1)) d\omega, i \in \{3, 6\}, \\
\mathcal{B}_{i2}(t, t') &= iB_{10}^{\mathfrak{R}}(t') \int_0^\infty (P_i^*(\omega, t') Q_{10}(\omega, t') (N(\omega) + 1) e^{-i\omega\tau} - P_i(\omega, t') Q_{10}^*(\omega, t') e^{i\omega\tau} N(\omega)) d\omega, i \in \{3, 6\}, \\
\mathcal{B}_{2i}(t, t') &= iB_{10}^{\mathfrak{R}}(t) \int_0^\infty (P_i^*(\omega, t') Q_{10}(\omega, t) N(\omega) e^{i\omega\tau} - P_i(\omega, t') Q_{10}^*(\omega, t) e^{-i\omega\tau} (N(\omega) + 1)) d\omega, i \in \{3, 6\}, \\
\zeta_{ij}(t, t') &= e^{i\mathfrak{S}} \left( \int_0^\infty \frac{(L_i(\omega) F_i(\omega, t) - L_j(\omega) F_j(\omega, t)) (L_i(\omega) F_i(\omega, t') - L_j(\omega) F_j(\omega, t'))^* e^{i\omega\tau}}{\omega^2} d\omega \right), \\
\xi_{ij}^\pm(t, t') &= e^{-\int_0^\infty \frac{|(L_i(\omega) F_i(\omega, t) - L_j(\omega) F_j(\omega, t)) e^{i\omega\tau} \pm (L_i(\omega) F_i(\omega, t') - L_j(\omega) F_j(\omega, t'))|^2}{2\omega^2} \coth\left(\frac{\beta\omega}{2}\right) d\omega}, \\
P_i(\omega, t) &= L_{i \bmod 2}(\omega) (1 - F_{i \bmod 2}(\omega, t)), \\
Q_{ij}(\omega, t) &= \frac{L_i(\omega) F_1(\omega, t) - L_j(\omega) F_j(\omega, t)}{\omega}, \\
a_i(\omega, t) &= \frac{\left(1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} (2\varepsilon_i(t) + 2R_i(t) - \varepsilon(t))\right)}{1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left(\varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega}{2}\right)}{\omega}\right)}, \\
b_i(\omega, t) &= \frac{2 \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \frac{1}{\omega} |B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega}{2}\right)}{1 - \frac{\tanh\left(\frac{\beta\eta(t)}{2}\right)}{\eta(t)} \left(\varepsilon(t) - 2(\varepsilon(t) - \varepsilon_i(t) - R_i(t)) - \frac{2|B_{10}(t)|^2 |V_{10}(t)|^2 \coth\left(\frac{\beta\omega}{2}\right)}{\omega}\right)}, \\
r_i(\omega, t) &= \frac{a_i(\omega, t)}{1 - b_0(\omega, t) b_1(\omega, t)}, \\
s_i(\omega, t) &= \frac{a_{(i+1) \bmod 2}(\omega, t) b_{i \bmod 2}(\omega, t)}{1 - b_0(\omega, t) b_1(\omega, t)}, \\
F_0(\omega, t) &= r_0(\omega, t) + \frac{L_1(\omega)}{L_0(\omega)} s_0(\omega, t), \\
F_1(\omega, t) &= \frac{L_0(\omega)}{L_1(\omega)} r_1(\omega, t) + s_1(\omega, t).
\end{aligned}$$

The eigenvalues of the Hamiltonian  $\overline{H_S}$  are given by the solution of the following algebraic equation:

$$\lambda^2 - \text{Tr}(\overline{H_S}) \lambda + \text{Det}(\overline{H_S}) = 0. \quad (899)$$

The solutions of this equation written in terms of  $\eta$  and  $\xi$  as defined in the previous section are given by  $\lambda_\pm = \frac{\xi \pm \eta}{2}$  and they satisfy  $H_S |\pm\rangle = \lambda_\pm |\pm\rangle$ . Using this notation is possible to write  $H_S = \lambda_+ |+\rangle\langle+| + \lambda_- |-\rangle\langle-|$ .

The time-dependence of the system operators  $\widetilde{A}_i(t)$  may be made explicit using the Fourier decomposition, in the case for time-independent  $\overline{H_S}$  we will obtain:

$$\widetilde{A}_i(\tau) = e^{i\overline{H_S}\tau} A_i e^{-i\overline{H_S}\tau} \quad (900)$$

$$= \sum_w e^{-i w \tau} \mathcal{A}_i(w). \quad (901)$$

Where the sum is defined on the set of all the differences between the eigenvalues of the system, in our case  $w \in \{0, \pm\eta\}$ .

In order to use the equation (901) to decompose the equation (370) we need to consider the time ordering operator  $\mathcal{T}$ , it's possible to write using the Dyson series or the expansion of the operator of the form  $U(t) \equiv \mathcal{T} \exp \left( -i \int_0^t dt' \overline{H_S}(t') \right)$  like:

$$U(t) \equiv \mathcal{T} \exp \left( -i \int_0^t dt' \overline{H_S}(t') \right) \quad (902)$$

$$= \mathbb{I} + \sum_{n=1}^{\infty} (-i)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} dt_n H(t_1) H(t_2) \dots H(t_n). \quad (903)$$

Here  $0 < t_1 < t_2 < \dots < t_{n-1} < t_n = t$  is a partition of the set  $[0, t]$ . We will use a perturbative solution to the exponential of a time-varying operator, this can be done if we write an effective hamiltonian  $H_E(t)$  such that  $\mathcal{T} \exp \left( -i \int_0^t dt' \overline{H_S}(t') \right) \equiv \exp(-it H_E(t))$ . The effective Hamiltonian is expanded in a series of terms of increasing order in time  $H_E(t) = H_E^{(0)}(t) + H_E^{(1)}(t) + H_E^{(2)}(t) + \dots$  so we can write:

$$U(t) = \exp \left( -it \left( H_E^{(0)}(t) + H_E^{(1)}(t) + H_E^{(2)}(t) + \dots \right) \right). \quad (904)$$

The terms can be found expanding  $\mathcal{T} \exp \left( -i \int_0^t dt' \overline{H_S}(t') \right)$  and  $U(t)$  then equating the terms of the same power. The lowest terms are:

$$H_E^{(0)}(t) = \frac{1}{t} \int_0^t \overline{H_S}(t') dt', \quad (905)$$

$$H_E^{(1)}(t) = -\frac{i}{2t} \int_0^t dt' \int_0^{t'} dt'' [\overline{H_S}(t'), \overline{H_S}(t'')], \quad (906)$$

$$H_E^{(2)}(t) = \frac{1}{6t} \int_0^t dt' \int_0^{t'} dt'' \int_0^{t''} dt''' ([\overline{H_S}(t'), \overline{H_S}(t'')], \overline{H_S}(t''')) + [[\overline{H_S}(t'''), \overline{H_S}(t'')], \overline{H_S}(t')]. \quad (907)$$

In this case the Fourier decomposition using the expansion of  $H_E(t)$  is:

$$\widetilde{A}_i(t) = U^\dagger(t) A_i(t) U(t) \quad (908)$$

$$\widetilde{A}_i(t) = e^{iH_E(t)t} A_i(t) e^{-iH_E(t)t} \quad (909)$$

$$= \sum_{w(t)} e^{-iw(t)t} \mathcal{A}_i(t, w(t)). \quad (910)$$

$w(t)$  belongs to the set of differences of eigenvalues of  $H_E(t)$  that depends of the time. As we can see the decomposition matrices are time-dependent as well.

Extending the Fourier decomposition to the matrix  $\widetilde{A}_j(t - \tau, t)$  using the Magnus expansion generates:

$$\widetilde{A}_j(t - \tau, t) = U(t) U^\dagger(t - \tau) A_j(t) U(t - \tau) U^\dagger(t) \quad (911)$$

$$= e^{-itH_E(t)} e^{i(t-\tau)H_E(t-\tau)} A_j(t) e^{-i(t-\tau)H_E(t-\tau)} e^{itH_E(t)} \quad (912)$$

$$= e^{-itH_E(t)} \left( \sum_{w'(t-\tau)} e^{-i(t-\tau)w(t-\tau)} \mathcal{A}_j(t, w(t-\tau)) \right) e^{itH_E(t)} \quad (913)$$

$$= \sum_{w(t), w'(t-\tau)} e^{iw'(t)t} e^{-i(t-\tau)w(t-\tau)} \mathcal{A}_j(t, w(t-\tau), w'(t)) \quad (914)$$

$$= \sum_{w(t), w'(t-\tau)} e^{iw'(t)t} e^{-i(t-\tau)w(t-\tau)} \mathcal{A}_j(t, w(t-\tau), w'(t)) \quad (915)$$

$$= \sum_{w(t), w'(t-\tau)} e^{i\tau w(t-\tau)} e^{-it(w(t-\tau) - w'(t))} \mathcal{A}_j(t, w(t-\tau), w'(t)) \quad (916)$$

where  $w'(t - \tau)$  and  $w(t)$  belongs to the set of the differences of the eigenvalues of the Hamiltonian  $\overline{H_E}(t - \tau)$  and  $\overline{H_E}(t)$  respectively.

In order to show the explicit form of the matrices present in the RHS of the equation (901) for a general  $2 \times 2$  matrix in a given time let's write the matrix  $A_i$  in the base  $V = \{|+\rangle, |-\rangle\}$  in the following way:

$$A_i = \sum_{\alpha, \beta \in V} \langle \alpha | A_i | \beta \rangle | \alpha \rangle \langle \beta |. \quad (917)$$

Given that  $[|+\rangle\langle+|, |-\rangle\langle-|] = 0$ , then using the Zassenhaus formula we obtain:

$$e^{i\overline{H_E}\tau} = e^{i(\lambda_+|+\rangle\langle+| + \lambda_-|-\rangle\langle-|)\tau} \quad (918)$$

$$= e^{i\lambda_+|+\rangle\langle+|\tau} e^{i\lambda_-|-\rangle\langle-|\tau} \quad (919)$$

$$= (|-\rangle\langle-| + e^{i\lambda_+\tau}|+\rangle\langle+|) (|+\rangle\langle+| + e^{i\lambda_-\tau}|-\rangle\langle-|) \quad (920)$$

$$= e^{i\lambda_+\tau}|+\rangle\langle+| + e^{i\lambda_-\tau}|-\rangle\langle-|. \quad (921)$$

Calculating the transformation (901) directly using the previous relationship we find that:

$$U^\dagger(\tau) A_i(\tau) U(\tau) = (e^{i\lambda_+\tau}|+\rangle\langle+| + e^{i\lambda_-\tau}|-\rangle\langle-|) \left( \sum_{\alpha, \beta \in V} \langle \alpha | A_i(\tau) | \beta \rangle | \alpha \rangle \langle \beta | \right) (e^{-i\lambda_+\tau}|+\rangle\langle+| + e^{-i\lambda_-\tau}|-\rangle\langle-|) \quad (922)$$

$$= \langle + | A_i(\tau) | + \rangle | + \rangle \langle + | + e^{i\eta\tau} \langle + | A_i(\tau) | - \rangle | + \rangle \langle - | + e^{-i\eta\tau} \langle - | A_i(\tau) | + \rangle | - \rangle \langle + | + \langle - | A_i(\tau) | - \rangle | - \rangle \langle - |. \quad (923)$$

$$= A_i(0) + A_i(-w) e^{iw\tau} + A_i(w) e^{-iw\tau} \quad (924)$$

Here  $w = \lambda_+ - \lambda_-$ . Comparing the RHS of the equations (901) and the explicit expression for  $\widetilde{A}_i(\tau)$  in (909), we obtain the form of the expansion matrices of the Fourier decomposition for a general  $2 \times 2$  matrix:

$$A_i(0) = \langle + | A_i(\tau) | + \rangle | + \rangle \langle + | + \langle - | A_i(\tau) | - \rangle | - \rangle \langle - |, \quad (925)$$

$$A_i(-w) = \langle + | A_i(\tau) | - \rangle | + \rangle \langle - |, \quad (926)$$

$$A_i(w) = \langle - | A_i(\tau) | + \rangle | - \rangle \langle + |. \quad (927)$$

For a decomposition of the interaction Hamiltonian in terms of Hermitian operators, i.e.  $\widetilde{A}_i(\tau) = \widetilde{A}_i^\dagger(\tau)$  and  $\widetilde{B}_i(\tau) = \widetilde{B}_i^\dagger(\tau)$  we can use the equation (901) to write the master equation in the following neater form:

$$\frac{d\overline{\rho_S}(t)}{dt} = -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ij} \int_0^t d\tau C_i(t) C_j(t-\tau) \left( \mathcal{B}_{ij}(\tau) \left[ A_i, \widetilde{A}_j(t-\tau, t) \overline{\rho_S}(t) \right] + \mathcal{B}_{ji}(-\tau) \left[ \overline{\rho_S}(t) \widetilde{A}_j(t-\tau, t), A_i \right] \right) \quad (928)$$

$$= -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijw} \int_0^t d\tau C_i(t) C_j(t-\tau) \left( \mathcal{B}_{ij}(\tau) \left[ A_i, e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_j(w(t-\tau), w'(t)) \overline{\rho_S}(t) \right] \right. \quad (929)$$

$$\left. - \mathcal{B}_{ji}(-\tau) \left[ A_i, \overline{\rho_S}(t) e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_j(w(t-\tau), w'(t)) \right] \right) \quad (930)$$

Given that  $\mathcal{A}_j(w(t-\tau), w'(t)) = \mathcal{A}_j^\dagger(-w(t-\tau), -w'(t))$  from the Fourier decomposition (901) then we can rearrange the precedent sum in the following way with the trace respect to the bath:

$$\mathcal{B}_{ij}(\tau) = \text{Tr}_B \left( \widetilde{B}_i(t) \widetilde{B}_j(s) \rho_B \right) \quad (931)$$

$$= \text{Tr}_B \left( \widetilde{B}_i(\tau) \widetilde{B}_j(0) \rho_B \right). \quad (932)$$

Let's define:

$$\mathcal{A}_j(w(t-\tau), w'(t)) = \mathcal{A}_{jww'}(t-\tau, t) \quad (933)$$

The master equation can be re-written in the following form:

$$\frac{d\overline{\rho_S}(t)}{dt} = -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \int_0^t d\tau C_i(t) C_j(t-\tau) \mathcal{B}_{ij}(\tau) \left[ A_i, e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \overline{\rho_S}(t) \right] \quad (934)$$

$$+ \sum_{ijww'} \mathcal{B}_{ji}(-\tau) \left[ A_i, \overline{\rho_S}(t) e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \right] \quad (935)$$

$$= -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \int_0^t d\tau C_i(t) C_j(t-\tau) \mathcal{B}_{ij}(\tau) \left[ A_i, e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \overline{\rho_S}(t) \right] \quad (936)$$

$$+ \sum_{ijww'} \mathcal{B}_{ji}(-\tau) \left[ A_i, \overline{\rho_S}(t) e^{-i\tau w(t-\tau)} e^{it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \right] \quad (937)$$

$$= -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \int_0^t d\tau C_i(t) C_j(t-\tau) \mathcal{B}_{ij}(\tau) \left[ A_i, e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \overline{\rho_S}(t) \right] \quad (938)$$

$$+ \sum_{ijww'} \mathcal{B}_{ji}(-\tau) \left[ A_i, \overline{\rho_S}(t) e^{-i\tau w(t-\tau)} e^{it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \right] \quad (939)$$

$$= -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \int_0^t d\tau C_i(t) C_j(t-\tau) \mathcal{B}_{ij}(\tau) \left[ A_i, e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \overline{\rho_S}(t) \right] \quad (940)$$

$$- \mathcal{B}_{ji}(-\tau) \left[ A_i, \overline{\rho_S}(t) e^{-i\tau w(t-\tau)} e^{it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \right] \quad (941)$$

$$= -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \int_0^t d\tau C_i(t) C_j(t-\tau) \text{Tr}_B \left( \left[ A_i, \widetilde{B}_i(\tau) \widetilde{B}_j(0) \rho_B e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \overline{\rho_S}(t) \right] \right) \quad (942)$$

$$- \left[ A_i, \widetilde{B}_j(-\tau) \widetilde{B}_i(0) \rho_B \overline{\rho_S}(t) e^{-i\tau w(t-\tau)} e^{it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \right] \quad (943)$$

Given that if we define:

$$D_{ijww'}(t-\tau, t) = C_i(t) C_j(t-\tau) \mathcal{B}_{ij}(\tau) e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \quad (944)$$

then

$$D_{ijww'}^\dagger(t-\tau, t) = \left( C_i(t) C_j(t-\tau) \mathcal{B}_{ij}(\tau) e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}(t-\tau, t) \right)^\dagger \quad (945)$$

$$= \mathcal{B}_{ij}^*(\tau) C_i(t) C_j(t-\tau) e^{-i\tau w(t-\tau)} e^{it(w(t-\tau)-w'(t))} \mathcal{A}_{jww'}^\dagger(t-\tau, t) \quad (946)$$

We used the fact that  $C_i(t), C_j(t-\tau)$  are real. Now let's consider the following trace recalling that  $\text{Tr}(A)^* = \text{Tr}(A^\dagger)$  so:

$$\text{Tr}_B \left( \widetilde{B}_j(-\tau) \widetilde{B}_i(0) \rho_B \right) = \text{Tr}_B \left( e^{-i\tau H_B(\tau)} B_j e^{i\tau H_B(\tau)} B_i \rho_B \right) \quad (947)$$

$$= \text{Tr}_B \left( B_j e^{i\tau H_B(\tau)} B_i \rho_B e^{-i\tau H_B(\tau)} \right) \text{ (by cyclic permutivity of trace)} \quad (948)$$

$$= \text{Tr}_B \left( B_j e^{i\tau H_B(\tau)} B_i e^{-i\tau H_B(\tau)} \rho_B \right) \text{ (by commutativity of } e^{-i\tau H_B(\tau)} \text{ and } \rho_B) \quad (949)$$

$$= \text{Tr}_B \left( B_j \widetilde{B}_i(\tau) \rho_B \right) \text{ (by definition of time evolution)} \quad (950)$$

$$= \text{Tr}_B \left( B_j \widetilde{B}_i(\tau) \rho_B \right) \quad (951)$$

$$= \text{Tr}_B \left( \rho_B B_j \widetilde{B}_i(\tau) \right) \quad (952)$$

$$= \text{Tr}_B \left( \left( \widetilde{B}_i(\tau) B_j \rho_B \right)^\dagger \right) \text{ (by definition of adjoint)} \quad (953)$$

$$= \text{Tr}_B \left( \widetilde{B}_i(\tau) B_j \rho_B \right)^* \quad (954)$$

$$= \mathcal{B}_{ij}^*(\tau) \quad (955)$$

So we can write the master equation like:

$$\frac{d\overline{\rho}_S(t)}{dt} = -i[\overline{H}_S(t), \overline{\rho}_S(t)] - \sum_{ijww'} \int_0^t d\tau C_i(t) C_j(t-\tau) \left( \mathcal{B}_{ij}(\tau) \left[ A_i, e^{i\tau w(t-\tau)} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_j(w(t-\tau), w'(t)) \overline{\rho}_S(t) \right] \right. \quad (956)$$

$$\left. - \mathcal{B}_{ij}^*(\tau) \left[ A_i, \overline{\rho}_S(t) e^{-i\tau w(t-\tau)} e^{it(w(t-\tau)-w'(t))} \mathcal{A}_j^\dagger(w(t-\tau), w'(t)) \right] \right) \quad (957)$$

$$= -i[\overline{H}_S(t), \overline{\rho}_S(t)] - \sum_{ijww'} \int_0^t d\tau \left( [A_i, D_{ijww'}(t-\tau, t) \overline{\rho}_S(t)] - [A_i, \overline{\rho}_S(t) D_{ijww'}^\dagger(t-\tau, t)] \right) \quad (958)$$

Let's define the response matrix in the following way.

$$\mathcal{D}_{ijww'}(t) = \int_0^t d\tau D_{ijww'}(t-\tau, t) \quad (959)$$

Then the master equation can be written as:

$$\frac{d\overline{\rho}_S(t)}{dt} = -i[\overline{H}_S(t), \overline{\rho}_S(t)] - \sum_{ijww'} \left( [A_i, \mathcal{D}_{ijww'}(t) \overline{\rho}_S(t)] - [A_i, \overline{\rho}_S(t) \mathcal{D}_{ijww'}^\dagger(t)] \right) \quad (960)$$

If we extend the upper limit of integration to  $\infty$  in the equation (959) then the system will be independent of any preparation at  $t = 0$ , so the evolution of the system will depend only on its present state as expected in the Markovian approximation.

Applying the inverse transformation we will obtain that:

$$e^{-V} \frac{d\overline{\rho}_S(t)}{dt} e^V = \frac{d(e^{-V} \overline{\rho}_S e^V)}{dt} \quad (961)$$

$$= \frac{d\rho_S}{dt} \quad (962)$$

$$= -ie^{-V} [\overline{H}_S(t), \overline{\rho}_S(t)] e^V - \sum_{ijww'} \int_0^t d\tau \left( e^{-V} [A_i, D_{ijww'}(t-\tau, t) \overline{\rho}_S(t)] e^V - e^{-V} [A_i, \overline{\rho}_S(t) D_{ijww'}^\dagger(t-\tau, t)] e^V \right). \quad (963)$$

For a product we have the following:



$$e^{-V} \overline{AB} e^V = e^{-V} \overline{A} \overline{B} e^V \quad (964)$$

$$= e^{-V} \overline{A} e^V e^{-V} \overline{B} e^V \quad (965)$$

$$= (e^{-V} \overline{A} e^V) (e^{-V} \overline{B} e^V) \quad (966)$$

$$= AB. \quad (967)$$

We can use this to prove the following property for the inverse transformation of a commutator:

$$e^{-V} [A, B] e^V = e^{-V} (\overline{AB} - \overline{BA}) e^V \quad (968)$$

$$= e^{-V} \overline{AB} e^V - e^{-V} \overline{BA} e^V \quad (969)$$

$$= AB - BA \quad (970)$$

$$= [A, B]. \quad (971)$$

So we will obtain that

$$\frac{d\rho_S}{dt} = -i e^{-V} [\overline{H_S}(t), \overline{\rho_S}(t)] e^V - e^{-V} \sum_{ijww'} \left( [A_i, \mathcal{D}_{ijww'}(t) \overline{\rho_S}(t)] - [A_i, \overline{\rho_S}(t) \mathcal{D}_{ijww'}^\dagger(t)] \right) e^V \quad (972)$$

$$= -i e^{-V} [\overline{H_S}(t), \overline{\rho_S}(t)] e^V - \sum_{ijww'} \left( e^{-V} [A_i, \mathcal{D}_{ijww'}(t) \overline{\rho_S}(t)] e^V - e^{-V} [A_i, \overline{\rho_S}(t) \mathcal{D}_{ijww'}^\dagger(t)] e^V \right) \quad (973)$$

$$= -i [H_S(t), \rho_S(t)] - \sum_{ijww'} \left( [e^{-V} A_i e^V, e^{-V} \mathcal{D}_{ijww'}(t) \overline{\rho_S}(t) e^V] - [e^{-V} A_i e^V, e^{-V} \overline{\rho_S}(t) \mathcal{D}_{ijww'}^\dagger(t) e^V] \right) \quad (974)$$

$$= -i [H_S(t), \rho_S(t)] - \sum_{ijww'} \left( [e^{-V} A_i e^V, e^{-V} \mathcal{D}_{ijww'}(t) e^V e^{-V} \overline{\rho_S}(t) e^V] - [e^{-V} A_i e^V, e^{-V} \overline{\rho_S}(t) e^V e^{-V} \mathcal{D}_{ijww'}^\dagger(t) e^V] \right) \quad (975)$$

$$= -i [H_S(t), \rho_S(t)] - \sum_{ijww'} \left( [e^{-V} A_i e^V, e^{-V} \mathcal{D}_{ijww'}(t) e^V \rho_S(t)] - [e^{-V} A_i e^V, \rho_S(t) e^{-V} \mathcal{D}_{ijww'}^\dagger(t) e^V] \right). \quad (976)$$

#### IV. LIMIT CASES

In order to show the plausibility of the master equation (960) for a time-dependent Hamiltonian we will show that this equation reproduces the following cases under certain limits conditions that will be pointed in each subsection.

##### A. Time-independent variational quantum master equation

At first let's show that the master equation (960) reproduces the results of the reference [1], for the latter case we have that  $i, j \in \{1, 2, 3\}$  and  $\omega \in (0, \pm\eta)$ . The Hamiltonian of the system considered in this reference written in the same basis than the Hamiltonian (1) is given by:

$$H = \left( \delta + \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \right) |1\rangle\langle 1| + \frac{\Omega}{2} \sigma_x + \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}. \quad (977)$$

After performing the transformation (24) on the Hamiltonian (977) it's possible to split that result in the following set of Hamiltonians:

$$\overline{H_S} = (\delta + R) |1\rangle\langle 1| + \frac{\Omega_r}{2} \sigma_x, \quad (978)$$

$$\overline{H_I} = B_z |1\rangle\langle 1| + \frac{\Omega}{2} (B_x \sigma_x + B_y \sigma_y), \quad (979)$$

$$H_B = \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}. \quad (980)$$

The Hamiltonian (978) differs from the transformed Hamiltonian  $H_S$  of the reference written like  $H_S = \frac{R}{2}\mathbb{I} + \frac{\epsilon}{2}\sigma_z + \frac{\Omega_r}{2}\sigma_x$  by a term proportional to the identity, this can be seen in the following way taking  $\epsilon = \delta + R$

$$(\delta + R) |1\rangle\langle 1| - \frac{\delta}{2}\mathbb{I} = \left(\frac{\delta}{2} + R\right) |1\rangle\langle 1| - \frac{\delta}{2} |0\rangle\langle 0| \quad (981)$$

$$= \frac{R}{2}\mathbb{I} + \frac{\delta + R}{2}\sigma_z \quad (982)$$

$$= \frac{R}{2}\mathbb{I} + \frac{\epsilon}{2}\sigma_z. \quad (983)$$

In this Hamiltonian we can write  $A_i = \sigma_x$ ,  $A_2 = \sigma_y$  and  $A_3 = \frac{I+\sigma_z}{2} = |1\rangle\langle 1|$  with  $\sigma_z = |1\rangle\langle 1| - |0\rangle\langle 0|$ . In order to find the decomposition matrices of the Fourier decomposition let's obtain the eigenvalues and eigenvectors of the matrix  $\overline{H_S}$ . Given that  $\overline{H_S} = \frac{R}{2}\mathbb{I} + \frac{\epsilon}{2}\sigma_z + \frac{\Omega_r}{2}\sigma_x$  then  $\text{Tr}(\overline{H_S}) = R$  and  $\text{Det}(\overline{H_S}) = \frac{R^2 - \epsilon^2}{4} - \frac{\Omega_r^2}{4}$  then by the Caley-Hamilton theorem then we will have that the equations of the eigenvalues and it's values are given by::

$$0 = \lambda^2 - R\lambda + \frac{R^2 - \epsilon^2 - \Omega_r^2}{4}, \quad (984)$$

$$\lambda_{\pm} = \frac{R \pm \sqrt{(-R)^2 - 4\left(\frac{R^2 - \epsilon^2 - \Omega_r^2}{4}\right)}}{2} \quad (985)$$

$$= \frac{R \pm \sqrt{R^2 - (R^2 - \epsilon^2 - \Omega_r^2)}}{2} \quad (986)$$

$$= \frac{R \pm \sqrt{\epsilon^2 + \Omega_r^2}}{2} \quad (987)$$

$$\eta = \sqrt{\epsilon^2 + \Omega_r^2}, \quad (988)$$

$$\lambda_{\pm} = \frac{R \pm \eta}{2}. \quad (989)$$

For  $\lambda_+ = \frac{R+\eta}{2}$  we will obtain the associated eigenvector like:

$$\begin{pmatrix} \frac{R}{2} - \frac{\epsilon}{2} - \frac{R+\eta}{2} & \frac{\Omega_r}{2} \\ \frac{\Omega_r}{2} & \frac{R}{2} + \frac{\epsilon}{2} - \frac{R+\eta}{2} \end{pmatrix} = \begin{pmatrix} -\frac{\epsilon}{2} - \frac{\eta}{2} & \frac{\Omega_r}{2} \\ \frac{\Omega_r}{2} & \frac{\epsilon}{2} - \frac{\eta}{2} \end{pmatrix}. \quad (990)$$

so the eigenvector  $|+\rangle = a|0\rangle + b|1\rangle$  satisfies  $-\frac{\epsilon+\eta}{2}a + \frac{\Omega_r}{2}b = 0$ , so  $a = \frac{\Omega_r}{\epsilon+\eta}b$  then the normalized eigenvector is  $|+\rangle = \frac{\Omega_r}{\sqrt{(\epsilon+\eta)^2 + \Omega_r^2}}|0\rangle + \frac{\epsilon+\eta}{\sqrt{(\epsilon+\eta)^2 + \Omega_r^2}}|1\rangle$  with  $\sin(\theta) = \frac{\Omega_r}{\sqrt{(\epsilon+\eta)^2 + \Omega_r^2}}$  and  $\cos(\theta) = \frac{\epsilon+\eta}{\sqrt{(\epsilon+\eta)^2 + \Omega_r^2}}$ . The vector is written in reduced way like  $|+\rangle = \sin(\theta)|0\rangle + \cos(\theta)|1\rangle$ .

For  $\lambda_- = \frac{R-\eta}{2}$  we will obtain the associated eigenvector like:

$$\begin{pmatrix} \frac{R}{2} - \frac{\epsilon}{2} - \frac{R-\eta}{2} & \frac{\Omega_r}{2} \\ \frac{\Omega_r}{2} & \frac{R}{2} + \frac{\epsilon}{2} - \frac{R-\eta}{2} \end{pmatrix} = \begin{pmatrix} -\frac{\epsilon}{2} + \frac{\eta}{2} & \frac{\Omega_r}{2} \\ \frac{\Omega_r}{2} & \frac{\epsilon}{2} + \frac{\eta}{2} \end{pmatrix}. \quad (991)$$

so the eigenvector  $|+\rangle = a|0\rangle + b|1\rangle$  satisfies  $\frac{\Omega_r}{2}a + \frac{\epsilon+\eta}{2}b = 0$ , so  $a = -\frac{\epsilon+\eta}{\Omega_r}b$  then the normalized eigenvector is  $|-\rangle = \frac{\epsilon+\eta}{\sqrt{(\epsilon+\eta)^2 + \Omega_r^2}}|0\rangle - \frac{\Omega_r}{\sqrt{(\epsilon+\eta)^2 + \Omega_r^2}}|1\rangle$ . The vector is written in reduced way like  $|-\rangle = \cos(\theta)|0\rangle - \sin(\theta)|1\rangle$ . Summarizing these results we can write:

$$\lambda_+ = \frac{\epsilon + \eta}{2}, \quad (992)$$

$$\lambda_- = \frac{\epsilon - \eta}{2}, \quad (993)$$

$$|+\rangle = \sin(\theta) |0\rangle + \cos(\theta) |1\rangle, \quad (994)$$

$$|-\rangle = \cos(\theta) |0\rangle - \sin(\theta) |1\rangle, \quad (995)$$

$$\sin(\theta) = \frac{\Omega_r}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}, \quad (996)$$

$$\cos(\theta) = \frac{\epsilon + \eta}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}. \quad (997)$$

This result is plausible because in the paper [1] we have that:

$$\theta = \frac{1}{2} \tan^{-1} \left( \frac{\Omega_r}{\epsilon} \right). \quad (998)$$

We can obtain the value of  $\tan(\theta)$  through the following trigonometry identity for  $x = \tan^{-1} \left( \frac{\Omega_r}{\epsilon} \right)$ .

$$\tan \left( \frac{x}{2} \right) = \frac{\sin(x)}{\cos(x) + 1}. \quad (999)$$

So the value of  $\tan(\theta)$  using (999) is equal to:

$$\tan(\theta) = \frac{\frac{\Omega_r}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}}{\frac{\epsilon}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}} + 1} \quad (1000)$$

$$= \frac{\frac{\Omega_r}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}}{\frac{\epsilon + \sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}} \quad (1001)$$

$$= \frac{\Omega_r}{\epsilon + \eta}. \quad (1002)$$

This proves our assertion.

Using this basis we can find the decomposition matrices using the equations (926)-(927) and the fact that  $|+\rangle = \sin(\theta) |0\rangle + \cos(\theta) |1\rangle = \begin{pmatrix} \sin(\theta) \\ \cos(\theta) \end{pmatrix}$  and  $|-\rangle = \cos(\theta) |0\rangle - \sin(\theta) |1\rangle = \begin{pmatrix} \cos(\theta) \\ -\sin(\theta) \end{pmatrix}$  with  $\sin(\theta) = \frac{\Omega_r}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}$  and  $\cos(\theta) = \frac{\epsilon + \eta}{\sqrt{(\epsilon + \eta)^2 + \Omega_r^2}}$ :

$$\langle +|\sigma_x|+\rangle = \begin{pmatrix} \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \sin(\theta) \\ \cos(\theta) \end{pmatrix} \quad (1003)$$

$$= 2 \sin(\theta) \cos(\theta) \quad (1004)$$

$$= \sin(2\theta), \quad (1005)$$

$$\langle -|\sigma_x|-\rangle = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \cos(\theta) \\ -\sin(\theta) \end{pmatrix} \quad (1006)$$

$$= -2 \sin(\theta) \cos(\theta) \quad (1007)$$

$$= -\sin(2\theta), \quad (1008)$$

$$\langle -|\sigma_x|+\rangle = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \sin(\theta) \\ \cos(\theta) \end{pmatrix} \quad (1009)$$

$$= \cos^2(\theta) - \sin^2(\theta) \quad (1010)$$

$$= \cos(2\theta), \quad (1011)$$

$$\langle +|\sigma_y|+\rangle = \begin{pmatrix} \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \begin{pmatrix} \sin(\theta) \\ \cos(\theta) \end{pmatrix} \quad (1012)$$

$$= i \sin(\theta) \cos(\theta) - i \sin(\theta) \cos(\theta) \quad (1013)$$

$$= 0, \quad (1014)$$

$$\langle -|\sigma_y|-\rangle = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \end{pmatrix} \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \begin{pmatrix} \cos(\theta) \\ -\sin(\theta) \end{pmatrix} \quad (1015)$$

$$= i \sin(\theta) \cos(\theta) - i \sin(\theta) \cos(\theta) \quad (1016)$$

$$= 0, \quad (1017)$$

$$\langle -|\sigma_y|+\rangle = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \end{pmatrix} \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \begin{pmatrix} \sin(\theta) \\ \cos(\theta) \end{pmatrix} \quad (1018)$$

$$= i \cos^2(\theta) + i \sin^2(\theta) \quad (1019)$$

$$= i. \quad (1020)$$

$$\langle +|\frac{1+\sigma_z}{2}|+\rangle = \begin{pmatrix} \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \sin(\theta) \\ \cos(\theta) \end{pmatrix} \quad (1021)$$

$$= \cos(\theta) \cos(\theta) \quad (1022)$$

$$= \cos^2(\theta), \quad (1023)$$

$$\langle -|\frac{1+\sigma_z}{2}|-\rangle = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos(\theta) \\ -\sin(\theta) \end{pmatrix} \quad (1024)$$

$$= \sin(\theta) \sin(\theta) \quad (1025)$$

$$= \sin^2(\theta), \quad (1026)$$

$$\langle -|\frac{1+\sigma_z}{2}|+\rangle = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \sin(\theta) \\ \cos(\theta) \end{pmatrix} \quad (1027)$$

$$= -\sin(\theta) \cos(\theta) \quad (1028)$$

$$= -\sin(\theta) \cos(\theta). \quad (1029)$$

Composing the parts shown give us the Fourier decomposition matrices for this case:

$$A_1(0) = \sin(2\theta) (|+\chi+\rangle - |-\chi-\rangle), \quad (1030)$$

$$A_1(\eta) = \cos(2\theta) |-\chi+\rangle, \quad (1031)$$

$$A_2(0) = 0, \quad (1032)$$

$$A_2(\eta) = i|-\chi+\rangle, \quad (1033)$$

$$A_3(0) = \cos^2(\theta) |+\chi+\rangle + \sin^2(\theta) |-\chi-\rangle, \quad (1034)$$

$$A_3(\eta) = -\sin(\theta) \cos(\theta) |-\chi+\rangle. \quad (1035)$$

Now to prove the fact that the model of the “Time-independent variational quantum master equation” is a special case the master equation (963) we need to take account of the time-independence of the hamiltonian of this system. From this perspective is possible to show that for the equation (944) is equivalent to:

$$\mathcal{D}_{ijww'}(t) = \int_0^t d\tau D_{ijww'}(t-\tau, t) \quad (1036)$$

$$= \int_0^t d\tau C_i(t) C_j(t-\tau) \Lambda_{ij}(\tau) e^{i\tau w} e^{-it(w(t-\tau)-w'(t))} \mathcal{A}_j(w(t-\tau), w'(t)) \quad (1037)$$

$$= \int_0^t d\tau C_i(t) C_j(t-\tau) \Lambda_{ij}(\tau) e^{i\tau w} e^{-it(w-w')} \mathcal{A}_j(w, w'). \quad (1038)$$

Now to make comparisons between the model obtained and the model of the system under discussion we will define that the correlation functions of the reference [1] denoted by  $\Lambda'_{ij}(\tau)$  relate with the correlation functions defined in the equation (411) in the following way:

$$\Lambda'_{ij}(\tau) = C_i(t) C_j(t-\tau) \Lambda_{ij}(\tau). \quad (1039)$$

So the response matrix can be rewritten as:

$$\mathcal{D}_{ijww'}(t) = \left( \int_0^t d\tau \Lambda'_{ij}(\tau) e^{i\tau w} e^{-it(w-w')} \right) \mathcal{A}_j(w, w') \quad (1040)$$

Let's define the response function like:

$$K_{ij}(w, w', t) = \int_0^t C_i(t) C_j(t-\tau) \Lambda_{ij}(\tau) e^{i\tau w} e^{-it(w-w')} d\tau \quad (1041)$$

$$= \int_0^t \Lambda'_{ij}(\tau) e^{i\tau w} e^{-it(w-w')} d\tau \quad (1042)$$

$$= K_{ijww'}(t). \quad (1043)$$

Then we have the following equivalence:

$$\mathcal{D}_{ijww'}(t) = K_{ijww'}(t) \mathcal{A}_j(w, w') \quad (1044)$$

$$= K_{ijww'}(t) \mathcal{A}_{jww'} \quad (1045)$$

We can proof that

$$\frac{d\overline{\rho_S}(t)}{dt} = -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \left( [A_i, \mathcal{D}_{ijww'}(t) \overline{\rho_S}(t)] - [A_i, \overline{\rho_S}(t) \mathcal{D}_{ijww'}^\dagger(t)] \right) \quad (1046)$$

$$= -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \left( [A_i, K_{ijww'}(t) \mathcal{A}_{jww'} \overline{\rho_S}(t)] - [A_i, \overline{\rho_S}(t) K_{ijww'}^*(t) \mathcal{A}_{jww'}^\dagger] \right) \quad (1047)$$

$$= -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \left( K_{ijww'}(t) [A_i, \mathcal{A}_{jww'} \overline{\rho_S}(t)] - K_{ijww'}^*(t) [A_i, \overline{\rho_S}(t) \mathcal{A}_{jww'}^\dagger] \right) \quad (1048)$$

$$= -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} \left( (K_{ijww'}^{\Re}(t) + i K_{ijww'}^{\Im}(t)) [A_i, \mathcal{A}_{jww'} \overline{\rho_S}(t)] - (K_{ijww'}^{\Re}(t) - i K_{ijww'}^{\Im}(t)) [A_i, \overline{\rho_S}(t) \mathcal{A}_{jww'}^\dagger] \right) \quad (1049)$$

$$= -i [\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijww'} K_{ijww'}^{\Re}(t) [A_i, \mathcal{A}_{jww'} \overline{\rho_S}(t) - \overline{\rho_S}(t) \mathcal{A}_{jww'}^\dagger] - i \sum_{ijww'} K_{ijww'}^{\Im}(t) [A_i, \mathcal{A}_{jww'} \overline{\rho_S}(t) + \overline{\rho_S}(t) \mathcal{A}_{jww'}^\dagger] \quad (1050)$$

Using the notation of the master equation (960), we can say that  $C_1(t) = \frac{\Omega}{2} = C_2(t)$  and  $C_3(t) = 1$ , being  $\Omega$  a constant. Furthermore given that  $\overline{H_S}$  is time-independent then  $B(t) = B$ . Taking the equations(??)-(??) we find that the correlation functions of the reference [1] written in terms of the RHS of the equation (411) are equal to:

$$\langle \widetilde{B}_{1z}(t) \widetilde{B}_{1z}(s) \rangle_B = \sum_{\mathbf{k}} ((g_{1\mathbf{k}} - v_{1\mathbf{k}}) (g_{1\mathbf{k}} - v_{1\mathbf{k}})^* e^{i\omega_{\mathbf{k}}\tau} N_{\mathbf{k}} + (g_{1\mathbf{k}} - v_{1\mathbf{k}})^* (g_{1\mathbf{k}} - v_{1\mathbf{k}}) e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1)) \quad (1051)$$

$$= \sum_{\mathbf{k}} |g_{1\mathbf{k}} - v_{1\mathbf{k}}|^2 (e^{i\omega_{\mathbf{k}}\tau} N_{\mathbf{k}} + e^{-i\omega_{\mathbf{k}}\tau} (N_{\mathbf{k}} + 1)) \quad (1052)$$

$$\approx \int_0^\infty J_1(\omega) (1 - F_1(\omega))^2 (e^{i\omega\tau} N(\omega) + e^{-i\omega\tau} (N(\omega) + 1)) d\omega \quad (1053)$$

$$G_{\pm}(\omega, \tau) = e^{i\omega\tau} N(\omega) + e^{-i\omega\tau} (N(\omega) + 1) \quad (1054)$$

$$\langle \widetilde{B}_{1z}(t) \widetilde{B}_{1z}(s) \rangle_B \approx \int_0^\infty J_1(\omega) (1 - F_1(\omega))^2 G_+(\omega, t) d\omega \quad (1055)$$

$$\chi_{10}(t) = 0 \text{ (because } v_{0\mathbf{k}}(t) = 0 \text{ for all } \mathbf{k}) \quad (1056)$$

$$U_{10}(t, s) = \prod_{\mathbf{k}} \exp \left( i \left( \frac{(v_{1\mathbf{k}}(t) - v_{0\mathbf{k}}(t)) (v_{1\mathbf{k}}(s) - v_{0\mathbf{k}}(s))^* e^{i\omega_{\mathbf{k}}\tau}}{\omega_{\mathbf{k}}^2} \right)^{\Im} \right) \quad (1057)$$

$$= \prod_{\mathbf{k}} \exp \left( i \left( \frac{v_{1\mathbf{k}}^2(t) e^{i\omega_{\mathbf{k}}\tau}}{\omega_{\mathbf{k}}^2} \right)^{\Im} \right) \quad (1058)$$

$$= \prod_{\mathbf{k}} \exp \left( i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} \right) \quad (1059)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_x(s) \rangle_B = \frac{1}{2} \left( \prod_{\mathbf{k}} \exp \left( i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} \right) \prod_{\mathbf{k}} \exp \left( - \frac{|v_{1\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}|^2}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) + \prod_{\mathbf{k}} \exp \left( - i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} \right) \prod_{\mathbf{k}} \exp \left( - \frac{|v_{1\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} - v_{1\mathbf{k}}|^2}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) \quad (1060)$$

$$- \left( \exp \left( - \frac{1}{2} \sum_{\mathbf{k}} \left( \left| \frac{v_{1\mathbf{k}}}{\omega_{\mathbf{k}}} \right|^2 \right) \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) \left( \exp \left( - \frac{1}{2} \sum_{\mathbf{k}} \left( \left| \frac{v_{1\mathbf{k}}}{\omega_{\mathbf{k}}} \right|^2 \right) \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) \quad (1061)$$

$$= \frac{1}{2} \left( \prod_{\mathbf{k}} \exp \left( i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{|v_{1\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}|^2}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) + \prod_{\mathbf{k}} \exp \left( - i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{|v_{1\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} - v_{1\mathbf{k}}|^2}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) \quad (1062)$$

$$- \left( \exp \left( - \sum_{\mathbf{k}} \frac{v_{1\mathbf{k}}^2}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) \quad (1063)$$

$$|v_{1\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} \pm v_{1\mathbf{k}}|^2 = v_{1\mathbf{k}}^2 |e^{i\omega_{\mathbf{k}}\tau} \pm 1|^2 \quad (1064)$$

$$= v_{1\mathbf{k}}^2 |\cos(\omega_{\mathbf{k}}\tau) + i \sin(\omega_{\mathbf{k}}\tau) \pm 1|^2 \quad (1065)$$

$$= v_{1\mathbf{k}}^2 \left( (1 \pm \cos(\omega_{\mathbf{k}}\tau))^2 + \sin^2(\omega_{\mathbf{k}}\tau) \right) \quad (1066)$$

$$= 2v_{1\mathbf{k}}^2 (1 \pm \cos(\omega_{\mathbf{k}}\tau)) \quad (1067)$$

$$B \equiv \exp \left( - \frac{1}{2} \sum_{\mathbf{k}} \frac{v_{1\mathbf{k}}^2}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \quad (1068)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_x(s) \rangle_B = \frac{1}{2} \left( \exp \left( \sum_{\mathbf{k}} i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{|v_{1\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} + v_{1\mathbf{k}}|^2}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) + \exp \left( \sum_{\mathbf{k}} - i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{|v_{1\mathbf{k}} e^{i\omega_{\mathbf{k}}\tau} - v_{1\mathbf{k}}|^2}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) \quad (1069)$$

$$- \left( \exp \left( - \sum_{\mathbf{k}} \frac{v_{1\mathbf{k}}^2}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) \quad (1070)$$

$$\phi(\tau) = \sum_{\mathbf{k}} \frac{v_{1\mathbf{k}}^2}{\omega_{\mathbf{k}}^2} \left( -i \sin(\omega_{\mathbf{k}}\tau) + \cos(\omega_{\mathbf{k}}\tau) \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \quad (1071)$$

$$\approx \int_0^\infty \frac{J_1(\omega) F_1^2(\omega)}{\omega^2} \left( -i \sin(\omega\tau) + \cos(\omega\tau) \coth \left( \frac{\beta\omega}{2} \right) \right) d\omega \quad (1072)$$

$$= \int_0^\infty \frac{J_1(\omega) F_1^2(\omega)}{\omega^2} G_+(\omega, \tau) d\omega \quad (1073)$$

$$\langle \widetilde{B}_x(t) \widetilde{B}_x(s) \rangle_B = \frac{1}{2} \left( \exp \left( \sum_{\mathbf{k}} i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{2v_{1\mathbf{k}}^2(1+\cos(\omega_{\mathbf{k}}\tau))}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) + \exp \left( \sum_{\mathbf{k}} - i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{2v_{1\mathbf{k}}^2(1-\cos(\omega_{\mathbf{k}}\tau))}{2\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) - B^2 \quad (1074)$$

$$= \frac{1}{2} \left( \exp \left( \sum_{\mathbf{k}} i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{v_{1\mathbf{k}}^2(1+\cos(\omega_{\mathbf{k}}\tau))}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) + \exp \left( \sum_{\mathbf{k}} - i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{v_{1\mathbf{k}}^2(1-\cos(\omega_{\mathbf{k}}\tau))}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) - B^2 \quad (1075)$$

$$= \frac{1}{2} \left( \exp \left( - \sum_{\mathbf{k}} \frac{v_{1\mathbf{k}}^2}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \exp \left( \sum_{\mathbf{k}} i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{v_{1\mathbf{k}}^2 \cos(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) + \exp \left( - \sum_{\mathbf{k}} \frac{v_{1\mathbf{k}}^2}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \exp \left( \sum_{\mathbf{k}} - i \frac{v_{1\mathbf{k}}^2 \sin(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} - \frac{v_{1\mathbf{k}}^2 \cos(\omega_{\mathbf{k}}\tau)}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right) \right) - B^2 \quad (1076)$$

$$B^2 = \exp \left( - \sum_{\mathbf{k}} \frac{v_{1\mathbf{k}}^2}{\omega_{\mathbf{k}}^2} \coth \left( \frac{\beta\omega_{\mathbf{k}}}{2} \right) \right)$$

$$\Lambda'_{11}(\tau) = \left(\frac{\Omega}{2}\right)^2 \text{Tr}_B \left( \widetilde{B}_1(\tau) \widetilde{B}_1(0) \rho_B \right) \quad (1118)$$

$$= \frac{\Omega_r^2}{8} \left( e^{\phi(\tau)} + e^{-\phi(\tau)} - 2 \right) \quad (1119)$$

$$= \frac{\Omega_r^2}{4} (\cosh(\phi(\tau)) - 1) \quad (1120)$$

$$\Lambda'_{22}(\tau) = \left(\frac{\Omega}{2}\right)^2 \text{Tr}_B \left( \widetilde{B}_2(\tau) \widetilde{B}_2(0) \rho_B \right) \quad (1121)$$

$$= \frac{\Omega_r^2}{8} \left( e^{\phi(\tau)} - e^{-\phi(\tau)} \right), \quad (1122)$$

$$\Lambda'_{33}(\tau) = \int_0^\infty d\omega J(\omega) (1 - F(\omega))^2 G_+(\tau), \quad (1123)$$

$$\Lambda'_{32}(\tau) = \frac{\Omega_r}{2} \int_0^\infty d\omega \frac{J(\omega)}{\omega} F(\omega) (1 - F(\omega)) iG_-(\tau), \quad (1124)$$

$$\Lambda'_{32}(\tau) = -\Lambda'_{23}(\tau), \quad (1125)$$

$$\Lambda'_{12}(\tau) = \Lambda'_{21}(\tau) \quad (1126)$$

$$= \Lambda'_{13}(\tau) \quad (1127)$$

$$= \Lambda'_{31}(\tau) \quad (1128)$$

$$= 0. \quad (1129)$$

Finally taking the Hamiltonian (977) and given that to reproduce this Hamiltonian we need to impose in (5) that  $V_{10}(t) = \frac{\Omega}{2}$ ,  $\varepsilon_0(t) = 0$  and  $\varepsilon_1(t) = \delta$ , then we obtain that  $\text{Det}(\overline{H}_S) = -\frac{\Omega_r^2}{4}$ ,  $\text{Tr}(\overline{H}_S) = \epsilon$ . Now  $\eta = \sqrt{\epsilon^2 + \Omega_r^2}$  and using the equation (335) we have that:

$$f_k = \frac{g_k \left( 1 - \frac{\epsilon \tanh\left(\frac{\beta\eta}{2}\right)}{\eta} \right)}{1 - \frac{\tanh\left(\frac{\beta\eta}{2}\right)}{\eta} \left( \epsilon - \frac{\Omega_r^2 \coth\left(\frac{\beta\omega_k}{2}\right)}{2\omega_k} \right)} \quad (1130)$$

$$= \frac{g_k \left( 1 - \frac{\epsilon \tanh\left(\frac{\beta\eta}{2}\right)}{\eta} \right)}{1 - \frac{\epsilon \tanh\left(\frac{\beta\eta}{2}\right)}{\eta} \left( 1 - \frac{\Omega_r^2 \coth\left(\frac{\beta\omega_k}{2}\right)}{2\epsilon\omega_k} \right)}. \quad (1131)$$

This shows that the expression obtained reproduces the variational parameters of the time-independent model of the reference. In general we can see that the time-independent model studied can be reproduced using the master equation (447) under a time-independent approach providing similar results.

Given that the Hamiltonian of this system is time-independent, then  $U(t)U^\dagger(t-\tau) = U(\tau)$ . From the equation (960) and using the fact that

$$\widetilde{A}_j(t-\tau, t) = U(\tau) A_j U(-\tau) \quad (1132)$$

$$= \sum_w e^{iw\tau} \mathcal{A}_j(-w) \quad (1133)$$

$$= \sum_w e^{-iw\tau} \mathcal{A}_j(w). \quad (1134)$$

because the matrices  $U(t)$  and  $U(t-\tau)$  commute from the fact that  $H_S(t)$  and  $H_S(t-\tau)$  commute as well for time independent Hamiltonians. The master equation is equal to:



$$\frac{d\overline{\rho_S}(t)}{dt} = -i[H_S(t), \overline{\rho_S}(t)] - \frac{1}{2} \sum_{ij} \sum_w \gamma_{ij}(w, t) [A_i, \mathcal{A}_j(w) \overline{\rho_S}(t) - \overline{\rho_S}(t) \mathcal{A}_j^\dagger(w)] \quad (1135)$$

$$- \sum_{ij} \sum_w S_{ij}(w, t) [A_i, \mathcal{A}_j(w) \overline{\rho_S}(t) + \overline{\rho_S}(t) \mathcal{A}_j^\dagger(w)]. \quad (1136)$$

where  $\mathcal{A}_j^\dagger(w) = \mathcal{A}_j(-w)$ , as we can see the equation (1136) contains the rates and energy shifts  $\gamma_{ij}(w, t) = 2K_{ij}^{\Re}(w, t)$  and  $S_{ij}(w, t) = K_{ij}^{\Im}(w, t)$ , respectively, defined in terms of the response functions

$$K_{ij}^{\Im}(w, t) = \int_0^t \Lambda'_{ij}(\tau) e^{iw\tau} d\tau.$$

The fact  $\mathcal{A}_j^\dagger(w) = \mathcal{A}_j(-w)$  can be verified directly for a  $2 \times 2$  matrix. given that  $\overline{H_S}$  is independent of time then we have that:

$$e^{i\overline{H_S}(t-\tau)} = e^{i(\lambda_+|+\rangle\langle+| + \lambda_-|-\rangle\langle-|)(t-\tau)} \quad (1137)$$

$$= e^{i\lambda_+|+\rangle\langle+|(t-\tau)} e^{i\lambda_-|-\rangle\langle-|(t-\tau)} \quad (1138)$$

$$= \left(|-\rangle\langle-| + e^{i\lambda_+(t-\tau)}|+\rangle\langle+|\right) \left(|+\rangle\langle+| + e^{i\lambda_-(t-\tau)}|-\rangle\langle-|\right) \quad (1139)$$

$$= e^{i\lambda_+(t-\tau)}|+\rangle\langle+| + e^{i\lambda_-(t-\tau)}|-\rangle\langle-|. \quad (1140)$$

Where  $\lambda_+, \lambda_-$  are the eigenvalues associated to the eigenvectors  $|+\rangle\langle+|, |-\rangle\langle-|$  of  $\overline{H_S}$ . Calculating the transformation (901) of (925)-(927) directly using the previous relationship we find that:

$$\widetilde{A_i(0)(t-\tau)} = (e^{i\lambda_+(t-\tau)}|+\rangle\langle+| + e^{i\lambda_-(t-\tau)}|-\rangle\langle-|)(\langle+|A_i|+\rangle|+\rangle\langle+| + \langle-|A_i|-\rangle|-\rangle\langle-|)(e^{-i\lambda_+(t-\tau)}|+\rangle\langle+| + e^{-i\lambda_-(t-\tau)}|-\rangle\langle-|) \quad (1141)$$

$$= \langle+|A_i|+\rangle|+\rangle\langle+| + \langle-|A_i|-\rangle|-\rangle\langle-|, \quad (1142)$$

$$\widetilde{A_i(w)(t-\tau)} = (e^{i\lambda_+(t-\tau)}|+\rangle\langle+| + e^{i\lambda_-(t-\tau)}|-\rangle\langle-|)(\langle+|A_i|-\rangle|+\rangle\langle-|)(e^{-i\lambda_+(t-\tau)}|+\rangle\langle+| + e^{-i\lambda_-(t-\tau)}|-\rangle\langle-|) \quad (1143)$$

$$= \langle+|A_i|-\rangle|+\rangle\langle-|e^{iw(t-\tau)}, \quad (1144)$$

$$\widetilde{A_i(-w)(t-\tau)} = (e^{i\lambda_+(t-\tau)}|+\rangle\langle+| + e^{i\lambda_-(t-\tau)}|-\rangle\langle-|)(\langle-|A_i|+\rangle|-\rangle\langle+|)(e^{-i\lambda_+(t-\tau)}|+\rangle\langle+| + e^{-i\lambda_-(t-\tau)}|-\rangle\langle-|) \quad (1145)$$

$$= \langle-|A_i|+\rangle|-\rangle\langle+|e^{-iw(t-\tau)}. \quad (1146)$$

Here  $w = \lambda_+ - \lambda_-$ . So we can see that for the equation (911) it's possible to deduce for this case of time-independent matrix  $\overline{H_S}$  if  $w \neq w'$  then  $A'_j(w, w') = 0$  so:

$$\widetilde{A}_j(t - \tau, t) = U(t) U^\dagger(t - \tau) A_j(t) U(t - \tau) U^\dagger(t) \quad (1147)$$

$$= U(t) \left( \sum_{w(t-\tau)} e^{-i(t-\tau)w(t-\tau)} A_j(w(t-\tau)) \right) U^\dagger(t) \quad (1148)$$

$$= \sum_{w(t-\tau)} e^{-i(t-\tau)w(t-\tau)} U(t) A_j(w(t-\tau)) U^\dagger(t) \quad (1149)$$

$$= \sum_{w'(t), w(t-\tau)} e^{-i(t-\tau)w(t-\tau)} e^{itw'(t)} A_j(w(t-\tau), w'(t)) \quad (1150)$$

$$= \sum_{w'(t), w(t-\tau)} e^{-i(t-\tau)w(t-\tau)} e^{itw'(t)} A_{jww'} \quad (1151)$$

$$= \sum_{w'(t), w(t-\tau)} e^{-i(t-\tau)w(t-\tau)} e^{itw'(t)} A_j(w) \delta_{ww'} \quad (1152)$$

$$= \sum_w e^{-i(t-\tau)w} e^{itw} A_j(w) \quad (1153)$$

$$= \sum_w e^{i\tau w} A_j(w) \quad (1154)$$

$$= U^\dagger(-\tau) A_j U(-\tau) \quad (1155)$$

So using now as reference the equation (1050) and  $A'_j(w, w') = 0$  we can deduce that:

$$\frac{d\overline{\rho_S}(t)}{dt} = -i[\overline{H_S}(t), \overline{\rho_S}(t)] - \sum_{ijw} K_{ij}^{\Re}(w, t) [A_i, A_j(w) \overline{\rho_S}(t) - \overline{\rho_S}(t) A_j^\dagger(w)] - i \sum_{ijw} K_{ij}^{\Im}(w, t) [A_i, A_j(w) \overline{\rho_S}(t) + \overline{\rho_S}(t) A_j^\dagger(w)] \quad (1156)$$

## B. Time-dependent polaron quantum master equation

Following the reference [1], when  $\Omega_k \ll \omega_k$  then  $f_k \approx g_k$  so we recover the full polaron transformation. It means from the equation (106) that  $B_z = 0$ . The Hamiltonian studied is given by:

$$H = \left( \delta + \sum_{\mathbf{k}} \left( g_{\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{\mathbf{k}}^* b_{\mathbf{k}} \right) \right) |1\rangle\langle 1| + \frac{\Omega(t)}{2} \sigma_x + \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}. \quad (1157)$$

If  $v_{\mathbf{k}} \approx g_{\mathbf{k}}$  then  $B(\tau) = B$ , so  $B$  is independent of the time. In order to reproduce the Hamiltonian of the equation (1157) using the Hamiltonian of the equation (1) we can say that  $\delta = \varepsilon_1(t)$ ,  $\varepsilon_0(t) = 0$ ,  $V_{10}(t) = \frac{\Omega(t)}{2}$ . Now given that  $v_{\mathbf{k}} \approx g_{\mathbf{k}}$  then, in this case and using the equation (??) and (??) we obtain the following transformed Hamiltonians:

$$\overline{H_S} = (\delta + R_1) |1\rangle\langle 1| + \frac{B\sigma_x}{2} \Omega(t), \quad (1158)$$

$$\overline{H_I} = \frac{\Omega(t)}{2} (B_x \sigma_x + B_y \sigma_y). \quad (1159)$$

In this case  $R_1 = \sum_{\mathbf{k}} \left( \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - 2 \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} g_{\mathbf{k}} \right)$  from (27) and given that  $v_{\mathbf{k}} \approx g_{\mathbf{k}}$  and  $\frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} = g_{\mathbf{k}}/\omega_{\mathbf{k}}$  then  $R_1 = \sum_{\mathbf{k}} (-\omega_{\mathbf{k}}^{-1} |g_{\mathbf{k}}|^2) = \sum_{\mathbf{k}} (-\omega_{\mathbf{k}} | \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} |^2)$  as expected, take  $\delta + R_1 = \delta'$ . If  $F(\omega_{\mathbf{k}}) = 1$  and using the equations (1118)-(1126) we can deduce that the only terms that survive are  $\Lambda_{11}(\tau)$  and  $\Lambda_{22}(\tau)$ . The phonon propagator for this case is:

$$\phi(\tau) = \int_0^\infty \frac{J(\omega)}{\omega^2} G_+(\tau) d\omega. \quad (1160)$$

Writing  $G_+(\tau) = \coth\left(\frac{\beta\omega}{2}\right) \cos(\omega\tau) - i \sin(\omega\tau)$  so (1160) can be written as:

$$\phi(\tau) = \int_0^\infty \frac{J(\omega)}{\omega^2} \left( \coth\left(\frac{\beta\omega}{2}\right) \cos(\omega\tau) - i \sin(\omega\tau) \right) d\omega. \quad (1161)$$

Writing the interaction Hamiltonian (1159) in the similar way to the equation (??) allow us to write  $A_1 = \sigma_x$ ,  $A_2 = \sigma_y$ ,  $B_1(t) = B_x$ ,  $B_2(t) = B_y$  and  $C_1(t) = \frac{\Omega(t)}{2} = C_2(t)$ . Now taking the equation (??) with  $\delta'|1\rangle\langle 1| = \frac{\delta'}{2}\sigma_z + \frac{\delta'}{2}\mathbb{I}$  help us to reproduce the hamiltonian of the reference [2]. Then  $\overline{H_S}$  is equal to:

$$\overline{H_S} = \frac{\delta'}{2}\sigma_z + \frac{B\sigma_x}{2}\Omega(t). \quad (1162)$$

As we can see the function  $B$  is a time-independent function because we consider that  $g_k$  doesn't depend of the time. In this case the relevant correlation functions are given by:

$$\Lambda_{11}(\tau) = \text{Tr}_B \left( \widetilde{B}_1(\tau) \widetilde{B}_1(0) \rho_B \right) \quad (1163)$$

$$= \frac{B^2}{2} \left( e^{\phi(\tau)} + e^{-\phi(\tau)} - 2 \right), \quad (1164)$$

$$\Lambda_{22}(\tau) = \text{Tr}_B \left( \widetilde{B}_2(\tau) \widetilde{B}_2(0) \rho_B \right) \quad (1165)$$

$$= \frac{B^2}{2} \left( e^{\phi(\tau)} + e^{-\phi(\tau)} \right). \quad (1166)$$

These functions match with the equations  $\Lambda_x(\tau)$  and  $\Lambda_y(\tau)$  of the reference [2] and  $\Lambda_i(\tau) = \Lambda_i(-\tau)$  for  $i \in \{x, y\}$  respectively. The master equation for this section based on the equation(447) is:

$$\frac{d\rho_S(t)}{dt} = -i \left[ \frac{\delta'}{2}\sigma_z + \frac{\Omega_r(t)\sigma_x}{2}, \rho_S(t) \right] - \sum_{i=1}^2 \int_0^t d\tau \left( C_i(t) C_i(t-\tau) \Lambda_{ii}(\tau) \left[ A_i, \widetilde{A}_i(t-\tau, t) \rho_S(t) \right] \right) \quad (1167)$$

$$+ C_i(t) C_i(t-\tau) \Lambda_{ii}(-\tau) \left[ \rho_S(t) \widetilde{A}_i(t-\tau, t), A_i \right]. \quad (1168)$$

Replacing  $C_i(t) = \frac{\Omega(t)}{2}$  and  $\widetilde{A}_i(t-\tau, t) = \widetilde{\sigma}_i(t-\tau, t)$ , also using the equations (1163) and (1166) on the equation (1168) we obtain that:

$$\frac{d\rho_S(t)}{dt} = -\frac{i}{2} [\delta'\sigma_z + \Omega_r(t)\sigma_x, \rho_S(t)] - \frac{\Omega(t)}{4} \int_0^t d\tau \Omega(t-\tau) ([\sigma_x, \widetilde{\sigma}_x(t-\tau, t) \rho_S(t)] \Lambda_x(\tau) \quad (1169)$$

$$+ [\sigma_y, \widetilde{\sigma}_y(t-\tau, t) \rho_S(t)] \Lambda_y(\tau) + [\rho_S(t) \widetilde{\sigma}_x(t-\tau, t), \sigma_x] \Lambda_x(\tau) + [\rho_S(t) \widetilde{\sigma}_y(t-\tau, t), \sigma_y] \Lambda_y(\tau)). \quad (1170)$$

As we can see  $[A_j, \widetilde{A}_i(t-\tau, t) \rho_S(t)]^\dagger = [\rho_S(t) \widetilde{A}_i(t-\tau, t), A_j]$ ,  $\Lambda_x(\tau) = \Lambda_x(-\tau)$  and  $\Lambda_y(\tau) = \Lambda_y(-\tau)$ , so the result obtained is the same master equation (21) of the reference [2] extended in the hermitian conjugate.

### C. Time-Dependent Weak-Coupling Limit

In order to prove that the master equation deduced reproduces the equation (S17) of the reference [3] we will impose that  $F(\omega) = 0$ , so there is no transformation in this case. As we can see from the definition (411) the only term that survives is  $\Lambda_{33}(\tau)$ . Taking  $\hbar = 1$  the Hamiltonian of the reference can be written in the form:

$$H = \Delta|1\rangle\langle 1| + \frac{\Omega(t)}{2} (|1\rangle\langle 0| + |0\rangle\langle 1|) + \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + |1\rangle\langle 1| \sum_{\mathbf{k}} \left( g_{\mathbf{k}} b_{\mathbf{k}}^\dagger + g_{\mathbf{k}}^* b_{\mathbf{k}} \right). \quad (1171)$$

Using the equation (960), from the fact that the Hamiltonian is time-independent in the evolution time allow us to write:

$$\frac{d\rho_S}{dt} = -i [H_S(t), \rho_S(t)] - \frac{1}{2} \sum_w \gamma_{33}(w, t) \left[ A_3, A_3(w) \rho_S(t) - \rho_S(t) A_3^\dagger(w) \right] \quad (1172)$$

$$- \sum_w S_{33}(w, t) \left[ A_3, A_3(w) \rho_S(t) + \rho_S(t) A_3^\dagger(w) \right] \Bigg). \quad (1173)$$

The correlation functions are relevant if  $F(\omega) = 0$  for the weak-coupling approximation are:

$$\Lambda_{33}(\tau) = \int_0^\infty d\omega J(\omega) G_+(\tau), \quad (1174)$$

$$\Lambda_{33}(-\tau) = \int_0^\infty d\omega J(\omega) G_+(-\tau). \quad (1175)$$

In our case  $A_3 = \frac{\mathbb{I} + \sigma_z}{2}$ , the equation (1173) can be transformed in

$$\frac{d\rho_S}{dt} = -i [H_S(t), \rho_S(t)] - \sum_w (K_{33}(w, t) [A_3, A_3(w) \rho_S(t)] + K_{33}^*(w, t) [\rho_S(t) A_3(w), A_3]). \quad (1176)$$

As the paper suggest we will consider that the quantum system is in resonance, so  $\Delta = 0$  and furthermore, the relaxation time of the bath is less than the evolution time to be considered, so the frequency of the Rabi frequency of the laser can be taken as constant and equal to  $\tilde{\Omega}$ . To find the matrices  $A_3(w)$ , we have to remember that  $H_S = \frac{\Omega(t)}{2} (|1\rangle\langle 0| + |0\rangle\langle 1|)$ , this Hamiltonian using the approximation  $\tilde{\Omega}$  have the following eigenvalues and eigenvectors:

$$\lambda_+ = \frac{\tilde{\Omega}}{2}, \quad (1177)$$

$$|+\rangle = \frac{1}{\sqrt{2}} (|1\rangle + |0\rangle), \quad (1178)$$

$$\lambda_- = -\frac{\tilde{\Omega}}{2}, \quad (1179)$$

$$|-\rangle = \frac{1}{\sqrt{2}} (|0\rangle - |1\rangle). \quad (1180)$$

The elements of the decomposition matrices are:

$$\langle + | \frac{1 + \sigma_z}{2} | + \rangle = \frac{1}{2} \begin{pmatrix} 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (1181)$$

$$= \frac{1}{2}, \quad (1182)$$

$$\langle - | \frac{1 + \sigma_z}{2} | - \rangle = \frac{1}{2} \begin{pmatrix} 1 & -1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad (1183)$$

$$= \frac{1}{2}, \quad (1184)$$

$$\langle - | \frac{1 + \sigma_z}{2} | + \rangle = \frac{1}{2} \begin{pmatrix} 1 & -1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (1185)$$

$$= -\frac{1}{2}. \quad (1186)$$

The decomposition matrices are

$$A_3(0) = \frac{1}{2}|+\rangle\langle+| + \frac{1}{2}|-\rangle\langle-| \quad (1187)$$

$$= \frac{\mathbb{I}}{2}, \quad (1188)$$

$$A_3(\eta) = -\frac{1}{2}|-\rangle\langle+| \quad (1189)$$

$$= \frac{1}{4}(\sigma_z + i\sigma_y), \quad (1190)$$

$$A_3(-\eta) = -\frac{1}{2}|+\rangle\langle-| \quad (1191)$$

$$= \frac{1}{4}(\sigma_z - i\sigma_y). \quad (1192)$$

Neglecting the term proportional to the identity in the Hamiltonian we obtain that:

$$\frac{d\rho_S(t)}{dt} = -i\frac{\tilde{\Omega}}{2}[\sigma_x, \rho_S(t)] - K_{33}(\tilde{\Omega}, t) \left[ \frac{\sigma_z}{2}, \frac{1}{4}(\sigma_z + i\sigma_y)\rho_S(t) \right] - K_{33}(-\tilde{\Omega}, t) \left[ \frac{\sigma_z}{2}, \frac{1}{4}(\sigma_z - i\sigma_y)\rho_S(t) \right] \quad (1193)$$

$$- K_{33}^*(\tilde{\Omega}, t) \left[ \rho_S(t) \frac{1}{4}(\sigma_z + i\sigma_y), \frac{\sigma_z}{2} \right] - K_{33}^*(-\tilde{\Omega}, t) \left[ \rho_S(t) \frac{1}{4}(\sigma_z - i\sigma_y), \frac{\sigma_z}{2} \right]. \quad (1194)$$

Calculating the response functions extending the upper limit of  $\tau$  to  $\infty$ , we obtain:

$$K_{33}(\tilde{\Omega}) = \int_0^\infty \int_0^\infty J(\omega) G_+(\tau) e^{i\tilde{\Omega}\tau} d\tau d\omega \quad (1195)$$

$$= \int_0^\infty \int_0^\infty J(\omega) e^{i\tilde{\Omega}\tau} ((n(\omega) + 1)e^{-i\tau\omega} + n(\omega)e^{i\tau\omega}) d\tau d\omega \quad (1196)$$

$$= \int_0^\infty \int_0^\infty J(\omega) e^{i\tilde{\Omega}\tau} (n(\omega) + 1)e^{-i\tau\omega} d\tau d\omega \quad (1197)$$

$$= \int_0^\infty \int_0^\infty J(\omega) (n(\omega) + 1)e^{i\tilde{\Omega}\tau - i\tau\omega} d\tau d\omega \quad (1198)$$

$$= \int_0^\infty J(\omega) (n(\omega) + 1) \pi \delta(\tilde{\Omega} - \omega) d\omega \quad (1199)$$

$$= \pi J(\tilde{\Omega}) (n(\tilde{\Omega}) + 1), \quad (1200)$$

$$K_{33}(-\tilde{\Omega}) = \int_0^\infty \int_0^\infty J(\omega) G_+(\tau) e^{-i\tilde{\Omega}\tau} d\tau d\omega \quad (1201)$$

$$= \int_0^\infty \int_0^\infty J(\omega) e^{-i\tilde{\Omega}\tau} ((n(\omega) + 1)e^{-i\tau\omega} + n(\omega)e^{i\tau\omega}) d\tau d\omega \quad (1202)$$

$$= \int_0^\infty \int_0^\infty J(\omega) e^{-i\tilde{\Omega}\tau} n(\omega) e^{i\tau\omega} d\tau d\omega \quad (1203)$$

$$= \int_0^\infty \int_0^\infty J(\omega) n(\omega) e^{-i\tilde{\Omega}\tau + i\tau\omega} d\tau d\omega \quad (1204)$$

$$= \int_0^\infty J(\omega) n(\omega) \pi \delta(-\tilde{\Omega} + \omega) d\omega \quad (1205)$$

$$= \pi J(\tilde{\Omega}) n(\tilde{\Omega}). \quad (1206)$$

Here we have used  $\int_0^\infty ds e^{\pm i\epsilon s} = \pi \delta(\epsilon) \pm i \frac{\text{V.P.}}{\epsilon}$ , where V.P. denotes the Cauchy's principal value. These principal values are ignored because they lead to small renormalizations of the Hamiltonian. Furthermore we don't take

account of value associated to the matrix  $A_3(0)$  because the spectral density  $J(\omega)$  is equal to zero when  $\omega = 0$ . Replacing in the equation (1193) lead us to obtain:

$$\frac{d\rho_S(t)}{dt} = -i\frac{\tilde{\Omega}}{2} [\sigma_x, \rho_S(t)] - \frac{\pi}{8} J(\tilde{\Omega}) \left( (n(\tilde{\Omega}) + 1) [\sigma_z, (\sigma_z + i\sigma_y) \rho_S(t)] + n(\tilde{\Omega}) [\sigma_z, (\sigma_z - i\sigma_y) \rho_S(t)] \right) \quad (1207)$$

$$- \frac{\pi}{8} J(\tilde{\Omega}) \left( (n(\tilde{\Omega}) + 1) [\rho_S(t) (\sigma_z + i\sigma_y), \sigma_z] + n(\tilde{\Omega}) [\rho_S(t) (\sigma_z - i\sigma_y), \sigma_z] \right). \quad (1208)$$

This is the same result than the equation (S17), so we have proved that our general master equation allows to reproduce the results of the weak-coupling time-dependent. Now the master equation in the evolution time is given by

$$\frac{d\rho_S(t)}{dt} = -i\frac{\Omega(t)}{2} [\sigma_x, \rho_S(t)] - \frac{\pi}{8} J(\Omega(t)) \left( (n(\Omega(t)) + 1) [\sigma_z, (\sigma_z + i\sigma_y) \rho_S(t)] + n(\Omega(t)) [\sigma_z, (\sigma_z - i\sigma_y) \rho_S(t)] \right) \quad (1209)$$

$$- \frac{\pi}{8} J(\Omega(t)) \left( (n(\Omega(t)) + 1) [\rho_S(t) (\sigma_z + i\sigma_y), \sigma_z] + n(\Omega(t)) [\rho_S(t) (\sigma_z - i\sigma_y), \sigma_z] \right). \quad (1210)$$

## V. TIME-DEPENDENT MULTI-SITE MODEL WITH V BATHS COUPLING

Let's consider the following Hamiltonian for a system of m-level system coupled to v-baths. We start with a time-dependent Hamiltonian of the form:

$$H(t) = H_S(t) + H_I + H_B, \quad (1211)$$

$$H_S(t) = \sum_n \varepsilon_n(t) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m|, \quad (1212)$$

$$H_I = \sum_{nuk} |n\rangle\langle n| \left( g_{nuk} b_{uk}^\dagger + g_{nuk}^* b_{uk} \right), \quad (1213)$$

$$H_B = \sum_{uk} \omega_{uk} b_{uk}^\dagger b_{uk}. \quad (1214)$$

### A. Variational Transformation

We consider the following operator:

$$V = \sum_{nuk} |n\rangle\langle n| \omega_{uk}^{-1} \left( f_{nuk} b_{uk}^\dagger - f_{nuk}^* b_{uk} \right) \quad (1215)$$

At first let's obtain  $e^{\pm V}$  under the transformation (1215), consider  $\hat{\varphi}_n = \sum_{uk} \omega_{uk}^{-1} \left( f_{nuk} b_{uk}^\dagger - f_{nuk}^* b_{uk} \right)$ , so the equation (1215) can be written as  $V = \sum_n |n\rangle\langle n| \hat{\varphi}_n$ , then we have:

$$e^{\pm V} = e^{\pm \sum_n |n\rangle\langle n| \hat{\varphi}_n} \quad (1216)$$

$$= \mathbb{I} \pm \sum_n |n\rangle\langle n| \hat{\varphi}_n + \frac{(\sum_n |n\rangle\langle n| \hat{\varphi}_n)^2}{2!} + \dots \quad (1217)$$

$$= \mathbb{I} \pm \sum_n |n\rangle\langle n| \hat{\varphi}_n + \frac{\sum_n |n\rangle\langle n| \hat{\varphi}_n^2}{2!} + \dots \quad (1218)$$

$$= \sum_n |n\rangle\langle n| \pm \sum_n |n\rangle\langle n| \hat{\varphi}_n + \frac{\sum_n |n\rangle\langle n| \hat{\varphi}_n^2}{2!} + \dots \quad (1219)$$

$$= \sum_n |n\rangle\langle n| \left( \mathbb{I} \pm \hat{\varphi}_n + \frac{\hat{\varphi}_n^2}{2!} + \dots \right) \quad (1220)$$

$$= \sum_n |n\rangle\langle n| e^{\pm \hat{\varphi}_n} \quad (1221)$$

Given that  $\left[ f_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger - f_{nu\mathbf{k}}^* b_{u\mathbf{k}}, f_{nu'\mathbf{k}'} b_{u'\mathbf{k}'}^\dagger - f_{nu'\mathbf{k}'}^* b_{u'\mathbf{k}'} \right] = 0$  for all  $\mathbf{k}', \mathbf{k}$  and  $u, u'$  then we can proof using the Zassenhaus formula and defining  $D(\pm \alpha_{nu\mathbf{k}}) = e^{\pm (\alpha_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger - \alpha_{nu\mathbf{k}}^* b_{u\mathbf{k}})}$  in the same way than (23) with  $\alpha_{nu\mathbf{k}} = \frac{f_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}}$ :

$$e^{\pm \sum_{u\mathbf{k}} \omega_{u\mathbf{k}}^{-1} (f_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger - f_{nu\mathbf{k}}^* b_{u\mathbf{k}})} = \prod_u e^{\pm \sum_{\mathbf{k}} \omega_{u\mathbf{k}}^{-1} (f_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger - f_{nu\mathbf{k}}^* b_{u\mathbf{k}})} \quad (1222)$$

$$= \prod_u \left( \prod_{\mathbf{k}} e^{\pm \omega_{u\mathbf{k}}^{-1} (f_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger - f_{nu\mathbf{k}}^* b_{u\mathbf{k}})} \right) \quad (1223)$$

$$= \prod_u \left( \prod_{\mathbf{k}} D(\pm \alpha_{nu\mathbf{k}}) \right) \quad (1224)$$

$$= \prod_{u\mathbf{k}} D(\pm \alpha_{nu\mathbf{k}}) \quad (1225)$$

$$= \prod_u B_{nu\pm} \quad (1226)$$

$$B_{nu\pm} \equiv \prod_{\mathbf{k}} D(\pm \alpha_{nu\mathbf{k}}) \quad (1227)$$

As we can see  $e^{-V} = \sum_n |n\rangle\langle n| \prod_u B_{nu-}$  and  $e^V = \sum_n |n\rangle\langle n| \prod_u B_{nu+}$  this implies that  $e^{-V} e^V = \mathbb{I}$ . This allows us to write the canonical transformation in the following explicit way:

$$e^V A e^{-V} = \left( \sum_n |n\rangle\langle n| \prod_u B_{nu+} \right) A \left( \sum_n |n\rangle\langle n| \prod_u B_{nu-} \right) \quad (1228)$$

Now let's obtain the canonical transformation of the principal elements of the Hamiltonian (1211):

$$\overline{|0\rangle\langle 0|} = \left( \sum_n |n\rangle\langle n| \prod_u B_{nu+} \right) |0\rangle\langle 0| \left( \sum_n |n\rangle\langle n| \prod_u B_{nu-} \right), \quad (1229)$$

$$= \prod_u B_{0u+} |0\rangle\langle 0| \prod_u B_{0u-}, \quad (1230)$$

$$= |0\rangle\langle 0| \prod_u B_{0u+} \prod_u B_{0u-}, \quad (1231)$$

$$= |0\rangle\langle 0| \prod_u B_{0u+} B_{0u-} \quad (1232)$$

$$= |0\rangle\langle 0| \prod_u \mathbb{I} \quad (1233)$$

$$= |0\rangle\langle 0|. \quad (1234)$$

$$\overline{|m\rangle\langle n|} = \left( \sum_n |n\rangle\langle n| \prod_u B_{nu+} \right) |m\rangle\langle n| \left( \sum_n |n\rangle\langle n| \prod_u B_{nu-} \right), \quad (1235)$$

$$= |m\rangle\langle m| \prod_u B_{mu+} |m\rangle\langle n| \prod_u B_{nu-}, \quad (1236)$$

$$= |m\rangle\langle n| \prod_u B_{mu+} \prod_u B_{nu-}, \quad (1237)$$

$$= |m\rangle\langle n| \prod_u (B_{mu+} B_{nu-}), \quad m \neq n, \quad (1238)$$

$$= |m\rangle\langle n| \prod_u \left( \prod_{\mathbf{k}} D(\alpha_{muk}) \prod_{\mathbf{k}} D(-\alpha_{nuk}) \right), \quad (1239)$$

$$= |m\rangle\langle n| \prod_u \prod_{\mathbf{k}} (D(\alpha_{muk}) D(-\alpha_{nuk})), \quad (1240)$$

$$= |m\rangle\langle n| \prod_{u\mathbf{k}} \left( D(\alpha_{muk} - \alpha_{nuk}) \exp \left( \frac{1}{2} (-\alpha_{muk} \alpha_{nuk}^* + \alpha_{muk}^* \alpha_{nuk}) \right) \right). \quad (1241)$$

$$\Pi_u(B_{mu+} B_{nu-}) = \prod_{u\mathbf{k}} \left( D(\alpha_{muk} - \alpha_{nuk}) \exp \left( \frac{1}{2} (-\alpha_{muk} \alpha_{nuk}^* + \alpha_{muk}^* \alpha_{nuk}) \right) \right). \quad (1242)$$

$$\overline{\sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}}} = (\sum_n |n\rangle\langle n| \prod_u B_{nu+}) \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} (\sum_n |n\rangle\langle n| \prod_u B_{nu-}), \quad (1243)$$

$$= (|0\rangle\langle 0| \prod_u B_{0u+} + |1\rangle\langle 1| \prod_u B_{1u+} + \dots) (\sum_n |n\rangle\langle n| \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}}) (|0\rangle\langle 0| \prod_u B_{0u-} + |1\rangle\langle 1| \prod_u B_{1u-} + \dots), \quad (1244)$$

$$= |0\rangle\langle 0| \prod_u B_{0u+} + \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} \prod_u B_{0u-} + |1\rangle\langle 1| \prod_u B_{1u+} + \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} \prod_u B_{1u-} + \dots, \quad (1245)$$

$$= |0\rangle\langle 0| \prod_u B_{0u+} + (\sum_{\mathbf{k}} \omega_{0\mathbf{k}} b_{0\mathbf{k}}^\dagger b_{0\mathbf{k}} + \sum_{\mathbf{k}} \omega_{1\mathbf{k}} b_{1\mathbf{k}}^\dagger b_{1\mathbf{k}} + \dots) \prod_u B_{0u-} + |1\rangle\langle 1| \prod_u B_{1u+} + (\sum_{\mathbf{k}} \omega_{0\mathbf{k}} b_{0\mathbf{k}}^\dagger b_{0\mathbf{k}} + \sum_{\mathbf{k}} \omega_{1\mathbf{k}} b_{1\mathbf{k}}^\dagger b_{1\mathbf{k}} + \dots) \prod_u B_{1u-} + \dots \quad (1246)$$

$$= |0\rangle\langle 0| (\prod_u B_{0u+} + \sum_{\mathbf{k}} \omega_{0\mathbf{k}} b_{0\mathbf{k}}^\dagger b_{0\mathbf{k}} \prod_u B_{0u-} + \prod_u B_{0u+} + \sum_{\mathbf{k}} \omega_{1\mathbf{k}} b_{1\mathbf{k}}^\dagger b_{1\mathbf{k}} \prod_u B_{0u-} + \dots) \quad (1247)$$

$$+ |1\rangle\langle 1| (\prod_u B_{1u+} + \sum_{\mathbf{k}} \omega_{0\mathbf{k}} b_{0\mathbf{k}}^\dagger b_{0\mathbf{k}} \prod_u B_{1u-} + \prod_u B_{1u+} + \sum_{\mathbf{k}} \omega_{1\mathbf{k}} b_{1\mathbf{k}}^\dagger b_{1\mathbf{k}} \prod_u B_{1u-} + \dots) + \dots \quad (1248)$$

$$= |0\rangle\langle 0| \left( \sum_{\mathbf{k}} \omega_{0\mathbf{k}} \left( b_{0\mathbf{k}}^\dagger - \frac{v_{00\mathbf{k}}}{\omega_{0\mathbf{k}}} \right) \left( b_{0\mathbf{k}} - \frac{v_{00\mathbf{k}}}{\omega_{0\mathbf{k}}} \right) + \sum_{\mathbf{k}} \omega_{1\mathbf{k}} \left( b_{1\mathbf{k}}^\dagger - \frac{v_{01\mathbf{k}}}{\omega_{1\mathbf{k}}} \right) \left( b_{0\mathbf{k}} - \frac{v_{01\mathbf{k}}}{\omega_{1\mathbf{k}}} \right) + \dots \right) \quad (1249)$$

$$+ |1\rangle\langle 1| \left( \sum_{\mathbf{k}} \omega_{0\mathbf{k}} \left( b_{0\mathbf{k}}^\dagger - \frac{v_{10\mathbf{k}}}{\omega_{0\mathbf{k}}} \right) \left( b_{0\mathbf{k}} - \frac{v_{10\mathbf{k}}}{\omega_{0\mathbf{k}}} \right) + \sum_{\mathbf{k}} \omega_{1\mathbf{k}} \left( b_{1\mathbf{k}}^\dagger - \frac{v_{11\mathbf{k}}}{\omega_{1\mathbf{k}}} \right) \left( b_{0\mathbf{k}} - \frac{v_{11\mathbf{k}}}{\omega_{1\mathbf{k}}} \right) + \dots \right) + \dots \quad (1250)$$

$$= |0\rangle\langle 0| \left( \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} \left( b_{u\mathbf{k}}^\dagger - \frac{v_{0u\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \left( b_{u\mathbf{k}} - \frac{v_{0u\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) + |1\rangle\langle 1| \left( \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} \left( b_{u\mathbf{k}}^\dagger - \frac{v_{1u\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \left( b_{u\mathbf{k}} - \frac{v_{1u\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) + \dots \quad (1251)$$

$$= \sum_{n\mathbf{u}\mathbf{k}} |n\rangle\langle n| \left( \omega_{u\mathbf{k}} \left( b_{u\mathbf{k}}^\dagger - \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \left( b_{u\mathbf{k}} - \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) \quad (1252)$$

$$= \sum_{n\mathbf{u}\mathbf{k}} |n\rangle\langle n| \left( \omega_{u\mathbf{k}} \left( b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} - \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} b_{u\mathbf{k}}^\dagger - \frac{v_{nu\mathbf{k}}^*}{\omega_{u\mathbf{k}}} b_{u\mathbf{k}} + \left| \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} \right|^2 \right) \right) \quad (1253)$$

$$= \sum_{n\mathbf{u}\mathbf{k}} |n\rangle\langle n| \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} + \sum_{n\mathbf{u}\mathbf{k}} |n\rangle\langle n| \left( \frac{|v_{nu\mathbf{k}}|^2}{\omega_{u\mathbf{k}}} - v_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger - v_{nu\mathbf{k}}^* b_{u\mathbf{k}} \right) \quad (1254)$$

$$= \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} + \sum_{n\mathbf{u}\mathbf{k}} |n\rangle\langle n| \left( \frac{|v_{nu\mathbf{k}}|^2}{\omega_{u\mathbf{k}}} - (v_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger + v_{nu\mathbf{k}}^* b_{u\mathbf{k}}) \right) \quad (1255)$$

The transformed Hamiltonians of the equations (1212) to (1214) written in terms of (1229) to (1253) are:



$$\overline{H_S(t)} = \overline{\sum_n \varepsilon_n(t) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m|} \quad (1256)$$

$$= \overline{\sum_n \varepsilon_n(t) |n\rangle\langle n|} + \overline{\sum_{n \neq m} V_{nm}(t) |n\rangle\langle m|} \quad (1257)$$

$$= \sum_n \varepsilon_n(t) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| \prod_u (B_{mu+} B_{nu-}) \quad (1258)$$

$$\overline{H_I} = \left( \sum_n |n\rangle\langle n| \prod_u B_{nu+} \right) \left( \sum_{n \neq m} |n\rangle\langle m| (g_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger + g_{nu\mathbf{k}}^* b_{u\mathbf{k}}) \right) \left( \sum_n |n\rangle\langle n| \prod_u B_{nu-} \right) \quad (1259)$$

$$= \left( \sum_n |n\rangle\langle n| \prod_u B_{nu+} \right) \left( \sum_{u\mathbf{k}} |0\rangle\langle 0| (g_{0u\mathbf{k}} b_{u\mathbf{k}}^\dagger + g_{0u\mathbf{k}}^* b_{u\mathbf{k}}) + \dots \right) \left( \sum_n |n\rangle\langle n| \prod_u B_{nu-} \right) \quad (1260)$$

$$= \prod_u B_{0u+} \sum_{u\mathbf{k}} |0\rangle\langle 0| (g_{0u\mathbf{k}} b_{u\mathbf{k}}^\dagger + g_{0u\mathbf{k}}^* b_{u\mathbf{k}}) \prod_u B_{0u-} + \prod_u B_{1u+} \sum_{u\mathbf{k}} |1\rangle\langle 1| (g_{1u\mathbf{k}} b_{u\mathbf{k}}^\dagger + g_{1u\mathbf{k}}^* b_{u\mathbf{k}}) \prod_u B_{1u-} + \dots \quad (1261)$$

$$= \sum_{u\mathbf{k}} |0\rangle\langle 0| (g_{0u\mathbf{k}} \prod_u B_{0u+} b_{u\mathbf{k}}^\dagger \prod_u B_{0u-} + g_{0u\mathbf{k}}^* \prod_u B_{0u+} b_{u\mathbf{k}} \prod_u B_{0u-}) + \sum_{u\mathbf{k}} |1\rangle\langle 1| (g_{1u\mathbf{k}} \prod_u B_{1u+} b_{u\mathbf{k}}^\dagger \prod_u B_{1u-} + g_{1u\mathbf{k}}^* \prod_u B_{1u+} b_{u\mathbf{k}} \prod_u B_{1u-}) + \dots \quad (1262)$$

$$= \sum_{u\mathbf{k}} |0\rangle\langle 0| \left( g_{0u\mathbf{k}} \left( b_{u\mathbf{k}}^\dagger - \frac{v_{0u\mathbf{k}}^*}{\omega_{u\mathbf{k}}} \right) + g_{0u\mathbf{k}}^* \left( b_{u\mathbf{k}} - \frac{v_{0u\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) + \sum_{u\mathbf{k}} |1\rangle\langle 1| \left( g_{1u\mathbf{k}} \left( b_{u\mathbf{k}}^\dagger - \frac{v_{1u\mathbf{k}}^*}{\omega_{u\mathbf{k}}} \right) + g_{1u\mathbf{k}}^* \left( b_{u\mathbf{k}} - \frac{v_{1u\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) + \dots \quad (1263)$$

$$= \sum_{n \neq u\mathbf{k}} |n\rangle\langle n| \left( g_{nu\mathbf{k}} \left( b_{u\mathbf{k}}^\dagger - \frac{v_{nu\mathbf{k}}^*}{\omega_{u\mathbf{k}}} \right) + g_{nu\mathbf{k}}^* \left( b_{u\mathbf{k}} - \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) \quad (1264)$$

$$= \sum_{n \neq u\mathbf{k}} |n\rangle\langle n| \left( g_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger + g_{nu\mathbf{k}}^* b_{u\mathbf{k}} - \left( g_{nu\mathbf{k}} \frac{v_{nu\mathbf{k}}^*}{\omega_{u\mathbf{k}}} + g_{nu\mathbf{k}}^* \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) \quad (1265)$$

$$\overline{H_B} = \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} + \sum_{n \neq u\mathbf{k}} |n\rangle\langle n| \left( \frac{|v_{nu\mathbf{k}}|^2}{\omega_{u\mathbf{k}}} - (v_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger + v_{nu\mathbf{k}}^* b_{u\mathbf{k}}) \right) \quad (1266)$$

Joining this terms allow us to write the transformed Hamiltonian as:

$$\overline{H} = \sum_n \varepsilon_n(t) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| \prod_u (B_{mu+} B_{nu-}) + \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} + \sum_{n \neq u\mathbf{k}} |n\rangle\langle n| \left( \frac{|v_{nu\mathbf{k}}|^2}{\omega_{u\mathbf{k}}} - (v_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger + v_{nu\mathbf{k}}^* b_{u\mathbf{k}}) \right) \quad (1267)$$

$$+ \sum_{n \neq u\mathbf{k}} |n\rangle\langle n| \left( g_{nu\mathbf{k}} b_{u\mathbf{k}}^\dagger + g_{nu\mathbf{k}}^* b_{u\mathbf{k}} - \left( g_{nu\mathbf{k}} \frac{v_{nu\mathbf{k}}^*}{\omega_{u\mathbf{k}}} + g_{nu\mathbf{k}}^* \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) \quad (1268)$$

Let's define the following functions:

$$R_n(t) = \sum_{u\mathbf{k}} \left( \frac{|v_{nu\mathbf{k}}|^2}{\omega_{u\mathbf{k}}} - \left( g_{nu\mathbf{k}} \frac{v_{nu\mathbf{k}}^*}{\omega_{u\mathbf{k}}} + g_{nu\mathbf{k}}^* \frac{v_{nu\mathbf{k}}}{\omega_{u\mathbf{k}}} \right) \right) \quad (1269)$$

$$B_{z,n}(t) = \sum_{u\mathbf{k}} \left( (g_{nu\mathbf{k}} - v_{nu\mathbf{k}}) b_{u\mathbf{k}}^\dagger + (g_{nu\mathbf{k}} - v_{nu\mathbf{k}})^* b_{u\mathbf{k}} \right) \quad (1270)$$

Using the previous functions we have that (1267) can be re-written in the following way:

$$\overline{H} = \sum_n \varepsilon_n(t) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| \prod_u (B_{mu+} B_{nu-}) + \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} + \sum_n R_n(t) |n\rangle\langle n| + \sum_n B_{z,n}(t) |n\rangle\langle n| \quad (1271)$$

$$(1272)$$

Now in order to separate the elements of the hamiltonian (1272) let's follow the references of the equations (??) and (??) to separate the hamiltonian, before proceeding to do this we need to consider the term of the form:

$$\langle \Pi_u (B_{mu+} B_{nu-}) \rangle_{\overline{H_0}} = \langle \Pi_{uk} (D(\alpha_{muk} - \alpha_{nuk}) \exp(\frac{1}{2}(-\alpha_{muk} \alpha_{nuk}^* + \alpha_{muk}^* \alpha_{nuk})) \rangle_{\overline{H_0}} \quad (1273)$$

$$= (\Pi_{uk} \exp(\frac{1}{2}(-\alpha_{muk} \alpha_{nuk}^* + \alpha_{muk}^* \alpha_{nuk}))) \langle \Pi_{uk} D(\alpha_{muk} - \alpha_{nuk}) \rangle_{\overline{H_0}} \quad (1274)$$

$$= \left( \Pi_{uk} \exp\left(\frac{(v_{muk}^* v_{nuk} - v_{nuk} v_{muk}^*)}{2\omega_{uk}^2}\right) \right) \Pi_u \exp\left(-\frac{1}{2} \sum_{\mathbf{k}} \frac{|v_{muk} - v_{nuk}|^2}{\omega_{uk}^2} \coth\left(\frac{\beta\omega_{uk}}{2}\right)\right) \quad (1275)$$

$$\equiv B_{nm} \quad (1276)$$

$$\langle \Pi_u (B_{nu+} B_{mu-}) \rangle_{\overline{H_0}} = \left( \Pi_{uk} \exp\left(\frac{(v_{nuk}^* v_{muk} - v_{muk} v_{nuk}^*)}{2\omega_{uk}^2}\right) \right) \Pi_u \exp\left(-\frac{1}{2} \sum_{\mathbf{k}} \frac{|v_{muk} - v_{nuk}|^2}{\omega_{uk}^2} \coth\left(\frac{\beta\omega_{uk}}{2}\right)\right) \quad (1277)$$

$$= B_{nm}^* \quad (1278)$$

Following the reference [4] we define:

$$J_{nm} = \prod_u (B_{mu+} B_{nu-}) - B_{nm} \quad (1279)$$

As we can see:

$$J_{nm}^\dagger = \left( \prod_u (B_{mu+} B_{nu-}) - B_{nm} \right)^\dagger \quad (1280)$$

$$= \prod_u (B_{nu+} B_{mu-}) - B_{nm}^* \quad (1281)$$

$$= \prod_u (B_{nu+} B_{mu-}) - B_{mn} \quad (1282)$$

$$= J_{mn} \quad (1283)$$

We can separate the Hamiltonian (1272) on the following way using similar arguments to the precedent sections to obtain:

$$\overline{H_{\bar{S}}}(t) = \sum_n (\varepsilon_n(t) + R_n) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| B_{nm} \quad (1284)$$

$$\overline{H_{\bar{I}}}(t) = \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| J_{nm} + \sum_n B_{z,n}(t) |n\rangle\langle n|, \quad (1285)$$

$$\overline{H_{\bar{B}}}(t) = \sum_{u\mathbf{k}} \omega_{u\mathbf{k}} b_{u\mathbf{k}}^\dagger b_{u\mathbf{k}} \quad (1286)$$

## B. Free-energy minimization

As first approach let's consider the minimization of the free-energy through the Feynman-Bogoliubov inequality

$$A \leq A_B \equiv -\frac{1}{\beta} \ln \left( \text{Tr} \left( e^{-\beta(\overline{H_{\bar{S}}}(t) + \overline{H_{\bar{B}}})} \right) \right) + \langle \overline{H_{\bar{I}}} \rangle_{\overline{H_{\bar{S}}}(t) + \overline{H_{\bar{B}}}} + O \left( \left\langle \overline{H_{\bar{I}}}^2 \right\rangle_{\overline{H_{\bar{S}}}(t) + \overline{H_{\bar{B}}}} \right). \quad (1287)$$

Taking the equations (244)-(252) and given that  $\text{Tr} \left( e^{-\beta \overline{H_{\bar{S}}}(t)} \right) = C(R_0, R_1, \dots, R_{d-1}, B_{01}, \dots, B_{0(d-1)}, \dots, B_{(d-2)(d-1)})$ , where each  $R_i$  and  $B_{kj}$  depend of the set of variational parameters  $\{v_{nuk}\}$ . Given that the numbers  $v_{nuk}$  are complex then we can separate them as  $v_{nuk} = v_{nuk}^{\Re} + i v_{nuk}^{\Im}$ . So our approach will be based on the derivation respect to  $v_{nuk}^{\Re}$  and  $v_{nuk}^{\Im}$ . The Hamiltonian  $\overline{H_{\bar{S}}}(t)$  can be written like:

$$\overline{H_S(t)} = \sum_n \left( \varepsilon_n(t) + \sum_{uk} \left( \frac{|v_{nuk}|^2}{\omega_{uk}} - \left( g_{nuk} \frac{v_{nuk}^*}{\omega_{uk}} + g_{nuk}^* \frac{v_{nuk}}{\omega_{uk}} \right) \right) \right) |n\rangle\langle n| \quad (1288)$$

$$+ \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| \left( \Pi_{uk} \exp \left( \frac{(v_{muk}^* v_{nuk} - v_{muk} v_{nuk}^*)}{2\omega_{uk}^2} \right) \right) \Pi_u \exp \left( -\frac{1}{2} \sum_k \frac{|v_{muk} - v_{nuk}|^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1289)$$

$$= \sum_n \left( \varepsilon_n(t) + \sum_{uk} \left( \frac{|v_{nuk}|^2}{\omega_{uk}} - \frac{g_{nuk} v_{nuk}^* + g_{nuk}^* v_{nuk}}{\omega_{uk}} \right) \right) |n\rangle\langle n| \quad (1290)$$

$$+ \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| \left( \Pi_{uk} \exp \left( \frac{(v_{muk}^* v_{nuk} - v_{muk} v_{nuk}^*)}{2\omega_{uk}^2} \right) \right) \Pi_u \exp \left( -\frac{1}{2} \sum_k \frac{|v_{muk} - v_{nuk}|^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1291)$$

$$= \sum_n \left( \varepsilon_n(t) + \sum_{uk} \left( \frac{(v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2}{\omega_{uk}} - \frac{(g_{nuk} + g_{nuk}^*) v_{nuk}^{\Re} + i v_{nuk}^{\Im} (g_{nuk}^* - g_{nuk})}{\omega_{uk}} \right) \right) |n\rangle\langle n| \quad (1292)$$

$$+ \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| \left( \Pi_{uk} \exp \left( \frac{(v_{muk}^* v_{nuk} - v_{muk} v_{nuk}^*)}{2\omega_{uk}^2} \right) \right) \Pi_u \exp \left( -\frac{1}{2} \sum_k \frac{|v_{muk} - v_{nuk}|^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1293)$$

$$v_{muk}^* v_{nuk} - v_{muk} v_{nuk}^* = (v_{muk}^{\Re} - i v_{muk}^{\Im}) (v_{nuk}^{\Re} + i v_{nuk}^{\Im}) - (v_{muk}^{\Re} + i v_{muk}^{\Im}) (v_{nuk}^{\Re} - i v_{nuk}^{\Im}) \quad (1294)$$

$$= (v_{muk}^{\Re} v_{nuk}^{\Re} + i v_{nuk}^{\Im} v_{muk}^{\Re} - i v_{muk}^{\Re} v_{nuk}^{\Im} + v_{muk}^{\Im} v_{nuk}^{\Im}) \quad (1295)$$

$$- (v_{muk}^{\Re} v_{nuk}^{\Re} - i v_{nuk}^{\Im} v_{muk}^{\Re} + i v_{muk}^{\Re} v_{nuk}^{\Im} + v_{muk}^{\Im} v_{nuk}^{\Im}) \quad (1296)$$

$$= 2i (v_{nuk}^{\Im} v_{muk}^{\Re} - v_{muk}^{\Im} v_{nuk}^{\Re}) \quad (1297)$$

$$\overline{H_S(t)} = \sum_n \left( \varepsilon_n(t) + \sum_{uk} \left( \frac{(v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2}{\omega_{uk}} - \frac{(g_{nuk} + g_{nuk}^*) v_{nuk}^{\Re} + i v_{nuk}^{\Im} (g_{nuk}^* - g_{nuk})}{\omega_{uk}} \right) \right) |n\rangle\langle n| \quad (1298)$$

$$+ \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| \left( \Pi_{uk} \exp \left( \frac{i(v_{nuk}^{\Im} v_{muk}^{\Re} - v_{muk}^{\Im} v_{nuk}^{\Re})}{\omega_{uk}^2} \right) \right) \Pi_u \exp \left( -\frac{1}{2} \sum_k \frac{|v_{muk} - v_{nuk}|^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1299)$$

$$|v_{muk} - v_{nuk}|^2 = (v_{muk} - v_{nuk})(v_{muk} - v_{nuk})^* \quad (1300)$$

$$= |v_{muk}|^2 + |v_{nuk}|^2 - (v_{nuk} v_{muk}^* + v_{muk} v_{nuk}^*) \quad (1301)$$

$$= (v_{muk}^{\Re})^2 + (v_{muk}^{\Im})^2 + (v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2 - (v_{nuk}^{\Re} + i v_{nuk}^{\Im})(v_{muk}^{\Re} - i v_{muk}^{\Im}) \quad (1302)$$

$$- (v_{nuk}^{\Re} - i v_{nuk}^{\Im})(v_{muk}^{\Re} + i v_{muk}^{\Im}) \quad (1303)$$

$$= (v_{muk}^{\Re})^2 + (v_{muk}^{\Im})^2 + (v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2 - 2(v_{nuk}^{\Re} v_{muk}^{\Re} + v_{nuk}^{\Im} v_{muk}^{\Im}) \quad (1304)$$

$$= (v_{muk}^{\Re} - v_{nuk}^{\Re})^2 + (v_{muk}^{\Im} - v_{nuk}^{\Im})^2 \quad (1305)$$

$$R_n(t) = \sum_{uk} \left( \frac{|v_{nuk}|^2}{\omega_{uk}} - \left( g_{nuk} \frac{v_{nuk}^*}{\omega_{uk}} + g_{nuk}^* \frac{v_{nuk}}{\omega_{uk}} \right) \right) \quad (1306)$$

$$= \sum_{uk} \left( \frac{(v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2 - (g_{nuk} + g_{nuk}^*) v_{nuk}^{\Re} - i v_{nuk}^{\Im} (g_{nuk}^* - g_{nuk})}{\omega_{uk}} \right) \quad (1307)$$

$$= \sum_{uk} \left( \frac{(v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2 - 2g_{nuk}^{\Re} v_{nuk}^{\Re} - 2g_{nuk}^{\Im} v_{nuk}^{\Im}}{\omega_{uk}} \right) \quad (1308)$$

$$B_{mn} = \left( \Pi_{uk} \exp \left( \frac{(v_{muk}^* v_{nuk} - v_{muk} v_{nuk}^*)}{2\omega_{uk}^2} \right) \right) \Pi_u \exp \left( -\frac{1}{2} \sum_k \frac{|v_{muk} - v_{nuk}|^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1309)$$

$$= \left( \Pi_{uk} \exp \left( \frac{i(v_{nuk}^{\Im} v_{muk}^{\Re} - v_{muk}^{\Im} v_{nuk}^{\Re})}{\omega_{uk}^2} \right) \right) \Pi_u \exp \left( -\frac{1}{2} \sum_k \frac{(v_{muk}^{\Re} - v_{nuk}^{\Re})^2 + (v_{muk}^{\Im} - v_{nuk}^{\Im})^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1310)$$

Then we can obtain using the chain rule that:

$$\frac{\partial R_{n'}}{\partial v_{nuk}^{\Re}} = \frac{\partial}{\partial v_{nuk}^{\Re}} \sum_{uk} \left( \frac{(v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2 - 2g_{nuk}^{\Re} v_{nuk}^{\Re} - 2g_{nuk}^{\Im} v_{nuk}^{\Im}}{\omega_{uk}} \right) \quad (1311)$$

$$= \frac{2v_{nuk}^{\Re} - 2g_{nuk}^{\Re}}{\omega_{uk}} \delta_{nn'} \quad (1312)$$

$$= 2 \frac{v_{nuk}^{\Re} - g_{nuk}^{\Re}}{\omega_{uk}} \delta_{nn'} \quad (1313)$$

$$\frac{\partial R_{n'}}{\partial v_{nuk}^{\Im}} = \frac{\partial}{\partial v_{nuk}^{\Im}} \sum_{uk} \left( \frac{(v_{nuk}^{\Re})^2 + (v_{nuk}^{\Im})^2 - 2g_{nuk}^{\Re} v_{nuk}^{\Re} - 2g_{nuk}^{\Im} v_{nuk}^{\Im}}{\omega_{uk}} \right) \quad (1314)$$

$$= \frac{2v_{nuk}^{\Im} - 2g_{nuk}^{\Im}}{\omega_{uk}} \delta_{nn'} \quad (1315)$$

$$= 2 \frac{v_{nuk}^{\Im} - g_{nuk}^{\Im}}{\omega_{uk}} \delta_{nn'} \quad (1316)$$

Given that:

$$\ln B_{mn} = \ln \left( \left( \prod_{uk} \exp \left( \frac{i(v_{nuk}^{\Im} v_{muk}^{\Re} - v_{muk}^{\Im} v_{nuk}^{\Re})}{\omega_{uk}^2} \right) \right) \prod_u \exp \left( -\frac{1}{2} \sum_{\mathbf{k}} \frac{(v_{muk}^{\Re} - v_{nuk}^{\Re})^2 + (v_{muk}^{\Im} - v_{nuk}^{\Im})^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \right) \quad (1317)$$

$$= \sum_{uk} \ln \exp \left( \frac{i(v_{nuk}^{\Im} v_{muk}^{\Re} - v_{muk}^{\Im} v_{nuk}^{\Re})}{\omega_{uk}^2} \right) + \sum_u \ln \exp \left( -\frac{1}{2} \sum_{\mathbf{k}} \frac{(v_{muk}^{\Re} - v_{nuk}^{\Re})^2 + (v_{muk}^{\Im} - v_{nuk}^{\Im})^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1318)$$

$$= \sum_{uk} \left( \frac{i(v_{nuk}^{\Im} v_{muk}^{\Re} - v_{muk}^{\Im} v_{nuk}^{\Re})}{\omega_{uk}^2} \right) + \sum_{uk} \left( -\frac{1}{2} \frac{(v_{muk}^{\Re} - v_{nuk}^{\Re})^2 + (v_{muk}^{\Im} - v_{nuk}^{\Im})^2}{\omega_{uk}^2} \coth \left( \frac{\beta_u \omega_{uk}}{2} \right) \right) \quad (1319)$$

$$\frac{\partial \ln B_{mn}}{\partial v_{nuk}^{\Re}} = \frac{-i v_{muk}^{\Im} - (v_{nuk}^{\Re} - v_{muk}^{\Re}) \coth \left( \frac{\beta_u \omega_{uk}}{2} \right)}{\omega_{uk}^2} \quad (1320)$$

$$\frac{\partial \ln B_{mn}}{\partial v_{nuk}^{\Im}} = \frac{i v_{muk}^{\Re} - (v_{nuk}^{\Im} - v_{muk}^{\Im}) \coth \left( \frac{\beta_u \omega_{uk}}{2} \right)}{\omega_{uk}^2} \quad (1321)$$

$$\frac{\partial \ln B_{mn}}{\partial a} = \frac{1}{B_{mn}} \frac{\partial B_{mn}}{\partial a} \quad (1322)$$

$$\frac{\partial B_{mn}}{\partial a} = B_{mn} \frac{\partial \ln B_{mn}}{\partial a} \quad (1323)$$

$$\frac{\partial B_{mn}}{\partial a} = \frac{\partial (B_{nm})^{\dagger}}{\partial a} \quad (1324)$$

Then the principal derivates are given by:

$$\frac{\partial B_{mn}}{\partial v_{n\mathbf{k}}^{\Re}} = B_{mn} \frac{\partial \ln B_{mn}}{\partial v_{n\mathbf{k}}^{\Re}} \quad (1325)$$

$$= B_{mn} \left( \frac{-iv_{m\mathbf{k}}^{\Im} - (v_{n\mathbf{k}}^{\Re} - v_{m\mathbf{k}}^{\Re}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \quad (1326)$$

$$= B_{mn} \left( \frac{-iv_{m\mathbf{k}}^{\Im} + (v_{m\mathbf{k}}^{\Re} - v_{n\mathbf{k}}^{\Re}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \quad (1327)$$

$$\frac{\partial B_{nm}}{\partial v_{n\mathbf{k}}^{\Re}} = \left( \frac{\partial B_{mn}}{\partial v_{n\mathbf{k}}^{\Re}} \right)^{\dagger} \quad (1328)$$

$$= \left( B_{mn} \left( \frac{-iv_{m\mathbf{k}}^{\Im} + (v_{m\mathbf{k}}^{\Re} - v_{n\mathbf{k}}^{\Re}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \right)^{\dagger} \quad (1329)$$

$$= B_{nm} \left( \frac{iv_{m\mathbf{k}}^{\Im} + (v_{m\mathbf{k}}^{\Re} - v_{n\mathbf{k}}^{\Re}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \quad (1330)$$

$$\frac{\partial B_{mn}}{\partial v_{n\mathbf{k}}^{\Im}} = B_{mn} \frac{\partial \ln B_{mn}}{\partial v_{n\mathbf{k}}^{\Im}} \quad (1331)$$

$$= B_{mn} \left( \frac{iv_{m\mathbf{k}}^{\Re} - (v_{n\mathbf{k}}^{\Im} - v_{m\mathbf{k}}^{\Im}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \quad (1332)$$

$$= B_{mn} \left( \frac{iv_{m\mathbf{k}}^{\Re} + (v_{m\mathbf{k}}^{\Im} - v_{n\mathbf{k}}^{\Im}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \quad (1333)$$

$$\frac{\partial B_{nm}}{\partial v_{n\mathbf{k}}^{\Im}} = \left( \frac{\partial B_{mn}}{\partial v_{n\mathbf{k}}^{\Im}} \right)^{\dagger} \quad (1334)$$

$$= (B_{mn})^{\dagger} \quad (1335)$$

$$= B_{nm} \left( \frac{-iv_{m\mathbf{k}}^{\Re} + (v_{m\mathbf{k}}^{\Im} - v_{n\mathbf{k}}^{\Im}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \quad (1336)$$

Introducing this derivatives in the equation (1311) give us:

$$\frac{\partial A_{\mathbf{B}}}{\partial v_{n\mathbf{k}}^{\Re}} = \frac{\partial A_{\mathbf{B}}}{\partial R_n} \left( 2 \frac{v_{n\mathbf{k}}^{\Re} - g_{u\mathbf{k}}^{\Re}}{\omega_{u\mathbf{k}}} \right) + \sum_{n < m} \left( \frac{\partial A_{\mathbf{B}}}{\partial B_{nm}} B_{nm} \left( \frac{iv_{m\mathbf{k}}^{\Im} + (v_{m\mathbf{k}}^{\Re} - v_{n\mathbf{k}}^{\Re}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \right) \quad (1337)$$

$$+ \frac{\partial A_{\mathbf{B}}}{\partial B_{mn}} B_{mn} \left( \frac{-iv_{m\mathbf{k}}^{\Im} + (v_{m\mathbf{k}}^{\Re} - v_{n\mathbf{k}}^{\Re}) \coth\left(\frac{\beta_u \omega_{u\mathbf{k}}}{2}\right)}{\omega_{u\mathbf{k}}^2} \right) \quad (1338)$$

$$= 0 \quad (1339)$$

We can obtain the variational parameters:

$$-2 \frac{\partial A_B}{\partial R_n} \frac{v_{nuk}^{\Re}}{\omega_{uk}} + \sum_{n < m} \left( \frac{\partial A_B}{\partial B_{nm}} B_{nm} \frac{v_{nuk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} + \frac{\partial A_B}{\partial B_{mn}} B_{mn} \frac{v_{nuk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right) \quad (1340)$$

$$= -\frac{\partial A_B}{\partial R_n} \frac{2g_{nuk}^{\Re}}{\omega_{uk}} + \sum_{n < m} \left( \frac{\partial A_B}{\partial B_{nm}} B_{nm} \left( \frac{iv_{muk}^{\Im} + v_{muk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right) + \frac{\partial A_B}{\partial B_{mn}} B_{mn} \left( \frac{-iv_{muk}^{\Im} + v_{muk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right) \right) \quad (1341)$$

$$v_{nuk}^{\Re} = \frac{\frac{\partial A_B}{\partial R_n} \frac{2g_{nuk}^{\Re}}{\omega_{uk}} - \sum_{n < m} \left( \frac{\partial A_B}{\partial B_{nm}} B_{nm} \left( \frac{iv_{muk}^{\Im} + v_{muk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right) + \frac{\partial A_B}{\partial B_{mn}} B_{mn} \left( \frac{-iv_{muk}^{\Im} + v_{muk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right) \right)}{2 \frac{\partial A_B}{\partial R_n} \frac{1}{\omega_{uk}} - \sum_{n \neq m} \left( \frac{\partial A_B}{\partial B_{nm}} B_{nm} \frac{\coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} + \frac{\partial A_B}{\partial B_{mn}} B_{mn} \frac{\coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right)} \quad (1342)$$

$$= \frac{2g_{nuk}^{\Re} \omega_{uk} \frac{\partial A_B}{\partial R_n} - \sum_{n < m} \left( \frac{\partial A_B}{\partial B_{nm}} B_{nm} (iv_{muk}^{\Im} + v_{muk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)) + \frac{\partial A_B}{\partial B_{mn}} B_{mn} (-iv_{muk}^{\Im} + v_{muk}^{\Re} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)) \right)}{2\omega_{uk} \frac{\partial A_B}{\partial R_n} - \sum_{n \neq m} \left( \frac{\partial A_B}{\partial B_{nm}} B_{nm} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right) + \frac{\partial A_B}{\partial B_{mn}} B_{mn} \coth\left(\frac{\beta_u \omega_{uk}}{2}\right) \right)} \quad (1343)$$

Let's consider the imaginary part of the variation parameters

$$\frac{\partial A_B}{\partial v_{nuk}^{\Im}} = \frac{\partial A_B}{\partial R_n} \left( 2 \frac{v_{nuk}^{\Im} - g_{nuk}^{\Im}}{\omega_{uk}} \right) + \sum_{n < m} \left( \frac{\partial A_B}{\partial B_{nm}} B_{nm} \left( \frac{-iv_{muk}^{\Re} - (v_{nuk}^{\Im} - v_{muk}^{\Im}) \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right) \right) \quad (1344)$$

$$+ \frac{\partial A_B}{\partial B_{mn}} B_{mn} \left( \frac{iv_{muk}^{\Re} - (v_{nuk}^{\Im} - v_{muk}^{\Im}) \coth\left(\frac{\beta_u \omega_{uk}}{2}\right)}{\omega_{uk}^2} \right) \quad (1345)$$

$$= 0 \quad (1346)$$

Rearranging we obtain



$$\tilde{O} \equiv U^\dagger(t) O U(t) \quad (1363)$$

$$U(t) \equiv \mathcal{T} \exp \left( -i \int_0^t dt' \overline{H_S}(t') \right). \quad (1364)$$

Therefore:

$$\widetilde{\overline{\rho_S}}(t) = U^\dagger(t) \overline{\rho_S}(t) U(t), \text{ where} \quad (1365)$$

$$\overline{\rho_S}(t) = \text{Tr}_B(\bar{\rho}(t)) \quad (1366)$$

We can re-write the transformed interaction Hamiltonian operator using the following matrices:

$$\sigma_{nm,x} = |n\rangle\langle m| + |m\rangle\langle n| \quad (1367)$$

$$\sigma_{nm,y} = i(|n\rangle\langle m| - |m\rangle\langle n|) \quad (1368)$$

$$B_{nm,x} = \frac{B_{nm} + B_{mn}}{2} \quad (1369)$$

$$B_{nm,x} = \frac{B_{nm} - B_{mn}}{2i} \quad (1370)$$

We can proof that  $B_{nm} = B_{mn}^\dagger$

$$B_{mn}^\dagger = (B_{m+}B_{n-} - B_m B_n)^\dagger \quad (1371)$$

$$= B_{n-}^\dagger B_{m+}^\dagger - B_n B_m \quad (1372)$$

$$= B_{n+} B_{m-} - B_n B_m \quad (1373)$$

$$= B_{nm} \quad (1374)$$

So we can say that the set of matrices (1367) are hermetic. Re-writing the transformed interaction Hamiltonian using the set (1367) give us.

$$\overline{H_I} = \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m| B_{nm} + \sum_n B_{z,n}(t) |n\rangle\langle n|, \quad (1375)$$

$$= \sum_n B_{z,n}(t) |n\rangle\langle n| + \sum_{n < m} (V_{nm}(t) |n\rangle\langle m| B_{nm} + V_{mn}(t) |m\rangle\langle n| B_{mn}) \quad (1376)$$

$$= \sum_n B_{z,n}(t) |n\rangle\langle n| + \sum_{n < m} \left( \Re(V_{nm}(t)) B_{nm} \left( \frac{\sigma_{nm,x} - i\sigma_{nm,y}}{2} \right) + iV_{nm}^\Im(t) B_{nm} \left( \frac{\sigma_{nm,x} - i\sigma_{nm,y}}{2} \right) \right) \quad (1377)$$

$$+ \Re(V_{nm}(t)) B_{mn} \left( \frac{\sigma_{nm,x} + i\sigma_{nm,y}}{2} \right) - iV_{nm}^\Im(t) B_{mn} \left( \frac{\sigma_{nm,x} + i\sigma_{nm,y}}{2} \right) \quad (1378)$$

$$= \sum_n B_{z,n}(t) |n\rangle\langle n| + \sum_{n < m} \left( \Re(V_{nm}(t)) \sigma_{nm,x} \left( \frac{B_{nm} + B_{mn}}{2} \right) + \Re(V_{nm}(t)) \sigma_{nm,y} \frac{i(B_{mn} - B_{nm})}{2} \right) \quad (1379)$$

$$+ i\Im(V_{nm}(t)) \sigma_{nm,x} \left( \frac{B_{nm} - B_{mn}}{2} \right) + \Im(V_{nm}(t)) \sigma_{nm,y} \left( \frac{B_{nm} + B_{mn}}{2} \right) \quad (1380)$$

$$= \sum_n B_{z,n}(t) |n\rangle\langle n| + \sum_{n < m} (\Re(V_{nm}(t)) \sigma_{nm,x} B_{nm,x} - \Im(V_{nm}(t)) \sigma_{nm,x} B_{nm,y} + \Re(V_{nm}(t)) \sigma_{nm,y} B_{nm,y} \quad (1381)$$

$$+ \Im(V_{nm}(t)) \sigma_{nm,y} B_{nm,x}) \quad (1382)$$

Let's define the set

$$P = \{(n, m) \in \mathbb{N}^2 | 0 \leq n, m \leq d-1 \wedge (n = m \vee n < m)\} \quad (1383)$$



Now consider the following set of operators,

$$A_{1,nm}(t) = \sigma_{nm,x} (1 - \delta_{mn}) \quad (1384)$$

$$A_{2,nm}(t) = \sigma_{nm,y} (1 - \delta_{mn}) \quad (1385)$$

$$A_{3,nm}(t) = \delta_{mn} |n\rangle\langle m| \quad (1386)$$

$$A_{4,nm}(t) = A_{2,mn}(t) \quad (1387)$$

$$A_{5,nm}(t) = A_{1,nm}(t) \quad (1388)$$

$$B_{1,nm}(t) = B_{nm,x} \quad (1389)$$

$$B_{2,nm}(t) = B_{nm,y} \quad (1390)$$

$$B_{3,nm}(t) = B_{z,n}(t) \quad (1391)$$

$$B_{4,nm}(t) = B_{1,nm}(t) \quad (1392)$$

$$B_{5,nm}(t) = B_{2,nm}(t) \quad (1393)$$

$$C_{1,nm}(t) = \Re(V_{nm}(t)) \quad (1394)$$

$$C_{2,nm}(t) = C_{1,nm}(t) \quad (1395)$$

$$C_{3,nm}(t) = 1 \quad (1396)$$

$$C_{4,nm}(t) = \Im(V_{nm}(t)) \quad (1397)$$

$$C_{5,nm}(t) = -\Im(V_{nm}(t)) \quad (1398)$$

The previous notation allows us to write the interaction Hamiltonian in  $\overline{H_I}(t)$  as:

$$\overline{H_I} = \sum_{j \in J, p \in P} C_{jp}(t) (A_{jp} \otimes B_{jp}(t)) \quad (1399)$$

Here  $J = \{1, 2, 3, 4, 5\}$  and  $P$  the set defined in (1383).

We write the interaction Hamiltonian transformed under (1363) as:

$$\widetilde{H_I}(t) = \sum_{j \in J, p \in P} C_{jp}(t) \left( \widetilde{A_{jp}}(t) \otimes \widetilde{B_{jp}}(t) \right) \quad (1400)$$

$$\widetilde{A_{jp}}(t) = U^\dagger(t) A_{jp} U(t) \quad (1401)$$

$$\widetilde{B_{jp}}(t) = e^{iH_B t} B_{jp}(t) e^{-iH_B t} \quad (1402)$$

Taking as reference state  $\rho_B$  and truncating at second order in  $H_I(t)$ , we obtain our master equation in the interaction picture:

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \text{Tr}_B \left[ \widetilde{H_I}(t), \left[ \widetilde{H_I}(s), \widetilde{\rho_S}(t) \rho_B \right] \right] ds \quad (1403)$$

Replacing the equation (1400) in (1403) we can obtain:

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \text{Tr}_B \left[ \widetilde{H}_I(t), \left[ \widetilde{H}_I(s), \widetilde{\rho_S}(t) \rho_B \right] \right] ds \quad (1404)$$

$$= - \int_0^t \text{Tr}_B \left[ \sum_{j \in J, p \in P} C_{jp}(t) \left( \widetilde{A}_{jp}(t) \otimes \widetilde{B}_{jp}(t) \right), \left[ \sum_{j' \in J, p' \in P} C_{j'p'}(s) \left( \widetilde{A}_{j'p'}(s) \otimes \widetilde{B}_{j'p'}(s) \right), \widetilde{\rho_S}(t) \rho_B \right] \right] ds \quad (1405)$$

$$= - \int_0^t \text{Tr}_B \left[ \sum_{j \in J, p \in P} C_{jp}(t) \left( \widetilde{A}_{jp}(t) \otimes \widetilde{B}_{jp}(t) \right), \sum_{j' \in J, p' \in P} C_{j'p'}(s) \left( \widetilde{A}_{j'p'}(s) \otimes \widetilde{B}_{j'p'}(s) \right) \widetilde{\rho_S}(t) \rho_B \right] \quad (1406)$$

$$- \widetilde{\rho_S}(t) \rho_B \sum_{j' \in J, p' \in P} C_{j'p'}(s) \left( \widetilde{A}_{j'p'}(s) \otimes \widetilde{B}_{j'p'}(s) \right) \Big] ds \quad (1407)$$

$$= - \int_0^t \text{Tr}_B \left( \sum_{j \in J, p \in P} C_{jp}(t) \left( \widetilde{A}_{jp}(t) \otimes \widetilde{B}_{jp}(t) \right) \sum_{j' \in J, p' \in P} C_{j'p'}(s) \left( \widetilde{A}_{j'p'}(s) \otimes \widetilde{B}_{j'p'}(s) \right) \widetilde{\rho_S}(t) \rho_B \right. \quad (1408)$$

$$\left. - \sum_{j \in J, p \in P} C_{jp}(t) \left( \widetilde{A}_{jp}(t) \otimes \widetilde{B}_{jp}(t) \right) \widetilde{\rho_S}(t) \rho_B \sum_{j' \in J, p' \in P} C_{j'p'}(s) \left( \widetilde{A}_{j'p'}(s) \otimes \widetilde{B}_{j'p'}(s) \right) \right) \quad (1409)$$

$$- \sum_{j' \in J, p' \in P} C_{j'p'}(s) \left( \widetilde{A}_{j'p'}(s) \otimes \widetilde{B}_{j'p'}(s) \right) \widetilde{\rho_S}(t) \rho_B \sum_{j \in J, p \in P} C_{jp}(t) \left( \widetilde{A}_{jp}(t) \otimes \widetilde{B}_{jp}(t) \right) \quad (1410)$$

$$\left. + \widetilde{\rho_S}(t) \rho_B \sum_{j' \in J, p' \in P} C_{j'p'}(s) \left( \widetilde{A}_{j'p'}(s) \otimes \widetilde{B}_{j'p'}(s) \right) \sum_{j \in J, p \in P} C_{jp}(t) \left( \widetilde{A}_{jp}(t) \otimes \widetilde{B}_{jp}(t) \right) \right) ds \quad (1411)$$

In order to calculate the correlation functions we define:

$$\Lambda_{jpj'p'}(\tau) = \left\langle \widetilde{B}_{jp}(t) \widetilde{B}_{j'p'}(s) \right\rangle_B \quad (1412)$$

$$= \left\langle \widetilde{B}_{jp}(\tau) \widetilde{B}_{j'p'}(0) \right\rangle_B \quad (1413)$$

Here  $s \rightarrow t - \tau$  and  $\text{Tr}_B \left( \widetilde{B}_{jp}(t) \widetilde{B}_{j'p'}(s) \right) = \left\langle \widetilde{B}_{jp}(t) \widetilde{B}_{j'p'}(s) \right\rangle_B$ . To evaluate the trace respect to the bath we need to recall that our master equation depends of elements related to the bath and represented by the operators  $\widetilde{B}_{jp}(t)$  and elements related to the system given by  $\widetilde{A}_{jp}(t)$ . The systems considered are in different Hilbert spaces so  $\text{Tr} \left( \widetilde{A}_{jp}(t) \widetilde{B}_{j'p'}(t) \right) = \text{Tr} \left( \widetilde{A}_{jp}(t) \right) \text{Tr} \left( \widetilde{B}_{j'p'}(t) \right)$ . The correlation functions relevant of the master equation (1411) are:

$$\text{Tr}_B \left( \widetilde{B_{jp}}(t) \widetilde{B_{j'p'}}(s) \rho_B \right) = \left\langle \widetilde{B_{jp}}(t) \widetilde{B_{j'p'}}(s) \right\rangle_B \quad (1414)$$

$$= \left\langle \widetilde{B_{jp}}(0) \widetilde{B_{j'p'}}(0) \right\rangle_B \quad (1415)$$

$$= \Lambda_{jpj'p'}(\tau) \quad (1416)$$

$$\text{Tr}_B \left( \widetilde{B_{jp}}(t) \rho_B \widetilde{B_{j'p'}}(s) \right) = \text{Tr}_B \left( \widetilde{B_{j'p'}}(s) \widetilde{B_{jp}}(t) \rho_B \right) \quad (1417)$$

$$= \left\langle \widetilde{B_{j'p'}}(s) \widetilde{B_{jp}}(t) \right\rangle_B \quad (1418)$$

$$= \left\langle \widetilde{B_{j'p'}}(-\tau) \widetilde{B_{jp}}(0) \right\rangle_B \quad (1419)$$

$$= \Lambda_{j'p'jp}(-\tau) \quad (1420)$$

$$\text{Tr}_B \left( \widetilde{B_{j'p'}}(s) \rho_B \widetilde{B_{jp}}(t) \right) = \text{Tr}_B \left( \widetilde{B_{jp}}(t) \widetilde{B_{j'p'}}(s) \rho_B \right) \quad (1421)$$

$$= \left\langle \widetilde{B_{jp}}(t) \widetilde{B_{j'p'}}(s) \right\rangle_B \quad (1422)$$

$$= \left\langle \widetilde{B_{jp}}(\tau) \widetilde{B_{j'p'}}(0) \right\rangle_B \quad (1423)$$

$$= \Lambda_{jpj'p'}(\tau) \quad (1424)$$

$$\text{Tr}_B \left( \rho_B \widetilde{B_{j'p'}}(s) \widetilde{B_{jp}}(t) \right) = \text{Tr}_B \left( \widetilde{B_{j'p'}}(s) \widetilde{B_{jp}}(t) \rho_B \right) \quad (1425)$$

$$= \left\langle \widetilde{B_{j'p'}}(s) \widetilde{B_{jp}}(t) \right\rangle_B \quad (1426)$$

$$= \left\langle \widetilde{B_{j'p'}}(-\tau) \widetilde{B_{jp}}(0) \right\rangle_B \quad (1427)$$

$$= \Lambda_{j'p'jp}(-\tau) \quad (1428)$$

We made use of the cyclic property for the trace to evaluate the correlation functions, from the equations obtained in (1404) and (1411) and using the equations (1414)-(1428) we can re-write:

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \sum_{j,j',p,p'} \left( C_{jp}(t) C_{j'p'}(s) \left( \Lambda_{jpj'p'}(\tau) \widetilde{A_{jp}}(t) \widetilde{A_{j'p'}}(s) \widetilde{\rho_S}(t) - \Lambda_{j'p'jp}(-\tau) \widetilde{A_{jp}}(t) \widetilde{\rho_S}(t) \widetilde{A_{j'p'}}(s) \right) \right. \quad (1429)$$

$$\left. + C_{jp}(t) C_{j'p'}(s) \left( \Lambda_{j'p'jp}(-\tau) \widetilde{\rho_S}(t) \widetilde{A_{j'p'}}(s) \widetilde{A_{jp}}(t) - \Lambda_{jpj'p'}(\tau) \widetilde{A_{j'p'}}(s) \widetilde{\rho_S}(t) \widetilde{A_{jp}}(t) \right) \right) ds \quad (1430)$$

$$= - \int_0^t \sum_{jj'pp'} \left( C_{jp}(t) C_{j'p'}(s) \left( \Lambda_{jpj'p'}(\tau) \left[ \widetilde{A_{jp}}(t), \widetilde{A_{j'p'}}(s) \widetilde{\rho_S}(t) \right] + \Lambda_{j'p'jp}(-\tau) \left[ \widetilde{\rho_S}(t) \widetilde{A_{j'p'}}(s), \widetilde{A_{jp}}(t) \right] \right) \right) \quad (1431)$$

Rearranging and identifying the commutators allow us to write a more simplified version

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \sum_{jj'pp'} \left( C_{jp}(t) C_{j'p'}(t-\tau) \left( \Lambda_{jpj'p'}(\tau) \left[ A_{jp}(t), A_{j'p'}(t-\tau, t) \widetilde{\rho_S}(t) \right] + \Lambda_{j'p'jp}(-\tau) \left[ \widetilde{\rho_S}(t) A_{j'p'}(t-\tau, t), A_{jp}(t) \right] \right) \right) d\tau - i [H_S(t), \widetilde{\rho_S}(t)] \quad (1432)$$

For this case we used that  $A_{jp}(t-\tau, t) = U(t) U^\dagger(t-\tau) A_{jp}(t) U(t-\tau) U^\dagger(t)$ . This is a non-Markovian equation.

## VI. TIME-DEPENDENT MULTI-SITE MODEL WITH ONE BATH COUPLING

Let's consider the following Hamiltonian for a system of d-levels (qudit). We start with a time-dependent Hamiltonian of the form:

$$H(t) = H_S(t) + H_I + H_B, \quad (1433)$$

$$H_S(t) = \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m|, \quad (1434)$$

$$H_I = \left( \sum_{n=0} \mu_n(t) |n\rangle\langle n| \right) \left( \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \right), \quad (1435)$$

$$H_B = \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}. \quad (1436)$$

We will start with a system-bath coupling operator of the form  $\sum_{n=0} \mu_n(t) |n\rangle\langle n|$ .

### A. Variational Transformation

We consider the following operator:

$$V = \left( \sum_{n=1} |n\rangle\langle n| \right) \left( \sum_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger - b_{\mathbf{k}}) \right) \quad (1437)$$

At first let's obtain  $e^V$  under the transformation (1437), consider  $\hat{\varphi} = \sum_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger - b_{\mathbf{k}})$ :

$$e^V = e^{\sum_{n=1} |n\rangle\langle n| \hat{\varphi}} \quad (1438)$$

$$= \mathbb{I} + \sum_{n=1} |n\rangle\langle n| \hat{\varphi} + \frac{(\sum_{n=1} |n\rangle\langle n| \hat{\varphi})^2}{2!} + \dots \quad (1439)$$

$$= \mathbb{I} + \sum_{n=1} |n\rangle\langle n| \hat{\varphi} + \frac{\sum_{n=1} |n\rangle\langle n| \hat{\varphi}^2}{2!} + \dots \quad (1440)$$

$$= \mathbb{I} - \sum_{n=1} |n\rangle\langle n| + \sum_{n=1} |n\rangle\langle n| \left( \mathbb{I} + \hat{\varphi} + \frac{\hat{\varphi}^2}{2!} + \dots \right) \quad (1441)$$

$$= |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| e^{\hat{\varphi}} \quad (1442)$$

$$= |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \quad (1443)$$

Given that  $[b_{\mathbf{k}'}^\dagger - b_{\mathbf{k}'}^\dagger, b_{\mathbf{k}}^\dagger - b_{\mathbf{k}}] = 0$  if  $\mathbf{k}' \neq \mathbf{k}$  then we can proof using the Zassenhaus formula and defining  $D\left(\pm \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}}\right) = e^{\pm \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger - b_{\mathbf{k}})}$  in the same way than (23):

$$e^{\sum_{\mathbf{k}} \pm \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger - b_{\mathbf{k}})} = \prod_{\mathbf{k}} e^{\pm \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger - b_{\mathbf{k}})} \quad (1444)$$

$$= \prod_{\mathbf{k}} D\left(\pm \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}}\right) \quad (1445)$$

$$= B_{\pm} \quad (1446)$$

As we can see  $e^{-V} = |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B$ . because this form imposes that  $e^{-V} e^V = \mathbb{I}$  and the inverse of a operator is unique. This allows us to write the canonical transformation in the following explicit way:

$$e^V A e^{-V} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) A \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right) \quad (1447)$$

Now let's obtain the canonical transformation of the principal elements of the Hamiltonian (1433):

$$\overline{|0\rangle\langle 0|} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) |0\rangle\langle 0| \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right), \quad (1448)$$

$$= |0\rangle\langle 0|, \quad (1449)$$

$$\overline{|m\rangle\langle n|} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) |m\rangle\langle n| \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right), \quad (1450)$$

$$= |m\rangle\langle m| B^+ |m\rangle\langle n| n\rangle\langle n| B^-, \quad (1451)$$

$$= |m\rangle\langle n|, \quad m \neq 0, \quad n \neq 0, \quad (1452)$$

$$\overline{|0\rangle\langle m|} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) |0\rangle\langle m| \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right), \quad (1453)$$

$$= |0\rangle\langle m| B^- \quad m \neq 0, \quad (1454)$$

$$\overline{|m\rangle\langle 0|} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) |m\rangle\langle 0| \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right) \quad (1455)$$

$$= |0\rangle\langle m| B^+ \quad m \neq 0, \quad (1456)$$

$$\overline{\sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right) \quad (1457)$$

$$= |0\rangle\langle 0| \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} B^+ b_{\mathbf{k}}^\dagger b_{\mathbf{k}} B^- \quad (1458)$$

$$= |0\rangle\langle 0| \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} \left( B^+ b_{\mathbf{k}}^\dagger B^- \right) \left( B^+ b_{\mathbf{k}} B^- \right) \quad (1459)$$

$$= |0\rangle\langle 0| \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} \left( b_{\mathbf{k}}^\dagger - \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) \left( b_{\mathbf{k}} - \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) \quad (1460)$$

$$= |0\rangle\langle 0| \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} \left( b_{\mathbf{k}}^\dagger b_{\mathbf{k}} - \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) + \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) \quad (1461)$$

$$= \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} \left( \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \right) \quad (1462)$$

$$= \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - \sum_{n=1} |n\rangle\langle n| \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1463)$$

The transformed Hamiltonians of the equations (1434) to (1436) written in terms of (1448) to (1463) are:

$$\overline{H_S(t)} = \overline{\sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n \neq m} V_{nm}(t) |n\rangle\langle m|} \quad (1464)$$

$$= \overline{\sum_{n=0} \varepsilon_n(t) |n\rangle\langle n|} + \overline{\sum_{n \neq m} V_{nm}(t) |n\rangle\langle m|} \quad (1465)$$

$$= \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| + V_{n0}(t) |n\rangle\langle 0|) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| \quad (1466)$$

$$= \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n=1} (V_{0n}(t) \overline{|0\rangle\langle n|} + V_{n0}(t) \overline{|n\rangle\langle 0|}) + \sum_{m,n \neq 0} V_{mn}(t) \overline{|m\rangle\langle n|} \quad (1467)$$

$$= \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n=1} (V_{0n}(t) B^- |0\rangle\langle n| + V_{n0}(t) B^+ |n\rangle\langle 0|) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| \quad (1468)$$

$$= \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| B^- + V_{n0}(t) |n\rangle\langle 0| B^+) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| \quad (1469)$$

$$\overline{H_I} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) \left( \left( \sum_{n=0} \mu_n(t) |n\rangle\langle n| \right) \left( \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \right) \right) \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right) \quad (1470)$$

$$= \left( \mu_0(t) |0\rangle\langle 0| + \sum_{n=1} \mu_n(t) |n\rangle\langle n| B^+ \right) \left( \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \right) \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right) \quad (1471)$$

$$= \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) + \sum_{n=1} \mu_n(t) |n\rangle\langle n| \sum_{\mathbf{k}} g_{\mathbf{k}} B^+ (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) B^- \quad (1472)$$

$$= \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) + \sum_{n=1} \mu_n(t) |n\rangle\langle n| \sum_{\mathbf{k}} g_{\mathbf{k}} \left( b_{\mathbf{k}}^\dagger + b_{\mathbf{k}} - 2 \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) \quad (1473)$$

$$\overline{H_B} = \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - \sum_{n=1} |n\rangle\langle n| \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1474)$$

Joining this terms allow us to write:

$$\overline{H} = \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| B^- + V_{n0}(t) |n\rangle\langle 0| B^+) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| \quad (1475)$$

$$+ \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - \sum_{n=1} |n\rangle\langle n| \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1476)$$

$$+ \sum_{n=0} \mu_n(t) |n\rangle\langle n| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) - \sum_{n=1} \mu_n(t) |n\rangle\langle n| \sum_{\mathbf{k}} 2g_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \quad (1477)$$

$$= \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| B^- + V_{n0}(t) |n\rangle\langle 0| B^+) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| \quad (1478)$$

$$+ \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \left( \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - 2\mu_n(t) g_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) + \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1479)$$

$$+ \sum_{n=1} |n\rangle\langle n| \sum_{\mathbf{k}} \left( g_{\mathbf{k}} \mu_n(t) - \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1480)$$

Let's define the following functions:

$$R_n(t) = \sum_{\mathbf{k}} \left( \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - 2\mu_n(t) g_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) \quad (1481)$$

$$= \sum_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \left( \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} - 2\mu_n(t) g_{\mathbf{k}} \right) \quad (1482)$$

$$B_{z,n}(t) = \sum_{\mathbf{k}} \left( g_{\mathbf{k}} \mu_n(t) - \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1483)$$

Using the previous functions we have that (1480) can be re-written in the following way:

$$\overline{H} = \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| B^- + V_{n0}(t) |n\rangle\langle 0| B^+) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| \quad (1484)$$

$$+ \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \sum_{n=1} R_n |n\rangle\langle n| + \sum_{n=1} B_{z,n} |n\rangle\langle n| + \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1485)$$

Now in order to separate the elements of the hamiltonian (1485) let's follow the references of the equations (??) and (??) to separate the hamiltonian like:

$$\overline{H_S}(t) = \sum_{n=0} \varepsilon_n(t) |n\rangle\langle n| + B \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| + V_{n0}(t) |n\rangle\langle 0|) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| + \sum_{n=1} R_n |n\rangle\langle n| \quad (1486)$$

$$\overline{H_I} = \sum_{n=1} B_{z,n} |n\rangle\langle n| + \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) + \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| (B^- - B) + V_{n0}(t) |n\rangle\langle 0| (B^+ - B)), \quad (1487)$$

$$\overline{H_B} = \sum_{\mathbf{k}} \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \quad (1488)$$

Here B is given by:

$$\begin{aligned} B &= \langle B^+ \rangle \\ &= \langle B^- \rangle \end{aligned}$$

The transformed Hamiltonian can be written in function of the following set of hermitian operators:

$$\sigma_{nm,x} = |n\rangle\langle m| + |m\rangle\langle n| \quad (1489)$$

$$\sigma_{nm,y} = i(|n\rangle\langle m| - |m\rangle\langle n|) \quad (1490)$$

$$B_x = \frac{B^+ + B^- - 2B}{2} \quad (1491)$$

$$B_y = \frac{B^- - B^+}{2i} \quad (1492)$$

Using this set of hermitian operators to write the Hamiltonians (1434)-(1436)

$$\overline{H_S(t)} = \varepsilon_0(t) |0\rangle\langle 0| + \sum_{n=1} (\varepsilon_n(t) + R_n) |n\rangle\langle n| + B \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| + V_{n0}(t) |n\rangle\langle 0|) + \sum_{m,n \neq 0} V_{mn}(t) |m\rangle\langle n| \quad (1493)$$

$$= \varepsilon_0(t) |0\rangle\langle 0| + B \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| + V_{n0}(t) |n\rangle\langle 0|) + \sum_{0 < m < n} (V_{mn}(t) |m\rangle\langle n| + V_{nm}(t) |n\rangle\langle m|) \quad (1494)$$

$$+ \sum_{n=1} (\varepsilon_n(t) + R_n) |n\rangle\langle n| \quad (1495)$$

$$= \sum_{0 < m < n} ((\Re(V_{mn}(t)) + i\Im(V_{mn}(t))) |m\rangle\langle n| + (\Re(V_{mn}(t)) - i\Im(V_{mn}(t))) |n\rangle\langle m|) + \varepsilon_0(t) |0\rangle\langle 0| \quad (1496)$$

$$+ B \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| + V_{n0}(t) |n\rangle\langle 0|) + \sum_{n=1} (\varepsilon_n(t) + R_n) |n\rangle\langle n| \quad (1497)$$

$$= \sum_{0 < m < n} \left( (\Re(V_{nm}(t)) + i\Im(V_{mn}(t))) \frac{\sigma_{nm,x} - i\sigma_{nm,y}}{2} + (\Re(V_{nm}(t)) - i\Im(V_{mn}(t))) \frac{\sigma_{nm,x} + i\sigma_{nm,y}}{2} \right) \quad (1498)$$

$$+ B \sum_{n=1} \left( V_{0n}(t) \frac{\sigma_{0n,x} - i\sigma_{0n,y}}{2} + V_{n0}(t) \frac{\sigma_{0n,x} + i\sigma_{0n,y}}{2} \right) + \varepsilon_0(t) |0\rangle\langle 0| + \sum_{n=1} (\varepsilon_n(t) + R_n) |n\rangle\langle n| \quad (1499)$$

$$= \sum_{0 < m < n} (\Re(V_{nm}(t)) \sigma_{nm,x} + \Im(V_{nm}(t)) \sigma_{nm,y}) + B \sum_{n=1} (\Re(V_{0n}(t)) \sigma_{0n,x} + \Im(V_{mn}(t)) \sigma_{0n,y}) \quad (1500)$$

$$+ \varepsilon_0(t) |0\rangle\langle 0| + \sum_{n=1} (\varepsilon_n(t) + R_n) |n\rangle\langle n| \quad (1501)$$

$$\overline{H_I(t)} = \sum_{n=1} B_{z,n} |n\rangle\langle n| + \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) + \sum_{n=1} (V_{0n}(t) |0\rangle\langle n| (B^- - B) + V_{n0}(t) |n\rangle\langle 0| (B^+ - B)) \quad (1502)$$

$$= \sum_{n=1} \left( (\Re(V_{0n}(t)) + i\Im(V_{0n}(t))) (B^- - B) \frac{\sigma_{0n,x} - i\sigma_{0n,y}}{2} + (\Re(V_{0n}(t)) - i\Im(V_{0n}(t))) (B^+ - B) \frac{\sigma_{0n,x} + i\sigma_{0n,y}}{2} \right) \quad (1503)$$

$$+ \sum_{n=1} B_{z,n} |n\rangle\langle n| + \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1504)$$

$$= \sum_{n=1} B_{z,n} |n\rangle\langle n| + \sum_{n=1} \left( \frac{\sigma_{0n,x}}{2} ((B^- - B) (\Re(V_{0n}(t)) + i\Im(V_{0n}(t))) + (B^+ - B) (\Re(V_{0n}(t)) - i\Im(V_{0n}(t)))) \right) \quad (1505)$$

$$+ \frac{i\sigma_{0n,y}}{2} ((B^+ - B) (\Re(V_{0n}(t)) - i\Im(V_{0n}(t))) - (B^- - B) (\Re(V_{0n}(t)) + i\Im(V_{0n}(t)))) \quad (1506)$$

$$+ \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1507)$$

$$= \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) + \sum_{n=1} \left( \frac{\sigma_{0n,x}}{2} (B^+ + B^- - 2B) \Re(V_{0n}(t)) + i(B^- - B - B^+ + B) \Im(V_{0n}(t)) \right) \quad (1508)$$

$$+ \frac{i\sigma_{0n,y}}{2} ((B^+ - B - B^- + B) \Re(V_{0n}(t)) + i(B - B^- + B - B^+) \Im(V_{0n}(t))) + \sum_{n=1} B_{z,n} |n\rangle\langle n| \quad (1509)$$

$$= \sum_{n=1} B_{z,n} |n\rangle\langle n| + \mu_0(t) |0\rangle\langle 0| \sum_{\mathbf{k}} g_{\mathbf{k}} (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) + \sum_{n=1} (\sigma_{0n,x} (B_x \Re(V_{0n}(t)) - B_y \Im(V_{0n}(t))) \quad (1510)$$

$$+ \sigma_{0n,y} (B_y \Re(V_{0n}(t)) + B_x \Im(V_{0n}(t)))) \quad (1511)$$

## B. Free-energy minimization

As first approach let's consider the minimization of the free-energy through the Feynman-Bogoliubov inequality



$$A \leq A_B \equiv -\frac{1}{\beta} \ln \left( \text{Tr} \left( e^{-\beta(\overline{H}_S + \overline{H}_B)} \right) \right) + \langle \overline{H}_I \rangle_{\overline{H}_S + \overline{H}_B} + O \left( \langle \overline{H}_I^2 \rangle_{\overline{H}_S + \overline{H}_B} \right). \quad (1512)$$

Taking the equations (244)-(252) and given that  $\text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right) = C(R_1, R_2, \dots, R_{d-1}, B)$ , where each  $R_i$  and  $B$  depend of the set of variational parameters  $\{v_{\mathbf{k}}\}$ . From (252) and using the chain rule we obtain that:

$$\frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial v_{\mathbf{k}}} = \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial B} \frac{\partial B}{\partial v_{\mathbf{k}}} + \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} \frac{\partial R_n}{\partial v_{\mathbf{k}}}, \quad (1513)$$

$$= 0 \quad (1514)$$

Let's recall the equations (1481) and (1483), we can write them in terms of the variational parameters

$$B = \exp \left( - (1/2) \sum_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}^2} \coth(\beta \omega_{\mathbf{k}}/2) \right) \quad (1515)$$

$$R_n = \sum_{\mathbf{k}} \omega_{\mathbf{k}}^{-1} (v_{\mathbf{k}} - 2\mu_n(t) g_{\mathbf{k}} v_{\mathbf{k}}) \quad (1516)$$

The derivates needed to obtain the set of variational parameter are given by:

$$\frac{\partial B}{\partial v_{\mathbf{k}}} = -\frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}^2} \coth(\beta \omega_{\mathbf{k}}/2) B \quad (1517)$$

$$\frac{\partial R_n}{\partial v_{\mathbf{k}}} = \omega_{\mathbf{k}}^{-1} (2v_{\mathbf{k}} - 2\mu_n(t) g_{\mathbf{k}}) \quad (1518)$$

Introducing this derivates in the equation (1513) give us:

$$\frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial v_{\mathbf{k}}} = \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial B} \left( -\frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}^2} \coth(\beta \omega_{\mathbf{k}}/2) B \right) + \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} \omega_{\mathbf{k}}^{-1} (2v_{\mathbf{k}} - 2\mu_n(t) g_{\mathbf{k}}) \quad (1519)$$

$$= v_{\mathbf{k}} \left( \frac{2}{\omega_{\mathbf{k}}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} - \frac{\coth(\beta \omega_{\mathbf{k}}/2) B}{\omega_{\mathbf{k}}^2} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial B} \right) - \frac{2g_{\mathbf{k}}}{\omega_{\mathbf{k}}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} \mu_n(t) \quad (1520)$$

We can obtain the variational parameters:

$$v_{\mathbf{k}} = \frac{\frac{2g_{\mathbf{k}}}{\omega_{\mathbf{k}}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} \mu_n(t)}{\frac{2}{\omega_{\mathbf{k}}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} - \frac{\coth(\beta \omega_{\mathbf{k}}/2) B}{\omega_{\mathbf{k}}^2} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial B}} \quad (1521)$$

$$= \frac{2g_{\mathbf{k}} \omega_{\mathbf{k}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} \mu_n(t)}{2\omega_{\mathbf{k}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} - B \coth(\beta \omega_{\mathbf{k}}/2) \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial B}} \quad (1522)$$

Now taking  $v_{\mathbf{k}} = g_{\mathbf{k}} v_{\mathbf{k}}$  then we can obtain  $v_{\mathbf{k}}$  like:

$$v_{\mathbf{k}} = \frac{2\omega_{\mathbf{k}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} \mu_n(t)}{2\omega_{\mathbf{k}} \sum_{n=1} \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial R_n} - B \coth(\beta \omega_{\mathbf{k}}/2) \frac{\partial \text{Tr} \left( e^{-\beta \overline{H}_S(t)} \right)}{\partial B}}. \quad (1523)$$

### C. Master Equation

Let's consider that the initial state of the system is given by  $\rho(0) = |0\rangle\langle 0| \otimes \rho_B$ , as we can see this state is independent of the variational transformation:

$$e^V \rho(0) e^{-V} = \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^+ \right) (|0\rangle\langle 0| \otimes \rho_B) \left( |0\rangle\langle 0| + \sum_{n=1} |n\rangle\langle n| B^- \right) \quad (1524)$$

$$0 = |0\rangle\langle 0| \otimes \rho_B \quad (1525)$$

$$0 = \rho(0) \quad (1526)$$

We transform any operator  $O$  into the interaction picture in the following way:

$$\tilde{O} \equiv U^\dagger(t) O U(t) \quad (1527)$$

$$U(t) \equiv \mathcal{T} \exp \left( -i \int_0^t dt' \overline{H_S}(t') \right). \quad (1528)$$

Therefore:

$$\widetilde{\overline{\rho_S}}(t) = U^\dagger(t) \overline{\rho_S}(t) U(t), \text{ where} \quad (1529)$$

$$\overline{\rho_S}(t) = \text{Tr}_B(\tilde{\rho}(t)) \quad (1530)$$

We can re-write the transformed interaction Hamiltonian operator like:

$$\overline{H_I}(t) = B_{z,0} |0\rangle\langle 0| + \sum_{n=1} (\Re(V_{0n}(t)) B_x \sigma_{0n,x} + \Re(V_{0n}(t)) B_y \sigma_{0n,y} + B_{z,n} |n\rangle\langle n|) \quad (1531)$$

$$+ \Im(V_{0n}(t)) B_x \sigma_{0n,y} - \Im(V_{0n}(t)) B_y \sigma_{0n,x} \quad (1532)$$

where

$$B_{z,0} = \sum_{\mathbf{k}} g_{\mathbf{k}} \mu_0(t) (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \quad (1533)$$

$$B_{z,n} = \sum_{\mathbf{k}} \left( g_{\mathbf{k}} \mu_n(t) - \omega_{\mathbf{k}} \frac{v_{\mathbf{k}}}{\omega_{\mathbf{k}}} \right) (b_{\mathbf{k}}^\dagger + b_{\mathbf{k}}) \text{ if } n \neq 0 \quad (1534)$$

Now consider the following set of operators:

$$A_{1n}(t) = \sigma_{0n,x} \quad (1535)$$

$$A_{2n}(t) = \sigma_{0n,y} \quad (1536)$$

$$A_{3n}(t) = |n\rangle\langle n| \quad (1537)$$

$$A_{4n}(t) = A_{2n}(t) \quad (1538)$$

$$A_{5n}(t) = A_{1n}(t) \quad (1539)$$

$$B_{1n}(t) = B_x \quad (1540)$$

$$B_{2n}(t) = B_y \quad (1541)$$

$$B_{3n}(t) = B_{z,n} \quad (1542)$$

$$B_{4n}(t) = B_{1n}(t) \quad (1543)$$

$$B_{5n}(t) = B_{2n}(t) \quad (1544)$$

$$C_{10}(t) = 0 \quad (1545)$$

$$C_{20}(t) = 0 \quad (1546)$$

$$C_{40}(t) = 0 \quad (1547)$$

$$C_{50}(t) = 0 \quad (1548)$$

$$C_{30}(t) = 1 \quad (1549)$$

$$C_{1n}(t) = \Re(V_{0n}(t)) \quad (1550)$$

$$C_{2n}(t) = C_{1n}(t) \quad (1551)$$

$$C_{3n}(t) = 1 \quad (1552)$$

$$C_{4n}(t) = \Im(V_{0n}(t)) \quad (1553)$$

$$C_{5n}(t) = -\Im(V_{0n}(t)) \quad (1554)$$

The previous notation allows us to write the interaction Hamiltonian in  $\overline{H}_I(t)$  as:

$$\overline{H}_I = \sum_{j \in J} \sum_{n=1} C_{jn}(t) (A_{jn} \otimes B_{jn}(t)) \quad (1555)$$

Here  $J = \{1, 2, 3, 4, 5\}$ .

We write the interaction Hamiltonian transformed under (1527) as:

$$\widetilde{H}_I(t) = \sum_{j \in J} \sum_{n=1} C_{jn}(t) \left( \widetilde{A}_{jn}(t) \otimes \widetilde{B}_{jn}(t) \right) \quad (1556)$$

$$\widetilde{A}_i(t) = U^\dagger(t) A_i U(t) \quad (1557)$$

$$\widetilde{B}_i(t) = e^{iH_B t} B_i(t) e^{-iH_B t} \quad (1558)$$

Taking as reference state  $\rho_B$  and truncating at second order in  $H_I(t)$ , we obtain our master equation in the interaction picture:

$$\frac{d\widetilde{\rho}_S(t)}{dt} = - \int_0^t \text{Tr}_B \left[ \widetilde{H}_I(t), \left[ \widetilde{H}_I(s), \widetilde{\rho}_S(t) \rho_B \right] \right] ds \quad (1559)$$

Replacing the equation (1556) in (1559) we can obtain:

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \text{Tr}_B \left[ \widetilde{H}_I(t), \left[ \widetilde{H}_I(s), \widetilde{\rho_S}(t) \rho_B \right] \right] ds \quad (1560)$$

$$= - \int_0^t \text{Tr}_B \left[ \sum_{j \in J} \sum_{n=1} C_{jn}(t) \left( \widetilde{A}_{jn}(t) \otimes \widetilde{B}_{jn}(t) \right), \left[ \sum_{j' \in J} \sum_{n'=1} C_{j'n'}(s) \left( \widetilde{A}_{j'n'}(s) \otimes \widetilde{B}_{j'n'}(s) \right), \widetilde{\rho_S}(t) \rho_B \right] \right] ds \quad (1561)$$

$$= - \int_0^t \text{Tr}_B \left[ \sum_{j \in J} \sum_{n=1} C_{jn}(t) \left( \widetilde{A}_{jn}(t) \otimes \widetilde{B}_{jn}(t) \right), \sum_{j' \in J} \sum_{n'=1} C_{j'n'}(s) \left( \widetilde{A}_{j'n'}(s) \otimes \widetilde{B}_{j'n'}(s) \right) \widetilde{\rho_S}(t) \rho_B \right] ds \quad (1562)$$

$$- \widetilde{\rho_S}(t) \rho_B \sum_{j' \in J} \sum_{n'=1} C_{j'n'}(s) \left( \widetilde{A}_{j'n'}(s) \otimes \widetilde{B}_{j'n'}(s) \right) \Big] ds \quad (1563)$$

$$= - \int_0^t \text{Tr}_B \left( \sum_{j \in J} \sum_{n=1} C_{jn}(t) \left( \widetilde{A}_{jn}(t) \otimes \widetilde{B}_{jn}(t) \right) \sum_{j' \in J} \sum_{n'=1} C_{j'n'}(s) \left( \widetilde{A}_{j'n'}(s) \otimes \widetilde{B}_{j'n'}(s) \right) \widetilde{\rho_S}(t) \rho_B \right) ds \quad (1564)$$

$$- \sum_{j \in J} \sum_{n=1} C_{jn}(t) \left( \widetilde{A}_{jn}(t) \otimes \widetilde{B}_{jn}(t) \right) \widetilde{\rho_S}(t) \rho_B \sum_{j' \in J} \sum_{n'=1} C_{j'n'}(s) \left( \widetilde{A}_{j'n'}(s) \otimes \widetilde{B}_{j'n'}(s) \right) \quad (1565)$$

$$- \sum_{j' \in J} \sum_{n'=1} C_{j'n'}(s) \left( \widetilde{A}_{j'n'}(s) \otimes \widetilde{B}_{j'n'}(s) \right) \widetilde{\rho_S}(t) \rho_B \sum_{j \in J} \sum_{n=1} C_{jn}(t) \left( \widetilde{A}_{jn}(t) \otimes \widetilde{B}_{jn}(t) \right) \quad (1566)$$

$$+ \widetilde{\rho_S}(t) \rho_B \sum_{j' \in J} \sum_{n'=1} C_{j'n'}(s) \left( \widetilde{A}_{j'n'}(s) \otimes \widetilde{B}_{j'n'}(s) \right) \sum_{j \in J} \sum_{n=1} C_{jn}(t) \left( \widetilde{A}_{jn}(t) \otimes \widetilde{B}_{jn}(t) \right) \Big) ds \quad (1567)$$

In order to calculate the correlation functions we define:

$$\Lambda_{jn j'n'}(\tau) = \left\langle \widetilde{B}_{jn}(t) \widetilde{B}_{j'n'}(s) \right\rangle_B \quad (1568)$$

$$= \left\langle \widetilde{B}_{jn}(\tau) \widetilde{B}_{j'n'}(0) \right\rangle_B \quad (1569)$$

Here  $s \rightarrow t - \tau$  and  $\text{Tr}_B \left( \widetilde{B}_{jn}(t) \widetilde{B}_{j'n'}(s) \rho_B \right) = \left\langle \widetilde{B}_{jn}(t) \widetilde{B}_{j'n'}(s) \right\rangle_B$ . To evaluate the trace respect to the bath we need to recall that our master equation depends of elements related to the bath and represented by the operators  $\widetilde{B}_{jn}(t)$  and elements related to the system given by  $\widetilde{A}_{jn}(t)$ . The systems considered are in different Hilbert spaces so  $\text{Tr} \left( \widetilde{A}_{jn}(t) \widetilde{B}_{j'n'}(t) \right) = \text{Tr} \left( \widetilde{A}_{jn}(t) \right) \text{Tr} \left( \widetilde{B}_{j'n'}(t) \right)$ . The correlation functions relevant of the master equation (1567) are:

$$\text{Tr}_B \left( \widetilde{B_{jn}}(t) \widetilde{B_{j'n'}}(s) \rho_B \right) = \left\langle \widetilde{B_{jn}}(t) \widetilde{B_{j'n'}}(s) \right\rangle_B \quad (1570)$$

$$= \left\langle \widetilde{B_{jn}}(0) \widetilde{B_{j'n'}}(0) \right\rangle_B \quad (1571)$$

$$= \Lambda_{jnj'n'}(\tau) \quad (1572)$$

$$\text{Tr}_B \left( \widetilde{B_{jn}}(t) \rho_B \widetilde{B_{j'n'}}(s) \right) = \text{Tr}_B \left( \widetilde{B_{j'n'}}(s) \widetilde{B_{jn}}(t) \rho_B \right) \quad (1573)$$

$$= \left\langle \widetilde{B_{j'n'}}(s) \widetilde{B_{jn}}(t) \right\rangle_B \quad (1574)$$

$$= \left\langle \widetilde{B_{j'n'}}(-\tau) \widetilde{B_{jn}}(0) \right\rangle_B \quad (1575)$$

$$= \Lambda_{j'n'jn}(-\tau) \quad (1576)$$

$$\text{Tr}_B \left( \widetilde{B_{j'n'}}(s) \rho_B \widetilde{B_{jn}}(t) \right) = \text{Tr}_B \left( \widetilde{B_{jn}}(t) \widetilde{B_{j'n'}}(s) \rho_B \right) \quad (1577)$$

$$= \left\langle \widetilde{B_{jn}}(t) \widetilde{B_{j'n'}}(s) \right\rangle_B \quad (1578)$$

$$= \left\langle \widetilde{B_{jn}}(\tau) \widetilde{B_{j'n'}}(0) \right\rangle_B \quad (1579)$$

$$= \Lambda_{jnj'n'}(\tau) \quad (1580)$$

$$\text{Tr}_B \left( \rho_B \widetilde{B_{j'n'}}(s) \widetilde{B_{jn}}(t) \right) = \text{Tr}_B \left( \widetilde{B_{j'n'}}(s) \widetilde{B_{jn}}(t) \rho_B \right) \quad (1581)$$

$$= \left\langle \widetilde{B_{j'n'}}(s) \widetilde{B_{jn}}(t) \right\rangle_B \quad (1582)$$

$$= \left\langle \widetilde{B_{j'n'}}(-\tau) \widetilde{B_{jn}}(0) \right\rangle_B \quad (1583)$$

$$= \Lambda_{j'n'jn}(-\tau) \quad (1584)$$

We made use of the cyclic property for the trace to evaluate the correlation functions, from the equations obtained in (1560) and (1567) and using the equations (1570)-(1584) we can re-write:

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \sum_{j,j',n,n'} \left( C_{jn}(t) C_{j'n'}(s) \left( \Lambda_{jnj'n'}(\tau) \widetilde{A_{jn}}(t) \widetilde{A_{j'n'}}(s) \widetilde{\rho_S}(t) - \Lambda_{j'n'jn}(-\tau) \widetilde{A_{jn}}(t) \widetilde{\rho_S}(t) \widetilde{A_{j'n'}}(s) \right) \right. \quad (1585)$$

$$\left. + C_{jn}(t) C_{j'n'}(s) \left( \Lambda_{j'n'jn}(-\tau) \widetilde{\rho_S}(t) \widetilde{A_{j'n'}}(s) \widetilde{A_{jn}}(t) - \Lambda_{jnj'n'}(\tau) \widetilde{A_{j'n'}}(s) \widetilde{\rho_S}(t) \widetilde{A_{jn}}(t) \right) \right) ds \quad (1586)$$

$$= - \int_0^t \sum_{j,j',n,n'} \left( C_{jn}(t) C_{j'n'}(s) \left( \Lambda_{jnj'n'}(\tau) \left[ \widetilde{A_{jn}}(t), \widetilde{A_{j'n'}}(s) \widetilde{\rho_S}(t) \right] + \Lambda_{j'n'jn}(-\tau) \left[ \widetilde{\rho_S}(t) \widetilde{A_{j'n'}}(s), \widetilde{A_{jn}}(t) \right] \right) \right) \quad (1587)$$

$$\frac{d\widetilde{\rho_S}(t)}{dt} = - \int_0^t \sum_{j,j',n,n'} \left( C_{jn}(t) C_{j'n'}(t-\tau) \left( \Lambda_{jnj'n'}(\tau) [A_{jn}(t), A_{j'n'}(t-\tau, t) \widetilde{\rho_S}(t)] + \Lambda_{j'n'jn}(-\tau) [\widetilde{\rho_S}(t) A_{j'n'}(t-\tau, t), A_{jn}(t)] \right) \right) d\tau - i [H_S(t), \widetilde{\rho_S}(t)] \quad (1588)$$

For this case we used that  $A_{jn}(t-\tau, t) = U(t) U^\dagger(t-\tau) A_{jn}(t) U(t-\tau) U^\dagger(t)$ . This is a non-Markovian equation and if we take  $n = 2$  (two sites),  $\mu_0(t) = 0$ ,  $\mu_1(t) = 1$  then we can reproduce a similar expression to (447) as expected.

## VII. BIBLIOGRAPHY

- [1] McCutcheon D P S, Dattani N S, Gauger E M, Lovett B W and Nazir A 2011 Phys. Rev. B 84 081305
- [2] Dara P S McCutcheon and Ahsan Nazir 2010 New J. Phys. 12 113042
- [3] Supplement: Theoretical model of phonon induced dephasing. A.J. Ramsay et al 2009.
- [4] Felix A Pollock et al 2013 New J. Phys. 15 075018

---

\* [n.dattani@cfa.harvard.edu](mailto:n.dattani@cfa.harvard.edu)  
† [edcchaparroso@unal.edu.co](mailto:edcchaparroso@unal.edu.co)