

# **Bulkley River Watershed Connectivity Restoration Plan: 2021 - 2031**

Canadian Wildlife Federation

20-08-2024

# Table of contents

<b>Acknowledgements</b>	<b>5</b>
<b>Project Overview</b>	<b>7</b>
Connectivity Plan Purpose and Approach . . . . .	7
Vision Statement . . . . .	8
Project Scope . . . . .	8
Target Species . . . . .	11
Barrier Types . . . . .	16
<b>Connectivity Status Assessment and Goals</b>	<b>19</b>
Key Ecological Attributes and Current Connectivity Status . . . . .	19
Goals . . . . .	20
<b>Barrier Prioritization</b>	<b>21</b>
Bulkley River Watershed Barrier Prioritization Summary . . . . .	21
<b>Work Planning</b>	<b>48</b>
Annual Progress Update . . . . .	48
Annual Work Plan . . . . .	48
Operational Plan . . . . .	52
<b>References</b>	<b>54</b>

<b>Appendices</b>	<b>55</b>
<b>Project Partners</b>	<b>55</b>
Planning Team . . . . .	55
Key Actors . . . . .	56
<b>Supplementary Information</b>	<b>59</b>
Situation Analysis . . . . .	59
Strategies & Actions . . . . .	59
Strategy 1: Barrier Remediation . . . . .	59
Strategy 2: Barrier Prevention . . . . .	60
Strategy 3: Strengthen Indigenous Connections to Land and Water . . . . .	60
Strategy 4: Planning and Progress Tracking . . . . .	60
Theories of Change & Objectives . . . . .	60
Funding Sources . . . . .	62
<b>Data Download and Methods</b>	<b>70</b>
Connectivity Status Assessment Methods . . . . .	70



# Acknowledgements



This plan represents the culmination of a collaborative planning process undertaken in the Bulkley River watershed over many months of work with a multi-partner planning team of individuals and groups passionate about the conservation and restoration of freshwater ecosystems and the species they support. Plan development was funded by the BC Salmon Restoration and Innovation Fund and the RBC Bluewater Project. We were fortunate to benefit from the feedback, guidance, and wisdom of many groups and individuals who volunteered their time throughout this process — this publication would not have been possible without the engagement of our partners and the planning team. (see Table 15).

We recognize the incredible fish passage and connectivity work that has occurred in the Bulkley River watershed to date, and we are excited to continue partnering with local groups and organizations to build up existing initiatives and provide a road map to push connectivity remediation forward over the next 10 years and beyond

The Canadian Wildlife Federation recognizes that the lands and waters that form the basis of this plan are the traditional unceded territory of the Wet'suwet'en and Gitxsan peoples. We are grateful for the opportunity to learn from the stewards of this land and work together to benefit Pacific Salmon and Steelhead. A special thank you to Dallas Nikal, Mike Ridsdale, and Elaine Sampson for sharing the traditional Witsuwit'en and Gitxsanimax names used in this plan.

# Project Overview

## Connectivity Plan Purpose and Approach

The following Watershed Connectivity Remediation Plan (WCRP) represents the culmination of a one-year collaborative planning effort for the Bulkley River watershed (excluding the Morice River, see Figure 1), the overall aim of which is to clarify and reduce the threat of aquatic barriers to anadromous salmonids and the livelihoods that they support, including the values and laws of First Nations, as well their continued sustenance, cultural, and ceremonial needs both now and into the future. This 10-year plan was developed to identify priority actions that the Bulkley River WCRP planning team (see Table 15 for a list of team members) will undertake between 2021-2031 to conserve and restore fish passage in the watershed through strategies aimed at barrier remediation, barrier prevention, and strengthening Indigenous connections to land and water.

WCRPs are long-term, actionable plans that blend local stakeholder and rightsholder knowledge with innovative GIS analyses to gain a shared understanding of where remediation efforts will have the greatest benefit for anadromous salmonids. The planning process is inspired by the [Conservation Standards](#) (v.4.0), which is a conservation planning framework that allows planning teams to systematically identify, implement, and monitor strategies to apply the most effective solutions to high priority conservation problems. There is a rich history of connectivity and fish passage planning and remediation work in the Bulkley River watershed that this WCRP builds upon, including the work undertaken by the BC Fish Passage Technical Working Group, the Skeena Fisheries Commission, the Office of the Wet'suwet'en, the Wet'suwet'en First Nation, and the Society for Ecosystem Restoration in Northern British Columbia (SERNbc) among others (Wilson and Rabnet. (2007), McCarthy and Fernando (2015), Smith (2018), Casselman and Stanley (2010), A. Irvine (2018)). The Canadian Wildlife Federation will continue to engage and coordinate with local partners and existing initiatives, in part through the Skeena Environmental Stewardship Initiative. SERNbc is also currently undertaking [fish passage](#) work in the Bulkley River watershed, with some overlap and some differences in scope compared to the work and processes described in this WCRP. The SERNbc project relies on expert knowledge and field assessments in both the Bulkley and Morice watersheds to improve passage for all fish, including fluvial and resident species, and focuses on stream crossings that act as barriers (Irvine (2018)). This WCRP focuses specifically on improving connectivity for

anadromous salmonid species and uses consensus-based planning exercises and spatial model implementation to develop watershed-scale status assessments, goal setting, and prioritization for multiple barrier types. SERNbc and the WCRP planning team are currently collaborating on the development of the [bcfishpass](#) connectivity model and will continue to work together to promote coordination and collaboration between the two initiatives moving forward.

The planning team compiled existing barrier location and assessment data, habitat data, and previously identified priorities, and combined this with local and Indigenous knowledge to create a strategic watershed-scale plan to improve connectivity. To expand on this work, the Bulkley River WCRP planning team applied the WCRP planning framework to define the “thematic” scope of freshwater connectivity and refine the “geographic” scope to identify those portions of the watershed where barrier prioritization will be conducted, and subsequent remediation efforts will take place. Additionally, the team selected target fish species, assessed their current connectivity status in the watershed, defined concrete goals for gains in connectivity, and developed a priority list of barriers for remediation to achieve those goals. While the current version of this plan is based on the best-available information at the time of publishing, WCRPs are intended to be “living plans” that are updated regularly as new information becomes available, or if local priorities and contexts change. As such, this document should be interpreted as a current “snap-shot” in time, and future iterations of this WCRP will build upon the material presented in this plan to continuously improve aquatic barrier remediation for migratory fish in the Bulkley River watershed. For more information on how WCRPs are developed, see N. Mazany-Wright et al. (2021c).

## **Vision Statement**

Healthy, well-connected streams and rivers within the Bulkley River watershed support thriving populations of migratory and resident fish. In turn, these fish provide the continued sustenance, cultural, and ceremonial needs of the Wet’suwet’en and Gitxsan peoples, as they have since time immemorial. First Nations, residents, and visitors to the watershed work together for environmental stewardship to clarify, implement, and assess the effectiveness of actions to mitigate the negative effects of aquatic barriers, improving the resiliency of streams and rivers for the benefit and appreciation of all.

## **Project Scope**

Connectivity is a critical component of freshwater ecosystems that encompasses a variety of factors related to ecosystem structure and function, such as the ability of aquatic organisms to disperse and/or migrate, the transportation of energy and matter (e.g., nutrient cycling and sediment flows), and temperature regulation (Seliger and Zeiringer (2018).). Though

each of these factors are important when considering the health of a watershed, for the purposes of this WCRP the term “connectivity” is defined as the degree to which aquatic organisms can disperse and/or migrate freely through freshwater systems. Within this context, connectivity is primarily constrained by physical barriers including anthropogenic infrastructure such as dams, weirs, and stream crossings, and natural features such as waterfalls and debris flows. This plan is intended to focus on the direct remediation and prevention of localized, physical barriers instead of the broad land-use patterns that are causing chronic connectivity issues in the watershed. The planning team decided that the primary focus of this WCRP is addressing barriers to longitudinal connectivity (i.e., along the upstream-downstream plane) due to the magnitude of the threat posed by linear development (i.e., road and rail lines) in the watershed.

The primary geographic scope of this WCRP is the Bulkley River watershed, located in the mid-eastern portion of the Skeena River drainage basin in northwestern British Columbia (Figure 1). The scope constitutes the Bulkley River “watershed group” as defined by the [British Columbia Freshwater Atlas](#) (FWA). A consistent spatial framework was necessary to undertake a watershed-selection process at the provincial scale to identify target watersheds to improve connectivity for salmonids. The Bulkley River watershed was identified by the BC Fish Passage Restoration Initiative as one of four target watersheds for WCRP development S. M. N. Mazany-Wright N. and Rebellato (2021b). The Bulkley River watershed has a drainage area of 776,200 ha, spanning from Bulkley Lake in the southeast to the confluence with the Skeena River in the northwest. The watershed is generally divided into the “lower” Bulkley River and the “upper” Bulkley River by the confluence with the Morice River near the town of Houston. Culturally and economically important populations of Chinook Salmon (*Oncorhynchus tshawytscha*), Coho Salmon (*Oncorhynchus kisutch*), Sockeye Salmon (*Oncorhynchus nerka*), and Steelhead (*Oncorhynchus mykiss*) are all found in the watershed, which historically supported Indigenous sustenance and trading economies (Table 1; Irvine (2018)).

Table 1: Target fish species in the Bulkley River watershed. The Gitxsanimax, Witsuwit'en, and Western common and scientific species names are provided.

Gitxsanimax	Witsuwit'en	Common Name	Scientific Name
Ya'aa	Ggis	Chinook Salmon	<i>Oncorhynchus tshawytscha</i>
Eek	Deedzex	Coho Salmon	<i>Oncorhynchus kisutch</i>
Mi'soo	Taalook	Sockeye Salmon	<i>Oncorhynchus nerka</i>
Milit	Tësdli	Steelhead	<i>Oncorhynchus mykiss</i>

The Bulkley River watershed comprises parts of the traditional territories of two matrilineal nations:




Figure 1: The primary geographic scope — the Bulkley River watershed, excluding the Morice River drainage.

- Gitxsan peoples— the traditional Gitxsan Laxyip spans the northern portion of the watershed, including the Suskwa River, and is governed by a hereditary system of 60 Wilps or House Groups who are represented by Simgigyat (hereditary chiefs). Each Wilp has jurisdiction over several Anaat, or fishing sites. The Wilp groups that have territory coinciding with the Bulkley River watershed include Djogaslee, Gyet'm Galdo'o, Luutkudziwas, Axtii Tsex, Yagosip, and Spookw (G. Sebastian pers. comm.). The Gitxsan steward the land and waters based on Ayookw (Gitxsan law) and Adaakw (oral histories; Government (2019), Irvine (2018)). It is necessary to receive permission from the individual Wilp chief for any work to occur on their territory.
- Wet'suwet'en peoples— the Wedzin Kwah (Bulkley River watershed) is part of the larger Wet'suwet'en traditional territory. The hereditary territory is governed by a system made up of five clans – Gilseyhu (Big Frog), Laksilyu (Small Frog), Tsayu (Beaver), Gitdumden (Wolf/Bear) and Laksamshu (Fireweed) – each of which comprises multiple Yikhs (House Groups) represented by hereditary chiefs. The Wet'suwet'en steward the land based on Inuk Nu'at'en (Wet'suwet'en law), and the principle of Yintahk, meaning everything is connected to the land (Office of the Wet'suwet'en (2013), Irvine (2018)). It is necessary to receive permission from the appropriate bands (Witset First Nation or Wet'suwet'en first Nation, Skin Tyee, Nee Tahí Buhn, or Burns Lake Band), nation representatives (Office of the Wet'suwet'en (2013)), and the individual Yikh chiefs for any work to occur on their territory.

The geographic scope of this WCRP was further refined by identifying “potentially accessible” stream segments, which are defined as streams that target species should be able to access in the absence of anthropogenic barriers (Figure 2). Potentially accessible stream segments were spatially delineated using fish species observation and distribution data, as well as data on “exclusionary points”, which are waterfalls greater than 5 m in height and gradient barriers based on species-specific swimming abilities. These maps were explored by the planning team to incorporate additional local knowledge, ensure accuracy, and finalize the constraints on potentially accessible stream segments. All other stream segments were removed from the scope for further consideration. The “constrained geographic scope” formed the foundation for all subsequent analyses and planning steps, including mapping and modelling useable habitat types, quantifying the current connectivity status, goal setting, and action planning N. Mazany-Wright et al. (2021a).

## **Target Species**

Target species represent the ecologically and culturally important species for which habitat connectivity is being conserved and/or restored in the watershed. In the Bulkley River watershed, the planning team selected Anadromous Salmonids as the target species group, which comprises Chinook Salmon, Coho Salmon, Sockeye Salmon, and Steelhead. Anadromous salmonids also include Pink Salmon (*Oncorhynchus gorbuscha*) and Chum Salmon (*Oncorhynchus keta*) as beneficiary




Figure 2: Potentially accessible stream segments within the Bulkley River watershed. These do not represent useable habitat types, but rather identifies the stream segments within which habitat modelling and barrier mapping and prioritization was undertaken.

species (i.e., species that are not actively targeted through the planning process but will also benefit from connectivity improvements for target anadromous species in the watershed). The selection of these target species was driven primarily by the target species of the primary fund supporting this planning work. The planning team also identified other culturally and ecologically important species within the watershed to consider for inclusion in future iterations of the WCRP, including Pacific Lamprey (*Entosphenus tridentatus*) and Bull Trout (*Salvelinus confluentus*).

## **Anadromous Salmonids**

Anadromous salmonids are cultural and ecological keystone species that contribute to productive ecosystems by contributing marine-derived nutrients to the watershed and forming an important food source for grizzly bears and other species Schindler and Quay (2003). Salmon have enduring food, social, and ceremonial value for the Gitxsan and Wet'suwet'en peoples and contribute significant economic value for recreational and commercial fisheries. Salmon have sustained the culture and economies of Indigenous peoples in the watershed since time immemorial – providing the primary food source for communities, supporting wide-ranging trade systems, and helping pass knowledge and ceremony to future generations through fishing and fish processing (SSAF (2021), Office of the Wet'suwet'en (2013), Rescan (2012)).

Anadromous salmonid populations in the Bulkley River watershed have declined significantly in recent decades, leading both the Gitxsan and Wet'suwet'en nations to declare harvest moratoriums or fishing bans in their territories (Office of the Wet'suwet'en (2013), Government (2019)). The stewardship of these resources in their territories are imbued in the spirit and culture of these nations through a symbiotic relationship with these fish species – threats to the fish are threats to the well-being of the Wet'suwet'en and Gitxsan peoples (SSAF (2021)). The stewardship of their waters continues through the work of the [Gitksan Watershed Authorities](#) and the [Wet'suwet'en Fisheries Program](#), as well as collaborative initiatives like the Skeena Environment Stewardship Initiative.

For the purposes of this WCRP, anadromous salmonid populations are defined using Fisheries and Oceans Canada's Conservation Units. A Conservation Unit (CU) is a group of wild Pacific salmon sufficiently isolated from other groups that, if extirpated, is very unlikely to recolonize naturally within an acceptable timeframe, such as a human lifetime or a specified number of salmon generations. Conservation Units are not defined for Steelhead, as such there is no assessment information to provide for the Bulkley River watershed population. See [Data Downloads and Methods](#) for maps of modelled anadromous salmonid spawning and rearing habitat in the Bulkley River watershed.

### **Chinook Salmon | Ya'aa | Ggís| *Oncorhynchus tshawytscha***

Table 2: Chinook Salmon Conservation Units assessments in the Bulkley River watershed. Assessments undertaken by [the Pacific Salmon Foundation \(2020\)](#).

Conservation Unit	Biological Status	Run timing	Trend in spawner abundance (all available data)	Trend in spawner abundance (last 3 generations)
Middle Skeena – Mainstem Tributaries Upper Bulkley River	Good Data Deficient	July-September Data Deficient	575% Data Deficient	-63% Data Deficient

The Middle Skeena – Mainstem Tributaries Chinook Salmon spawn in the mainstem Bulkley River (downstream of the confluence with the Morice River) and in key tributaries, including Telkwa River, Goathorn Creek, Howson Creek, Kathlyn Creek, Suskwa River, Harold Price Creek, and Natlan Creek. The Middle Skeena Chinook Salmon stocks have seen a decline in recent years, particularly over the last three generations of spawners [Pacific Salmon Explorer](#). The upper Bulkley River Chinook Salmon (upstream of the confluence with the Morice River) are the first salmon to return in the year, usually early-to-mid June, marking the start of the salmon fishery in the watershed Office of the Wet'suwet'en (2013). The upper Bulkley River population is known to spawn in the mainstem and tributaries of the Bulkley River, including Buck Creek, Byman Creek, Richfield Creek, Maxan Creek, and Foxy Creek. In some years, low water flows prevent adult Chinook Salmon from migrating past Bulkley Falls. The upper Bulkley Chinook Salmon stocks have been observed to be in decline and are threatened, in part, by habitat degradation, including linear development (e.g., highway, rail, and road infrastructure) that fragments tributaries (Office of the Wet'suwet'en (2013), [Pacific Salmon Explorer](#)).

### Coho Salmon | Eek | Deedzex | *Oncorhynchus kisutch*

Table 3: Coho Salmon Conservation Unit assessment in the Bulkley River Watershed. Assessments undertaken by [the Pacific Salmon Foundation \(2020\)](#).

Conservation Unit	Biological Status	Run timing	Trend in spawner abundance (all available data)	Trend in spawner abundance (last 3 generations)
Middle Skeena	Good	July-September	82%	-26%

Coho Salmon are the most widely dispersed anadromous salmonid species in the Bulkley River watershed due to their ability to move into smaller tributaries, including headwater streams. Coho Salmon spawning migration peaks in early-to-mid August, though traditionally the main Coho Salmon fishery occurs later in the season (Office of the Wet'suwet'en (2013)). Spawning and rearing of the Middle Skeena population is known to occur within the watershed in the mainstem channels of the Bulkley, Telkwa, and Suskwa Rivers, and key tributaries, including Buck Creek, Aitken Creek, McQuarrie Creek, Byman Creek, Richfield Creek, Alport Creek, and Maxan Creek. In recent decades, Coho Salmon distribution has

often been limited to areas downstream of Bulkley Falls, but in years with sufficient flow, tributaries upstream of the falls are well-used by rearing juveniles (M. Risdale pers. comm.). The Coho Salmon population in the watershed appeared to begin recovering around 1998 but has since declined over the last three generations of spawners (Office of the Wet'suwet'en (2013), PSF 2014). Additionally, since 1989, tens of thousands of Coho Salmon fry have been released into the upper Bulkley River mainstem from the Toboggan Hatchery on Toboggan Creek (Office of the Wet'suwet'en (2013)).

### **Sockeye Salmon | Mi'soo | Taalook | *Oncorhynchus nerka***

Table 4: Sockeye Salmon Conservation Unit Assessments in the Bulkley River watershed. There were not enough data to support assessments for Sockeye Salmon populations at the time of analysis. Assessments undertaken by [the Pacific Salmon Foundation \(2020\)](#).

Conservation Unit	Biological Status	Run timing	Trend in spawner abundance (all available data)	Trend in spawner abundance (last 3 generations)
Bulkley/Maxan (Lake type)	Data Deficient	Data Deficient	Data Deficient	Data Deficient
Skeena River (River-type)	Data Deficient	Data Deficient	Data Deficient	Data Deficient

Sockeye Salmon have cultural and commercial importance within the watershed, especially for First Nations communities, in part due to their fat content which is optimal for the smoke-drying process. Limitations on the Sockeye Salmon harvest in the watershed have hindered the ability of the Wet'suwet'en and Gitxsan to practice important cultural activities and the associated sharing of oral traditions and histories (SSAF (2021), Office of the Wet'suwet'en (2013)). The Sockeye Salmon runs generally follow the spring Chinook Salmon migrations in the Bulkley River system, but Bulkley Falls and flows in some parts of Maxan Creek can limit migration during low-flow years. Data are insufficient for population assessments; however, it is believed that the Bulkley/Maxan populations are at risk of extirpation. There are two Sockeye Salmon populations present in the watershed with distinct life histories – the lake-type and the river-type (Office of the Wet'suwet'en (2013), PSF 2014). Two lake-type Sockeye Salmon sub-populations spawn and rear in and around Bulkley Lake and Maxan Lake, and a third lake-type sub-population was extirpated from Toboggan Lake. River-type Sockeye distribution and habitat use within the Bulkley River watershed is not well documented; however, there are records of Sockeye Salmon river spawners in the mainstem Bulkley River around Richfield Creek, McQuarrie Creek, the Morice River confluence, and the Suskwa River mainstem near Natlan Creek.

### **Steelhead | Milit | Tësdłí | *Oncorhynchus mykiss***

Steelhead migrations coincide with the arrival of Coho Salmon in the watershed and are an important traditional food source to augment winter stores (Office of the Wet'suwet'en (2013)). Steelhead are known to spawn and rear in the mainstem Bulkley River and important tributaries, including the Telkwa River, Hubert Creek, Buck Creek, McQuarrie Creek, Byman Creek, Richfield Creek, Ailport Creek, Johnny David Creek, and Robert Hatch Creek. In the lower part of the watershed, Steelhead are known to spawn and rear throughout the Suskwa River system, all the way up through Harold Price Creek and Blunt Creek. Local knowledge indicates that Steelhead populations have been declining in

recent decades and are currently in poor condition throughout the entire watershed. In response, some stocking enhancement actions have been undertaken in an attempt to increase the population in the watershed (Office of the Wet'suwet'en (2013), Chudyk (1979)).

## Barrier Types

The following table highlights which barrier types pose the greatest threat to anadromous salmon in the watershed. The results of this assessment were used to inform the subsequent planning steps, as well as to identify knowledge gaps where there is little spatial data to inform the assessment for a specific barrier type.

Table 5: Barrier Types in the Bulkley River watershed and barrier rating assessment results. For each barrier type listed, Extent refers to the proportion of anadromous salmonid habitat that is being blocked by that barrier type, Severity is the proportion of structures for each barrier type that are known to block passage for target species based on field assessments, and Irreversibility is the degree to which the effects of a barrier type can be reversed and connectivity restored. The amount of habitat blocked used in this exercise is a representation of total amount of combined spawning and rearing habitat.

Barrier Types	Extent	Severity	Irreversibility	Overall Threat Rating:
Road-Stream Crossings	Low	Low	Medium	High
Rail-stream Crossings	Low	Low	Medium	High
Natural Barriers	Medium	Very High	Medium	Medium
Lateral Barriers	Medium	High	Medium	Medium
Large Dams(>3m height)	Low	Low	High	Low
Small Dams(<3m height)	Low	Low	Medium	Low
Trail-stream Crossings	Low	Low	Low	Low

### Road-stream Crossings

Road-stream crossings are the most abundant barrier type in the watershed, with 48 assessed and modelled crossings located on stream segments with modelled habitat. Demographic road crossings (highways, municipal, and paved roads) block 17.6 of habitat (1% of the total blocked habitat), with 0% of assessed crossings having been identified as barriers to fish passage. Resource roads block 17.6 km of habitat (1%), with 0% of assessed crossings have been identified as barriers. Significant land use and linear development throughout the valley bottom has disconnected the Bulkley River from important habitat in many tributaries, including Highway 16 which represents one of the main drivers of fragmentation in the watershed (see Table 9). The collective experience and input from the planning team resulted in a Medium irreversibility rating due to the technical complexity and resources required to remediate road-stream crossings, though it was noted that this differs considerably between resource roads and highway crossings.

## Rail-stream crossings

There are relatively few rail-stream crossings in the watershed (24 crossings on “potentially accessible” streams), but those that exist block significant amounts of habitat (17.6 km or 1% of the total habitat blocked), with more than half of assessed crossings (0%) acting as barriers to anadromous salmonids. All rail-stream crossings in the watershed are associated with the Canadian National (CN) railway running along the Bulkley River. With significant financial costs, technical challenges, and stakeholder engagement required with CN to remediate these barriers, the planning team decided on an overall pressure rating of High for this barrier type.

## Lateral Barriers

There are numerous types of lateral barriers that potentially occur in the watershed, including dykes, berms, and linear development (i.e., road and rail lines), all of which can restrict the ability of anadromous salmonids to move into floodplains, riparian wetlands, and other off-channel habitats. No comprehensive lateral barrier data exists within the watershed, so pressure ratings were based on qualitative local knowledge. Lateral barriers are not thought to be as prevalent as road- or rail-stream crossings but are likely very severe where they do exist. Highway 16 and the CN rail line that run along a significant stretch of the Bulkley River were identified as major lateral barriers that disconnect the mainstem river from its historic floodplain and off-channel habitat. Overall, the planning team decided that a Medium pressure rating adequately captured the effect that lateral barriers are having on connectivity in the watershed, while recognizing that the lack of data on lateral barriers in the watershed is an important knowledge gap to fill.

## Natural Barriers

Natural barriers to fish passage can include debris flows, log jams, sediment deposits, etc., but natural features that have always restricted fish passage (e.g., waterfalls) are not considered under this barrier type. Natural barriers are difficult to include in a spatial prioritization framework due to their transient nature. The planning team felt that the extent of natural barriers in the watershed is seasonal and fluctuates with freshet flow levels. Both current and historic land-use practices, including historic mining and current forest-harvesting impacts, have created sediment wedges that can act as significant barriers to anadromous salmonids. Due to the nature of these land-use practices, the severity of natural barriers was rated as High and the irreversibility as Medium, the latter due to the nature of what would be required to rectify poor land-use practices at a watershed scale. Overall, the planning team felt that a pressure rating of Medium adequately captured the effects of natural barriers.

## Large Dams (>3m height) and Small Dams (<3m height)

There are 24 mapped large and small dams on “potentially accessible” stream segments in the watershed, blocking 17.6 km (0.72% of the total blocked habitat) of modelled spawning and rearing habitat, resulting in a Low extent. The extent rating of these structures was confirmed by the planning team. There is only one known fish passage structure in the watershed and the remaining dams likely block passage for anadromous salmonids. Remediating these dams will require significant resources; however, due to the Low extent of dams in the watershed, a final pressure rating of Low was assigned.

## **Trail-stream crossings**

There is very little spatial data available on trail-stream crossings in the watershed, so the planning team was unable to quantify the true Extent and Severity of this barrier type. However, the planning team felt that trail-stream crossings are not prevalent within the watershed and that where they do exist, they rarely significantly block passage for anadromous salmonids. Given that most crossings will likely be fords or similar structures, the remediation costs associated with these barriers would be quite low. Overall, the planning team felt that the pressure rating for trail-stream crossings was likely Low.

# Connectivity Status Assessment and Goals

## Key Ecological Attributes and Current Connectivity Status

The planning team devised two Key Ecological Attributes (KEAs) and associated indicators to assess the current connectivity status of the watershed – Accessible Spawning Habitat and Accessible Rearing Habitat (Table 6). KEAs are the key aspects of anadromous salmonid ecology that are being targeted by this WCRP. The connectivity status for the Anadromous Salmonid KEAs were used to establish goals to improve habitat connectivity in the watershed and will be the baseline against which progress is tracked over time.

The current connectivity status assessment relies on GIS analyses to map known and modelled barriers to fish passage, identify stream reaches that have potential spawning and rearing habitat, estimate the proportion of habitat that is currently accessible to target species, and prioritize barriers for field assessment that would provide the greatest gains in connectivity. To support a flexible prioritization framework to identify priority barriers in the watershed, two assumptions are made: 1) any modelled (i.e., passability status is unknown) or partial barriers are treated as complete barriers to passage and 2) the habitat modelling is binary, it does not assign any habitat quality values. As such, the current connectivity status will be refined over time as more data on habitat and barriers are collected. For more detail on how the connectivity status assessments were conducted, see [Supplementary Information](#).

Table 6: Connectivity status assessment for spawning (a) and rearing (b) habitat in the Bulkley River watershed. The two KEAs - Accessible Spawning Habitat and Accessible Rearing Habitat - are evaluated by dividing the length of linear habitat (of each type) that is currently accessible to target species by the total length of all linear habitat (of each type) in the watershed.

Target Species	KEA	Indicator	Poor	Fair	Good	Very Good
Andromous Salmon	Available Spawning Habitat	% of total spawning habitat Current Status:	<50%	51-75%	76-90%	>90% 85

**Comments:** Indicator rating definitions are based on the consensus decisions of the planning team. The current status is based on the CWF Barrier Prioritization Model output, which is current as of October 2022.

Table 7

Target Species	KEA	Indicator	Poor	Fair	Good	Very Good
Andromous Salmon	Available Rearing Habitat	% of total rearing habitat Current Status:	<50%	51-75%	76-90%	>90%

**Comments:** Indicator rating definitions are based on the consensus decisions of the planning team. The current status is based on the CWF Barrier Prioritization Model output, which is current as of October 2022.

## Goals

Table 8: Goals to improve spawning (1) and rearing (2) habitat connectivity for target species in the Bulkley River watershed over the lifespan of the WCRP (2021-2031). The goals were established through discussions with the planning team and represent the resulting desired state of connectivity in the watershed. The goals are subject to change as more information and data are collected over the course of the plan timeline (e.g., the current connectivity status is updated based on barrier field assessments).

Goal #	Goal
1	By 2031, the percent (%) of total linear spawning habitat accessible to anadromous salmonids will increase from 85% to 95% within the Bulkley River watershed (i.e., reconnect at least 143 km of spawning habitat)
2	By 2031, the percent (%) of total linear rearing habitat accessible to anadromous salmonids will increase from 69% to 80% within the Bulkley River watershed (i.e., reconnect at least 599 km of rearing habitat)

# Barrier Prioritization

## Bulkley River Watershed Barrier Prioritization Summary

The primary conservation outcome of the WCRP is the remediation of barriers to connectivity in the Bulkley River watershed. To achieve Goals 1 and 2 in this plan, it is necessary to prioritize and identify a suite of barriers that, if remediated, will provide access to a minimum of 20.49 km of spawning habitat and 29.36 km of rearing habitat (Table 9):

Table 9: Spawning and rearing habitat connectivity gain requirements to meet WCRP goals in the Bulkley River watershed.  
The measures of currently accessible and total habitat values are derived from the Intrinsic Potential habitat model described in Supplementary Information.

Habitat Type	Currently accessible (km)	Total	Current Connectivity Status	Goal	Gain required (km)
Spawning	1160.860000	1358.130000	85.47%	95%	129.360000
Rearing	1566.740000	2255.610000	69.46%	80%	237.750000

The barrier prioritization process comprises three stages:

- Stage 1: preliminary barrier list
- Stage 2: intermediate barrier list
- Stage 3: priority barrier list

Initially, the barrier prioritization analysis ranked all barriers in the watershed by the amount of habitat blocked to produce a “preliminary barrier list”, which also accounted for assessing “sets” of barriers for which remediation could be coordinated to maximize connectivity gains. From this list, the top-ranking subset of barriers - comprising more barriers than are needed to achieve the goals - is selected to produce an “intermediate barrier list”. Barriers that did not rank highly in the model results, but were identified as priority barriers by the local partners were also added to the intermediate barrier list. A longer list of barriers is needed due to the inherent assumptions and uncertainty in the connectivity and habitat models and gaps in available data. Barriers that have been modelled (i.e., points where streams and road/rail networks intersect) are assumed to be barriers until field verification is undertaken and structures that have been assessed as “potential” barriers (e.g., may be passable at certain flow levels or for certain life history stages) require further investigation before a definitive remediation decision is made.

Additionally, the habitat model identifies stream segments that have the potential to support spawning or rearing habitat for target species but does not attempt to quantify habitat quality or suitability (see [Supplementary Information](#)), which will require additional field verification once barrier assessments have completed. As such, the intermediate barrier list below (Table 11) should be considered as a starting point in the prioritization process and represents structures that are a priority to evaluate further through barrier assessment and habitat confirmations because some structures will likely be passable, others will not be associated with usable habitat, and others may not be feasible to remediate because of logistic considerations. The intermediate barrier list was updated following the barrier assessments and habitat confirmations that were undertaken during the 2021 field season - some barriers were moved forward to the “priority barrier list” (see Table 12) and others were eliminated from consideration due to one or more of the considerations discussed above (see Table 10). The priority barrier list represents structures that were confirmed to be partial or full barriers to fish passage and that block access to confirmed habitat. Barriers on the priority list were reviewed by planning team members and selected for inclusion for proactive pursuit of remediation. For more details on the barrier prioritization model, please see N. Mazany-Wright et al. (2021a).

Table 10: List of crossings that were removed from the barrier prioritization list following field assessment. Crossings include those prioritized as part of the first iteration of the intermediate barrier list and additional sites selected by 2021 field crews that were removed following discussion with the planning team due to these structures not existing, being passable, not being associated with usable habitat, or deemed not feasible to remediate because of logistic considerations.

ID	Stream name	Reason for removal from prioritization	Comments
1001805665	Canyon Creek	Structure doesn't exist	Adjacent landowner indicated that this crossing does not exist. Quad access site that Dallas may try to get back to.
197957 (1001801969)	Tributary to Bulkley River	Passable	ATV bridge over historic culverts that have been washed out. No fish passage issues.
197905 (1001803682)	Crow Creek	Passable	Bridge
197900 (1001805532)	Crow Creek	Passable	Bridge
198041 (1001802760)	Deep Creek	Passable	Bridge

ID	Stream name	Reason for removal from prioritization	Comments
1001802820	Deep Creek	Structure doesn't exist access with permission obtained from land owner to assess.	Crossing does not exist. Private land
123776	Corya Creek	Passable	Bridge, fairly recent replacement
1001802044	Ailport Creek	Passable	Bridge on private land
197955 (1001800422)	Ailport Creek	Passable	Ford
197914 (1001802089)	Johnny David Creek	Passable	Ford crossing; no issues besides intense cattle-related impacts
197910 (1001804694)	Tributary to Maxan Lake	Passable	Bridge
197964 (1001805529)	Bulkley River	Passable	Embedded and backwatered structure on railway. Not likely causing issues for fish passage. Wetland type area.
197913 (1001802088)	Robert Hatch Creek	Passable	Ford crossing. No issues besides intense cattle related impacts.
198042 (1001805553)	Glass Creek	Passable	Likely closed bottom structures (concrete boxes) but look like bridges. Deep channel with lake upstream. Not likely presenting much of barrier due to deep slow flows.
198047 (1001801773)	Boulder Creek	Structure doesn't exist	

ID	Stream name	Reason for removal from prioritization	Comments
197956 (1001801133)	Tributary to Bulkley River	Structure doesn't exist at this location. Private land access.	No crossing present at this location. Private land access.
123375 (1001800131)	Trib to Thompson Creek	Not suitable habitat	Barrier, but low quality habitat
123775 (1001800161)	Witset Creek	Not suitable habitat	Noted as low habitat value during both assessments in 2013 and 2021
197903 (1001800180)	Trib to Bulkley River	Not suitable habitat	Low quality habitat
197902 (1001800360)	Trib to Bulkley River	Not suitable habitat	Low quality habitat
197904 (1001800372)	Trib to Bulkley River	Not suitable habitat	Low quality habitat
1001800752	Trib to Elwin	Not suitable habitat	Could not be accessed. Unlikely to be a barrier, unlikely to be good habitat (low gradients and flows)
197972 (1001802040)	Trib to Broman Lake	Not suitable habitat	Low quality habitat
197970 (1001803813)	Trib to Broman Lake	Not suitable habitat	Low quality habitat
197907 (1001802611)	Trib to Maxan Creek	Not suitable habitat	Low quality habitat
197908 (1001803697)	Trib to Maxan Creek	Not suitable habitat	Low quality habitat
197906 (1001803706)	Trib to Maxan Creek	Not suitable habitat	Low quality habitat
197963 (1001805531)	Trib to Bulkley River	Not suitable habitat	Low quality habitat
198044 (1001805573)	Trib to Toboggan Creek	Not suitable habitat	Low quality habitat
197969 (1001800583)	Bulkley River (Conrad Lake)	Passable	Bridge
1001802374	Causqua Creek	Structure doesn't exist	
1001804408	Trib to Klo Creek	Not suitable habitat	Unlikely to be suitable habitat

ID	Stream name	Reason for removal from prioritization	Comments
1001806000		Not suitable habitat	Unlikely to be suitable habitat

Table 11: Updated intermediate barrier list resulting from the second barrier prioritization analysis in the Bulkley River watershed. After assessing the potential barriers on the first iteration of the intermediate list (during 2021 field season) and either identifying them as remediation priorities or eliminating them from consideration (e.g., because they passed fish or did not have suitable habitat upstream), the remaining potential barriers in the watershed were re-prioritized. The barriers on this list were prioritized to exceed the connectivity goals of the plan. Barriers highlighted in the same colour represent sets of barriers that have been prioritized as a group. In the Barrier Status column, P = potential barrier and B = confirmed barrier. All barrier assessment data is compiled from the BC Provincial Stream Crossing Inventory System.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of downstream barriers	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
124422	Trib to Waterfall Cr	Municipal road crossing	Habitat confirmation	B	4	0	0	100% back-passage at this location	Continue to watered and monitor fish likely passable to most life stages and species
1001803294	Waterfall Cr	Resource Road	Modelled	B	1	0	0	Barrier assessment	Scheduled for 2022 field season

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers	Spawning habitat downstream blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
198066	Thompson (1001802488Cr)	Resource road crossing	Habitat confirmation	P	0	2.38	2.39	Consider re-routing stream to the north downstream of crossing, continue to monitor passage.	Appears passable for all life stages and species but may present slight barrier during high flows. Landowners adjacent to the stream report that the channel was historically relocated through the fields near the river and they would like to work together to relocate.
198048	Cesford (1001800048Creek)	Highway crossing	Habitat confirmation	B	1	0.67	0.68	Monitor for fish passage once	Two downstream culverts, railway fish crossing observed (198090) is upstream. remediated.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers downstream	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
198090	Cesford Creek	Railway	Barrier Assessment	B	0	2.36	2.36	Work with CN to identify crossing location and status.	No crossing located in the field. Possibly an older culvert that has become blocked and buried in sediment over time.
123446	Tyhee Creek	Demographic road crossing	Habitat confirmation	P	1	0	11.37	Monitor for passage barrier, but once likely downstream passable for barrier (123445) is replaced.	Scored as barrier, but most fish and life stages.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers	Spawning habitat downstream blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
58158	McDowell Creek	Highway crossing	Assessed	B	1	0	0.3	Newly replaced. Baffled and embedded closed-bottom structure  Continue to monitor fish natural passage at this location. Steep gradient (5.5%) and perched 0.36 cm. Likely not a barrier for most adults and flows.	

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers	Spawning habitat downstream blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
57944	Toboggan Creek	Highway crossing	Habitat confirmation	P	0	13.43	24.95	Monitor existing passage for continued functionality; aim for replacement at end of culvert life.	Massive closed-bottom structure. Past back-watering work conducted. Pink salmon present at fences upstream in 2021. High quality habitat with moderate cover.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers downstream	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
(1001800355)	Airport Creek	Highway crossing	Habitat confirmation	P	0197976	0.4	0.4	Monitor during migration to determine extent of barrier.	Scores as barrier, but not likely an issue for any life stage except potentially at high flows. Observed juvenile sized fish swimming through crossing (~140 mm).
197967 (1001800050)	Taman Creek/Bulkley River	Highway crossing	Habitat confirmation	B	0	5.2	16.73	Monitor new crossing for passage.	Replaced by MOTI, confirmed to be passable to fish. Dry at time of assessment with wetland-type area upstream. Unlikely to support salmon.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers	Spawning habitat downstream blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
123770	John Brown Creek	Highway crossing	Habitat confirmation	P	0	11.71	15.88	Monitor during migration to determine efficacy of fish passage.	Very large system. Dolly Varden, Rainbow Trout, and Coho captured upstream indicating partial barrier only. Chinook utilize reach above the culvert for spawning. Some baffles in culvert. Somewhat steep gradients downstream of crossing.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers downstream	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
197663	Johnny David Creek	Highway crossing	Habitat confirmation	P	0	10.28	14.01	Monitor to confirm continued functionality of fish passage; look at open-bottom replacement as long-term fix.	Good candidate for permanent fix and may align with maintenance requirements for the site (inlet is damaged). Backwatering conducted by WFN and LGL in ~2017. Coho upstream and downstream.
198065 (1001800191)	Watson Creek	Demographic road crossing	Assess	1001800255 B	1	0	0.99	Reassess following exam of Hwy 16 crossing (197974).	Very low flow, but channel is mostly watered. Abundant gravels and excellent vegetation cover.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of downstream barriers	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
197974	Watson Creek (1001800356Creek)	Highway crossing	Assessed	B	0	0	1.98	Reassess during higher flows, inspect dam upstream.	Presumed low quality habitat but needs re-assessment during higher flows.
1100001493	Coffin Creek	Dam	Modelled	B	3	0.9	9.02	Barrier assessment	Scheduled for 2022 field season.
123392	Lemieux Creek	Demographic road crossing	Assessed	B	1	1.67	7.14	Habitat confirmation	Scheduled for 2022 field season.
57793	Vallee Creek	Demographic road crossing	Assessed	B	1	1.78	5.6	Habitat confirmation	Scheduled for 2022 field season.
57978		Resource road crossing	Assessed	B	0	0	4.73	Habitat confirmation	Scheduled for 2022 field season.
195288	Gibson Creek	Demographic road crossing	Assessed	B	1	0	4.06	Habitat confirmation	Scheduled for 2022 field season.
58242	Kathlyn Creek	Demographic road crossing	Assessed	B	0	1.77	3.57	Habitat confirmation	Scheduled for 2022 field season.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of downstream barriers	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
58067	Gramophone Creek	Demographic road crossing	Assessed	B	0	3.37	3.43	Habitat confirmation	Scheduled for 2022 field season.
195559	Four Creek	Resource road crossing	Assessed	B	0	0	3.23	Habitat confirmation	Scheduled for 2022 field season.
123699	Driftwood Creek	Resource road crossing	Assessed	B	0	1.4	1.45	Habitat confirmation	Scheduled for 2022 field season.
124504	Coffin Creek	Demographic road crossing	Assessed	B	1	1.32	1.32	Habitat confirmation	Scheduled for 2022 field season.
123426	Robin Creek	Demographic road crossing	Assessed	B	0	1.28	1.28	Habitat confirmation	Scheduled for 2022 field season.
195290	Gibson Creek	Demographic road crossing	Assessed	B	0	0	1.25	Habitat confirmation	Scheduled for 2022 field season.
195943	Stock Creek	Demographic road crossing	Assessed	B	1	0	0.74	Habitat confirmation	Scheduled for 2022 field season.
197653	Perow Creek	Demographic road crossing	Assessed	B	1	0	0.34	Habitat confirmation	Scheduled for 2022 field season.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of downstream barriers	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
123393	Lemieux Creek	Demographic road crossing	Assessed	B	0	0.11	0.11	Habitat confirmation	Scheduled for 2022 field season.
195944	Stock Creek	Resource road crossing	Assessed	B	2	0	0.09	Habitat confirmation	Scheduled for 2022 field season.
197668	Coffin Creek	Railway crossing	Assessed	B	0	0.02	0.02	Habitat confirmation	Scheduled for 2022 field season.
1001800865		Resource road crossing	Modelled	P	0	0.37	15.95	Barrier assessment	Scheduled for 2022 field season.
123697	Driftwood Creek	Resource road crossing	Assessed	P	1	7.67	10.86	Habitat confirmation	Scheduled for 2022 field season.
1001801115	Watson Creek	Resource road crossing	Modelled	P	3	0	7.825	Barrier assessment	Scheduled for 2022 field season.
1001800255	Stock Creek	Demographic road crossing	Modelled	P	3	0	7.09	Barrier assessment	Scheduled for 2022 field season.
1001801071	Perow Creek	Resource road crossing	Modelled	P	2	0	7.05	Barrier assessment	Scheduled for 2022 field season.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of downstream barriers	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
1001801328	Robin Creek	Resource road crossing	Modelled	P	1	0.16	6.05	Barrier assessment	Scheduled for 2022 field season.
1001802047	Bulkley River	Resource road crossing	Modelled	P	2	0	5.37	Barrier assessment	Scheduled for 2022 field season.
1001804538		Resource road crossing	Modelled	P	2	0	5.015	Barrier assessment	Scheduled for 2022 field season.
1001801123	Watson Creek	Resource road crossing	Modelled	P	2	0	4.93	Barrier assessment	Scheduled for 2022 field season.
1001802069	Johnny David Creek	Resource road crossing	Modelled	P	1	0	4.31	Barrier assessment	Scheduled for 2022 field season.
1001806259	Maish Creek	Resource road crossing	Modelled	P	0	0.09	4.25	Barrier assessment	Scheduled for 2022 field season.
1001800670	Vanderven Creek	Resource road crossing	Modelled	P	2	0	3.98	Barrier assessment	Scheduled for 2022 field season.
1001800403		Demographic road crossing	Modelled	P	0	0	3.69	Barrier assessment	Scheduled for 2022 field season.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of downstream barriers	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
1001802798	Coffin Creek	Resource road crossing	Modelled	P	2	3.45	3.66	Barrier assessment	Scheduled for 2022 field season.
1001805019		Resource road crossing	Modelled	P	2	0	3.58	Barrier assessment	Scheduled for 2022 field season.
1001803607		Resource road crossing	Modelled	P	0	0	3.35	Barrier assessment	Scheduled for 2022 field season.
1001802239		Resource road crossing	Modelled	P	0	0	3.3	Barrier assessment	Scheduled for 2022 field season.
1001801039		Resource road crossing	Modelled	P	0	0	3.06	Barrier assessment	Scheduled for 2022 field season.
1001805507	Stock Creek	Railway crossing	Modelled	P	0	1.54	2.23	Barrier assessment	Scheduled for 2022 field season.
1001802482	Vallee Creek	Resource road crossing	Modelled	P	0	0.88	0.88	Barrier assessment	Scheduled for 2022 field season.
1001800067	Perow Creek	Demographic road crossing	Modelled	P	0	0	0.08	Barrier assessment	Scheduled for 2022 field season.

ID	Stream name	Barrier type	Assessment status (completed to date)	Barrier status	Number of barriers downstream	Spawning habitat blocked – all species (km)	Rearing habitat blocked – all species (km)	Next Step	Comments
						Total gain: 71.85	243.57		

Table 12: The Bulkley River watershed priority barrier list, which includes barriers that have undergone field assessment, been reviewed by the planning team, and selected to pursue for proactive remediation.

Aggregated Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
197912	Robert (100180210) Hatch	Private	Private	Design	Barrier	1	1.54	1.700	High	Remediate	Short	Collapsed bridge removed from channel September 2022.
123445	Tyhee Creek	Hwy 16 E MOTI	Design	Barrier	0	0.00	0.490	High	Currently in design phase, channel rehabilitation part of plans	Short	Blocks connectivity to Tyhee Lake	

Aggregated Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
124500	Helps Creek	Lawson Rd	MOTI	Design	Barrier	0	0.84	8.860	Medium	Currently in design phase	Short	Multiple braided channels and beaver ponds for rearing
197640	Trib to Buck Cr	Buck Flats Rd	MOTI	Design	Barrier	0	1.12	6.560	Medium	Currently in design phase	Short	71 ha of suitable wetland rearing habitat upstream
58159	McDowell Creek	Woodmere Nursery	Woodmere Nursery	Design	Barrier	0	0.00	0.450	Medium	Designs complete. No funding secured for replacement yet.	Short	Medium quality habitat. Barrier near mouth of creek; blocks access to high quality habitat upstream. Coho and Chinook juveniles present at time of assessment.

Aggregate Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
197665	Barren Creek	Railway	CN Rail	Design	Passable	0	0.29	0.289	Medium	Culvert at end of life and needs replacing	Short	Salmon known to spawn between railway and highway. Frequent dredging required due to undersized culverts and high bedload movement.
197664	Barren Creek	Hwy 16	MOTI	Design	Passable	1	0.25	4.000	High	Consider replacing once railway fixed	Medium	Abundant gravels suitable for Coho spawning, moderate cover, high rearing potential.

Aggregate Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
124420	Station Creek	Hwy 16	MOTI	Design	Barrier	0	2.96	6.260	Medium	Currently in design phase	Medium	Chicago Creek Restoration Society reports issue with unstable banks downstream of the crossing. Beaver dam removals upstream by CN have caused destruction of past restoration works.
124424	Trib to Waterfall Cr	Railway Crossing	CN Rail	Assessed	Barrier	2	0.00	0.000	Medium	Potential to partner with CN at same time as highway crossing replaced	Medium	Awaiting replacement of highway crossing downstream.

Aggregate Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
197975 (1001801120)	Airport Creek	Private	Private	Habitat confirmation	Partial barrier	1	9.26	18.830	Medium	Continue to work with landowner Short for potential solution	Private driveway. Coho captured upstream and downstream. Heavily dredged upstream.	
197960 (1024704566)	Corya Creek	Railway	CN Rail	Habitat confirmation	Barrier	0	9.07	12.340	High*	Highway crossing recently replaced downstream	Large culvert. Dolly Varden captured upstream and downstream. High quality habitat but glaciated and unlikely to support salmon.	Short

Aggregate Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
124421	Station Creek	11th Ave	Hazelton	Habitat confirmation	Barrier	3	0.00	0.000	Med-High	Hazelton to try to remediate	Work with New	Slow moving almost wetland like stream. Deep glides throughout. Moderate to high stagnant water often smelling of sulfur. Lots of healthy riparian veg. Appears to have very unstable flows.
124487	Porphyry Creek	Hwy 16	MOTI	Habitat confirmation	Barrier	0	4.99	5.710	Medium	Scheduled for replacement in next ~4 years	Short	Complete barrier to fish passage

Aggregated Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
197658	Byman Creek	Hwy 16 E MOTI	Habitat confirmation	Barrier	0	4.99	5.300	High	Culvert at end of life and needs replacing. Stream needs work downstream to prevent dewatering, DFO pursuing options with MOTI	Short	Dry at time of site visit in 2021 but appeared to be flowing again a few weeks later.	
197662	Richfield Cr	Hwy 16 E MOTI	Habitat confirmation	Partial	0	7.59	13.770	High	Major barrier, not currently on MOTI books, some flooding issues upstream also need addressing	Medium	High quality habitat. Coho captured upstream and downstream of crossing.	

Aggregate Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
123377	Thompson Cr	Walcott Rd	MOTI?	Habitat confirmation	Barrier	1	1.71	1.700	High	Relatively easy fix	Medium	High value habitat, frequent deep pools to 80 cm and abundant large woody debris and gravel. Channel hard to locate upstream of confluence with reed canary grass and willow dominated vegetation.

Aggregated Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Downstream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
123544	McDowell Creek	Unnamed	Private	Assessed	Barrier	2	0.00	1.770	Medium	Flashy system, needs riparian restoration, etc.	Long	Private crossing 200 m upstream of Hwy 19 crossing. Noted in 2013 assessment as high quality habitat.
100180019 (198049)	Cesford Creek	Hwy 118	MOTI	Habitat confirmation	Barrier	2	2.98	12.060	High	Look at remediating following two downstream crossings	Long	High quality habitat with good flow
58151	McDowell Creek	Woodmere Rd	MOTI?	Habitat confirmation	Barrier	3	0.00	0.610	Medium	Needs stream restoration work	Long	Medium quality habitat with abundant cover. Additional crossing 58157/123544 200 m upstream
			Total gain:	47.59	100.700							

Aggregate Crossing ID	Stream Name	Road Name	Barrier Owner	PSCIS Status	Barrier Status	No. Down- stream Barriers	Spawning habitat blocked (km)	Rearing habitat blocked (km)	Habitat Quality	Priority Status	Remediation Timeline (short <5 yrs/ Med (5-9 yrs)/ Long (10+ yrs))	Comments
-----------------------------	----------------	--------------	------------------	-----------------	-------------------	---------------------------------	--	---------------------------------------	--------------------	--------------------	--	----------

# Work Planning

## Annual Progress Update

One priority barrier in the Bulkley watershed was removed on McDowell Creek at Woodmere Nursery in summer 2023. This crossing was replaced with a clearspan bridge and opens up approximately 200 m of seasonal off-channel habitat for salmonids during periods of high flow. The Ministry of Transportation and Infrastructure has been working with local partners to advance designs for barriers on Trib to Buck Creek, Helps Creek, Thompson Creek and Gramophone Creek. The Office of the Wet'suwet'en has hired a part-time staff to help with fish passage coordination in the Bulkley watershed. In addition, efforts have been made by CWF and local partners to build a campaign around Station Creek, to convince the BC government to invest in an appropriate crossing at Highway 16.

## Annual Work Plan

Table 13: 2024-2025 Work Plan

Action	Lead	Support	Estimated Cost	Timeline
Identify traditional use areas to help inform where to improve connectivity, as and when information is readily available	OW	CWF		April-June 2024
Continue to work with OW to identify opportunities to expand WCRP planning within OW territory	CWF	OW		October-November 2024

Action	Lead	Support	Estimated CostTimeline
Engage with Wilp/House Groups for planned field work in territory/territories: provide yearly email to House Groups on planned activities, type of field work to be occurring and why, and provide opportunities for feedback.	OW	CWF	April-June 2024
Obtain consent for pre-construction and construction activities in Wilp/House Group territories (Thompson, Trib to Buck, Gramophone)	OW	CWF, MOTI	July-August 2024
CWF to work with OW to explore opportunities for developing objectives around fish passage and areas of fish passage restoration for OW the upper Bulkley and Morice as part of the Water Sustainability Act and long-term water sustainability plan.	OW	CWF	April-November 2024

Action	Lead	Support	Estimated Cost	Timeline
Review results of crossing assessments from 2022 and 2023, and updated connectivity model with WG partners, decide which structures to advance for further field investigation or rehabilitation	CWF	OW, Witset, Gitxsan, Skeena Watershed Coalition, SERNBC, SSAF/ESI		Apr-24
Update model with all current information related to aquatic connectivity in the Bulkley watershed	CWF	SERNBC, OW, Witset		April-May 2024
Update WCRP based on 2022-2023 field results and partner feedback	CWF	OW		April-May 2024
schedule barrier assessment and habitat confirmations for sites flagged for follow up	CWF	OW, Witset, Gitxsan		Aug-Sep 2024
Complete in-depth passage assessments on Ailport, Coffin (railway, highway), John Brown, Johnny David, McDowell Cr (highway), Toboggan Creek, Vallee Creek, Moan (*NEW), Trib to Buck 197646 (*NEW)	OW, Witset, Gitxsan?	CWF, Skeena Watershed Coalition		June, Aug-Oct 2024

Action	Lead	Support	Estimated Cost	Timeline
Engage with CN to determine if Cesford Creek crossing exists and current status of crossing	CWF	OW		May-June 2024
Engage CN Rail re: plans for replacing Barren Creek culvert	CWF	OW		May-June 2024
Engage with landowner at Airport Creek; advance to design if landowner consents	CWF	OW		May-June 2024
Finalize designs for Trib to Buck, Helps, Thompson, Gramophone, potentially advance 1 or more to construction.	MOTI	OW, CWF		April-November 2024
Develop campaign to promote restoration of Mission Creek and its tributaries for salmon; work with barrier owners to address crossings. Includes 1 MOTI crossing, 1 CN Rail crossing, 3 New Hazelton crossings.	Skeena Watershed Coalition	CWF, Gitxsan		April-September 2024
Review 2024 field results with program partners	CWF			Jan-Feb 2025
re-run connectivity models	CWF			Feb-25

Action	Lead	Support	Estimated Cost	Timeline
Develop 2025-2026 work plan	CWF			Feb-25
Update WCRP based on 2024 field results and partner feedback	CWF			Mar-25

## Operational Plan

The operational plan represents a preliminary exercise undertaken by the planning team to identify the potential leads, potential participants, and estimated cost for the implementation of each action in the Bulkley River watershed. The table below summarizes individuals, groups, or organizations that the planning team felt could lead or participate in the implementation of the plan and should be interpreted as the first step in on-going planning and engagement to develop more detailed and sophisticated action plans for each entry in the table. The individuals, groups, and organizations listed under the “Lead(s)” or “Potential Participants” columns are those that provisionally expressed interest in participating in one of those roles or were suggested by the planning team for further engagement (denoted in bold), for those that are not members of the planning team. The leads, participants, and estimated costs in the operational plan are not binding nor an official commitment of resources, but rather provide a roadmap for future coordination and engagement to work towards implementation of the WCRP.

Table 14: Operational plan to support the implementation of strategic species in the Bulkley River watershed.

Strategy / Actions	Lead(s)	Participants
<b>Strategy 1: Barrier Remediation</b>		
1.1 – Remove and decommission barriers	CWF, SERN	Office of the Wet'suwet'en, Skeena
1.2 – Upgrade and resize crossings	CWF, SERN	Office of the Wet'suwet'en, Skeena
1.3 – Install temporary mitigation	SERN	Office of the Wet'suwet'en, Skeena
1.4 – Raise funds to remediate barriers	CWF, SERN	Office of the Wet'suwet'en, BCTS
1.5 – Request regulatory action for non-compliant crossings	TBD	Office of the Wet'suwet'en, CWF
1.6 – Engage with CN to address their barriers	SERN	CWF, ESI/SSAF
1.7 – Knowledge Gap: Continue updating the barrier prioritization model	CWF	Witset First Nation, SERN
1.8 - Knowledge Gap: Adapt the provincial fish passage framework to account for adult fish and low flow conditions	CWF	TBD
1.9 – Knowledge Gap: Assess barriers by applying adapted version of the provincial fish passage framework	Fish Passage Technical Working Group, CWF, SERN	Witset First Nation, ESI/SSAF
<b>Strategy 2: Barrier Prevention</b>		
2.1 – Work with forestry licensees to improve their aquatic connectivity practices	SERN	Witset First Nation, ESI/SSAF
2.2 – Monitor new crossing installation compliance with regulations regarding fish passage	TBD	Witset First Nation, ESI/SSAF
<b>Strategy 3: Strengthen Indigenous Connections to Land and Water</b>		
3.1 – Develop a First Nations Youth Mentorship capacity building program	ESI/SSAF	CWF + CCC, Witset First Nation
3.2 – Engage with Wilp/House Groups for work to occur in their territory	ESI/SSAF	CWF, SERN
3.3 – First Nations guardianship program to monitor fish passage in the watershed	ESI/SSAF	Witset First Nation, SERN, Office
3.4 – Knowledge Gap: Identify traditional use areas to help inform where to improve connectivity	ESI/SSAF	CWF
<b>Strategy 4: Progress Tracking Plan</b>		
4.1– Implement the WCRP Progress Tracking Plan	CWF, SERN	TBD
4.2 – Develop a data catalogue on all current information related to aquatic connectivity in the Bulkley Watershed	TBD	CWF

Strategy / Actions	Lead(s)	Participants
4.3 – Expand this WCRP to include the Morice River Watershed	TBD	CWF
Total:		
Fundraising total:		
Proponent/government contribution total:		

## References

- Casselman, J., and D. Stanley. 2010. "Bulkley/Fulton Watershed Fish Passage Culvert Assessment Program."
- Chudyk, W. E. 1979. "Suskwa River Steelhead Trout: The Colonization of Harold-Price Creek with Hatchery-Reared Steelhead." *Skeena Fisheries Report* 79-1.
- Government, Gitxsan Hulwip. 2019. *Release: Gitxsan Chiefs Extend Fishing Ban to 2020 & Urge Canada, BC and Ministries to Step up.* <http://gitxsan.ca/release-gitxsan-chiefs-extend-fishing-ban-to-2020-season-urge-canada-bc-and-flnro-to-step-up/>.
- Irvine. 2018. *Bulkley River and Morice River Watershed Groups Fish Passage Restoration Planning.* [https://www.newgraphenvironment.com/fish\\_passage\\_bulkley\\_2020\\_](https://www.newgraphenvironment.com/fish_passage_bulkley_2020_)
- Irvine, A. 2018. *Analysis and Priority Identification of Existing Fish Passage Data: Bulkley River.* <http://a100.gov.bc.ca/pub/acat/public/viewReport.do?reportId=56648>.
- Mazany-Wright, N, S M Norris, N W R Lapointe, and B Rebellato. 2021a. "A Freshwater Connectivity Modelling Framework to Support Barrier Prioritization and Remediation in British Columbia." *Canadian Wildlife Federation*, 2021a.
- Mazany-Wright, N, J Noseworthy, S Sra, S M Norris, and N W Lapointe. 2021c. "Breaking down Barriers: A Practitioners' Guide to Watershed Connectivity Remediation Planning." *Canadian Wildlife Federation*, 2021c.
- Mazany-Wright, S. M. Norris, N., and B. Rebellato. 2021b. "B.c. Fish Passage Restoration Initiative Target Watershed Selection Process: Technical Documentation." *Canadian Wildlife Federation*, 2021b.
- McCarthy, M., and A. Fernando. 2015. "2015 Inventory of High Priority Culverted Fish Passage Barriers in the Lower/Middle Skeena, Bulkley, Morice, and Babine River Watersheds."
- Office of the Wet'suwet'en. 2013. *Wet'suwet'en Title and Rights Regarding Canada Department of Fisheries & Oceans and Pacific Trails Pipeline.* [http://www.wetsuweten.com/files/PTP\\_FHCP\\_Response\\_to\\_DFO-25Nov13-Final.pdf](http://www.wetsuweten.com/files/PTP_FHCP_Response_to_DFO-25Nov13-Final.pdf).
- Rescan. 2012. *KSM Project: Gitxsan Nation Traditional Knowledge and Use Desk-Based Research Report.* [https://www.ceaa.gc.ca/050/documents\\_staticpost/49262/89282\\_D\\_Gitxsan\\_Tradnl\\_Use\\_Desk\\_Based\\_Research\\_Report.pdf](https://www.ceaa.gc.ca/050/documents_staticpost/49262/89282_D_Gitxsan_Tradnl_Use_Desk_Based_Research_Report.pdf): Rescan Environmental Services Ltd.
- Schindler, C. Brock, D. E. P. R. Leavitt, and P. D. Quay. 2003. "The Importance of Marine-Derived Nutrients to Lake Productivity and Salmon Population Dynamics over the Last Five Centuries in Southwest Alaska."
- Seliger, Carina, and Bernhard Zeiringer. 2018. "River Connectivity, Habitat Fragmentation and Related Restoration Measures," 171–86.
- Sheer, M B, D S Busch, E Gilbert, J M Bayer, S Lanigan, J L Schei, K M Burnett, and D Miller. 2009. *Development and Management of Fish Intrinsic Potential Data and Methodologies: State of the IP 2008 Summary Report.* Pacific Northwest Aquatic Monitoring Partnership Series 2009—4, 56 pp.
- Smith, J. 2018. "Assessing Barriers to Fish Passage Within the Wet'suwet'en First Nation Traditional Territory."
- SSAF. 2021. "Skeena Sustainability Assessment Forum's State of the Value Report for Fish & Fish Habitat."
- Wilson, T., and K. Rabnet. 2007. *Fish Passage Assessment of Highway 16 and CN Rail in the Bulkley Watershed.* <https://data.skeenosalmon.info/dataset/fish-passage-assessment-highway-16-cn-rail-bulkley>.

# Project Partners

## Planning Team

Table 15: Bulkley River watershed WCRP planning team members. Planning team members contributed to the development of this plan by participating in a series of workshops and document and data review. The plan was generated based on the input and feedback of the local groups and organizations list in this table.

Name	Organization
Betty Rebellato	Canadian Wildlife Federation
Nick Mazany-Wright	Canadian Wildlife Federation
Justin Duncan	Canadian Wildlife Federation
Nicolas Lapointe	Canadian Wildlife Federation
Sarah Sra	Canadian Wildlife Federation
Simon Norris	Hillcrest Geographics
Natalie Newman	Department of Fisheries and Oceans Canada
Peter Dekoning	Department of Fisheries and Oceans Canada
Sandra Devcic	Department of Fisheries and Oceans Canada
Brian Williams	Gitxsan Nation
Elaine Sampson	Gitxsan Nation
Gordon Sebastian	Gitxsan Nation
John Degagne	Ministry of Forests, Lands and Natural Resource Operations
Al Irvine	Society for Ecosystem Restoration in Northern BC
Sean Mitchell	Skeena Sustainability Assessment Forum

Name	Organization
Jesse Stoeppler	Skeena Watershed Conservation Coalition
David Dewit	Office of the Wet'suwet'en
Mike Ridsdale	Office of the Wet'suwet'en
Dallas Nikal	Witset First Nation

## Key Actors

Table 16: Additional Key Actors in the Bulkley River watershed. Key Actors are the individuals, groups, and/or organizations, outside of the planning team, with influence and relevant experience in the watershed, whose engagement will be critical for the successful implementation of this WCRP.

Individual or Organization Name	Role and Primary Interest
B.C. Fish Passage Technical Working Group	The provincial government of B.C. can assist with local knowledge, data, expertise and can help facilitate barrier remediation work.
Cindy Verbeek/ A Rocha Canada, Upper Bulkley Streamkeepers	A Rocha educates and coordinates environmental stewardship activities for local people in the upper Bulkley watershed and can advise and support where needed. A Rocha Canada runs the Upper Bulkley River Streamkeepers project working with local environmental organizations, churches, schools, governments and citizens to ensure the watershed is healthy and flourishing. They have bio-inventory surveying data, water temperature data, and can help ensure that salmon reach their spawning ground.
Canadian National Rail (CN Rail)	CN Rail is a major proponent and barrier owner that can play a role in improving and replacing crossings in the valley bottom of the Bulkley River watershed.

Individual or Organization Name	Role and Primary Interest
David Wilford – Ministry of Forests, Lands, Natural Resource Operations and Rural Development (FLNRO)	Dave Wilford is a hydrologist and researcher working in the watershed.
Gitksan Watershed Authorities	Gitksan Watershed Authorities is a traditional and science-based body of fisheries professionals, biologists, field technicians, community leaders and support staff. This group is governed according to the traditional Gitksan house system. The GWA functions to represent the Gitksan for the management of fisheries on the Skeena and within the Gitksan territories.
Jeff Anderson	Jeff Anderson is a hydrologist at Geomorphic Consulting working in the watershed.
Skeena Knowledge Trust	The Skeena Knowledge Trust can help with data management.
Ken Rabnett	Ken Rabnett is a researcher, field lead and an excellent source of local knowledge.
Ministry of Transportation and Infrastructure (MOTI)	MOTI is a major proponent and barrier owner that can play a role in improving and replacing crossings in the valley bottom of the Bulkley River watershed.
Ministry of Forests, Lands, Natural Resource Operations, and Rural Development (FLNRORD)	FLNRORD can assist with engineering and design work for remediation projects.

Individual or Organization Name	Role and Primary Interest
Environmental Stewardship Initiative – Skeena Sustainability Assessment Forum (ESI:SSAF)	ESI:SSAF is a forum that allows first Nations and the Province to Fully collaborate to enhance environmental sustainability, and to address First Nation's long standing concerns with stewardship of the land and cumulative impacts in their traditional territories by creating meaningful space for traditional ecological knowledge alongside western science. ESI:SSAFF should be engaged to coordinate local implementation of the outcomes of the WCRP and support future planning updates.
SkeenaWild Conservation Trust	SkeenaWild engages in applied scientific research on salmon ecosystems and can help provide data.
Stu Barns/Skeena Fisheries Commission	The Skeena Fisheries Commission is an umbrella organization for First Nations and works directly with the Office of the Wet'suwet'en and the Gitxsan Watershed Authority. They can advise and support where needed.

# Supplementary Information

## Situation Analysis

The following situation model was developed by the WCRP partnership to “map” the project context and brainstorm potential actions for implementation. Green text is used to identify actions that were selected for implementation (see ‘Strategies & Actions’), and red text is used to identify actions that the project team has decided to exclude from the current iteration of the plan, given that they were either outside of the project scope or were deemed to be ineffective by the planning team.

## Strategies & Actions

Effectiveness evaluation of identified conservation strategies and associated actions to improve connectivity for target species in the Bulkley River watershed. The planning team identified four broad strategies to implement through this WCRP, 1) barrier remediation, 2) barrier prevention, 3) strengthen Indigenous connections to land and water, and 4) planning and progress tracking. Individual actions were qualitatively evaluated based on the anticipated effect each action will have on realizing on-the-ground gains in connectivity. Effectiveness ratings are based on a combination of “Feasibility and”Impact”, Feasibility is defined as the degree to which the project team can implement the action within realistic constraints (financial, time, ethical, etc.) and Impact is the degree to which the action is likely to contribute to achieving one or more of the goals established in this plan.

### Strategy 1: Barrier Remediation

ID	Actions	Details
1.1	Remove and decommission barriers	This action represents some projects that would be led by the planning team with conservation partners.
1.2	Upgrade and resize crossings	Examples include installing larger culverts, replacing closed- with open-bottom culverts, or
1.3	Install temporary mitigation	Examples may include installing fish ladders on barriers that cannot be remediated; however, this is a temporary measure.
1.4	Raise funds to remediate barriers	See “Funding Sources” for more information. Consider inviting potential funders to a funding roundtable.
1.5	Request regulatory action for non-compliant crossings	Request provincial and/or federal agencies to require that targeted, high-priority barriers be removed or modified.
1.6	Engage with CN to address their barriers	Build relationships with CN to open a two-way discussion on the scale, priority and impact of their barriers.
1.7	Knowledge Gap: Continue updating the barrier prioritization model	The model has been updated to reflect 2021 field assessments and intermediate barrier reviews.
1.8	Knowledge Gap: Adapt the provincial fish passage framework to account for adult fish and low-flow conditions	

ID	Actions	Details
1.9	Knowledge Gap: Assess barriers by applying adapted version of the provincial fish passage framework	The first three steps are, (1) barrier assessments, (2) habitat confirmations, and (3) remediation.

## Strategy 2: Barrier Prevention

Table 18: Strategy 2

ID	Actions	Details	Feasibility	Impact
2.1	Work with forestry licensees to improve their aquatic connectivity practices	This should include encouraging better consultation before crossings are installed in the first place.	High	Medium
2.2	Monitor new crossing installation compliance with regulations regarding fish passage	This action could be directly tied to action 3.3 - First Nations guardianship program.	Medium	Medium

## Strategy 3: Strengthen Indigenous Connections to Land and Water

ID	Actions	Details
3.1	Develop a First Nations Youth Mentorship capacity building program	The program should emphasize: (1) working with First Nations youth, with emphasis on females, to help build their capacity to work with Wilp/Yikh.
3.2	Engage with Wilp/Yikh for work to occur in their territory	Obtain permission from the appropriate Wilp/Yikh before conducting any work, including barrier assessments.
3.3	First Nations guardianship program to monitor fish passage in the watershed	
3.4	Knowledge Gap: Identify traditional use areas to help inform where to improve connectivity	Used as decision support to choose which projects to proceed with. This information is proprietary, can only be used internally.

## Strategy 4: Planning and Progress Tracking

Table 20: Strategy 4

ID	Actions	Details
4.1	Implement the WCRP Progress Tracking Plan	The WCRP Progress Tracking Plan will help the team to determine whether we are achieving our goals.
4.2	Develop a data catalogue on all current information related to aquatic connectivity in the Bulkley Watershed	
4.3	Explore expanding this WCRP to include the Morice River Watershed	

## Theories of Change & Objectives

Theories of Change are explicit assumptions around how the identified actions will achieve gains in connectivity and contribute towards reaching the goals of the plan. To develop Theories of Change, the planning team developed explicit assumptions for each strategy which helped to




Figure 3: Situation analysis developed by the planning team to identify factors that contribute to fragmentation (orange boxes), biophysical results (brown boxes), and potential strategies/actions to improve connectivity (yellow hexagons) for target species in the Bulkley River watershed.

clarify the rationale used for undertaking actions and provided an opportunity for feedback on invalid assumptions or missing opportunities. The Theories of Change are results oriented and clearly define the expected outcome. The following theory of change models were developed by the WCRP planning team to “map” the causal (“if-then”) progression of assumptions of how the actions within a strategy work together to achieve project goals.

## Funding Sources

Table 21: Potential funding sources for plan implementation in the Bulkley River watershed. The Canadian Wildlife Federation and the planning team can coordinate proposal submission through these sources.

Funding Source	Spending Restrictions and Other Consideration
Land Based Investment Strategy	Assessment and remediation of fish passage using provincial strategic approach. Primarily for remediation of Ministry-owned/orphaned barriers on forest service roads.
Environmental Enhancement Fund	Fish and wildlife passage improvements and restoration at stream and animal crossings at MOTI roads including culvert retrofits and replacement to restore Pacific salmon and trout access, and wildlife tunnels. Primarily for crossings linked to highway infrastructure.
Community Salmon Program	For projects supporting the protection, conservation and enhancement or rehabilitation of Pacific salmonids and their habitat. Funding for volunteer and not-for-profit community-based groups. Applicant must have a significant volunteer component to their group and to the project. Requires 50% match for funding (volunteer, in-kind, donation or other grants).

Funding Source	Spending Restrictions and Other Consideration
Southern Boundary Restoration and Enhancement Fund	<p>Supports 3 activities: (1) develop improved information for resource management; (2) Rehabilitate and restore marine and freshwater habitat; and (3) enhance wild stock production through low technology techniques. Emphasis for funding is on stocks of conservation concern, particularly those contributing to a fishery and stocks of bilateral fishery relevance.</p>
Habitat Conservation Trust Foundation Enhancement and Restoration Grants	<p>Projects that focus on freshwater wild fish, native wildlife species and their habitats, have the potential to achieve a significant conservation outcome, while maintaining or enhancing opportunities for fishing, hunting, trapping, wildlife viewing and associated outdoor recreational activities. Primary focus is on provincially managed fisheries such as Steelhead and Westslope Cutthroat Trout. Requires 50% funding match.</p>
Environmental Damages Fund	<p>Direct funds received from fines, court orders and voluntary payments to priority projects that will benefit Canada's natural environment, under 4 categories of improvement (in order of preference): (1) restoration, (2) environmental quality improvement, (3) research and development, and (4) education and awareness.</p>

Funding Source	Spending Restrictions and Other Consideration
Habitat Stewardship Program for Aquatic Species at Risk	<p>Program for non-profits, Indigenous governments, academic institutions for activities that align with recovery actions identified in SARA recovery documents and/or COSEWIC assessment documents. Project must address one or more of 3 broad categories: (1) Important habitat for aquatic species at risk is improved and/or managed to meet their recovery needs; (2) Threats to aquatic species at risk and/or their habitat are stopped, removed, and/or mitigated; (3) Collaboration and partnerships support the conservation and recovery of aquatic species at risk. Limited to at-risk species listed under COSEWIC and/or SARA as threatened, endangered or special concern.</p>
Canada Nature Fund for Aquatic Species at Risk	<p>Funding program aimed at addressing priority threats for aquatic species at risk listed as endangered, threatened or Special Concern by COSEWIC, as they align with existing federal, provincial or other local recovery plans. Limited to species in the Columbia and Fraser basins in BC, among other priority areas across Canada. Focus on multi-year, multi-partner initiatives that apply an ecosystem or multi-species approach and create a legacy by enabling recovery actions that carry beyond the life of the funding program. Amounts from \$100K-\$1M available per year.</p>

Funding Source	Spending Restrictions and Other Consideration
BC Salmon Restoration and Innovation Fund	<p>Funding for Indigenous enterprises, academia, industry associations, stewardship groups and commercial groups to support initiatives that support the protection and restoration of wild Pacific salmon and other BC fish stocks or ensure fish and seafood sector in BC is environmentally and economically sustainable. Five main priorities including species of concern rebuilding through habitat restoration with priority for projects that are part of a watershed-scale restoration plan/prioritization effort; build on successful previous restoration efforts; focus on critical habitat and/or the rehabilitation of natural ecosystem processes.</p>
Aboriginal Fund for Species at Risk	<p>Program for Indigenous groups for activities that align with recovery actions identified in SARA recovery documents and/or COSEWIC assessment documents for species listed as Endangered, Threatened, or Special Concern by SARA or COSEWIC. Project must address one or more of 4 broad categories: (1) Habitat for species at risk is improved and/or managed to meet their recovery needs; (2) Threats to species at risk and/or their habitat are stopped, removed and/or mitigated; (3) Collaboration, information sharing and partnership between Indigenous communities, governments and organizations and other interested parties (e.g. federal/provincial/territorial governments, academia, industry, private sector) is enhanced; and (4) Capacity within Indigenous communities, to lead in the stewardship of species at risk and contribute to broader SARA implementation, is strengthened.</p>

Funding Source	Spending Restrictions and Other Consideration
Federal Gas Tax Fund - Community Works Fund	Funding available to local governments from federal gas tax, with funds to be allocated for a variety of municipal projects/initiatives, including local roads/bridges and disaster mitigation.
Disaster Mitigation and Adaptation Fund	For those projects where flood risk is high: Funding available to local, regional and provincial governments, private sector, non-profit organizations, and Indigenous groups for projects aimed at reducing the socio-economic, environmental and cultural impacts triggered by natural hazards and extreme weather events and taking into consideration current and future impacts of climate change in communities and infrastructure at high risk. Includes both new construction of public infrastructure and modification/reinforcement of existing infrastructure. Projects must have a minimum of \$20 M in eligible expenditures and can be bundled together.
Community Gaming Grants	Funding for non-profit organizations (check funding program guidelines for specific eligibility requirements) for programs that help to protect and improve the environment by: (1) Conserving or revitalizing local ecosystems, (2) Reducing greenhouse gas emissions, (3) Providing community education or engagement opportunities related to the environment and agriculture or (4) Supporting the welfare of domestic animals and/or wildlife. Grants range from \$100K-250K per year.

Funding Source	Spending Restrictions and Other Consideration
Sitka Foundation	Funding for registered charities, universities, and government agencies (qualified Canadian organizations) for projects related to coastline and watershed conservation and climate change in 4 key areas: (1) land, water, and ocean conservation, (2) scientific research for nature and the environment, (3) public engagement around the importance of a healthy environment, (4) innovative conservation efforts in Canadian communities, at the local, provincial, and federal levels
TULA Foundation	Supports various environmental programs of interest to the Foundation on a case-by-case basis.
Vancouver Foundation	Granting agency for community, social and environmental initiatives for qualified Canadian organizations (charitable organizations, universities, government agencies). Granting programs change on an annual basis.
BC Conservation Foundation Small Project Fund	Funding available to Non-profits, fish and wildlife clubs (sportsmen's associations), businesses, local/regional governments, public organizations and First Nations for projects with demonstrated positive impact for fish, wildlife and habitat, including outreach programs. Preference given to projects where BCCF is not the sole funder.
Real Estate Foundation of BC General Grantslocal and regional governments, and social enterprises registered as C3s for sustainable land use and real estate practices in BC. Funds up to 50% of cash portion of a project.	Funding for First Nations, charities and societies, non-governmental organizations, universities and colleges, trade associations,




Figure 4: Theory of change developed by the planning team for the actions identified under Strategy 1: Barrier Remediation in the Bulkley River watershed.




Figure 5: Theory of change developed by the planning team for the actions identified under Strategy 2: Barrier Prevention in the Bulkley River watershed.

# Data Download and Methods

## Connectivity Status Assessment Methods

The connectivity status assessment for anadromous salmonids in the Bulkley River watershed builds on existing connectivity modelling work undertaken by the BC Fish Passage Technical Working Group, resulting in a flexible, customizable open-source spatial model called “[bcfishpass](#)”. The model spatially locates known and modelled barriers to fish passage, identifies potential spawning and rearing habitat for target species, and estimates the amount of habitat that is currently accessible to target species. The model uses an adapted version of the Intrinsic Potential (IP) fish habitat modelling framework (see Sheer et al. (2009) for an overview of the IP framework). The habitat model uses two geomorphic characteristics of the stream network — channel gradient and mean annual discharge — to identify potential spawning habitat and rearing habitat for each target species. The habitat model does not attempt to definitively map each habitat type nor estimate habitat quality, but rather identifies stream segments that have high potential to support spawning or rearing habitat for each species based on the geomorphic characteristics of the segment. For more details on the connectivity and habitat model structure and parameters, please see N. Mazany-Wright et al. (2021a). The variables and thresholds used to model potential spawning and rearing habitat for each target species are summarized in Table 15. The quantity of modelled habitat for each species was aggregated for each habitat type to inform the two KEAs — Accessible Spawning Habitat and Accessible Rearing Habitat — and represents a linear measure of potential habitat. To recognize the rearing value provided by features represented by polygons for certain species (e.g., wetlands for Coho Salmon and lakes for Sockeye Salmon) a multiplier of 1.5x the length of the stream segments flowing through the polygons was applied.

Table 22: Parameters and thresholds used to inform the Intrinsic Potential habitat model for spawning and rearing habitat for each target species in the Bulkley River watershed.

Species	Channel Gradient (%) (Spawning Habitat)	Mean annual discharge (m <sup>3</sup> /s) (Spawning Habitat)	Channel Gradient (%) (Rearing Habitat)	Mean annual discharge (m <sup>3</sup> /s) (Rearing Habitat)	Minimum area (ha) (Rearing Habitat)	Lake Multiplier (1.5x) (Rearing Habitat)
Chinook Salmon	0-3	0.46-322.5	0-5	0.28-100		

Species	Channel Gradient (%) (Spawning Habitat)	Mean annual discharge (m <sup>3</sup> /s) (Spawning Habitat)	Channel Gradient (%) (Rearing Habitat)	Mean annual discharge (m <sup>3</sup> /s) (Rearing Habitat)	Minimum Lake area (ha) (Rearing Habitat)	Multiplier (1.5x) (Rearing Habitat)
Coho Salmon	0-5  (Busch et al. 2011, Cooney and Holzer 2006)	0.164-59.15  (Bjornn and Reiser 1991, Newcombe 1977, Woll et al. 2017, Roberge et al. 2002, Raleigh and Miller 1986)	0-5  (Bjornn and Reiser 1991, Sloat et al. 1977, Woll et al. 2017, Neuman and Newcombe 1977, Roberge et al. 2002, Sloat et al. 2002, McMahon 1983)	0.03-40  (Woll et al. 2017, Porter et al. 2008, Rosenfeld et al. 2000)	200	Wetland  (Agrawal et al. 2005, Burnett et al. 2007)
Sockeye Salmon	0-2  (Lake 1999, Hoopes 1972)	0.175-65  (Bjornn and Reiser 1991, Woll et al. 2017, Neuman and Newcombe 1977, Roberge et al. 2002)	0-5  (Bjornn and Reiser 1991, Woll et al. 2017, Neuman and Newcombe 1977, Roberge et al. 2002)	0.02-60  (Woll et al. 2017)	200	Lake
Steelhead	0-4	0.447-75	0-7.4	0.02-60		

Species	Channel Gradient (%) (Spawning Habitat)	Mean annual discharge (m <sup>3</sup> /s) (Spawning Habitat)	Channel Gradient (%) (Rearing Habitat)	Mean annual discharge (m <sup>3</sup> /s) (Rearing Habitat)	Minimum Lake area (ha) (Rearing Habitat)	Multiplier (1.5x) (Rearing Habitat)
	(Sheer and Steel 2006, Cooney and Holzer 2006)	(Bjornn and Reiser 1991, Neuman and Newcombe 1977, Roberge et al. 2002)	(Porter et al. 2008)	(Agrawal et al. 2005, Burnett et al. 2007)		