

Math Notes

Math Notes



Canine-Table

Github

POSIX Nexus serves as a comprehensive cross-language reference hub that explores the implementation and behavior of POSIX-compliant functionality across a diverse set of programming environments. Built atop the foundational IEEE Portable Operating System Interface (POSIX) standards, this project emphasizes compatibility, portability, and interoperability between operating systems.

Abstract

Contents

I	Algebra	III
I	Algebraic Identities	III
I	Nomials	V
I	Polynomial Exponents	VI
I	Dividing Polynomials	IX
I	Multiplying Polynomials	X
I	Combining Polynomials	XI
I	FOIL Method	XIV
I	Factoring Polynomials	XV
I	Expanding Polynomials	XXII
I	Lowest Common Denominator	XXIII
I	Brackets	XXV
I	Dividing	XXVIII
I	Multiply	XXIX
I	Radicals	XXX
I	Linear Algebra	XXXIII
I	Linear Equations	XXXIII
I	Coordinate Plane	XXXV
I	Distance Formula	XXXVIII
II	Sets	XLI
II	Complex	XLI
II	Finite Fields	XLI
II	Integers	XLI
II	Natural Numbers	XLII
II	Primes	XLII
II	Quaternions	XLII
II	Rationals	XLII
II	Reals	XLIII
II	Vector Spaces	XLIII
III	Definitions	XLIII
III	Pairs	XLIII
III	Notation	XLVI
III	Delta in Mathematics	XLVI
III	Theta in Mathematics	XLVII
III	Epsilon in Mathematics	XLVII
III	Delta in Mathematics	XLVIII
III	Proportional Symbol	XLIX
III	Infinity in Mathematics	XLIX
III	Perpendicular Symbol	L
III	Plus-Minus and Minus-Plus	LI
III	Parallel Symbol	LII

III	Angle Symbol	LII
III	Similar vs Equivalent	LIII
III	Approximate Symbol	LIV
III	Similar or Equal Symbol	LV
III	Congruent Symbol	LVI
III	Limit Symbol (lim)	LVI



I Algebra

I Algebraic Identities

Difference of Squares

- ➔ **Definition** \Rightarrow Product of sum and difference equals difference of squares
- ➔ **General Formula** $\Rightarrow a^2 - b^2 = (a - b)(a + b)$
- ➔ **Pattern** \Rightarrow Always collapses into two linear factors
- ➔ **Example** $\Rightarrow x^2 - 9 = (x - 3)(x + 3)$
- ➔ **Extension** \Rightarrow Used in rationalizing denominators and factoring quadratics

Sum of Cubes

- ➔ **Definition** \Rightarrow Sum of cubes factors into binomial and trinomial
- ➔ **General Formula** $\Rightarrow a^3 + b^3 = (a + b)(a^2 - ab + b^2)$
- ➔ **Pattern** \Rightarrow Binomial carries the sum, trinomial balances with alternating signs
- ➔ **Example** $\Rightarrow x^3 + 8 = (x + 2)(x^2 - 2x + 4)$
- ➔ **Extension** \Rightarrow Pairs with difference of cubes for full cubic factorization

Difference of Cubes

- ➔ **Definition** \Rightarrow Difference of cubes factors into binomial and trinomial
- ➔ **General Formula** $\Rightarrow a^3 - b^3 = (a - b)(a^2 + ab + b^2)$
- ➔ **Pattern** \Rightarrow Binomial carries the difference, trinomial balances with all positives
- ➔ **Example** $\Rightarrow x^3 - 27 = (x - 3)(x^2 + 3x + 9)$
- ➔ **Extension** \Rightarrow Complements sum of cubes in cubic identities



Perfect Square Trinomial

- ➔ **Definition** \Rightarrow Squaring a binomial produces a trinomial pattern
- ➔ **General Formula** $\Rightarrow (a + b)^2 = a^2 + 2ab + b^2$
- ➔ **Negative Case** $\Rightarrow (a - b)^2 = a^2 - 2ab + b^2$
- ➔ **Pattern** \Rightarrow Always: square + double product + square
- ➔ **Example** $\Rightarrow (x + 3)^2 = x^2 + 6x + 9$
- ➔ **Extension** \Rightarrow Recognizing this pattern speeds up factoring and simplification

Exponent and Radical Identities

- ➔ **Power of a Power** $\Rightarrow (a^m)^n = a^{mn}$
- ➔ **Negative Exponent** $\Rightarrow a^{-n} = \frac{1}{a^n}$
- ➔ **Fractional Exponent** $\Rightarrow \sqrt[n]{a^m} = a^{m/n}$
- ➔ **Product Rule** $\Rightarrow a^m \cdot a^n = a^{m+n}$
- ➔ **Quotient Rule** $\Rightarrow \frac{a^m}{a^n} = a^{m-n}$
- ➔ **Extension** \Rightarrow Links radicals, reciprocals, and powers into one unified system

Binomial Expansion

- ➔ **Definition** \Rightarrow Expansion of $(a + b)^n$ into a sum of terms
- ➔ **General Formula** $\Rightarrow (a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$
- ➔ **Binomial Coefficient** $\Rightarrow \binom{n}{k} = \frac{n!}{k!(n-k)!}$
- ➔ **Pattern** \Rightarrow Exponents of a decrease, exponents of b increase
- ➔ **Symmetry** \Rightarrow Coefficients are symmetric: $\binom{n}{k} = \binom{n}{n-k}$
- ➔ **Connection** \Rightarrow Coefficients form Pascal's Triangle
- ➔ **Example** $\Rightarrow (x + 1)^4 = x^4 + 4x^3 + 6x^2 + 4x + 1$



I Nomials

Nomial Lineage

- ➔ **Monomial** ⇒ One term only, e.g. $7x^2$
- ➔ **Binomial** ⇒ Two terms, e.g. $x + 2$
- ➔ **Trinomial** ⇒ Three terms, e.g. $x^2 + 3x + 2$
- ➔ **Polynomial** ⇒ Many terms, general family
- ➔ **Technique** ⇒ Prefix indicates number of terms
- ➔ **Outcome** ⇒ Classification helps organize algebraic expressions

Monomial: ax^n

(1)

Binomial: $ax^n + bx^m$

(2)

Trinomial: $ax^n + bx^m + cx^k$

(3)

Polynomial Forms

- ➔ **Monomial** ⇒ One term only, e.g. $7x^2$
- ➔ **Binomial** ⇒ Two unlike terms, e.g. $x + 2$
- ➔ **Trinomial** ⇒ Three terms, e.g. $x^2 + 3x + 2$
- ➔ **Polynomial** ⇒ General family with many terms
- ➔ **Technique** ⇒ Prefix indicates number of terms
- ➔ **Outcome** ⇒ Classification helps organize algebraic expressions

Like Terms

- ➔ **Definition** ⇒ Expressions with identical variable parts, e.g. $3x^2$ and $-5x^2$
- ➔ **Variable Match** ⇒ Same variables with same exponents
- ➔ **Coefficient** ⇒ Numbers in front may differ
- ➔ **Technique** ⇒ Combine by adding or subtracting coefficients
- ➔ **Outcome** ⇒ Simplifies polynomials by reducing to fewer terms

**Coefficient**

- ➔ **Definition** \Rightarrow The number in front of a variable, e.g. in $7x$ the coefficient is 7
- ➔ **Variable Match** \Rightarrow It scales the variable part without changing its type
- ➔ **Examples** $\Rightarrow 3x^2$ has coefficient 3, $-5y$ has coefficient -5
- ➔ **Constants** \Rightarrow A constant term like 4 can be seen as coefficient 4 of x^0
- ➔ **Outcome** \Rightarrow Coefficients tell how strongly each variable contributes to the polynomial

I Polynomial Exponents**Definition of Negative Exponent**

- ➔ **Positive Exponent** $\Rightarrow x^n$ means multiply x by itself n times
- ➔ **Negative Exponent** $\Rightarrow x^{-n}$ means reciprocal of x^n
- ➔ **Rule** $\Rightarrow x^{-n} = \frac{1}{x^n}$
- ➔ **Example** $\Rightarrow x^{-3} = \frac{1}{x^3}$
- ➔ **Use** \Rightarrow Negative exponents express division or reciprocals in algebra and calculus

$$x^3 = x \cdot x \cdot x \Rightarrow x^{-3} = \frac{1}{x^3}$$

Using Negative Exponents

- ➔ **Step 1** \Rightarrow Recall exponent rules: $x^a \cdot x^b = x^{a+b}$
- ➔ **Step 2** \Rightarrow Set $a = 3, b = -3$: $x^3 \cdot x^{-3} = x^0$
- ➔ **Step 3** \Rightarrow But $x^0 = 1$
- ➔ **Step 4** \Rightarrow So x^{-3} must equal $\frac{1}{x^3}$
- ➔ **Outcome** \Rightarrow Negative exponent means reciprocal of positive power



Exponent Flip Examples

$$x^5 = \frac{1}{x^{-5}}$$

$$x^{17} = \frac{1}{x^{-17}}$$

$$28x^5 = \frac{28}{x^{-5}}$$

$$x^{-3} = \frac{1}{x^3}$$

$$x^{-3} = \frac{1}{x^3}$$

$$x^{12} = \frac{1}{x^{-12}}$$

$$15x^9 = \frac{15}{x^{-9}}$$

$$x^5 = \frac{1}{x^{-5}}$$

$$x^3 = \frac{1}{x^{-3}}$$

(4)

(5)

(6)

(7)

(8)

Polynomial Exponent Rules Applied

- ➔ **Power of a Power** $\Rightarrow (a^m)^n = a^{mn}$
- ➔ **Nested Powers** \Rightarrow Combine: $8 \cdot 3 = 24$
- ➔ **Outer Flip** \Rightarrow Apply (-9) : $x^{768y^{5z^3}} \rightarrow x^{-6912y^{5z^3}}$
- ➔ **Technique** \Rightarrow Multiply all exponents carefully, preserve inner structure
- ➔ **Outcome** \Rightarrow Final simplified form: $x^{-6912y^{5z^3}}$

$$\begin{aligned} (((x^{32y^{5z^3}})^8)^3)^{-9} &= (x^{32y^{5z^3}})^{8 \cdot 3} \Rightarrow x^{768y^{5z^3}} \\ &= (x^{768y^{5z^3}})^{-9} \Rightarrow x^{-9 \cdot 768y^{5z^3}} \\ &= x^{-6912y^{5z^3}} \end{aligned}$$

Exponent Rules Applied

- ➔ **Power of a Power** $\Rightarrow (a^m)^n = a^{mn}$
- ➔ **Nested Powers** \Rightarrow Combine: $8 \cdot 3 = 24$
- ➔ **Outer Flip** \Rightarrow Apply (-9) : $x^{768y^{5z^3}} \rightarrow x^{-6912y^{5z^3}}$
- ➔ **Technique** \Rightarrow Multiply all exponents carefully, preserve inner structure
- ➔ **Outcome** \Rightarrow Final simplified form: $x^{-6912y^{5z^3}}$



$$x^2 \cdot x^4 = x^{2+4} \Rightarrow x^6$$

$$x^7 \cdot x^5 = x^{5+7} \Rightarrow x^{12}$$

$$x^8 \cdot x^9 = x^{8+9} \Rightarrow x^{17}$$

$$(3x^3)(5x^6) = (3 \cdot 5)x^{3+6} \Rightarrow 15x^9$$

$$(4x^2)(7x^3) = (4 \cdot 7)x^{2+3} \Rightarrow 28x^5$$

$$(4xy^2)(8x^2y^3) = (4 \cdot 8)x^{1+2}(1 \cdot 1)y^{2+3} \Rightarrow 32x^3y^5$$

$$(5x^2y^3)(6x^3y^4) = (5 \cdot 6)x^{2+3}(1 \cdot 1)y^{3+4} \Rightarrow 30x^5y^7$$

$$(7x^3y^4)(8x^5y^7) = (7 \cdot 8)x^{3+5}(1 \cdot 1)y^{4+7} \Rightarrow 56x^8y^{11}$$

$$(x^3)^4 = x^{3 \cdot 4} = x^{12}$$

$$(x^4)^6 = x^{4 \cdot 6} = x^{24}$$

$$(x^3)^5 = x^{3 \cdot 5} = x^{15}$$

$$(3x^2)^4 = 3^{1 \cdot 4}x^{2 \cdot 4} = 3^4x^8 = 81x^8$$

$$(2x^3)^3 = 2^{1 \cdot 3}x^{3 \cdot 3} = 2^3x^9 = 8x^9$$

$$(3x^2)^2(2x^3)^3 = 3^{1 \cdot 2}x^{2 \cdot 2}2^{1 \cdot 3}x^3 \cdot 3$$

$$= 3^2x^42^3x^9$$

$$= 9 \cdot 8x^{4+9}$$

$$= 72x^{13}$$

$$(3^2x^3y^4)^2(2^3x^2y^5)^3 = (3^{2 \cdot 2}x^{3 \cdot 2}y^{4 \cdot 2})(2^{3 \cdot 3}x^{2 \cdot 3}y^{5 \cdot 3})$$

$$= (3^4x^6y^8)(2^9x^6y^{15})$$

$$= (81x^6y^8)(512x^6y^{15})$$

$$= 81 \cdot 512x^6+6y^{8+15}$$

$$= 41472x^{12}y^{23}$$



$$\begin{aligned} -2^3 &= -2 \cdot -2 \cdot -2 = -8 \\ (-2)^3 &= -2 \cdot -2 \cdot -2 = -8 \\ -(-2)^3 &= -2 \cdot -2 \cdot -2 = 8 \end{aligned}$$

$$\begin{aligned} (-7x^2y^3)^0 &= -7^{2 \cdot 0} x^{2 \cdot 0} y^{3 \cdot 0} \\ &= -7^0 x^0 y^0 \\ &= 1 \cdot 1 \cdot 1 \\ &= 1 \end{aligned}$$

$$\begin{aligned} 3x(5x + 8) &= 15x^2 + 24x \\ 4x(x^2 - 2x + 3) &= 4x^3 - 8x^2 + 12x \end{aligned}$$

I Dividing Polynomials

$$\frac{x^8}{x^3} = x^{8-3} \Rightarrow x^5 \quad (9)$$

$$\frac{x^5}{x^2} = x^{5-2} \Rightarrow x^3 \quad (10)$$

$$\frac{x^5}{x^8} = x^{5-8} \Rightarrow x^{-3} \quad (11)$$

$$\frac{x^4}{x^7} = x^{4-7} \Rightarrow x^{-3} \quad (12)$$

$$(13)$$

$$\begin{aligned} \frac{24x^9y^5}{8x^3y^{12}} &= \frac{24}{8} x^{9-3} \frac{1}{1} y^{5-12} \\ &= 3x^6 y^{-7} \\ &= \frac{3x^6}{y^7} \end{aligned}$$



$$\begin{aligned}\frac{12x^5y^{-3}z^4}{36x^8y^{-4}z^{-8}} &= \frac{\frac{12}{3}x^{5-8}}{\frac{36}{3}} \frac{1}{1}y^{-3-(-4)} \frac{1}{1}z^{4-(-8)} \\ &= \frac{x^{-3}y^1z^{12}}{3} \\ &= \frac{yz^{12}}{3x^3}\end{aligned}$$

I Multiplying Polynomials

Multiplication Symbols

- ➔ **Dot** \Rightarrow \cdot is clean, algebraic, avoids confusion with x
- ➔ **Times** \Rightarrow \times is bold, arithmetic, or cross product
- ➔ **Context** \Rightarrow Use \cdot in algebra, \times in arithmetic or vectors
- ➔ **Technique** \Rightarrow Choose based on clarity and audience
- ➔ **Outcome** \Rightarrow Both mean multiplication, but notation signals intent

$$x^2 \cdot x^4 = x^{2+4} \Rightarrow x^6$$

$$x^7 \cdot x^5 = x^{5+7} \Rightarrow x^{12}$$

$$x^8 \cdot x^9 = x^{8+9} \Rightarrow x^{17}$$

$$(3x^3)(5x^6) = (3 \cdot 5)x^{3+6} \Rightarrow 15x^9$$

$$(4x^2)(7x^3) = (4 \cdot 7)x^{2+3} \Rightarrow 28x^5$$





$$(4x^2y^2)(8x^2y^3) = (4 \cdot 8)x^{1+2}(1 \cdot 1)y^{2+3} \Rightarrow 32x^3y^5$$

$$(5x^2y^3)(6x^3y^4) = (5 \cdot 6)x^{2+3}(1 \cdot 1)y^{3+4} \Rightarrow 30x^5y^7$$

$$(7x^3y^4)(8x^5y^7) = (7 \cdot 8)x^{3+5}(1 \cdot 1)y^{4+7} \Rightarrow 56x^8y^{11}$$

I Combining Polynomials

$$x + 4 = 7 \Rightarrow x + 4 - 4 = 7 - 4 \Rightarrow x + 0 = 3$$

$$x + 9 = 15 \Rightarrow x + 9 - 9 = 15 - 9 \Rightarrow x + 0 = 6$$

$$6 + x = 13 \Rightarrow 6 + x - 6 = 13 - 6 \Rightarrow 0 + x - 0 = 7$$

$$x - 3 = 9 \Rightarrow x - 3 + 3 = 9 + 3 \Rightarrow x - 0 = 12 \Rightarrow x = 12$$

$$x - 8 = 7 \Rightarrow x + 8 - 8 = 7 + 8 \Rightarrow x - 0 = 15 \Rightarrow x = 15$$



$$3x + 5 = 11 \Rightarrow 3x + 5 - 5 = 11 - 5 \Rightarrow \frac{3x}{3} = \frac{6}{3} \Rightarrow x = 2$$

$$6.3 = -2 + x \Rightarrow 6.3 + 2 = -2 + 2 + x \Rightarrow 6.3 = 0 + x \Rightarrow x = 6.3$$

$$5 = x - 8 \Rightarrow 5 + 8 = x - 8 + 8 \Rightarrow 13 = x - 0 \Rightarrow \\ \hookrightarrow x = 13$$

$$5 - x = 12 \Rightarrow 5 - 12 - x + x = 12 - 12 + x \Rightarrow \\ \hookrightarrow -7 = 0 + x \Rightarrow x = -7$$

$$-8 = 5 - x \Rightarrow -8 + 8 = 5 + 8 - x \Rightarrow 0 + x = 13 - x + x \Rightarrow \\ \hookrightarrow x = 13$$

$$3x = 12 \Rightarrow \frac{3x}{3} = \frac{12}{3} \Rightarrow \frac{x}{1} = 4 \Rightarrow x = 4$$

$$7x = 14 \Rightarrow \frac{7x}{7} = \frac{14}{7} \Rightarrow \frac{x}{1} = 2 \Rightarrow x = 2$$

$$-6x = -30 \Rightarrow \frac{-6x}{-6} = \frac{-30}{-6} \Rightarrow \frac{-x}{-1} = 5 \Rightarrow x = 5$$



$$-8x = 48 \Rightarrow \frac{-8x}{-8} = \frac{48}{-8} \Rightarrow \frac{-x}{-1} = -6 \Rightarrow x = -6$$

$$7x = -56 \Rightarrow \frac{7x}{7} = \frac{-56}{7} \Rightarrow \frac{x}{1} = -8 \Rightarrow x = -8$$

$$-8x = -72 \Rightarrow \frac{-8x}{-8} = \frac{-72}{-8} \Rightarrow \frac{x}{1} = 9 \Rightarrow x = 9$$

$$4x + 3 = 6x - 15 \Rightarrow 4x - 4x + 3 + 15 = 6x - 4x - 15 + 15 \Rightarrow \\ \hookrightarrow \frac{18}{2} = \frac{2x}{2}; \Rightarrow x = 9$$

$$3(2x - 4) = 5(3x + 2) - 3 \Rightarrow 6x - 12 = 15x + 10 - 3 \Rightarrow \\ \hookrightarrow 6x - 6x - 12 - 7 = 15x - 6x + 7 - 7 \Rightarrow \\ \hookrightarrow \frac{-19}{9} = \frac{9x}{9} \Rightarrow x = \frac{-19}{9}$$

$$\frac{3}{4}x - \frac{2}{3} = 12 \Rightarrow (\frac{3}{4}x \cdot 4)3 - (\frac{2}{3} \cdot 3)4 = 12 \cdot 12 \\ \hookrightarrow (3x)3 - (2)4 = 144 \Rightarrow 9x - 8 + 8 = 144 + 8 \Rightarrow \\ \hookrightarrow \frac{9x}{9} = \frac{152}{9} \Rightarrow x = \frac{152}{9}$$

$$\frac{2}{3}x + 5 = 8 \Rightarrow (\frac{2}{3}x + 5 = 8)3 \Rightarrow 2x + 15 - 15 = 24 - 15 \Rightarrow \\ \hookrightarrow \frac{2x}{2} = \frac{9}{2} \Rightarrow x = \frac{9}{2}$$



$$\begin{aligned}(3x + 5) + (4x - 2) &= 7x + 3 \\(4x^2 + 3x + 9) + (5x^2 + 7x - 4) &= 9x^2 + 10x + 5 \\(5x^2 - 6x - 12) - (7x^2 + 4x - 13) &= 5x^2 - 6x - 12 - 7x^2 - 4x + 13 \\&= 5x^2 - 7x^2 - 12 + 13 - 6x - 4x \\&= -2x^2 + 1 - 10x\end{aligned}$$

I FOIL Method

$$(a + b)(c + d) = ac + ad + bc + bd \quad (14)$$

FOIL Method

- ➔ First ⇒ Multiply first terms: $a \cdot c$
- ➔ Outer ⇒ Multiply outer terms: $a \cdot d$
- ➔ Inner ⇒ Multiply inner terms: $b \cdot c$
- ➔ Last ⇒ Multiply last terms: $b \cdot d$
- ➔ Outcome ⇒ Sum all four products to get the expanded expression

$$\begin{aligned}(2x + 5)(4x^2 - 3x + 6) &= (2 \cdot 4)x^{1+2} + (2 \cdot -3)x^{1+1} + (2 \cdot 6)x \\&\quad \hookrightarrow +(5 \cdot 4)x^2 + (5 \cdot -3)x + (5 \cdot 6) \\&= 8x^3 + (-6 + 20)x + (12 + -15)x + 30 \\&= 8x^3 + 14x^2 - 3x + 30\end{aligned}$$



$$\begin{aligned}
 (3x^2 - 2x + 4)(4x^2 + 5x - 6) &= \\
 \hookrightarrow (3 \cdot 4)x^{2+2} + (3 \cdot 5)x^{2+1} + (3 \cdot -6)x^{2+0} \\
 \hookrightarrow +(-2 \cdot 4)x^{1+2} + (-2 \cdot 5)x^{1+1} + (-2 \cdot -6)x^{1+0} \\
 \hookrightarrow +(4 \cdot 4)x^{0+2} + (4 \cdot 5)x^{0+1} + (4 \cdot -6)x^{0+0} \\
 &= 12x^4 + (15 + -8)x^3 + (-18 + -10 + 16)x^2 + (12 + 20)x + -24 \\
 &= 12x^4 + 7x^3 + -12x^2 + 32x + -24
 \end{aligned}$$

I Factoring Polynomials

$$4a^2 + 2ab - 3a^2b + 5$$

Terms	Factors	Prime Factors
$4a^2$	$4, a^2$	$2, 2, a, a$
$2ab$	$3, a, b$	$2, a, b$
$-3a^2b$	$-3, a^2, b$	$-3, a, a, b$
5	5	5

$$xy^2 - 3x^2y^2 - 6y + z$$

Terms	Factors	Prime Factors
xy^2	$4, a^2$	$2, 2, a, a$
$-3x^2y$	$3, a, b$	$2, a, b$
$-6y$	$-6, y$	$-2, 3, y$
z	z	z

$$-5 + 2(3a^2 - 3t)$$

Terms	Factors	Prime Factors
-5	-5	-5
$2(3a^2 - 3t)$	$2 \cdot 3a^2 - 3t$	$2 \cdot 3a^2 - 3t$
$6t$	$6, t$	$3, 3, t$



$$3x^2 + 5x - 2$$

Terms	Factors	Prime Factors
$3x^2$	$3, x^2$	$3 \cdot x \cdot x$
$5x$	$5, x$	$5, x$
-2	-2	-2

Definition of Prime Factor

- ➔ **Prime** \Rightarrow A number greater than 1 divisible only by 1 and itself
- ➔ **Factor** \Rightarrow A number that divides another evenly
- ➔ **Prime Factor** \Rightarrow A prime number that divides another number exactly
- ➔ **Example** $\Rightarrow 60 = 2^2 \cdot 3 \cdot 5$; prime factors are 2, 3, 5
- ➔ **Use** \Rightarrow Prime factors are the building blocks of integers, used in LCM, GCD, and simplification

Find LCM of 12 and 18

$$12 = 2^2 \cdot 3 \Rightarrow$$

$$18 = 2 \cdot 3^2 \Rightarrow$$

$$\text{LCM} = 2^2 \cdot 3^2 \hookrightarrow 36$$

Using Prime Factors for LCM

- ➔ **Step 1** \Rightarrow Prime factorize each number
- ➔ **Step 2** \Rightarrow Collect all distinct primes
- ➔ **Step 3** \Rightarrow Take the highest power of each prime
- ➔ **Step 4** \Rightarrow Multiply them together
- ➔ **Outcome** $\Rightarrow \text{LCM}(12, 18) = 36$

Euclidean Modulus

```
1  define nx_pt_mod(x, y) {  
2      x = nx_abs(x)  
3      if (x == 0)  
4          return 0  
5      y = nx_abs(y)
```



```

6         if (y > 0)
7             return x - y * nx_pt_trunc(x / y)
8     print "<nx:impurity/>"
9     return -1
10 }

```

The Greatest Common Factor

```

1  define nx_euc(x, y) {
2      auto n
3      if (x == y)
4          return x
5      while (x > 0 && y > 0) {
6          n = x
7          x = nx_pt_mod(y, x)
8          y = n
9      }
10     return n
11 }

```

$$(8, 12) \mapsto \gcd(8, 12) = 4$$

$$8x + 12 \Rightarrow 4\left(\frac{8x}{2} + \frac{12}{4}\right) \Rightarrow 4(2x + 3)$$

$$(4, 2) \mapsto \gcd(4, 2) = 2$$

$$4x^2 + 2x \Rightarrow 2x\left(\frac{4x^2}{2x} + \frac{2x}{2x}\right) \Rightarrow 2x(2x + 1)$$

$$(12, 18) \mapsto \gcd(12, 18) = 6$$

$$12ab^2 + 18a^2b^3 \Rightarrow 6ab^2(2 + 3ab)$$



Definition of Perfect Square

- ➔ **Perfect Square** \Rightarrow A number that can be expressed as n^2 for some integer n
- ➔ **Integer Squared** \Rightarrow Formed by multiplying an integer by itself
- ➔ **Examples** $\Rightarrow 1, 4, 9, 16, 25, 36, \dots$
- ➔ **Non-Examples** $\Rightarrow 2, 3, 5, 6, 7, 10, \dots$
- ➔ **Use** \Rightarrow Perfect squares appear in factoring, radicals, and Pythagorean identities

Check if 49 is a perfect square

$$\begin{aligned} 49 &= 7 \cdot 7 \Rightarrow \\ &= 7^2 \hookrightarrow \end{aligned}$$

Therefore, 49 is a perfect square.

Using Perfect Squares

- ➔ **Step 1** \Rightarrow Identify the number
- ➔ **Step 2** \Rightarrow Ask if it can be written as n^2
- ➔ **Step 3** \Rightarrow If yes, it is a perfect square
- ➔ **Step 4** \Rightarrow If no, it is not
- ➔ **Outcome** \Rightarrow 49 is a perfect square since $49 = 7^2$

Difference of Squares Applied

- ➔ **Identity** $\Rightarrow (a^2 - b^2) = (a - b)(a + b)$
- ➔ **Example** $\Rightarrow x^2 - 9$
- ➔ **Factorization** \Rightarrow Apply rule: $x^2 - 3^2 = (x - 3)(x + 3)$
- ➔ **Technique** \Rightarrow Recognize perfect squares and subtract
- ➔ **Outcome** \Rightarrow Final factored form: $(x - 3)(x + 3)$

$$x^2 - 9 \Rightarrow x^2 - 3^2 \Rightarrow (x - 3)(x + 3)$$



Perfect Squares Difference

- ➔ Identity $\Rightarrow (a^2 - b^2) = (a - b)(a + b)$
- ➔ Example $\Rightarrow x^2 - 9$
- ➔ Factorization $\Rightarrow (x - 3)(x + 3)$
- ➔ Technique \Rightarrow Spot the squares, apply the difference rule
- ➔ Outcome \Rightarrow Factored polynomial form

$$(25) \mapsto \sqrt{(25)} = 5$$

$$x^2 - 25 \Rightarrow (x + 5)(x - 5)$$

$$(x - 5)(x + 5) \Rightarrow x^2 + 5x + -5x + -25 \Rightarrow$$

$$\hookrightarrow x^2 + (5x + -5x \Rightarrow 0) - 25 \Rightarrow x^2 - 25$$

$$(9) \mapsto \sqrt{(9)} = 3$$

$$x^2 - 9 \Rightarrow (x + 3)(x - 3)$$

$$(4) \mapsto \sqrt{(4)} = 2$$

$$x^2 - 4 \Rightarrow (x + 2)(x - 2)$$

$$4x^2 - 25 \Rightarrow (2x + 5)(2x - 5)$$



$$\begin{aligned}(81) &\mapsto \sqrt{(81)} = 9 \\ (16) &\mapsto \sqrt{(16)} = 4 \\ 16x^2 - 25 &\Rightarrow (4x + 9)(4x - 9)\end{aligned}$$

$$25x^2 - 16y^2 \Rightarrow (5x + 4y)(5x - 4y)$$

$$81x^4 - 16y^8 \Rightarrow (9x^2 + 4y^4)(9x^2 - 4y^4) \Rightarrow (3x + 2y^2)(3x - 2y^2)$$

Factor by Grouping Applied

- ➔ Setup \Rightarrow Polynomial with 4 terms
- ➔ Grouping \Rightarrow Split into two pairs
- ➔ Inner Factor \Rightarrow Factor each pair separately
- ➔ Common Binomial \Rightarrow Extract the shared binomial
- ➔ Outcome \Rightarrow Final factored form

$$\begin{aligned}x^3 + 3x^2 + 2x + 6 &\Rightarrow (x^3 + 3x^2) + (2x + 6) \Rightarrow \\ &\hookrightarrow x^2(x + 3) + 2(x + 3) \Rightarrow (x^2 + 2)(x + 3)\end{aligned}$$

$$\begin{aligned}x^3 - 4x^2 + 3x - 12 &\Rightarrow x^2(x - 4) + 3(x - 4) \Rightarrow \\ &\hookrightarrow \frac{x^2(x - 4)}{x - 4} + \frac{3(x - 4)}{x - 4} \Rightarrow \\ &\hookrightarrow (x - 4)(x^2 + 3)\end{aligned}$$



$$\begin{aligned}
 2x^3 - 6x^2 + 4x - 12 &\Rightarrow 2x^2(x - 3) + 4(x - 3) \Rightarrow \\
 &\hookrightarrow \frac{2x^2(x - 3)}{x - 3} + \frac{4(x - 3)}{x - 3} \Rightarrow \\
 &\hookrightarrow (x - 3)(2x^2 + 4)
 \end{aligned}$$

$$\begin{aligned}
 3x^3 + 8x^2 - 6x - 16 &\Rightarrow x^2(3x + 8) - 2(3x + 8) \Rightarrow \\
 &\hookrightarrow \frac{x^2(3x + 8)}{3x + 8} + \frac{-2(3x + 8)}{3x + 8} \Rightarrow \\
 &\hookrightarrow (x^2 - 2)(3x + 8) \Rightarrow \\
 \hookrightarrow 3x^3 - 6x + 8x^2 - 16 &\Rightarrow 3x(x^2 - 2) + 8(x^2 - 2) \Rightarrow \\
 &\hookrightarrow \frac{3x(x^2 - 2)}{x^2 - 2} + \frac{8(x^2 - 2)}{x^2 - 2} \Rightarrow \\
 &\hookrightarrow (x^2 - 2)(3x + 8)
 \end{aligned}$$

Perfect Square Trinomial

- ➔ **Definition** \Rightarrow A trinomial formed by squaring a binomial
- ➔ **General Form** $\Rightarrow (a + b)^2 = a^2 + 2ab + b^2$
- ➔ **Expansion** \Rightarrow First term squared, double product of terms, last term squared
- ➔ **Example** $\Rightarrow (x + 3)^2 = x^2 + 6x + 9$
- ➔ **Negative Case** $\Rightarrow (a - b)^2 = a^2 - 2ab + b^2$
- ➔ **Pattern** \Rightarrow Always: square + double product + square



I Expanding Polynomials

Binomial Expansion

- ➔ **Definition** \Rightarrow Expansion of $(a + b)^n$ into a sum of terms
- ➔ **General Formula** $\Rightarrow (a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$
- ➔ **Binomial Coefficient** $\Rightarrow \binom{n}{k} = \frac{n!}{k!(n-k)!}$
- ➔ **Pattern** \Rightarrow Exponents of a decrease, exponents of b increase
- ➔ **Symmetry** \Rightarrow Coefficients are symmetric: $\binom{n}{k} = \binom{n}{n-k}$
- ➔ **Connection** \Rightarrow Coefficients form Pascal's Triangle
- ➔ **Example** $\Rightarrow (x + 1)^4 = x^4 + 4x^3 + 6x^2 + 4x + 1$

Pascal's Triangle Connection

- ➔ **Structure** \Rightarrow Pascal's Triangle is built by starting with 1 at the top, each entry below is the sum of the two above
- ➔ **Coefficients** \Rightarrow Row n of Pascal's Triangle gives the coefficients for $(a + b)^n$
- ➔ **Symmetry** \Rightarrow Coefficients are symmetric: $\binom{n}{k} = \binom{n}{n-k}$
- ➔ **Example Row 4** \Rightarrow Coefficients 1, 4, 6, 4, 1 expand $(a + b)^4$
- ➔ **Combinatorics** \Rightarrow Each coefficient counts the number of ways to choose k items from n
- ➔ **Extension** \Rightarrow Pascal's Triangle also encodes identities: hockey-stick pattern, Fibonacci connections, binomial sums

Hockey-Stick Identity

- ➔ **Pattern** \Rightarrow Pick a diagonal in Pascal's Triangle, sum entries until a row
- ➔ **Result** \Rightarrow The sum equals the entry just below and to the right (like a hockey stick)
- ➔ **Formula** $\Rightarrow \binom{r}{r} + \binom{r+1}{r} + \cdots + \binom{n}{r} = \binom{n+1}{r+1}$
- ➔ **Example** $\Rightarrow \binom{2}{2} + \binom{3}{2} + \binom{4}{2} = \binom{5}{3}$
- ➔ **Visual** \Rightarrow The diagonal entries form the "shaft," the final entry is the "blade"



Fibonacci Connection

- ➔ **Pattern** \Rightarrow Sum shallow diagonals of Pascal's Triangle
- ➔ **Result** \Rightarrow The sums produce Fibonacci numbers
- ➔ **Example** \Rightarrow Row sums: 1, 1, 2, 3, 5, 8, 13, ...
- ➔ **Formula** $\Rightarrow \text{Fib}(n) = \sum \binom{n-k}{k}$
- ➔ **Visual** \Rightarrow Diagonal "paths" through Pascal's Triangle trace Fibonacci lineage

Binomial Sums

- ➔ **Row Sum** $\Rightarrow \sum_{k=0}^n \binom{n}{k} = 2^n$
- ➔ **Alternating Sum** $\Rightarrow \sum_{k=0}^n (-1)^k \binom{n}{k} = 0$
- ➔ **Weighted Sum** $\Rightarrow \sum_{k=0}^n k \binom{n}{k} = n \cdot 2^{n-1}$
- ➔ **Combinatorial Meaning** \Rightarrow Row sum counts all subsets of an n -element set
- ➔ **Probability Link** \Rightarrow Binomial distribution probabilities sum to 1 using these identities

I Lowest Common Denominator

refer to I to see the euc function.

The Lowest Common Denominator

```

1  define nx_lcd(x, y) {
2      return x * y / nx_euc(x, y)
3  }
```

$$\frac{3}{4} + \frac{2}{5} \Rightarrow \frac{3 \cdot 5}{4 \cdot 5} + \frac{2 \cdot 4}{5 \cdot 4} \Rightarrow \frac{15}{20} + \frac{8}{20} \Rightarrow \frac{15 + 8}{20} \Rightarrow \frac{23}{20}$$

$$\frac{5}{6} + \frac{4}{7} \Rightarrow \frac{5 \cdot 7}{6 \cdot 7} + \frac{4 \cdot 6}{7 \cdot 6} \Rightarrow \frac{35}{42} + \frac{24}{42} \Rightarrow \frac{35 + 24}{42} \Rightarrow \frac{59}{42}$$



$$\frac{7}{5} \cdot \frac{4}{3} \Rightarrow \frac{7 \cdot 4}{5 \cdot 3} \Rightarrow \frac{28}{15} \Rightarrow 1 \frac{13}{15}$$

$$(18, 20) \mapsto \gcd(18, 20) = 2$$
$$\frac{3}{5} \cdot \frac{6}{4} \Rightarrow \frac{3 \cdot 6}{5 \cdot 4} \Rightarrow \frac{18}{20} \Rightarrow \frac{\frac{18}{2}}{\frac{20}{2}} \Rightarrow \frac{9}{10}$$

$$(28, 63) \mapsto \gcd(28, 63) = 7$$
$$(56, 35) \mapsto \gcd(56, 35) = 7$$
$$\frac{28}{63} \cdot \frac{56}{35} \Rightarrow \frac{\frac{28}{7}}{\frac{63}{7}} \cdot \frac{\frac{56}{7}}{\frac{35}{7}} \Rightarrow \frac{4}{9} \cdot \frac{8}{5} \Rightarrow \frac{4 \cdot 8}{9 \cdot 5} \Rightarrow \frac{32}{45}$$

Definition of Keep-Change-Flip

- ➔ **Keep** \Rightarrow Keep the first fraction exactly as it is
- ➔ **Change** \Rightarrow Change the division sign to multiplication
- ➔ **Flip** \Rightarrow Flip the second fraction (take its reciprocal)
- ➔ **Example** $\Rightarrow \frac{3}{4} \div \frac{2}{5}$
- ➔ **Outcome** $\Rightarrow \frac{3}{4} \times \frac{5}{2} = \frac{15}{8}$

$$\frac{3}{4} \div \frac{2}{5} \Rightarrow \frac{3}{4} \times \frac{5}{2}$$



Using Keep-Change-Flip

- ➔ **Step 1** ⇒ Write the division problem
- ➔ **Step 2** ⇒ Keep the first fraction
- ➔ **Step 3** ⇒ Change division to multiplication
- ➔ **Step 4** ⇒ Flip the second fraction
- ➔ **Step 5** ⇒ Multiply across numerators and denominators
- ➔ **Outcome** ⇒ Simplified fraction result

I Brackets

Curly Braces {}

- ➔ **Definition** ⇒ Used to denote sets or grouping in mathematics, e.g. $\{1, 2, 3\}$
- ➔ **Programming** ⇒ Common in code to enclose blocks of statements
- ➔ **TeX** ⇒ Used to group arguments for commands
- ➔ **Example** ⇒ $\{x \mid x > 0\}$ means the set of all positive x
- ➔ **Outcome** ⇒ Curly braces signal structured grouping or set notation

Square Brackets []

- ➔ **Definition** ⇒ Used for intervals, optional elements, or matrices
- ➔ **Interval** ⇒ $[a, b]$ means all values between a and b , inclusive
- ➔ **Matrix** ⇒ Brackets often enclose arrays of numbers
- ➔ **Example** ⇒ $[2, 5]$ includes 2 and 5
- ➔ **Outcome** ⇒ Square brackets emphasize inclusion or structured arrays

$$\begin{aligned}
 8[-6 + 8(-2 + 4)] &= 8[-6 + 8 \cdot 2] \\
 &= 8[-6 + 16] \\
 &= 8 \cdot 10 \\
 &= 80
 \end{aligned}$$

**Parentheses ()**

- ➔ **Definition** \Rightarrow Used for grouping, order of operations, or function arguments
- ➔ **Math** $\Rightarrow (a + b)c$ ensures addition happens before multiplication
- ➔ **Functions** $\Rightarrow f(x)$ shows input to a function
- ➔ **Interval** $\Rightarrow (a, b)$ means values strictly between a and b
- ➔ **Outcome** \Rightarrow Parentheses control grouping and precedence in math and logic

$$\begin{aligned} -(-(1 - 2) + (5)) &= -(-(-1) + 5) \\ &= -(1 + 5) \\ &= -(6) \\ &= -6 \end{aligned}$$

$$\begin{aligned} 3 \cdot (6 - 3 + 1) - 4^2 &= 3 \cdot 4 - 16 \\ &= 12 - 16 \\ &= -4 \end{aligned}$$

$$\begin{aligned} (10, 4) &\mapsto \gcd(10, 4) = 2 \\ \frac{2 \cdot (5 - 1) + 2}{4 \cdot (2 - 1)} &\Leftrightarrow (2 \cdot (5 - 1) + 2) \div (4 \cdot (2 - 1)) \\ &= \frac{2 \cdot 4 + 2}{4 \cdot 1} \\ &= \frac{8 + 2}{4} \\ &= \frac{10}{4} \\ &= \frac{\frac{10}{2}}{\frac{4}{2}} \\ &= \frac{5}{2} \end{aligned}$$



$$\begin{aligned}
 5 + (1 \cdot 4 - (2 - 5) - 2) &= 4 - (-3) - 2 \\
 &= 7 - 2 \\
 &= 5
 \end{aligned}$$

Double Vertical Bars ||

- ➔ **Definition** ⇒ Used to denote absolute value or norm
- ➔ **Absolute Value** ⇒ $|x|$ is distance of x from 0
- ➔ **Norm** ⇒ $\|v\|$ is length of vector v
- ➔ **Example** ⇒ $|-5| = 5$, $\|(3, 4)\| = 5$
- ➔ **Outcome** ⇒ Double bars measure magnitude or distance

$$\begin{aligned}
 4 + |3 - 9| &= 4 + |-6| & (15) \\
 &= 4 + 6 & (16) \\
 &= 10 & (17)
 \end{aligned}$$

Floor Brackets ⌊ ⌋

- ➔ **Definition** ⇒ Used for the floor function
- ➔ **Floor** ⇒ $\lfloor x \rfloor$ = greatest integer less than or equal to x
- ➔ **Example** ⇒ $\lfloor 3.7 \rfloor = 3$
- ➔ **Use** ⇒ Rounds down to nearest integer
- ➔ **Outcome** ⇒ Floor brackets capture downward rounding



Ceiling Brackets []

- ➔ **Definition** ⇒ Used for the ceiling function
- ➔ **Ceiling** ⇒ $\lceil x \rceil$ = smallest integer greater than or equal to x
- ➔ **Example** ⇒ $\lceil 3.2 \rceil = 4$
- ➔ **Use** ⇒ Rounds up to nearest integer
- ➔ **Outcome** ⇒ Ceiling brackets capture upward rounding

I Dividing

Fraction Components

- ➔ **Numerator** ⇒ Top of the fraction, counts selected parts, e.g. in $\frac{3}{4}$ the numerator is 3
- ➔ **Denominator** ⇒ Bottom of the fraction, defines total equal parts, e.g. in $\frac{3}{4}$ the denominator is 4
- ➔ **Relationship** ⇒ Fraction = Numerator ÷ Denominator
- ➔ **Technique** ⇒ Numerator changes with quantity chosen, denominator fixes the partition size
- ➔ **Outcome** ⇒ Understanding both clarifies fraction meaning and operations

$$\frac{6 - 2^3}{2} = \frac{6 - 8}{2} \quad (18)$$

$$= \frac{-2}{2} \quad (19)$$

$$= -1 \quad (20)$$

Verbose Floating-Point Multiplication

- ➔ **Step 1** ⇒ Choose two floating-point numbers, e.g. 3.25 and 2.5
- ➔ **Step 2** ⇒ Express them as fractions: $3.25 = \frac{325}{100}$, $2.5 = \frac{25}{10}$
- ➔ **Step 3** ⇒ Multiply numerators: $325 \times 25 = 8125$
- ➔ **Step 4** ⇒ Multiply denominators: $100 \times 10 = 1000$
- ➔ **Step 5** ⇒ Form product fraction: $\frac{8125}{1000}$
- ➔ **Step 6** ⇒ Simplify fraction: $\frac{8125}{1000} = 8.125$
- ➔ **Step 7** ⇒ Restore decimal form: product is 8.125



$$\begin{aligned}
 10.4 &\Rightarrow \frac{10.4}{1} \frac{10.4}{1} \cdot 10 && \Rightarrow \Rightarrow \frac{104}{10} \\
 1.3 &\Rightarrow \frac{1.3}{1} && \Rightarrow \frac{1.3}{1} \cdot 10 \Rightarrow \frac{13}{10} \\
 1.5 &\Rightarrow \frac{1.5}{1} && \Rightarrow \frac{1.5}{1} \cdot 10 \Rightarrow \frac{15}{10} \\
 -2.35 &\Rightarrow \frac{-2.35}{1} \Leftrightarrow \frac{2.35}{-1} && \Rightarrow \frac{-2.35}{1} \cdot 100 \Rightarrow \frac{-235}{100}
 \end{aligned}$$

$$(10056, 1950) \mapsto \gcd(10056, 1950) = 26$$

$$\begin{aligned}
 \frac{10.3^2 - (-2.35)^2}{1.3(1.5)} &= \frac{106.09 - 5.522}{1.95} \\
 &= \frac{100.568}{1.95} \\
 &= \frac{\frac{100568}{26}}{\frac{1950}{26}} \\
 &= \frac{3868}{75}
 \end{aligned}$$

I Multiply

$$\left(\frac{103}{10}\right)^2 \Rightarrow \frac{103}{10} \cdot \frac{103}{10}$$

 \Rightarrow

	1	0	3
×	1	0	3
	3	0	9
	0	0	0
+	1	0	3
	1	0	6

$$\frac{10609}{100} \Rightarrow 106.09$$



$$\left(\frac{-235}{100}\right)^2 \Rightarrow \frac{-235}{10} \cdot \frac{-235}{10}$$

$$\frac{57665}{100} \Rightarrow 576.65$$

⇒

		−	2	3	5
	×	−	2	3	5
	1	0	5	1	5
		7	1	5	0
+	4	7	0	0	0
	5	7	6	6	5

I Radicals

Identity of $\sqrt[n]{a^n}$

- ➔ **Expression** ⇒ $\sqrt[n]{a^n}$
- ➔ **Meaning** ⇒ The n -th root of a^n
- ➔ **Simplification** ⇒ Cancels the root and exponent, yielding a
- ➔ **Condition** ⇒ Valid when $a \geq 0$ for real roots, or in complex domain otherwise
- ➔ **Example** ⇒ $\sqrt[3]{2^3} = \sqrt[3]{8} = 2$
- ➔ **Extension** ⇒ $\sqrt[n]{a^m} = a^{m/n}$

$$\sqrt[n]{a^n} = a$$

Definition of Radical

- ➔ **Radical** ⇒ An expression that uses the root symbol $\sqrt{}$
- ➔ **Square Root** ⇒ The most common radical, \sqrt{x} , meaning the number which squared gives x
- ➔ **Index** ⇒ The small number above the radical, e.g. $\sqrt[3]{x}$ is the cube root
- ➔ **Radicand** ⇒ The number or expression inside the radical sign
- ➔ **Example** ⇒ $\sqrt{25} = 5$, $\sqrt[3]{8} = 2$
- ➔ **Use** ⇒ Radicals are used to express roots, simplify algebraic expressions, and solve equations





$$\sqrt{50} \Rightarrow \sqrt{25 \cdot 2} \Rightarrow \sqrt{25} \cdot \sqrt{2} \Rightarrow 5\sqrt{2}$$

Using Radicals

- ➔ **Step 1** \Rightarrow Identify the radicand
- ➔ **Step 2** \Rightarrow Factor the radicand into perfect powers and leftovers
- ➔ **Step 3** \Rightarrow Simplify by taking the root of the perfect power
- ➔ **Step 4** \Rightarrow Leave the leftover inside the radical
- ➔ **Outcome** $\Rightarrow \sqrt{50} = 5\sqrt{2}$

Overview of `nx_squares`

- ➔ **Purpose** \Rightarrow Computes square roots using iterative scaling and subtraction
- ➔ **Input** \Rightarrow A single value x
- ➔ **Initialization** \Rightarrow Takes absolute value, sets scaling factor, prepares working variables
- ➔ **Scaling** \Rightarrow Doubles a seed until it exceeds the radicand
- ➔ **Iteration** \Rightarrow Subtracts scaled values, updates root approximation step by step
- ➔ **Condition** \Rightarrow Handles zero and invalid cases gracefully, returns early if needed
- ➔ **Output** \Rightarrow Final square root approximation, or truncated value if iteration fails

Square Roots

```

1  define nx_nr_sqrt(x) {
2      auto y, p
3      if (nx_xy_breach(x, 1) == -1)
4          return -1
5      y = x / 2
6      p = 0
7      while (y != p) {
8          p = y
9          y = (y + x / y) / 2
10     }
11     return y
12 }
```




Square Roots

```
1  define nx_squares(x) {
2      auto a, s, b, y
3      a = nx_abs(x)
4      if (a == 0 || scale == 0)
5          return x
6      s = 1
7      while (s < a)
8          s = s * 2
9      b = nx_scale(1)
10     x = 0
11     y = 0
12     while (s > b) {
13         if (s <= a) {
14             a = a - s
15             x = y
16             y = nx_nr_sqrt(s)
17             if (y == -1)
18                 return x
19             if (x != 0)
20                 print x, ", "
21         }
22         s = s / 3
23     }
24     return y
25 }
```

Definition of Vinculum

- ➔ **Vinculum** ⇒ The horizontal bar drawn over the radicand in a radical
- ➔ **Scope** ⇒ Shows exactly which terms are included under the radical
- ➔ **Fraction Use** ⇒ Also used as the bar separating numerator and denominator in fractions
- ➔ **Repeating Decimal** ⇒ Used to mark repeating digits, e.g. $0.\overline{3}$
- ➔ **Example** ⇒ In $\sqrt{a+b}$, the vinculum extends over $a+b$
- ➔ **Use** ⇒ Ensures clarity of grouping inside radicals, fractions, and repeating decimals

Definition of Radicand

- ➔ **Radicand** ⇒ The number or expression placed under the radical sign $\sqrt{}$
- ➔ **Role** ⇒ It is the quantity from which a root is extracted
- ➔ **Example** ⇒ In $\sqrt{25}$, the radicand is 25
- ➔ **Extended** ⇒ In $\sqrt[3]{8}$, the radicand is 8, and the index is 3
- ➔ **Scope** ⇒ The vinculum (bar) shows exactly which terms belong to the radicand
- ➔ **Outcome** ⇒ Identifying the radicand clarifies what is being rooted in the expression





Square Root Expansion

- ➔ **Radical** $\Rightarrow \sqrt{72}$
- ➔ **Factor** \Rightarrow Break into perfect square and leftover: $72 = 36 \cdot 2$
- ➔ **Expand** $\Rightarrow \sqrt{72} = \sqrt{36 \cdot 2} = \sqrt{36} \cdot \sqrt{2}$
- ➔ **Simplify** $\Rightarrow \sqrt{36} = 6$
- ➔ **Outcome** $\Rightarrow \sqrt{72} = 6\sqrt{2}$

Square Root Truncation

- ➔ **Radical** $\Rightarrow \sqrt{72}$
- ➔ **Approximation** \Rightarrow Leave as decimal: $\sqrt{72} \approx 8.485$
- ➔ **Truncation** \Rightarrow Cut after two decimals: 8.48
- ➔ **Outcome** \Rightarrow Truncated radical value ≈ 8.48

Types of Square Roots

- ➔ **Integer Square Root** \Rightarrow The root of a whole number. It may simplify to a rational (e.g. $\sqrt{25} = 5$) or remain irrational (e.g. $\sqrt{2}$)
- ➔ **Fraction Square Root** \Rightarrow The root of a ratio $\frac{p}{q}$, defined as $\sqrt{\frac{p}{q}} = \frac{\sqrt{p}}{\sqrt{q}}$
- ➔ **Rational Square Root** \Rightarrow Occurs when both numerator and denominator are perfect squares, e.g.

$$\sqrt{\frac{9}{16}} = \frac{3}{4}$$
- ➔ **Irrational Square Root** \Rightarrow Occurs when either numerator or denominator is not a perfect square, e.g.

$$\sqrt{\frac{2}{9}} = \frac{\sqrt{2}}{3}$$
- ➔ **Impurity Case** \Rightarrow When the denominator equals 0, the expression is undefined, e.g. $\sqrt{\frac{p}{0}}$

I Linear Algebra

I Linear Equations

$$x + 4 = 7 \Rightarrow x + 4 - 4 = 7 - 4 \Rightarrow x + 0 = 3$$



$$x + 9 = 15 \Rightarrow x + 9 - 9 = 15 - 9 \Rightarrow x + 0 = 6$$

$$6 + x = 13 \Rightarrow 6 + x - 6 = 13 - 6 \Rightarrow 0 + x - 0 = 7$$

$$x - 3 = 9 \Rightarrow x - 3 + 3 = 9 + 3 \Rightarrow x - 0 = 12 \Rightarrow x = 12$$

$$x - 8 = 7 \Rightarrow x + 8 - 8 = 7 + 8 \Rightarrow x - 0 = 15 \Rightarrow x = 15$$

$$6.3 = -2 + x \Rightarrow 6.3 + 2 = -2 + 2 + x \Rightarrow 6.3 = 0 + x \Rightarrow x = 6.3$$

$$5 = x - 8 \Rightarrow 5 + 8 = x - 8 + 8 \Rightarrow 13 = x - 0 \Rightarrow x = 13$$

$$\begin{aligned} 5 - x = 12 &\Rightarrow 5 - 12 - x + x = 12 - 12 + x \Rightarrow \\ &\Leftrightarrow -7 = 0 + x \Rightarrow x = -7 \end{aligned}$$

$$\begin{aligned} -8 = 5 - x &\Rightarrow -8 + 8 = 5 + 8 - x \Rightarrow \\ &\Leftrightarrow 0 + x = 13 - x + x \Rightarrow x = 13 \end{aligned}$$

$$3x = 12 \Rightarrow \frac{3x}{3} = \frac{12}{3} \Rightarrow \frac{3}{1}x = 4 \Rightarrow x = 4$$



$$7x = 14 \Rightarrow \frac{7x}{7} = \frac{14}{7} \Rightarrow \frac{x}{1} = 2 \Rightarrow x = 2$$

$$-6x = -30 \Rightarrow \frac{-6x}{-6} = \frac{-30}{-6} \Rightarrow \frac{-x}{-1} = 5 \Rightarrow x = 5$$

$$-8x = 48 \Rightarrow \frac{-8x}{-8} = \frac{48}{-8} \Rightarrow \frac{-x}{-1} = -6 \Rightarrow x = -6$$

$$7x = -56 \Rightarrow \frac{7x}{7} = \frac{-56}{7} \Rightarrow \frac{x}{1} = -8 \Rightarrow x = -8$$

$$-8x = -72 \Rightarrow \frac{-8x}{-8} = \frac{-72}{-8} \Rightarrow \frac{x}{1} = 9 \Rightarrow x = 9$$

I Coordinate Plane

Domain

- ➔ **Definition** \Rightarrow The set of all possible input values x for a function
- ➔ **Coordinate Plane** \Rightarrow Represents the horizontal axis (x-axis)
- ➔ **Example** \Rightarrow For $f(x) = \sqrt{x}$, the domain is $x \geq 0$
- ➔ **Relation** \Rightarrow Domain specifies where the function is defined
- ➔ **Outcome** \Rightarrow Determines the allowable values you can plug into the function



Range

- ➔ **Definition** \Rightarrow The set of all possible output values y from a function
- ➔ **Coordinate Plane** \Rightarrow Represents the vertical axis (y-axis)
- ➔ **Example** \Rightarrow For $f(x) = \sqrt{x}$, the range is $y \geq 0$
- ➔ **Relation** \Rightarrow Range shows the values the function can produce
- ➔ **Outcome** \Rightarrow Determines the spread of results plotted on the plane

Interval Notation

- ➔ **$[a, b]$** \Rightarrow Closed interval: includes both endpoints a and b
- ➔ **(a, b)** \Rightarrow Open interval: excludes both endpoints a and b
- ➔ **$[a, b)$** \Rightarrow Half-open interval: includes a but excludes b
- ➔ **$(a, b]$** \Rightarrow Half-open interval: excludes a but includes b
- ➔ **$[0, x)$** \Rightarrow All real numbers from 0 up to but not including x
- ➔ **$(0, x)$** \Rightarrow All real numbers greater than 0 and less than x

Relation of Ordered Pairs


- ➔ **Ordered Pair** \Rightarrow A pair written as (x, y) , where x is the input and y is the output
- ➔ **Relation** \Rightarrow Any set of ordered pairs that connects elements from one set (domain) to another (range)
- ➔ **Domain** \Rightarrow The collection of all first elements x in the ordered pairs
- ➔ **Range** \Rightarrow The collection of all second elements y in the ordered pairs
- ➔ **Function** \Rightarrow A special type of relation where each x is paired with exactly one y
- ➔ **Example** \Rightarrow Relation: $\{(1, 2), (2, 3), (3, 4)\}$; Domain: $\{1, 2, 3\}$; Range: $\{2, 3, 4\}$



Ordered Pairs

Input (a, b)	Output Value
$(0, 0)$	5
$(1, 0)$	9
$(1, 1)$	8
$(2, 1)$	15
$(2, 2)$	17
$(3, 1)$	26

Ordered Pairs





I Distance Formula

$$\text{2D points} \quad d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad (21)$$

$$\text{3D points} \quad d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2} \quad (22)$$

Distance Formula

- ➔ **Definition** ⇒ Gives the length of the line segment between two points in the plane
- ➔ **Points** ⇒ Two points (x_1, y_1) and (x_2, y_2)
- ➔ **Formula** ⇒ $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- ➔ **Origin** ⇒ Derived from the Pythagorean Theorem
- ➔ **Domain** ⇒ Applies to all real coordinates in 2D space
- ➔ **Extension** ⇒ Generalizes to 3D: $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$

Identity of $\sqrt[n]{a^n}$

- ➔ **Expression** ⇒ $\sqrt[n]{a^n}$
- ➔ **Meaning** ⇒ The n -th root of a^n
- ➔ **Simplification** ⇒ Cancels the root and exponent, yielding a
- ➔ **Condition** ⇒ Valid when $a \geq 0$ for real roots, or in complex domain otherwise
- ➔ **Example** ⇒ $\sqrt[3]{2^3} = \sqrt[3]{8} = 2$
- ➔ **Extension** ⇒ $\sqrt[n]{a^m} = a^{m/n}$

refer to I to see the sqrt function.

Distance Formula Functions

```

1  define nx_fma_dist1(x1, x2) {
2      return nx_nr_sqrt((x2 - x1)^2)
3  }
4
5  define nx_fma_dist2(x1, x2, y1, y2) {
6      return nx_nr_sqrt((x2 - x1)^2 + (y2 - y1)^2)
7  }
8
9  define nx_fma_dist3(x1, x2, y1, y2, z1, z2) {
10     return nx_nr_sqrt((x2 - x1)^2 + (y2 - y1)^2 + (z2 - z1)^2)

```





11

}


$$P_1(1, 2) \quad P_2(4, 6)$$

$$x_1 = 1 \quad x_2 = 4$$

$$y_1 = 2 \quad y_2 = 6$$

$$d = \sqrt{(4 - 1)^2 + (6 - 2)^2} \Rightarrow \sqrt{3^2 + 4^2} \Rightarrow \sqrt{9 + 16} \Rightarrow \sqrt{25} \Rightarrow 5$$

$$d = \sqrt{(4 - 1)^2 + (6 - 2)^2}$$






$$P_1(-3, 4) \quad P_2(9, -1)$$

$$x_1 = -3 \quad x_2 = 9$$

$$y_1 = 4 \quad y_2 = -1$$

$$d = \sqrt{(9 - (-3))^2 + ((-1) - 4)^2} \Rightarrow \sqrt{12^2 + -5^2} \Rightarrow \sqrt{144 + 25} \Rightarrow \sqrt{169} \Rightarrow 13$$

$$d = \sqrt{(9 - (-3))^2 + ((-1) - 4)^2}$$





II Sets

$$\mathbb{P} \subset \mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C} \subset \mathbb{H} \subset \mathbb{R}^n \quad \text{and branches into } \mathbb{F}_p$$

II Complex

Complex Numbers \mathbb{C}

- ➔ **Definition** \Rightarrow All numbers of the form $a + bi$, where $a, b \in \mathbb{R}$ and $i = \sqrt{-1}$
- ➔ **Examples** $\Rightarrow 3 + 2i, -1 - i, 0 + 4i$
- ➔ **Relation** \Rightarrow Contains all reals as the special case $b = 0$

II Finite Fields

Finite Fields \mathbb{F}_p

- ➔ **Definition** \Rightarrow A set of integers modulo a prime p , with addition and multiplication defined mod p
- ➔ **Examples** $\Rightarrow \mathbb{F}_2 = \{0, 1\}, \mathbb{F}_5 = \{0, 1, 2, 3, 4\}$
- ➔ **Relation** \Rightarrow Finite fields are algebraic systems distinct from \mathbb{R} , but essential in number theory
- ➔ **Use** \Rightarrow Cryptography, coding theory, error correction, and algebraic geometry

II Integers

Integers \mathbb{Z}

- ➔ **Definition** \Rightarrow All whole numbers, both positive and negative, including zero
- ➔ **Examples** $\Rightarrow \{-3, -2, -1, 0, 1, 2, 3, \dots\}$
- ➔ **Relation** \Rightarrow Every integer can be expressed as a rational $\frac{n}{1}$, so $\mathbb{Z} \subset \mathbb{Q}$



II Natural Numbers

Natural Numbers \mathbb{N}

- ➔ **Definition** \Rightarrow The counting numbers, starting from 1, sometimes including 0 depending on convention
- ➔ **Examples** $\Rightarrow \{1, 2, 3, 4, \dots\}$ or $\{0, 1, 2, 3, \dots\}$
- ➔ **Relation** $\Rightarrow \mathbb{N} \subset \mathbb{Z}$, since naturals are a subset of the integers
- ➔ **Use** \Rightarrow Foundation for counting, arithmetic, and building larger number sets

II Primes

Primes \mathbb{P}

- ➔ **Definition** \Rightarrow Natural numbers greater than 1 that have no divisors other than 1 and themselves
- ➔ **Examples** $\Rightarrow \{2, 3, 5, 7, 11, 13, \dots\}$
- ➔ **Relation** $\Rightarrow \mathbb{P} \subset \mathbb{N}$, primes are a special subset of the naturals
- ➔ **Use** \Rightarrow Foundation of number theory, factorization, and cryptography

II Quaternions

Quaternions \mathbb{H}

- ➔ **Definition** \Rightarrow Numbers of the form $a + bi + cj + dk$, where $a, b, c, d \in \mathbb{R}$ and $i^2 = j^2 = k^2 = ijk = -1$
- ➔ **Examples** $\Rightarrow 1 + 2i + 3j + 4k, -i + j$
- ➔ **Relation** $\Rightarrow \mathbb{C} \subset \mathbb{H}$, since complex numbers are a special case with $c = d = 0$
- ➔ **Use** \Rightarrow Applied in 3D rotations, computer graphics, and physics for representing orientation

II Rationals

Rationals \mathbb{Q}

- ➔ **Definition** \Rightarrow Numbers expressible as a fraction $\frac{p}{q}$ with integers p, q and $q \neq 0$
- ➔ **Examples** $\Rightarrow \frac{2}{3}, -\frac{5}{4}, 7 = \frac{7}{1}$
- ➔ **Relation** \Rightarrow All rationals are contained in the reals, so $\mathbb{Q} \subset \mathbb{R}$



II Reals

Reals \mathbb{R}

- ➔ **Definition** \Rightarrow All rationals plus irrationals (non-repeating, non-terminating decimals)
- ➔ **Examples** $\Rightarrow \pi, \sqrt{2}, 0.333\dots$
- ➔ **Relation** \Rightarrow Every real is a complex number with imaginary part 0, so $\mathbb{R} \subset \mathbb{C}$

II Vector Spaces

Vector Spaces \mathbb{R}^n

- ➔ **Definition** \Rightarrow Ordered tuples of real numbers, representing points or vectors in n -dimensional space
- ➔ **Examples** $\Rightarrow \mathbb{R}^2 = \{(x, y) \mid x, y \in \mathbb{R}\}, \mathbb{R}^3 = \{(x, y, z)\}$
- ➔ **Relation** \Rightarrow Built on \mathbb{R} , extending reals into higher dimensions
- ➔ **Use** \Rightarrow Geometry, physics, linear algebra, and computer graphics for modeling multidimensional systems

III Definitions

III Pairs

Factorial ($n!$)

Concept	Factorial ($n!$)
Definition	The product of all positive integers up to n . Defined as $n! = n \times (n - 1) \times \dots \times 1$.
Core Idea	Factorial counts permutations and combinations — it grows extremely fast.
Example	$5! = 120$.
Applications	Used in combinatorics, probability, and series expansions.
Pair	Inverse gamma function (not elementary).

**Logarithm Base 2 (\log_2)**

Concept	Logarithm Base 2 ($\log_2(x)$)
Definition	The inverse of the power of 2. Defined as the exponent y such that $2^y = x$.
Core Idea	$\log_2(x)$ measures how many times you multiply 2 to reach x .
Example	$\log_2(8) = 3$.
Applications	Widely used in computer science, information theory, and binary systems.
Pair	Power of 2 function (2^x).

Natural Logarithm (\ln)

Concept	Natural Logarithm ($\ln(x)$)
Definition	The inverse of the exponential function. Defined as the power to which e must be raised to equal x .
Core Idea	$\ln(x)$ undoes exponentiation with base e .
Example	$\ln(e^3) = 3$.
Applications	Used in calculus, growth/decay models, and solving exponential equations.
Pair	Exponential function (e^x).

Logarithm Base 2 (\log_2)

Concept	Logarithm Base 2 ($\log_2(x)$)
Definition	The inverse of the power of 2. Defined as the exponent y such that $2^y = x$.
Core Idea	$\log_2(x)$ measures how many times you multiply 2 to reach x .
Example	$\log_2(8) = 3$.
Applications	Widely used in computer science, information theory, and binary systems.
Pair	Power of 2 function (2^x).



Natural Logarithm (ln)

Concept	Natural Logarithm ($\ln(x)$)
Definition	The inverse of the exponential function. Defined as the power to which e must be raised to equal x .
Core Idea	$\ln(x)$ undoes exponentiation with base e .
Example	$\ln(e^3) = 3$.
Applications	Used in calculus, growth/decay models, and solving exponential equations.
Pair	Exponential function (e^x).

Tangent and Cotangent

Concept	Tangent ($\tan(x)$) and Cotangent ($\cot(x)$)
Definition	$\tan(x) = \frac{\sin(x)}{\cos(x)}$, while $\cot(x) = \frac{\cos(x)}{\sin(x)}$. They are reciprocals: $\cot(x) = \frac{1}{\tan(x)}$.
Core Idea	Tangent measures slope (rise/run). Cotangent flips that slope (run/rise).
Example	At 45° , $\tan(45^\circ) = 1$ and $\cot(45^\circ) = 1$.
Applications	Used in trigonometry, calculus, and geometry — especially for slope and angle analysis.
Pair	Reciprocal functions: $\tan(x) \leftrightarrow \cot(x)$.

Factorial (n!)

Concept	Factorial ($n!$)
Definition	The product of all positive integers up to n . Defined as $n! = n \times (n-1) \times \dots \times 1$.
Core Idea	Factorial counts permutations and combinations — it grows extremely fast.
Example	$5! = 120$.
Applications	Used in combinatorics, probability, and series expansions.
Pair	Inverse gamma function (not elementary).

**Factorial ($n!$)**

Concept	Factorial ($n!$)
Definition	The product of all positive integers up to n . Defined as $n! = n \times (n - 1) \times \dots \times 1$.
Core Idea	Factorial counts permutations and combinations — it grows extremely fast.
Example	$5! = 120$.
Applications	Used in combinatorics, probability, and series expansions.
Pair	Inverse gamma function (not elementary).

III Notation**III Delta in Mathematics****Finite Difference**

- ➔ **Definition** \Rightarrow Delta denotes change or difference between two values
- ➔ **Formula** $\Rightarrow \Delta x = x_{\text{final}} - x_{\text{initial}}$
- ➔ **Usage** \Rightarrow Discrete counterpart to derivative: $\Delta y / \Delta x \approx dy/dx$
- ➔ **Example** $\Rightarrow \Delta y = f(x_2) - f(x_1)$
- ➔ **Extension** \Rightarrow Forms the basis of finite difference methods in numerical analysis

Discriminant

- ➔ **Definition** \Rightarrow In quadratic equations, Δ denotes the discriminant
- ➔ **Formula** $\Rightarrow \Delta = b^2 - 4ac$
- ➔ **Usage** \Rightarrow Determines the nature of roots of $ax^2 + bx + c = 0$
- ➔ **Example** $\Rightarrow \Delta > 0$: two real roots; $\Delta = 0$: one real root; $\Delta < 0$: complex roots
- ➔ **Extension** \Rightarrow Generalized discriminants exist for higher-degree polynomials



Triangle Symbol

- ➔ **Definition** \Rightarrow Δ also denotes a triangle in geometry
- ➔ **Usage** $\Rightarrow \Delta ABC$ means triangle with vertices A, B, C
- ➔ **Connection** \Rightarrow Links algebraic glyph with geometric figure lineage
- ➔ **Extension** \Rightarrow Area of a triangle often denoted by Δ

III Theta in Mathematics

Angle Representation

- ➔ **Definition** \Rightarrow Greek letter θ used to denote an angle
- ➔ **Trigonometry** \Rightarrow Appears in sine, cosine, tangent: e.g. $\sin(\theta)$, $\cos(\theta)$, $\tan(\theta)$
- ➔ **Geometry** \Rightarrow Used in right triangles to relate sides and angles
- ➔ **Polar Coordinates** \Rightarrow Point (r, θ) defined by radius and angle
- ➔ **Example** \Rightarrow In a right triangle, $\cos(\theta) = \frac{\text{adjacent}}{\text{hypotenuse}}$

Other Mathematical Uses

- ➔ **Statistics** \Rightarrow θ often denotes parameters in probability distributions
- ➔ **Complex Numbers** \Rightarrow Angle of rotation in Euler's formula: $e^{i\theta} = \cos(\theta) + i \sin(\theta)$
- ➔ **Calculus** \Rightarrow Variable of integration in polar coordinates
- ➔ **Numerical Value** \Rightarrow In Greek numerals, θ has value 9
- ➔ **Connection** \Rightarrow Symbol of rotation, periodicity, and parameterization across math

III Epsilon in Mathematics

Infinitesimal Bound

- ➔ **Definition** \Rightarrow Greek letter ε used to denote a very small positive quantity
- ➔ **Limit Definition** \Rightarrow Appears in ε - δ proofs: for every $\varepsilon > 0$, there exists $\delta > 0$
- ➔ **Usage** \Rightarrow Measures closeness of a function to a limit
- ➔ **Example** $\Rightarrow |f(x) - L| < \varepsilon$ whenever $|x - c| < \delta$
- ➔ **Connection** \Rightarrow Core of rigorous calculus and real analysis



Error and Approximation

- ➔ **Definition** $\Rightarrow \varepsilon$ often denotes error tolerance or margin
- ➔ **Numerical Analysis** \Rightarrow Used to bound approximation error
- ➔ **Example** \Rightarrow If $|x - x_0| < \varepsilon$, then x is within tolerance
- ➔ **Connection** \Rightarrow Links exact mathematics with numerical computation

Other Uses

- ➔ **Set Theory** $\Rightarrow \varepsilon$ sometimes used as membership symbol, though \in is standard
- ➔ **Complexity** $\Rightarrow \varepsilon$ denotes arbitrarily small constants in algorithm analysis
- ➔ **Probability** $\Rightarrow \varepsilon$ used in inequalities like Chebyshev's or ε -nets
- ➔ **Connection** \Rightarrow Universal glyph for “smallness” across math disciplines

III Delta in Mathematics

Infinitesimal Change

- ➔ **Definition** \Rightarrow Greek letter δ denotes a very small positive quantity
- ➔ **Limit Proofs** \Rightarrow Appears in ε - δ definitions of limits
- ➔ **Formula** \Rightarrow For every $\varepsilon > 0$, there exists $\delta > 0$
- ➔ **Usage** \Rightarrow Controls how close x must be to c for $f(x)$ to be within ε of L
- ➔ **Example** \Rightarrow If $|x - c| < \delta$, then $|f(x) - L| < \varepsilon$
- ➔ **Connection** $\Rightarrow \delta$ measures input closeness, ε measures output closeness

Variation and Error

- ➔ **Definition** $\Rightarrow \delta$ often denotes small variation or tolerance
- ➔ **Numerical Analysis** \Rightarrow Used to bound input error
- ➔ **Example** \Rightarrow If $|x - x_0| < \delta$, then x is within tolerance of x_0
- ➔ **Connection** \Rightarrow Pairs with ε to formalize precision in analysis and computation



Other Mathematical Uses

- ➔ **Geometry** \Rightarrow δ sometimes used for small angles
- ➔ **Statistics** \Rightarrow δ may denote deviation or perturbation
- ➔ **Complexity** \Rightarrow δ used for small constants in algorithm analysis
- ➔ **Connection** \Rightarrow Universal glyph for “small input change” across math disciplines

III Proportional Symbol

Definition

- ➔ **Glyph** \Rightarrow \propto resembles the left half of ∞
- ➔ **Meaning** \Rightarrow Denotes proportionality between two quantities
- ➔ **Formula** \Rightarrow $y \propto x$ means $y = kx$ for some constant k
- ➔ **Usage** \Rightarrow Used in algebra, physics, and statistics to show direct proportionality
- ➔ **Example** \Rightarrow Gravitational force: $F \propto \frac{1}{r^2}$

Key Properties

- ➔ **Constant of Proportionality** \Rightarrow Always exists: $y = kx$
- ➔ **Direct Proportionality** \Rightarrow If one doubles, the other doubles
- ➔ **Inverse Proportionality** \Rightarrow Written as $y \propto \frac{1}{x}$
- ➔ **Scaling** \Rightarrow Proportionality preserves ratios
- ➔ **Connection** \Rightarrow Symbol bridges ratios, scaling laws, and functional dependence

III Infinity in Mathematics

Concept of Infinity

- ➔ **Definition** \Rightarrow ∞ denotes an unbounded quantity, larger than any real number
- ➔ **Calculus** \Rightarrow Appears in limits: $\lim_{x \rightarrow \infty} f(x)$
- ➔ **Set Theory** \Rightarrow Represents cardinalities of infinite sets
- ➔ **Geometry** \Rightarrow Used to mark points at infinity in projective geometry
- ➔ **Connection** \Rightarrow Symbol of endlessness, beyond finite measurement



Infinity in Calculus

- ➔ **Improper Integrals** $\Rightarrow \int_1^\infty \frac{1}{x^2} dx$
- ➔ **Limits** $\Rightarrow \lim_{x \rightarrow \infty} \frac{1}{x} = 0$
- ➔ **Series** \Rightarrow Infinite sums: $\sum_{n=1}^\infty \frac{1}{n^2}$
- ➔ **Connection** $\Rightarrow \infty$ marks the boundary of convergence and divergence
- ➔ **Example** \Rightarrow Harmonic series diverges: $\sum_{n=1}^\infty \frac{1}{n}$

Infinity in Set Theory

- ➔ **Countable Infinity** \Rightarrow Size of natural numbers, denoted \aleph_0
- ➔ **Uncountable Infinity** \Rightarrow Size of real numbers, larger than \aleph_0
- ➔ **Comparison** \Rightarrow Not all infinities are equal
- ➔ **Connection** $\Rightarrow \infty$ as a concept differs from cardinal numbers
- ➔ **Example** $\Rightarrow |\mathbb{N}| = \aleph_0$, but $|\mathbb{R}| > \aleph_0$

III Perpendicular Symbol

Definition

- ➔ **Glyph** $\Rightarrow \perp$ is the mathematical symbol for perpendicularity
- ➔ **Meaning** \Rightarrow Two lines, segments, or planes meet at a right angle (90°)
- ➔ **Notation** $\Rightarrow AB \perp CD$ means line AB is perpendicular to line CD
- ➔ **Geometry** \Rightarrow Used to denote orthogonality in Euclidean space
- ➔ **Example** \Rightarrow In a square, adjacent sides are \perp to each other

Linear Algebra Connection

- ➔ **Orthogonality** $\Rightarrow \perp$ denotes vectors with dot product zero
- ➔ **Formula** $\Rightarrow \vec{u} \perp \vec{v} \iff \vec{u} \cdot \vec{v} = 0$
- ➔ **Usage** \Rightarrow Defines orthogonal bases and projections
- ➔ **Example** $\Rightarrow (1, 0) \perp (0, 1)$ in \mathbb{R}^2
- ➔ **Extension** \Rightarrow Orthogonality generalizes perpendicularity to higher dimensions





Other Uses

- ➔ **Logic** \Rightarrow \perp sometimes denotes contradiction or falsity
- ➔ **Probability** \Rightarrow \perp used to denote independence in some texts
- ➔ **Connection** \Rightarrow Symbol bridges geometry, algebra, and logic
- ➔ **Visual** \Rightarrow Always evokes the right-angle lineage

III Plus-Minus and Minus-Plus

Plus-Minus (\pm)

- ➔ **Definition** \Rightarrow Symbol \pm means “plus or minus”
- ➔ **Usage** \Rightarrow Represents two possible values: $a + b$ or $a - b$
- ➔ **Example** \Rightarrow Quadratic formula: $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
- ➔ **Connection** \Rightarrow Encodes duality in solutions, symmetry in expansions
- ➔ **Extension** \Rightarrow Used in error bounds and approximations: $x \pm \varepsilon$

Minus-Plus (\mp)

- ➔ **Definition** \Rightarrow Symbol \mp means “minus or plus,” paired with \pm
- ➔ **Usage** \Rightarrow Ensures opposite choice when \pm is used earlier
- ➔ **Example** \Rightarrow If first term is $+$, second term takes $-$; if first is $-$, second takes $+$
- ➔ **Connection** \Rightarrow Keeps expressions consistent in paired signs
- ➔ **Extension** \Rightarrow Common in trigonometric identities and vector formulas

Combined Expression

- ➔ **Notation** $\Rightarrow a \pm b \mp c$
- ➔ **Meaning** \Rightarrow Two cases: $a + b - c$ or $a - b + c$
- ➔ **Pattern** \Rightarrow \pm and \mp always paired to flip signs consistently
- ➔ **Example** \Rightarrow In trig: $\sin(x \pm y) = \sin x \cos y \pm \cos x \sin y$
- ➔ **Connection** \Rightarrow Encodes dual solutions in compact symbolic form



III Parallel Symbol

Definition

- ➔ **Glyph** \Rightarrow \parallel is the mathematical symbol for parallelism
- ➔ **Meaning** \Rightarrow Two lines, segments, or planes never intersect and remain equidistant
- ➔ **Notation** $\Rightarrow AB \parallel CD$ means line AB is parallel to line CD
- ➔ **Geometry** \Rightarrow Used in Euclidean geometry to denote parallel lines and planes
- ➔ **Example** \Rightarrow In a rectangle, opposite sides are \parallel to each other

Linear Algebra Connection

- ➔ **Vectors** \Rightarrow \parallel denotes vectors that are scalar multiples of each other
- ➔ **Formula** $\Rightarrow \vec{u} \parallel \vec{v} \iff \vec{u} = k\vec{v}$
- ➔ **Usage** \Rightarrow Defines direction equivalence in vector spaces
- ➔ **Example** $\Rightarrow (2, 4) \parallel (1, 2)$ since $(2, 4) = 2(1, 2)$
- ➔ **Extension** \Rightarrow Parallelism generalizes beyond geometry into linear algebra and physics

Other Mathematical Uses

- ➔ **Analysis** $\Rightarrow \|x\|$ sometimes denotes norm of a vector
- ➔ **Logic** $\Rightarrow \parallel$ used in some texts for “parallel execution” or independence
- ➔ **Connection** \Rightarrow Symbol bridges geometry, algebra, and analysis
- ➔ **Visual** \Rightarrow Always evokes equidistant, non-intersecting lineage

III Angle Symbol

Definition

- ➔ **Glyph** $\Rightarrow \angle$ is the mathematical symbol for an angle
- ➔ **Meaning** \Rightarrow Represents the measure of rotation between two intersecting lines or rays
- ➔ **Notation** $\Rightarrow \angle ABC$ means the angle formed at vertex B by rays BA and BC
- ➔ **Units** \Rightarrow Measured in degrees ($^\circ$) or radians
- ➔ **Example** $\Rightarrow \angle ABC = 90^\circ$ denotes a right angle





Geometry Usage

- ➔ Right Angle $\Rightarrow \angle = 90^\circ$
- ➔ Acute Angle $\Rightarrow \angle < 90^\circ$
- ➔ Obtuse Angle $\Rightarrow 90^\circ < \angle < 180^\circ$
- ➔ Straight Angle $\Rightarrow \angle = 180^\circ$
- ➔ Reflex Angle $\Rightarrow 180^\circ < \angle < 360^\circ$

Other Mathematical Uses

- ➔ Trigonometry $\Rightarrow \angle$ used inside sine, cosine, tangent functions
- ➔ Polar Coordinates \Rightarrow Point (r, θ) defined by radius and angle
- ➔ Complex Numbers \Rightarrow Angle defines argument of a complex number
- ➔ Vector Analysis \Rightarrow Angle between vectors via dot product
- ➔ Connection \Rightarrow Symbol bridges geometry, trigonometry, and analysis

III Similar vs Equivalent

Similar Symbol (\sim)

- ➔ Glyph $\Rightarrow \sim$ is the symbol for similarity
- ➔ Geometry $\Rightarrow \triangle ABC \sim \triangle DEF$ means triangles have equal angles and proportional sides
- ➔ Algebra \Rightarrow Sometimes used to denote asymptotic equivalence: $f(x) \sim g(x)$ as $x \rightarrow \infty$
- ➔ Pattern \Rightarrow Similarity preserves shape but not necessarily size
- ➔ Example $\Rightarrow \triangle ABC \sim \triangle DEF$ if $\angle A = \angle D, \angle B = \angle E, \angle C = \angle F$
- ➔ Connection \Rightarrow Symbol of proportionality in geometry and asymptotics in analysis



Equivalent Symbol (\equiv)

- **Glyph** \Rightarrow \equiv is the symbol for equivalence
- **Congruence** $\Rightarrow \triangle ABC \equiv \triangle DEF$ means triangles are identical in shape and size
- **Number Theory** \Rightarrow Used for modular congruence: $a \equiv b \pmod{n}$
- **Logic** \Rightarrow Denotes logical equivalence: $P \equiv Q$
- **Pattern** \Rightarrow Equivalence preserves both shape and size, or exact relation
- **Example** $\Rightarrow 17 \equiv 5 \pmod{12}$
- **Connection** \Rightarrow Symbol of exact sameness across geometry, algebra, and logic

III Approximate Symbol

Definition

- **Glyph** $\Rightarrow \approx$ is the mathematical symbol for approximation
- **Meaning** \Rightarrow Indicates two values are close but not exactly equal
- **Notation** $\Rightarrow a \approx b$ means a is approximately equal to b
- **Usage** \Rightarrow Common in numerical analysis, applied math, and physics
- **Example** $\Rightarrow \pi \approx 3.1416$

Contexts of Use

- **Numerical** \Rightarrow Used when rounding or truncating decimals
- **Physics** \Rightarrow Marks measured values close to theoretical ones
- **Statistics** \Rightarrow Denotes approximate probabilities or estimates
- **Analysis** \Rightarrow Signals asymptotic closeness in limits
- **Connection** \Rightarrow Symbol bridges exact math with practical computation



Related Symbols

- ➔ Equal Sign \Rightarrow $=$ denotes exact equality
- ➔ Tilde \Rightarrow \sim denotes similarity or asymptotic equivalence
- ➔ Congruence \Rightarrow \equiv denotes exact equivalence or modular congruence
- ➔ Approx \Rightarrow \approx specifically signals numerical closeness
- ➔ Extension \Rightarrow Each glyph stages a different level of sameness

III Similar or Equal Symbol

Definition

- ➔ Glyph \Rightarrow \simeq is the symbol for “similar or equal”
- ➔ Meaning \Rightarrow Indicates two quantities are nearly equal and share structural similarity
- ➔ Usage \Rightarrow Common in analysis, approximation, and asymptotic notation
- ➔ Example \Rightarrow $f(x) \simeq g(x)$ means functions are close in value and form
- ➔ Connection \Rightarrow Bridges similarity (\sim) and equality ($=$)

Contexts of Use

- ➔ Approximation \Rightarrow Used when values are not exactly equal but very close
- ➔ Asymptotics \Rightarrow Signals functions behave similarly as $x \rightarrow \infty$
- ➔ Geometry \Rightarrow Sometimes used to denote figures nearly congruent
- ➔ Physics \Rightarrow Marks quantities equal within experimental tolerance
- ➔ Extension \Rightarrow \simeq is less strict than \equiv (equivalent) but stronger than \approx (approximate)



III Congruent Symbol

Definition

- ➔ **Glyph** \Rightarrow \cong is produced in LaTeX with `\cong`
- ➔ **Meaning** \Rightarrow Denotes congruence: “is congruent to”
- ➔ **Geometry** $\Rightarrow \triangle ABC \cong \triangle DEF$ means triangles are identical in shape and size
- ➔ **Pattern** \Rightarrow Congruence preserves both angles and side lengths
- ➔ **Example** \Rightarrow \triangle with sides 3,4,5 is \cong to another 3,4,5 triangle
- ➔ **Connection** \Rightarrow Stronger than similarity (\sim), exact match without scaling

Other Mathematical Uses

- ➔ **Number Theory** \Rightarrow Sometimes used interchangeably with $=$ for modular congruence
- ➔ **Analysis** \Rightarrow Can denote “is approximately congruent” in some texts
- ➔ **Logic** \Rightarrow Rarely used for structural equivalence
- ➔ **Extension** \Rightarrow \cong bridges geometry congruence with algebraic congruence
- ➔ **Visual** \Rightarrow Glyph resembles equality with a tilde, staging sameness plus shape relation

III Limit Symbol (lim)

Definition

- ➔ **Glyph** \Rightarrow \lim denotes the limit of a function or sequence
- ➔ **Meaning** \Rightarrow Describes the value a function approaches as the input approaches some point
- ➔ **Notation** $\Rightarrow \lim_{x \rightarrow c} f(x)$
- ➔ **Example** $\Rightarrow \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$
- ➔ **Connection** \Rightarrow Core concept in calculus, analysis, and continuity



Types of Limits

- ➔ **Finite Limit** \Rightarrow Function approaches a finite value as input approaches a point
- ➔ **Infinite Limit** \Rightarrow Function grows without bound: $\lim_{x \rightarrow 0^+} \frac{1}{x} = \infty$
- ➔ **One-Sided Limit** $\Rightarrow \lim_{x \rightarrow c^-} f(x)$ or $\lim_{x \rightarrow c^+} f(x)$
- ➔ **At Infinity** $\Rightarrow \lim_{x \rightarrow \infty} f(x)$
- ➔ **Sequence Limit** $\Rightarrow \lim_{n \rightarrow \infty} a_n$

Formal Definition

- ➔ **ϵ - δ Definition** \Rightarrow For every $\epsilon > 0$, there exists $\delta > 0$
- ➔ **Condition** \Rightarrow If $|x - c| < \delta$, then $|f(x) - L| < \epsilon$
- ➔ **Meaning** $\Rightarrow f(x)$ gets arbitrarily close to L as x approaches c
- ➔ **Connection** \Rightarrow Defines continuity and rigor in calculus
- ➔ **Example** $\Rightarrow \lim_{x \rightarrow 2} (3x + 1) = 7$

Limits

- ➔ **Usage** \Rightarrow Appears in limit notation: $\lim_{x \rightarrow c} f(x)$
- ➔ **Meaning** \Rightarrow “to” means the variable approaches a value
- ➔ **Example** $\Rightarrow \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$
- ➔ **Connection** \Rightarrow Glyph of approach, not exact arrival
- ➔ **Extension** \Rightarrow Also used in one-sided limits: $x \rightarrow c^+$, $x \rightarrow c^-$

Mappings

- ➔ **Usage** \Rightarrow Appears in function notation: $f : A \rightarrow B$
- ➔ **Meaning** \Rightarrow “to” denotes mapping from domain to codomain
- ➔ **Example** $\Rightarrow f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$
- ➔ **Connection** \Rightarrow Glyph of transformation, linking sets
- ➔ **Extension** \Rightarrow Used in category theory: arrows $A \rightarrow B$



Ranges and Intervals

- ➔ Usage ⇒ Appears in describing ranges: “from ... to ...”
- ➔ Meaning ⇒ Marks boundaries of intervals or sums
- ➔ Example ⇒ $\sum_{i=1}^n a_i$ means i runs from 1 to n
- ➔ Connection ⇒ Glyph of span, marking start and end
- ➔ Extension ⇒ Used in integrals: $\int_a^b f(x) dx$