


# Lemma 1 Proof/Example

Canyon Foot

October 18, 2019

For binary random variables  $X, Y, S$ , we wish to construct two causal graphs,  $G_1, G_2 = G_1 \setminus \{Y \rightarrow S\}$  with corresponding probability functions  $P_1, P_2$  such that  $P_2(y = 1|x = 1) = P_1(y = 1|x = 1, s = 1)$  but  $P_2(y = 1|x = 1) \neq P_1(y = 1|x = 1)$ . This demonstrates that the 'selection biased' conditional distribution  $P_1(y = 1|x = 1, s = 1)$  is consistent with a conditional  $P_2(y = 1|x = 1)$  that is different from the 'true' conditional  $P_1(y = 1|x = 1)$ . This is a violation of the definition of  $s$ -recoverability, therefore giving the result that  $G_1$  is unrecoverable.



We begin by using the definition of conditional probability to expand  $P_1(y = 1|x = 1, s = 1)$  as:

$$P_1(y = 1|x = 1, s = 1) = \frac{P_1(y = 1, x = 1, s = 1)}{P_1(x = 1, s = 1)}$$

Now, the definition of Markov factorization property for an SCM gives that  $P_1(y = 1, x = 1, s = 1) = P_1(s = 1|y = 1)p(y = 1|x = 1)$ . Then, expanding the denominator using the law of total probability, we have:

$$P_1(y = 1|x = 1, s = 1) = \frac{P_1(s = 1|y = 1)p(y = 1|x = 1)}{P_1(s = 1|y = 1)p(y = 1|x = 1) + P_1(s = 1|y = 0)p(y = 1|x = 1)}$$

Given this formulation, we can assign explicit values to our probabilities. Let  $P_1(y = 1|x = 1) = \frac{1}{2}$ , therefore giving  $P_1(y = 0|x = 1) = \frac{1}{2}$ . Additionally, let  $P_1(s = 1|y = 1) = \frac{3}{4}$  and  $P_1(s = 1|y = 0) = \frac{1}{4}$ . This means that we are more likely to 'sample' pairs  $(x, y)$  when  $y = 1$  than when  $y = 0$ . Then

$$P_1(y = 1|x = 1, s = 1) = \frac{\frac{3}{4} \frac{1}{2}}{\frac{3}{4} \frac{1}{2} + \frac{1}{4} \frac{1}{2}} = \frac{3}{4}.$$

By our earlier condition, we make  $P_2(y = 1|x = 1) = P_1(y = 1|x = 1, s = 1) = \frac{3}{4}$ , and here we get the result since  $P_2(y = 1|x = 1) \neq P_1(y = 1|x = 1) = \frac{1}{2}$ .