Solution for HW5 20161102

Taper

November 3, 2016

Abstract

陈鸿翔(11310075)

1 Uncertainty of electron velocity

2 Proof

Proof. Assuming an orthonormal basis labeled by n: $|n\rangle$. Using Einstein Summation convention, one finds:

$$\begin{split} \operatorname{Tr}(AB) &= \langle n|AB|n\rangle = \langle n|A|m\rangle \, \langle m|B|n\rangle = \langle m|B|n\rangle \, \langle n|A|m\rangle \\ &= \langle m|BA|m\rangle = \operatorname{Tr}(BA) \end{split}$$

Hence

$$\operatorname{Tr}(XYZ)=\operatorname{Tr}((XY)Z)=\operatorname{Tr}(Z(XY))=\operatorname{Tr}((ZX)Y)=\operatorname{Tr}(YZX)$$

3 Proof

Proof.

$$\langle [A, B] \rangle = \langle AB - BA \rangle = \langle AB \rangle - \langle BA \rangle = \langle AB \rangle - \langle AB \rangle^* = 2i \operatorname{Im}(\langle AB \rangle)$$

Hence it is imaginary or zero. Similar, by replacing the - sign above with + sign, one easily finds:

$$\langle \{A, B\} \rangle = 2 \operatorname{Re}(\langle AB \rangle)$$

So it is real. $\hfill\Box$

4 Diagonalization

A is real and symmetric, hence it is diagonalizable:

$$\det \begin{pmatrix} 1 - \lambda & 1 & 3 \\ 1 & 5 - \lambda & 1 \\ 3 & 1 & 1 - \lambda \end{pmatrix} = \det \begin{pmatrix} 1 - \lambda & 1 & 3 \\ 1 & 5 - \lambda & 1 \\ 0 & -14 + 3\lambda & -\lambda - 2 \end{pmatrix}$$
$$= (1 - \lambda) [(5 - \lambda)(-\lambda - 2) + 14 - 3\lambda] - (-10\lambda + 4) = -\lambda^3 + 7\lambda^2 - 36$$

The roots are $\lambda_1 = -2$, $\lambda_2 = 3$, $\lambda_3 = 6$. For $\lambda = -2$, we have

$$\left(\begin{array}{ccc} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{array}\right) \left(\begin{array}{c} x \\ y \\ z \end{array}\right) = -2 \left(\begin{array}{c} x \\ y \\ z \end{array}\right)$$

Or

$$\left(\begin{array}{ccc} 3 & 1 & 3 \\ 1 & 7 & 1 \\ 3 & 1 & 3 \end{array}\right) \left(\begin{array}{c} x \\ y \\ z \end{array}\right) = 0$$

Or

$$\left(\begin{array}{ccc} 0 & -20 & 0\\ 1 & 7 & 1\\ 0 & 0 & 0 \end{array}\right) \left(\begin{array}{c} x\\ y\\ z \end{array}\right) = 0$$

Obviously the corresponding eigenvector is $\alpha \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$, where α is any

nonzero complex number. The case for $\lambda=3$ and $\lambda=6$ can be similar solved by examing the following two equations:

$$\begin{pmatrix} -2 & 1 & 3 \\ 1 & 2 & 1 \\ 3 & 1 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 0, \quad \begin{pmatrix} -5 & 1 & 3 \\ 1 & -1 & 1 \\ 3 & 1 & -5 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 0$$

and the result is summarized as