

ECE770-T14/QIC 885: Quantum Electronics & Photonics

Problem Set 1, Winter 2014, Instructor: A. Hamed Majedi

Problem 1- Consider two-slit experiment with sources of monoenergetic electrons where detectors are placed along the vertical screen, e.g. x direction, to monitor the diffraction pattern of electrons.

When only one slit is open, the amplitude of detected electrons can be written as

$$\Psi_1(x, t) = A_1 \frac{e^{-i(kx - \omega t)}}{1 + x^2} \text{ and when only the other one is open, the amplitude is } \Psi_2(x, t) = A_2 \frac{e^{-i(kx - \omega t + \pi x)}}{1 + x^2}.$$

A_1 and A_2 are constants to be found.

Calculate the density of the detected electrons on the screen when

- a) both slits are open and a light source is used to determine which of the slits the electron went through.
- b) both slits are open and no light source is used.

Plot the intensity registered on the screen as a function of x for both cases in parts a) and b).

Problem 2- Consider a simplest wave function that potentially describes a surface bound state of the form of $\Psi(z, t) = Ae^{-\alpha|z|}e^{-i\omega t}$

where A , α and ω are positive real constants.

- a) Find A .
- b) Find the uncertainty in z .
- c) Find the momentum wave function associated with $\Psi(z, t = 0)$.
- d) If $\Psi(z, t = 0)$ is the initial wave function of a free electron, construct $\Psi(z, t)$, in the form of integral.
- e) Discuss the limiting cases (α very large and α very small).

Problem 3- A free object is located at $z = a$ at $t = 0$ and its wavefunction is given by $\Psi(z, t = 0) = \delta(z - a)$.

Find the wavefunction for $t > 0$, namely $\Psi(z, t)$. This solution is called the **free object propagator**.

Hint: To evaluate an integral of the form $\int_{-\infty}^{\infty} \exp[i\lambda z^2 + i\beta z]dz$, pretend that λ is $\lambda + i\epsilon$ with $\epsilon > 0$

so that the integral is convergent. Then complete the square in the exponent of the exponential and change variables. The $i\epsilon$ in your answer will allow you to decide whether to take the positive or negative square root. Finally let $i\epsilon$ goes to zero. This technique is called *regularization*.

Problem 4- Consider an object with mass m in the one-dimensional short range Dirac Delta function potential, $V(z) = V_o\delta(z)$.

- a) Find the criterion for V_o for having a bound state.
- b) What is the binding energy?
- c) Find the value of z_o such that the probability of finding the object within $|z| < z_o$ is equal to 0.9.

Problem 5- Consider an object of mass m that is constrained to move in a circle of radius a in the $x - y$ plane.

Determine the quantized energy levels for the following cases:

- a) The motion of the object is nonrelativistic with a constant linear momentum.
- b) The motion of the object is relativistic, namely $E^2 = |\mathbf{p}|^2c^2 + m_o^2c^4$, where m_o is the rest mass and c is the speed of light.

Due: Wednesday Jan 29, 2014. (before starting the class)