

Summary on Classical Mechanics

Yuyang Songsheng

September 21, 2016

1 Lagrangian Formulation

$$S = \int_{t_1}^{t_2} L(q_i, \dot{q}_i, t) dt, \quad \delta q_i(t_1) = \delta q_i(t_2) = 0$$

$$\delta S = 0 \rightarrow \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_i} \right) - \frac{\partial L}{\partial q_i} = 0$$

1.1 Example

The form of Lagrangian for a system of particles in inertial frame:

$$L = \sum_a \frac{1}{2} m_a v_a^2 - U(\vec{r}_1, \vec{r}_2, \dots,)$$

To get the form of Lagrangian for a system of interacting particles, we must assume:

- (1) Space and time are homogeneous and isotropic in inertial frame;
- (2) Galileo's relativity principle and Galilean transformation;
- (3) Spontaneous interaction between particles;

2 Symmetry and Conservation Laws(1)

2.1 Nother's theorem

For $q_i \rightarrow q_i + \delta q_i$ and $L \rightarrow L + \delta L$, if $\delta L = \frac{df(q, \dot{q}, t)}{dt}$, then we get

$$\frac{d}{dt} (p^i \delta q_i - f) = 0 \quad (p^i = \frac{\partial L}{\partial \dot{q}_i})$$

2.2 Homogeneity of time

If $\frac{\partial L}{\partial t} = 0$, then we get

$$\frac{dH}{dt} = 0 \quad (H = \sum_i \dot{q}_i p^i - L)$$

3 Hamilton formulation

$$p^i = \frac{\partial L}{\partial \dot{q}_i}$$

$$H(q, p, t) = \sum_i p^i \dot{q}_i - L$$

$$\dot{p}^i = -\frac{\partial H}{\partial q_i} \quad \dot{q}_i = \frac{\partial H}{\partial p^i}$$

3.1 Poisson Brackets

First, we assume the bracket operation has the following properties:

$$[f, g] = -[g, f]$$

$$[\alpha_1 f_1 + \alpha_2 f_2, \beta_1 g_1 + \beta_2 g_2] = \alpha_1 \beta_1 [f_1, g_1] + \alpha_1 \beta_2 [f_1, g_2] + \alpha_2 \beta_1 [f_2, g_1] + \alpha_2 \beta_2 [f_2, g_2]$$

$$[f_1 f_2, g_1 g_2] = f_1 [f_2, g_1] g_2 + f_1 g_1 [f_2, g_2] + g_1 [f_1, g_2] f_2 + [f_1, f_2] g_2 f_2$$

$$[f, [g, h]] + [g, [h, f]] + [h, [f, g]] = 0$$

Here, f, g, h are functions of p^i, q_i, t . Then, if we assume that

$$[q_i, p^k] = \delta_i^k$$

we can deduce that

$$[f, g] = \sum_k \left(\frac{\partial f}{\partial q_k} \frac{\partial g}{\partial p^k} - \frac{\partial f}{\partial p^k} \frac{\partial g}{\partial q_k} \right)$$

The Hamilton equation can be written as

$$\dot{p}^i = [p^i, H] \quad \dot{q}_i = [q_i, H] \tag{1}$$

And we can also derive that $\frac{df}{dt} = [f, H]$.

4 Symmetry and Conservation Laws(2)

Suppose g is a function of p and q . If the transformation of q and p can be described as

$$q \rightarrow q + \epsilon[q, g]$$

$$p \rightarrow p + \epsilon[p, g]$$

We can prove that

$$H \rightarrow H + \epsilon[H, g]$$

So if H is invariant under the transformation, then $[H, g] = 0$, that means $\frac{dg}{dt} = 0$, i.e. g is a conserved quantity of the motion.

5 Hamilton-Jacobi equation

We define

$$S(q, t) = \left(\int_{q_0, t_0}^{q, t} L dt \right) |_{extremum}$$

We can prove that

$$p = \frac{\partial S}{\partial q}, \quad H = -\frac{\partial S}{\partial t}$$

So, we have

$$-\frac{\partial S}{\partial t} = H(q, \frac{\partial S}{\partial q})$$

This is called Hamiltonian-Jacobi equation.

6 Symmetry and Conservation Laws(3)

If S is invariant under transformation $q_i \rightarrow q_i + \delta q_i$, then

$$\delta S = \left(\sum_i p^i \delta q_i \right) |_{q_0, t_0}^{q, t} = 0$$

So, we have

$$\frac{d}{dt}(p^i \delta q_i) = 0$$

Further more, if

$$\delta S = \left(\sum_i p^i \delta q_i \right) |_{q_0, t_0}^{q, t} = f(q_i, \dot{q}_i, t) |_{q_0, t_0}^{q, t}$$

we will have conserved quantity

$$\frac{d}{dt}(p^i \delta q_i - f) = 0$$