Worksheet 1: Quantum mechanical approaches: Hückel approximation and DFT methods

April Cooper, Patrick Kreissl und Sebastian Weber

April 28, 2013 University of Stuttgart

Contents

1	Computational Task: DFT calculations with Siesta						
	1.1	Geometry optimization of adenine	2				
	1.2	Theoretical Prediction of Watson-Crick Hydrogen-bond length in adenine-					
		thymine base pair	3				

1 Computational Task: DFT calculations with Siesta

1.1 Geometry optimization of adenine

Distance	Here	Ref [20]	Ref [7]	Exp	Angle	Here	Ref [20]	Ref [7]	Exp
C2-N3	•	•	•	•	•	•	•	•	•
N1-C2	•	•	•	•	•	•	•	•	•
C6-N1	•	•	•	•	•	•	•	•	•
C5-C6	•	•	•	•	•	•	•	•	•
C4-C5	•	•	•	•	•	•	•	•	•
N3-C4	•	•	•	•	•	•	•	•	•

1.2 Theoretical Prediction of Watson-Crick Hydrogen-bond length in adenine-thymine base pair




Figure 1: Results for two flavours of XC functional. Blue numbers show bond lengths and red ones show bond angles.

Angle in Å	LDA	GGA	Literature ¹
O4-H61-N6	176.7	175.6	175.8
N1-H3-N3	178.1	179.4	178.1

Table 1: Bond angles

Distance in Å	LDA	GGA	Experiment 1 ¹	Experiment 2 ¹
O4-N6	2.68	2.89	2.95	2.93
N1-N3	2.60	2.79	2.82	2.85

Table 2: Bond lengths

TODO: Write some text. Write down number of steps.

¹source: Attachment1b.pdf