

Worksheet 5: Monte-Carlo

April Cooper, Patrick Kreissl und Sebastian Weber

January 22, 2013

University of Stuttgart

Contents

1	Simple Sampling- Intrgration	2
2	Importance Sampling- Metropolis-Hastings-Algorithm	4

1 Simple Sampling- Intrgration

First we were asked to program a function `runge(x)` that computes the runge function $f(x) = \frac{1}{1+x^2}$ and to plot it on the interval $[-5, 5]$.




Figure 1: Runge function $f(x)$ on the interval $[-5, 5]$

Then we had to write a function that computes the exact integral of the runge function. The exact result is:

$$\int_{-5}^5 \frac{1}{1+x^2} = \arctan(5) - \arctan(-5) \approx 2.7468$$

After that we were asked to program a function `simple_sampling(f,a,b,N)` that performs N steps of a simple sampling Monte-Carlo integration and to use this function to compute the Integral for $N = 2^i$ with $2 \leq i \leq 20$. Furthermore we were asked to determine the actual and statistical error and to plot them against the number of integration steps N . In the following are two plots of the error from two different runs.





Figure 2: Statistical and actual error of the Integration with the simple sampling method

As it can be seen due to the randomness of the points in which f is evaluated the actual error behaves quite different from run to run. Furthermore it also varies for small numbers of integration steps quite a lot from the statistical error. In contrast to this the statistical error has a much more comparable behavior in different runs.

2 Importance Sampling- Metropolis-Hastings-Algorithm

In this task we were asked to implement the Metropolis-Hastings-algorithm and to use that function to generate a given number of samples for several Δx being used in the trial move. The resulting histograms can be found in the following plots




Figure 3: Distribution histograms of the Metropolis samples with 100 bins compared to the normalized runge function (red)

By eye it is hard to say whether $dx = 1.0$ or $dx = 10.0$ is the better choice. Having a look at the acceptance rates - $\approx 84.8\%$ for $dx = 1.0$, $\approx 32.5\%$ for $dx = 10.0$ - and taking into account that it was stated on the worksheet that an acceptance rate of about 30% is optimal, it can be said that $dx = 10.0$ is a better choice than $dx = 1.0$.