

Quantum Physics 1

Notes-4

The wavefunction and its interpretation

Probability current

The wave packet

Space and momentum representations

Heisenberg Uncertainty Principle

Important results so far

Einstein– deBroglie relations: $p = \hbar k = \frac{h}{\lambda}$; $E = \hbar\omega = hf$

The probability interpretation: $P(x)dx = \Psi^*\Psi dx$

$$\langle q \rangle = \int \Psi^* q(x) \Psi dx$$

The standard deviation of a distribution is:

$$\sigma = \sqrt{\langle x^2 \rangle - \langle x \rangle^2}$$

The general form for a wave packet using waves with

well-defined momentum: $\Psi(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} A(k) e^{i(kx - \omega t)} dk$

where $A(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Psi(x, 0) e^{-ikx} dx$ and $P(k)dk = A^* A dk$

Class 4 In-Class Activity

- a) Assume that the wavefunction of a square pulse at time zero is

$$\Psi(x, 0) = \begin{cases} 0 & \text{for } x < -a \\ \frac{1}{\sqrt{2a}} & \text{for } -a < x < a \\ 0 & \text{for } x > a \end{cases}$$

- b) Find the standard deviation $\sqrt{\langle x^2 \rangle - \langle x \rangle^2}$ of this distribution.
- c) Find $A(k)$
- d) Find the standard deviation of $A(k)$.
- e) Does this wavefunction obey the Heisenberg Uncertainty Principle?

Conservation of Probability

It is important that the total probability of finding a particle does not change with time.

Is this consistent with the Schrodinger equation?

- First calculate the time derivative of the probability density:

$$\frac{\partial P}{\partial t} = \frac{\partial \Psi^* \Psi}{\partial t} = \Psi^* \frac{\partial \Psi}{\partial t} + \Psi \frac{\partial \Psi^*}{\partial t} \quad (1)$$

- And from the SE:

$$\frac{\partial \Psi}{\partial t} = \frac{1}{i\hbar} \left(-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x) \Psi \right) \text{ and } \frac{\partial \Psi^*}{\partial t} = \frac{-1}{i\hbar} \left(-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi^*}{\partial x^2} + V(x) \Psi^* \right)$$

and substituting space derivatives for time derivatives in (1):

$$\begin{aligned} \frac{\partial P}{\partial t} &= \Psi^* \frac{1}{i\hbar} \left(-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x) \Psi \right) - \Psi \frac{1}{i\hbar} \left(-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi^*}{\partial x^2} + V(x) \Psi^* \right) \\ \frac{\partial P}{\partial t} &= \frac{i\hbar}{2m} \left(\Psi^* \frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} - \Psi \frac{\hbar^2}{2m} \frac{\partial^2 \Psi^*}{\partial x^2} \right) = \frac{i\hbar}{2m} \frac{\partial}{\partial x} \left(\Psi^* \frac{\partial \Psi}{\partial x} - \Psi \frac{\partial \Psi^*}{\partial x} \right) \end{aligned}$$

Conservation of probability

From the previous page: $\frac{\partial P}{\partial t} = \frac{i\hbar}{2m} \frac{\partial}{\partial x} \left(\Psi^* \frac{\partial \Psi}{\partial x} - \Psi \frac{\partial \Psi^*}{\partial x} \right)$

So the rate of change of the total probability is:

$$\frac{\partial}{\partial t} \int P dx = \int \frac{\partial P}{\partial t} dx = \frac{i\hbar}{2m} \int_{-\infty}^{\infty} \frac{\partial}{\partial x} \left(\Psi^* \frac{\partial \Psi}{\partial x} - \Psi \frac{\partial \Psi^*}{\partial x} \right) dx$$

$$\frac{\partial}{\partial t} \int P dx = \frac{i\hbar}{2m} \left(\Psi^* \frac{\partial \Psi}{\partial x} - \Psi \frac{\partial \Psi^*}{\partial x} \right) \Big|_{-\infty}^{\infty} = 0$$

The final step is because in order to be normalizable the wavefunction must go to zero at $\pm\infty$.

THUS, IF THE WAVEFUNCTION IS A SOLUTION TO THE SE AND IS NORMALIZABLE, THEN PROBABILITY IS CONSERVED.

Probability Current

- The probability current density

$$-\frac{i\hbar}{2m} \left(\Psi^* \frac{\partial \Psi}{\partial x} - \Psi \frac{\partial \Psi^*}{\partial x} \right) \equiv j_x(x)$$

is a useful concept that helps in thinking about traveling particles.

Probability Current for a Pure Momentum Function

$$\begin{aligned}\Psi_{p_0}(x, t) &= A e^{i(p_0 x - Et)/\hbar} \\ j_x(x) &= -\frac{i\hbar}{2m} \left(\Psi^* \frac{\partial \Psi}{\partial x} - \Psi \frac{\partial \Psi^*}{\partial x} \right) \\ &= -\frac{i\hbar}{2m} \left(\frac{A^* A i p_0}{\hbar} - \left(-\frac{A^* A i p_0}{\hbar} \right) \right) \\ &= \frac{|A|^2 p_0}{m} = |A|^2 v_0 \\ &\quad \text{(Makes sense.)}\end{aligned}$$

Note that $\frac{\partial}{\partial t} P([a, b], t) = j_x(a) - j_x(b) = 0$ for this case.

In class activity (not to be handed in)

- Evaluate the probability current for the wave function $\Psi = Ae^{ikx+i\omega t} + Be^{-ikx+i\omega t}$

Particles and waves: the Gaussian wavepacket

We will simplify calculations by assuming that the spatial part of the wavefunction can be described at $t = 0$ by a Normal Gaussian function centered at x_0 and moving with average wavenumber k_0 ;

$$\Psi(x, 0) = \left(\frac{1}{\sigma_x \sqrt{2\pi}} \right)^{1/2} e^{ik_0 x} e^{-(x-x_0)^2/4\sigma_x^2}.$$


(We have already solved a similar problem in Notes 2.)

$$\text{So } A(k) = \frac{1}{\sqrt{2\pi}} \left(\frac{1}{\sigma_x \sqrt{2\pi}} \right)^{1/2} \int_{-\infty}^{\infty} e^{i(k_0-k)x} e^{-(x-x_0)^2/4\sigma_x^2} dx$$

Gaussian wavepacket continued

to simplify further let's take $x_0 = 0$,

$$\text{so } A(k) = \frac{1}{\sqrt{2\pi}} \left(\frac{1}{\sigma_x \sqrt{\pi}} \right)^{1/2} \int_{-\infty}^{\infty} e^{i(k_0 - k)x} e^{-x^2/2\sigma_x^2} dx$$



Then letting $k' = k - k_0$, we solve by completing the square:

$$-ik'x - \frac{1}{2\sigma_x^2} x^2 + \sigma_x^2 k'^2 - \sigma_x^2 k'^2 = \left(\frac{1}{\sqrt{2}\sigma_x} x - i\sigma_x k' \right)^2 + \sigma_x^2 k'^2$$

$$A(k) = \frac{1}{\sqrt{2\pi}} \left(\frac{1}{\sigma_x \sqrt{\pi}} \right)^{1/2} e^{-\sigma_x^2 k'^2} \int_{-\infty}^{\infty} e^{-\left(\frac{1}{\sqrt{2}\sigma_x} x - i\sigma_x k' \right)^2} dx$$

$$\text{let } x' = \frac{x}{\sqrt{2}\sigma_x} - i\sigma_x k', \text{ then } A(k) = \frac{1}{\sqrt{2\pi}} \left(\frac{1}{\sigma_x \sqrt{\pi}} \right)^{1/2} \sqrt{2}\sigma_x e^{-\sigma_x^2 k'^2} \int_{-\infty}^{\infty} e^{-x'^2} dx'$$

$$A(k) = \frac{1}{\sqrt{2\pi}} \left(\frac{1}{\sigma_x \sqrt{\pi}} \right)^{1/2} \sqrt{2}\sigma_x e^{-\sigma_x^2 k'^2} \sqrt{\pi} = \left(\frac{1}{\pi} \sigma_x^2 \right)^{\frac{1}{4}} e^{-\frac{\sigma_x^2 (k - k_0)^2}{2}}$$


Gaussian wavepacket continued

$$A(k) = \left(\frac{1}{\pi} \sigma_x^2 \right)^{\frac{1}{4}} e^{-\frac{\sigma_x^2 (k-k_0)^2}{2}}$$

This is a Normal Gaussian k – distribution,
centered at k_0 with width $\sigma_k = \frac{1}{\sigma_x}$.

- Although we specified a wave momentum k_0 we find that the wavefunction actually contains a spread of wave momenta, $\frac{1}{\sigma_x}$, which increases as the spatial Gaussian narrows. Remember that $\Delta k = \frac{\Delta p}{\hbar}$.
- Waves of differing k move at different speeds, so the original wavefunction will spread with time.

Inversely correlated widths



Note that if Ψ is properly normalized ($\int \Psi^* \Psi dx = 1$)

then the wavevector amplitude function is also normalized to 1:

$$\int |A(k)|^2 dk = 1$$

An aside on the definition of the width of a Gaussian




Figure 4.8 A cornucopia of definitions of the width w_x of a function $f(x)$. The function at hand is $f(x) = e^{-x^2}$. (a) The full-width-at-half-maximum, $w_x = 1.665$. (b) The extent of the function at the special point where $f(x) = [f(x)]_{\max}/e$. The function e^{-x^2} is equal to $1/e$ of its maximum value at $x = \pm 1$, so this definition gives $w_x = 1.738$. (c) My definition: the standard deviation of x ; for this function, $w_x = \Delta x = 0.560$.

The width of a Gaussian can be defined in various ways. Two conventional ways are by the Full Width at Half Maximum (FWHM) and twice the standard deviation. Morrison uses $\Delta L = \text{standard deviation of the position}$.

Time evolution of the wavefunction

The general form of the wavefunction is:

$$\Psi(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} A(k) e^{-i(kx - \omega t)} dk.$$

This means that if you are given $\Psi(x, 0)$ you can find $\Psi(x, t)$.

$$A(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Psi(x, 0) e^{-i(kx)} dx.$$

To do the top integral, you must know $\omega(k)$.

The simplest case is for a pulse of light, $\omega = kc$

$$\Psi(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} A(k) e^{-ik(x - ct)} dk.$$

which produces a travelling pulse that does not change shape as a function of time.

Group velocity for a wave-packet

$$\psi(x, t) = \int A(k) e^{-i(kx - \omega(k)t)} dk$$

Assuming that $A(k)$ has a fairly narrow range of important k 's.
we will allow ourselves to expand $\omega(k)$:

$$\omega(k) \cong \omega(k_0) + (k - k_0) \left(\frac{d\omega}{dk} \right)_{k_0} + \dots$$

Letting $k' = k - k_0$

$$\psi(x, t) \cong e^{i(k_0 x - \omega(k_0)t)} \int dk' e^{-ik' \left[x - \left\{ \frac{d\omega}{dk} \right\} t \right]_0}$$

Therefore in this simple expansion, the group velocity is just

$$v_g = \frac{d\omega}{dk}$$

Momentum Probability Amplitude

Note that $A(k)$ clearly contains information about momentum because $k = p/\hbar$.

It is useful (later) to be able to describe the state function directly in terms of momentum states.

$$\Phi(p) \equiv \frac{1}{\sqrt{\hbar}} A\left(\frac{p}{\hbar}\right) \text{ (the momentum probability amplitude)}$$

Then we can rewrite the wavefunction as;

$$\Psi(x, 0) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \Phi(p) e^{ipx/\hbar} dp \text{ and}$$

$$\Phi(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \Psi(x, 0) e^{-ipx/\hbar} dx.$$

Note that the factor of $1/\sqrt{\hbar}$ is required for proper normalization of Ψ and Φ .

Interpretation of the space and momentum state functions

$\Psi(x, t)$ = the wavefunction and directly gives information about the probability of finding a particle at a specific position, but it is difficult to directly calculate properties like the expected momentum because momentum is not written as a function of position.

$\Phi(p)$ = the momentum probability amplitude and directly gives information about the probability of finding a particle with a specific momentum.

They both describe the same quantum state.

If the members of an ensemble are in the quantum state $\Psi(x, t)$ with Fourier transform $\Phi(p) = F[\Psi(x, 0)]$ then

$$P(p)dp = \Phi^*(p)\Phi(p)dp$$

is the probability that in a momentum measurement at time t a particle's momentum will be found to be between p and $p + dp$.

Expectation values

Given the momentum amplitude function and the associated probability distribution, we can compute expectation values:

$$\langle p \rangle = \int_{-\infty}^{\infty} \Phi(p) * p \Phi(p) dp = \text{expectation value (average) of } p.$$

$$\Delta p = \text{momentum uncertainty} = [(p - \langle p \rangle)^2]^{1/2} = [\langle p^2 \rangle - \langle p \rangle^2]^{1/2}$$

$$\langle p^2 \rangle = \int_{-\infty}^{\infty} \Phi(p) * p^2 \Phi(p) dp$$

TABLE 4.1 POSITION AND MOMENTUM INFORMATION FOR A GAUSSIAN STATE FUNCTION WITH $L = 1.0$, $x_0 = 0$, AND $p_0 = 0$

Position	Momentum
$\Psi(x, 0) = \left(\frac{1}{2\pi L^2} \right)^{1/4} e^{-x^2/(4L^2)}$ $\langle x \rangle = 0$ $\Delta x = L$	$\Phi(p) = \left(\frac{2 L^2}{\pi \hbar^2} \right)^{1/4} e^{-p^2 L^2 / \hbar^2}$ $\langle p \rangle = 0$ $\Delta p = \frac{1}{2L} \hbar$