


i-Tree Canopy v7.0


Cover Assessment and Tree Benefits Report

Estimated using random sampling statistics on 11/29/2020



Google

Land Cover



Abbr.	Cover Class	Description	Points	% Cover ± SE	Area (ac) ± SE
H	Grass/Herbaceous		40	26.67 ± 3.61	0.66 ± 0.09
IB	Impervious Buildings		0	0.00 ± 0.00	0.00 ± 0.00
IO	Impervious Other		0	0.00 ± 0.00	0.00 ± 0.00
IR	Impervious Road		34	22.67 ± 3.42	0.56 ± 0.08
S	Soil/Bare Ground		0	0.00 ± 0.00	0.00 ± 0.00
T	Tree/Shrub		76	50.67 ± 4.08	1.25 ± 0.10
W	Water		0	0.00 ± 0.00	0.00 ± 0.00
Total			150	100.00	2.47

Tree Benefit Estimates: Carbon (English units)

Description	Carbon (T)	±SE	CO ₂ Equiv. (T)	±SE	Value (USD)	±SE
Sequestered annually in trees	1.71	±0.14	6.27	±0.50	\$146	±12
Stored in trees (Note: this benefit is not an annual rate)	42.91	±3.46	157.34	±12.68	\$3,659	±295

Currency is in USD and rounded. Standard errors of removal and benefit amounts are based on standard errors of sampled and classified points. Amount sequestered is based on 1.365 T of Carbon, or 5.005 T of CO₂, per ac/yr and rounded. Amount stored is based on 34.281 T of Carbon, or 125.697 T of CO₂, per ac and rounded. Value (USD) is based on \$85.28/T of Carbon, or \$23.26/T of CO₂ and rounded. (English units: T = tons (2,000 pounds), ac = acres)

Tree Benefit Estimates: Air Pollution (English units)

Abbr.	Description	Amount (oz)	±SE	Value (USD)	±SE
CO	Carbon Monoxide removed annually	15.89	±1.28	\$1	±0
NO2	Nitrogen Dioxide removed annually	126.91	±10.22	\$1	±0
O3	Ozone removed annually	775.22	±62.46	\$69	±6
SO2	Sulfur Dioxide removed annually	45.56	±3.67	\$0	±0
PM10*	Particulate Matter greater than 2.5 microns and less than 10 microns removed annually	266.24	±21.45	\$52	±4
PM2.5	Particulate Matter less than 2.5 microns removed annually	60.00	±4.83	\$274	±22
Total		1,289.83	±103.92	\$397	±32

Currency is in USD and rounded. Standard errors of removal and benefit amounts are based on standard errors of sampled and classified points. Air Pollution Estimates are based on these values in oz/ac/yr @ \$/oz/yr and rounded:
CO 12.697 @ \$0.04 | NO2 101.387 @ \$0.01 | O3 619.311 @ \$0.09 | SO2 36.400 @ \$0.00 | PM10* 212.699 @ \$0.20 | PM2.5 47.932 @ \$4.56 (English units: oz = ounces, ac = acres)

Tree Benefit Estimates: Hydrological (English units)

Abbr.	Benefit	Amount (Kgal)	±SE	Value (USD)	±SE
AVRO	Avoided Runoff	1.18	±0.09	\$11	±1
E	Evaporation	4.68	±0.38	N/A	N/A
I	Interception	4.71	±0.38	N/A	N/A
T	Transpiration	2.77	±0.22	N/A	N/A
PE	Potential Evaporation	15.34	±1.24	N/A	N/A
PET	Potential Evapotranspiration	13.01	±1.05	N/A	N/A

Currency is in USD and rounded. Standard errors of removal and benefit amounts are based on standard errors of sampled and classified points. Hydrological Estimates are based on these values in Kgal/ac/yr @ \$/Kgal/yr and rounded:
AVRO 0.941 @ \$8.94 | E 3.738 @ N/A | I 3.762 @ N/A | T 2.216 @ N/A | PE 12.255 @ N/A | PET 10.394 @ N/A (English units: Kgal = thousands of gallons, ac = acres)

About i-Tree Canopy

The concept and prototype of this program were developed by David J. Nowak, Jeffery T. Walton, and Eric J. Greenfield (USDA Forest Service). The current version of this program was developed and adapted to i-Tree by David Ellingsworth, Mike Binkley, and Scott Maco (The Davey Tree Expert Company)

Limitations of i-Tree Canopy

The accuracy of the analysis depends upon the ability of the user to correctly classify each point into its correct class. As the number of points increase, the precision of the estimate will increase as the standard error of the estimate will decrease. If too few points are classified, the standard error will be too high to have any real certainty of the estimate.



Use of this tool indicates acceptance of the [EULA](#).