Every Elliptic Curve over a CM Number Field Is Modular

First version 7 October 2026
Attribution revision 9 October 2026

Abstract

We prove that every elliptic curve over a CM number field is modular. The proof combines the covering construction of OpenAI with a mod-5 automorphy lifting argument that retains a residually ramified Steinberg place. Following a method of Taylor and Khare–Thorne, the local condition at this place excludes the exceptional cohomology classes for $\mathrm{SL}_2(\mathbb{F}_5)$. Partial framing then supplies Taylor–Wiles presentations with the required dimension, while integral local–global compatibility imposes the same Steinberg condition on the Hecke-valued representations. A specialization of the covering family produces a modular auxiliary elliptic curve and three congruences, in characteristics p, 3, and p, connecting it to the original curve. The construction works over non-Galois CM fields and includes fields containing fifth roots of unity.

1 Introduction

Modularity identifies the Galois representations of an elliptic curve with automorphic representations of GL_2. Over a CM field, automorphy lifting provides a way to propagate this correspondence through congruences. The problem is to construct a suitable starting curve and a sequence of congruences that preserves the necessary local conditions.

The covering construction of OpenAI [O] supplies such a sequence. Its character spaces connect an auxiliary elliptic curve to a quadratic twist of a prescribed curve. We combine this construction with a lifting argument that also applies when the base field contains μ_5.

Theorem 1.1. *Every elliptic curve E over a CM number field K is modular. If E has no geometric complex multiplication, there is a regular algebraic cuspidal automorphic representation π_E of $\mathrm{GL}_2(\mathbb{A}_K)$, of weight zero and with trivial unitary central character, whose compatible system is $H^1(E)$. Its finite local parameters, in the cohomological normalization, agree with those of E. In particular, its elliptic-normalized local L-factors and conductor agree with those of E at every finite place.*

If E has geometric complex multiplication, the corresponding automorphic representation is obtained from its Hecke characters. It is cuspidal when the endomorphisms are not all defined over K; otherwise it is an isobaric sum of two characters.

The characteristic-five argument uses the exceptional-cohomology method of Taylor [Tay03, §1, condition E3 and Lemma 1.2], developed further by Khare–Thorne [KT, Introduction and Propositions 6.6 and 6.10]. We formulate the retained-Steinberg variant needed here in Theorem 4.2. Our auxiliary family supplies a place with nontrivial unipotent residual inertia. Retaining the Steinberg condition at this place excludes the exceptional classes from dual Selmer before Taylor–Wiles primes are chosen. Sections 3–4 give the local calculation and its integration into the CM-field patching argument.

The protected place belongs to an auxiliary curve, so the theorem imposes no restriction on the reduction of E. The geometric family has a split multiplicative degeneration of width three. Specializing near it makes the auxiliary discriminant valuation prime to 5. At the same time, specialization near a different boundary preserves the reduction categories needed in the congruence argument.

Here is the resulting chain. For a sufficiently large prime p, let η and ψ have orders 3 and p. Over $K(\mu_{3p})$, the compatible systems satisfy

$$V_\eta \equiv V_{\eta\psi} \pmod{p}, \qquad V_{\eta\psi} \equiv V_\psi \pmod{3}, \qquad V_\psi \equiv V_1 \pmod{p}.$$

The first system is the cohomology of the auxiliary curve; the last is that of a quadratic twist of E. The retained-Steinberg theorem makes the first system automorphic. Three applications of [CN, Theorem 5.2] give the last, and solvable descent returns to K.

Notation

A CM field is a totally imaginary quadratic extension of a totally real field. We write F^+ for the maximal totally real subfield of F and c for complex conjugation. Let G_F denote the absolute Galois group of F, ε_ℓ its ℓ-adic cyclotomic character, and q_v the cardinality of the residue field at a finite place v.

We use the cohomological normalization of [CN]: H^1 of an elliptic curve has determinant ε_ℓ^{-1} and labelled Hodge–Tate weights $\{0, 1\}$. Automorphic representations of weight zero are normalized unitarily. We write $r_\iota(\pi)$ for the associated Galois representation. Residual representations are semisimplified unless a lattice is specified.

A residual representation in characteristic ℓ is *decomposed generic* if there is a rational prime q, split completely in F, at whose places it is unramified and has Frobenius eigenvalue ratio outside $\{1, q, q^{-1}\}$. We will choose $q \equiv 1$ modulo ℓ, so distinct eigenvalues suffice. This agrees with [CN, Definition 2.1.27].

For a finite extension of $\mathbb{Q}_5$, let O be its ring of integers, ϖ a uniformizer, and k its residue field. All deformation problems below have fixed determinant ε_5^{-1}. Framed local rings use GL_2 frames, including scalar frame directions.

2 The geometric congruence construction

Let E/K be an elliptic curve without geometric complex multiplication, written $y^2 = h(x)$. Choose a prime $p > 100$ and put $n = 3p$. Over a nonempty open subset U of the parameter line, set

$$B_t : h(t)y^2 = h(x), \qquad R_t = (t, 1), \qquad S_t = [n]R_t.$$

The divisor $(S_t) - (-S_t) - n(R_t) + n(-R_t)$ is principal. Normalize its defining function f_t by $f_t(O) = 1$ and, for m dividing n, form

$$\operatorname{div}(f_t) = (S_t) - (-S_t) - n(R_t) + n(-R_t), \qquad C_m(t) : Y^m = f_t(Z).$$

The involution $w(Z, Y) = (-Z, Y^{-1})$ is defined over K. Put $A_t = \operatorname{Jac}(C_3(t)/\langle w\rangle)$. The construction, its compact-type degeneration, its integral homology, and its geometric monodromy are given in [O, Lemma 3.1, Proposition 3.3, Proposition 4.2, and Theorem 4.5].

Proposition 2.1. *Over $F = K(\mu_n)$, every rotation character θ defines a semisimple rank-two compatible system V_θ. It has determinant ε_ℓ^{-1}, labelled weights $\{0, 1\}$, and is potentially semistable above ℓ. Moreover, A_t is an elliptic curve and*

$$V_\eta \simeq H^1(A_t)|_{G_F}, \qquad V_1 \simeq H^1(B_t),$$

where η has order 3. The three congruences displayed in the introduction hold.

Proof. The integral calculation of [O, Proposition 4.2] makes $H^1(C_n, \mathbb{Z}_\ell)$ free of rank two over the rotation group ring. Thus every character space has dimension two. Cup product, composed with w in the second variable, gives an alternating perfect pairing on a character space with values in $\mathbb{Q}_\ell(-1)$, proving the determinant assertion. The involution exchanges θ and θ^{-1}, and Hodge decomposition gives one copy of each Hodge weight. Smooth proper geometry supplies finite ramification and potential semistability. Semisimplicity follows from Faltings [F], and the good-reduction trace formula supplies compatibility.

The trivial rotation space is $H^1(B_t)$. On C_3, the involution acts as -1 on this space and exchanges the two nontrivial character spaces. Its invariant space therefore has dimension two, identifying the quotient Jacobian with the stated endpoint.

For the congruences, choose O containing all character values and use the evaluation lattice

$$T_\theta = H^1(C_n, \mathbb{Z}_\ell) \otimes_{\mathbb{Z}_\ell[\mu_n],\theta} O.$$

It is free of rank two, and its reduction depends only on the reduced character. A p-power root of unity reduces to 1 in characteristic p; a third root reduces to 1 in characteristic 3. This proves the three congruences without dividing by n. □

Local conditions

Proposition 2.2. *At every place v_0 dividing $3p$ there is a nonempty open subset Ω_{v_0} of $U(K_{v_0})$ on which all four systems have the reduction category determined by E: potentially good ordinary, potentially good supersingular, or potentially multiplicative. In the multiplicative case, every rank-two system has monodromy of rank one, at every coefficient prime.*

Proof. Move the marked point toward O on the fixed curve E. The compact-type degeneration has Jacobian E^n [O, Proposition 3.3]. The local stability results of [O, §7] provide neighborhoods with the stated ordinary, supersingular, or totally toric behavior. Intersect the neighborhoods needed for the finitely many completions of F above v_0.

At a point outside the ramification locus of x, the local inverse function theorem gives an analytic section $(t, y(t))$, with $y(t)^2 = h(t)$. The map $(x, y) \mapsto (x, y(t)y)$ identifies B_t with E and identifies their normalized covering constructions. This gives Ω_{v_0}.

The ordinary filtration respects the rotations and induces the required filtration on each character space. In the supersingular case, all Newton slopes are $1/2$, excluding an ordinary weight-zero line after finite extension. In the totally toric case, the full monodromy has rank n. It is the direct sum of n rank-two nilpotent operators, each of rank at most one; all have rank one. □

Simultaneous specialization

Proposition 2.3. *The prime p and a parameter $t \in U(K)$ can be chosen so that:*

1. *all the local conditions of Proposition 2.2 hold;*
2. *the residual representations used in the $p, 3, p$ chain are absolutely irreducible over the corresponding cyclotomic fields and decomposed generic over F;*
3. *$A_t[5]$ is decomposed generic over K, and its image is $\mathrm{SL}_2(\mathbb{F}_5)$ when $\mu_5 \subset K$;*
4. *finitely many further nonempty local parameter opens, away from $3p$, may be prescribed.*

Proof. Write $d_K = [K : \mathbb{Q}]$ and let N be the normal closure of K. Serre's open-image theorem [S72] allows us to choose $p > 100$ so that every conjugate E^σ has image containing $\mathrm{SL}_2(\mathbb{F}_p)$ over N and

$$\frac{2d_K p}{p^2 - 1} < 1.$$

Put $D = N(\mu_{15p})$. Since D/N is abelian and $\mathrm{SL}_2(\mathbb{F}_p)$ is perfect, the image of each $E^\sigma[p]$ over D is exactly $\mathrm{SL}_2(\mathbb{F}_p)$. There are p^2 matrices of each trace 2 and -2 in this group. Hence the proportion with repeated eigenvalues is $2p/(p^2 - 1)$. In any subgroup of the product surjecting onto every factor, the union bound gives an element with distinct eigenvalues in every coordinate. Chebotarev produces rational primes split completely in D with that simultaneous property. Scalar quadratic twists preserve eigenvalue ratios, so this proves the required assertions for V_1 modulo p.

For the varying systems, consider $(m, \ell) = (3, p), (3, 5), (p, 3)$. The common-basis monodromy calculation [O, Theorem 4.5] supplies opposite unipotents on each nontrivial λ-character space:

$$T_a = \begin{pmatrix} 1 & sc_\lambda \\ 0 & 1 \end{pmatrix}, \qquad T_b = \begin{pmatrix} 1 & 0 \\ uc_\lambda & 1 \end{pmatrix}, \qquad c_\lambda = 2 - \lambda - \lambda^{-1}, \quad s, u \in \{1, -1\}.$$

For $m = 3$, the coefficient is 3. These matrices generate $\mathrm{SL}_2(\mathbb{F}_5)$ in characteristic 5. The word $T_a T_b^2$ has trace 20 or -16, distinct from ± 2 in characteristics 5 and $p > 100$. For $(m, \ell) = (p, 3)$, use $T_a T_b$. A repeated eigenvalue would force λ to satisfy a polynomial of degree at most four over $\mathbb{F}_3$. Then p divides $3^e - 1$ for some $e \le 4$, impossible for $p > 100$. Thus one word is regular semisimple on every nontrivial character space for each pair.

Let $W = \mathrm{Res}(K/\mathbb{Q})U$. It is an open subset of affine d_K-space and, over D, is the product of the conjugate parameter curves. Record the finite monodromy representations of all three pairs on all factors in a single finite quotient H of $\pi_1(W)$, taking the normal core of their common kernel. Include the quotient $\mathrm{Gal}(D/\mathbb{Q})$. A loop supported on one factor acts trivially on the others, so H contains the simultaneous regular-semisimple elements and the individual opposite unipotents.

Hilbert irreducibility with weak approximation [SG, Proposition 3.3.1 and Theorem 3.5.3], [O, Lemma 8.3] gives $x \in W(\mathbb{Q})$ whose specialization onto H is surjective and which meets every prescribed local open. Its corresponding parameter is a single $t \in U(K)$. Since the specialization respects $\mathrm{Gal}(D/\mathbb{Q})$, the image of G_D is precisely the kernel H_D.

The opposite unipotents in H_D have distinct fixed lines, proving absolute irreducibility over G_D. Since D contains $F(\mu_\ell)$ for each coefficient prime $\ell \in \{3, p\}$ in the congruence chain, this implies absolute irreducibility over the required cyclotomic fields. The simultaneous regular-semisimple elements, and their conjugacy classes in H, yield decomposed genericity by Chebotarev: they lie in H_D, so the resulting rational primes split completely in D, and hence have residue cardinality 1 modulo the relevant coefficient prime. The use of every conjugate factor ensures the property at every place above the chosen rational prime, both over F for the congruence chain and over K for $A_t[5]$.

Here ℓ does not divide m, so character projectors on C_m commute with reduction. Their rational identifications with the corresponding spaces on C_n give the desired semisimplified reductions, even when ℓ divides n. This is separate from the evaluation-lattice argument of Proposition 2.1. □

3 A Steinberg place in characteristic five

After the cyclotomic twist described below, the local deformation condition used in our fixed-determinant application is the residually ramified form of Taylor's condition E3 [Tay03, §1, condition E3]. The exclusion of finite-group cohomology classes is closely related to Taylor's Lemma 1.2 [Tay03] and the dual-Selmer argument of Khare–Thorne, Propositions 6.6 and 6.10 [KT]. We give the local calculation and resulting Taylor–Wiles prime selection explicitly in the form needed for the CM application.

The local and global settings differ. Taylor's proof of Lemma 1.2 first removes exceptional classes from ordinary Selmer using a place where the residual representation is unramified and the projective Frobenius has order five. Khare–Thorne work over totally real fields with exceptional projective S_5 image; their relevant auxiliary local condition has unramified residual inertia and nontrivial unipotent residual Frobenius. Here the field is CM and contains μ_5, the residual image is $\mathrm{SL}_2(\mathbb{F}_5)$, and the retained place has nontrivial unipotent residual inertia and trivial residual Frobenius. We use their method, while keeping the local and CM-field patching arguments in Sections 3–4.

For the comparison with Taylor's determinant convention, twisting by ε_5 changes our determinant ε_5^{-1} to ε_5; since the field contains μ_5, the residual twist is trivial. The projective image is A_5. The central subgroup of order two acts trivially on the adjoint module, and its order is invertible in characteristic five, so passage to the projective image does not change the relevant finite-group cohomology.

We now work in characteristic 5. The essential local condition is that inertia remain unipotent and residually nontrivial.

Lemma 3.1. *Let F_v have residue characteristic different from 5 and contain μ_5. Suppose that, for tame inertia and arithmetic Frobenius generators,*

$$\bar{\rho}(\tau) = U = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \qquad \bar{\rho}(\phi) = 1, \qquad \phi\tau\phi^{-1} = \tau^{q_v}.$$

Fix an unramified determinant ψ lifting $\det\bar\rho$. *The framed deformation condition in which every inertia element has characteristic polynomial* $(X-1)^2$ *is formally smooth:*

$$R_v^{\square,\mathrm{St},\psi} \simeq O[[a,u,z]].$$

For $M = \mathrm{ad}^0\,\bar\rho$, *put* $e = E_{12}$, $f = E_{21}$, *and* $h = \mathrm{diag}(1,-1)$. *Then*

$$H^1(F_v, M) = k[(f,0)] \oplus k[(0,e)],$$
$$\mathcal{L}_v = k[(0,e)], \qquad \dim_k \mathcal{L}_v = h^0(F_v, M) = 1.$$

A pair records the cocycle values on (τ,ϕ), *and* $\mathcal{L}_v$ *is the tangent space of the Steinberg condition.*

Proof. For $D = \mathrm{Ad}(U)$, direct calculation gives $D(e) = e$, $D(h) = h - 2e$, and $D(f) = f + h - e$. Hence $M^D = ke$, $(D-1)M = ke \oplus kh$, and $(D-1)^3 = 0$. In characteristic 5, the norm $1 + D + \cdots + D^4$ is zero. Since $q_v \equiv 1$ modulo 5, the sum of the first q_v powers of D is the identity.

Applying a cocycle to the tame relation gives $(D-1)c(\phi) = 0$. Coboundaries change $c(\tau)$ by $(D-1)M$ and leave $c(\phi)$ unchanged. Wild inertia and the prime-to-5 tame subgroup contribute no cohomology. This proves the description of H^1.

For the integral statement, let A be an Artinian coefficient algebra and put $N = \rho(\tau) - 1$. Cayley–Hamilton gives $N^2 = 0$. Its upper-right entry is a unit, so

$$N = \begin{pmatrix} a & b \\ -a^2/b & -a \end{pmatrix}, \qquad a \in \mathfrak{m}_A, \quad b \in 1 + \mathfrak{m}_A.$$

The strict basis change (Ne_2, e_2) makes $N = E_{12}$. The tame relation then forces

$$\rho(\phi) = \begin{pmatrix} q_v d & z \\ 0 & d \end{pmatrix}, \qquad d^2 = \psi(\phi)/q_v.$$

The square root congruent to 1 is unique. Setting $u = b - 1$, we obtain the unrestricted parameters a, u, z in the displayed presentation. The lifted wild and prime-to-5 inertial subgroups are trivial, since they map to the pro-5 kernel of reduction.

For a first-order cocycle $x = c(\tau)$, the trace condition is $\mathrm{tr}(xU) = 0$. It removes exactly the f-coordinate. The remaining inertia variation is a coboundary; after removing it, $c(\phi)$ is an arbitrary multiple of e. This proves the tangent assertion.

Finally, $\ker N = \mathrm{im}\, N = A(Ne_2)$ is a rank-one direct summand. It is the unique such summand on which inertia acts trivially. This kernel description applies over nonreduced coefficient algebras. □

Lemma 3.2. *Choose a primitive fifth root of unity to identify* $\mathbb{F}_5(1)$ *with* $\mathbb{F}_5$, *and extend scalars to* k. *This identifies* $M(1)$ *with* M. *Under local Tate duality,* $\mathcal{L}_v$ *is its own annihilator. A class with nonzero restriction to inertia therefore fails the dual Steinberg condition.*

Proof. The trace form on ad^0 is invariant, symmetric, and nondegenerate. Local Tate duality and graded commutativity give a nondegenerate alternating pairing on the two-dimensional $H^1(F_v, M)$. Every line is its own annihilator, and Lemma 3.1 identifies $\mathcal{L}_v$ with the unramified line. □

Excluding the exceptional classes

Let F contain μ_5, and let $\bar\rho : G_F \to \mathrm{SL}_2(\mathbb{F}_5)$ be surjective. Let S contain its ramification and the places above 5. Fix a nonempty set $P \subset S$ at which Lemma 3.1 applies. Put $M = \mathrm{ad}^0\,\bar\rho$. Let D_{Sel} be a subspace of $H^1(G_{F,S}, M)$ satisfying the dual Steinberg condition at P.

Lemma 3.3. *If* L/F *is cut out by* $\bar\rho$ *and* $C_N = F(\mu_{5^N})$, *then restriction to* G_{LC_N} *is injective on* D_{Sel}, *for every* $N \geq 1$.

Proof. The group $G = \mathrm{SL}_2(\mathbb{F}_5)$ is perfect, whereas C_N/F is abelian. Thus $L \cap C_N = F$. Since $M^G = 0$, inflation identifies

$$H^1(\mathrm{Gal}(LC_N/F), M) \simeq H^1(G, M).$$

The subgroup $\langle U \rangle$ has index 24. Restriction–corestriction therefore makes restriction to this Sylow 5-subgroup injective. The cyclic cohomology calculation gives

$$H^1(\langle U \rangle, M) = \ker(1 + D + \cdots + D^4)/(D - 1)M = k[f].$$

At a protected place, residual inertia maps onto a conjugate of $\langle U \rangle$, and inflation to inertia is injective. Every nonzero inflated finite-group class consequently has nonzero inertial restriction, contrary to Lemma 3.2. Inflation–restriction for G_{LC_N} now proves the claim. □

The following prime-selection argument is the retained-Steinberg version of the dual-Selmer strategy in Khare–Thorne [KT, Propositions 6.6 and 6.10]; we retain the proof to make the hypotheses used here explicit.

Lemma 3.4. *At every cyclotomic depth N, Taylor–Wiles primes can be chosen to kill the relative dual Selmer group for any framing set disjoint from P. They can be chosen of degree one over $\mathbb{Q}$ and outside any prescribed finite set.*

Proof. Let κ be a nonzero relative dual Selmer class. By Lemma 3.3 its restriction to G_{LC_N} is a nonzero homomorphism. The k-span of its image is G-stable, by the cocycle identity, and equals M because the adjoint representation of $\mathrm{SL}_2(\mathbb{F}_5)$ is absolutely irreducible.

Take $g = \mathrm{diag}(2, 3)$, and choose $\sigma \in G_{C_N}$ with residual image g. The coinvariants $M/(\mathrm{Ad}(g) - 1)M$ are the diagonal line. Multiplying σ by an element of G_{LC_N} changes $\kappa(\sigma)$ by a value whose possible k-span is M. We can therefore make its image in the coinvariants nonzero.

Chebotarev in the finite quotient recording κ, $\bar{\rho}$, and C_N gives a positive-density set of detecting primes w with $q_w \equiv 1$ modulo 5^N and distinct residual eigenvalues. Exclude also every prime above a rational prime ramified in F. Primes of degree greater than one over $\mathbb{Q}$ have density zero among the primes of F, so w may have degree one. Its rational residue characteristic is then unramified and has residue degree one in every imaginary quadratic subfield of F, and therefore splits in each of them.

These evaluation maps separate the dual Selmer group. Choose enough independent evaluations and add further primes to obtain a fixed sufficiently large cardinality r. At a Taylor–Wiles prime, the fixed-determinant local condition has the full two-dimensional tangent space: unramified and tame diagonal variations. Its dual condition is zero. An augmented dual Selmer class is therefore unramified at each added prime, belongs to the original group, and vanishes by the selected evaluations. □

4 Automorphy lifting with a retained Steinberg place

We formulate the lifting result in the prepared setting needed for the application. This keeps the local conditions and the patching calculation explicit.

Let F be CM with $\mu_5 \subset F$, and put $d = [F^+ : \mathbb{Q}]$. Choose a decomposition

$$S = S_5 \sqcup R \sqcup P, \qquad S_5 = S_5^{\mathrm{cr}} \sqcup S_5^{\mathrm{st}},$$
$$P \neq \varnothing, \qquad P^c \cap S = \varnothing.$$

All places of $R \cup P$ are prime to 5. Impose the following field conditions:

- Every rational prime below S or ramifying in F splits in an imaginary quadratic subfield of F.
- For each place $\bar{v}$ above 5 of F^+ there is a different place $\bar{v}'$ above 5 such that

$$\sum_{\substack{\bar{w} \mid 5 \\ \bar{w} \neq \bar{v}, \bar{v}'}} [F^+_{\bar{w}} : \mathbb{Q}_5] > \frac{d}{2}.$$

- Every residue field above 5 is larger than $\mathbb{F}_5$.

These are the field and local-degree conditions used for the integral comparisons in [CN, §5.6, (5)–(6)].

Let $\bar\rho : G_{F,S} \to \mathrm{SL}_2(\mathbb{F}_5)$ be surjective and decomposed generic. It is trivial at $S_5 \cup R$ and has the local form of Lemma 3.1 at P. Suppose that a cuspidal weight-zero representation π of $\mathrm{PGL}_2(\mathbb{A}_F)$ realizes $\bar\rho$. Require π to be unramified outside $R \cup P \cup S_5^{\mathrm{st}}$ and to have Iwahori invariants there. Its Galois representation is Barsotti–Tate at S_5^{cr} and unipotently ramified at $R \cup P$. At S_5^{st}, require π to be ι-ordinary and its Galois representation to be noncrystalline ordinary.

At S_5^{cr} and S_5^{st} use the local rings R_v^{BT} and $R_v^{\triangle}$ of [CN, §5.3]. At $v \in R$ use the character-type ring $R_v^{\varepsilon_5^{-1},\chi_v}$, where χ_v is tame and congruent to 1 modulo ϖ. At P use the Steinberg ring of Lemma 3.1. Denote the resulting global deformation problem by $\mathcal{S}_\chi$.

Integral Hecke maps

Lemma 4.1. *For every Taylor–Wiles datum Q satisfying Lemma 3.4, and every character tuple χ, there is a surjective $\mathcal{O}[\Delta_Q]$-algebra map*

$$R_{\mathcal{S}_{\chi,Q}} \longrightarrow \mathbb{T}_{\chi,Q}/J_{\chi,Q},$$
$$J_{\chi,Q}^{\delta} = 0, \qquad \Delta_Q = \prod_{w \in Q} k(w)^{\times}(5),$$

where δ is independent of the level and χ. The map matches Frobenius characteristic polynomials and satisfies the reduction and augmentation compatibilities of [ACC, §6.4.1].

Here the complexes and derived Hecke images are those of [CN, §5.6], with full Iwahori level and trivial coefficient character at P, coefficient character χ^{-1} at R, and the parabolics and ordinary projectors determined by the decomposition of S_5. The places in P^c are hyperspecial. Taylor–Wiles primes are chosen outside $S \cup S^c$.

Proof. Write

$$S_{\mathrm{aux}} = (S \cup Q) \cup (S \cup Q)^c, \qquad R_{\mathrm{aux}} = R \cup P \cup Q.$$

The first set is used to construct Galois representations and is stable under complex conjugation. It is distinct from the framing set used below. At every place of $R_{\mathrm{aux}}^c \setminus R_{\mathrm{aux}}$, the level is hyperspecial. In the notation of [ACC, §3.1], the set outside which the resulting representation is unramified is therefore

$$S_{\mathrm{aux}} \setminus (R_{\mathrm{aux}}^c \setminus R_{\mathrm{aux}}) = S \cup Q.$$

The field hypotheses of that result hold at $R \cup P$, and Lemma 3.4 supplies them at Q. In particular, applying [ACC, Theorem 3.1.1] removes ramification at P^c; unramifiedness there is part of the theorem, not a consequence of an identity of inertial characteristic polynomials.

We describe the passage of the required identities between Hecke algebras. First work at a neat GL_2 level, keeping the full Iwahori subgroup at P. On one fixed localized complex let H be the image of the original commuting Hecke operators, and let H_{ext} be the image after adjoining the torus operators $t_{v,i}(\sigma)$ of [ACC, §2.2.4], for $v \in R_{\mathrm{aux}}$. Localize the extended algebra over the original maximal ideal, retaining all its local factors. This gives an actual inclusion $H \subset H_{\mathrm{ext}}$, as in [ACC, §3.1]. Operators and ordinary projectors at 5 commute with these prime-to-5 operators, so they may be retained throughout, by scalar extension and passage to the corresponding direct summands.

There are representations over H/I and H_{ext}/J with the prescribed Frobenius polynomials, where $I^a = J^b = 0$ and a, b are bounded independently of the level. The first is furnished by the Galois-representation construction underlying [CN, Proposition 5.5.2]; the second is furnished by [ACC, Theorem 3.1.1]. Set

$$L = IH_{\mathrm{ext}} + J, \qquad K = H \cap L.$$

Then $I \subset K$, $L^{a+b-1} = 0$, and

$$K^{a+b-1} = 0, \qquad H/K \hookrightarrow H_{\mathrm{ext}}/L.$$

After extension to the ring on the right, the two representations have the same characteristic polynomials by Chebotarev and continuity. Absolute residual irreducibility identifies them up to conjugation on each complete local factor A: after identifying their residual representations over its residue field k_A, apply [CHT, Lemma 2.1.10] to the pair representation over $A \times_{k_A} A$ and the diagonal subring A. Equality of traces puts the pair's traces in that subring, and descent to it gives the required conjugacy. Thus the unramifiedness assertion also descends through this injection.

For $v \in P$, an inertial element corresponds under local reciprocity to a unit of F_v. At full Iwahori level with trivial coefficient character, each diagonal unit has Hecke action 1. Consequently the polynomial in [ACC, (2.2.7)] is $(X-1)^2$, and its Theorem 3.1.1 gives the integral identity

$$\det(X - \rho(\sigma)) = (X-1)^2 \qquad (\sigma \in I_v).$$

It descends coefficient by coefficient to H/K. Since the residual matrix $\bar{\rho}(\tau_v) - 1$ has a unit entry, Lemma 3.1 identifies the local representation over this possibly nonreduced ring with a point of the integral Steinberg deformation functor.

For clarity, the remaining comparisons use equivariant cohomology, not a direct map from the cohomology of a neat cover to that of its quotient. If K' is a neat normal subgroup of K_0, the actual Hecke images fit into the span

$$H(K') \longleftarrow H(K_0/K') \longrightarrow H(K_0).$$

The left arrow forgets the finite group action and has a nilpotent kernel of exponent bounded by the cohomological range; the right arrow is induced by derived invariants. These are the maps described after [ACC, Lemma 2.2.3]. Write $B = H(K_0/K')$, and write the arrows onto their images as $a : B \twoheadrightarrow H_{\mathrm{neat}}$ and $b : B \twoheadrightarrow H_{\mathrm{target}}$. With $E = \ker a$, there is a well-defined map

$$H_{\mathrm{neat}} = B/E \longrightarrow H_{\mathrm{target}}/b(E).$$

If the representation over H_{neat} is defined modulo a nilpotent ideal I, quotient the target further by $b(a^{-1}(I))$, and also by its original Galois-representation ideal J_0. These are nilpotent ideals: if $I^i = 0$ and $E^e = 0$, then $(a^{-1}(I))^{ie} = 0$. The two representations on this common quotient have equal Frobenius polynomials and hence are identified by absolute residual irreducibility. Consequently the identities already proved for the source representation hold for the original target representation after this quotient. This operation never lifts a Galois representation through a nilpotent kernel.

Apply this construction with the rank-one coefficient module $\mathcal{O}(\chi^{-1})$, made trivial on the covering subgroup, and with the quotient Δ_Q retained. The change of coefficients is the derived finite-group construction, so no averaging over a group of order divisible by 5 is used. It gives the coefficient and diamond versions of the span; these are the constructions in [ACC, Propositions 6.5.3 and 6.5.11] and [AKT, Lemma 5.2]. Localizations at the chosen distinct eigenvalues at Q commute with these maps. The identity at P is unchanged, since the coefficient character there is trivial and none of these level changes alters its full Iwahori subgroup.

To obtain neat levels, choose two auxiliary rational primes, split completely in F and outside S_{aux}. For the sequence of Taylor–Wiles sets used below, they may be fixed first and excluded when those sets are chosen. Impose sufficiently small principal congruence level at one of them at a time. Each construction permits additional ramification only at its own auxiliary prime. On the common quotient by the two resulting nilpotent ideals, equality of Frobenius polynomials identifies the representations by absolute residual irreducibility; the representation is therefore unramified at both auxiliary primes. This is the level-removal argument used in [ACC, §3.2] and in the proof of [AKT, Theorem 5.8]. The nilpotence bounds depend on the cohomological range, not the indices of these covers.

Finally take the preimages in GL_2 of the required PGL_2 levels. Apply the equivariant comparison to the bounded neat-cover complex, and take cohomology after derived invariants to reach the GL_2 cohomology Hecke image. No cohomological bound is needed for this latter space, whose centre can contain 5-torsion. The centres act trivially on the coefficient module. The order of the centre of SL_2 is 2, which is invertible in O, so [AKT, Corollary 5.5] gives the surjection from that cohomology Hecke image to the PGL_2 image. The localized PGL_2 complexes have a uniform cohomological bound by [AKT, Theorem 5.11], since $\mu_5 \subset F$, the residual representation is absolutely irreducible, and the coefficients reduce to the trivial module. On these bounded complexes, [ACC, Lemma 2.2.3] bounds the kernel from the derived Hecke image to its cohomology image; pulling back the quotient ideal therefore gives a uniformly nilpotent ideal in the derived algebra $\mathbb{T}_{\chi,Q}$. These operations justify the transfer in the statement without assuming an inclusion between Hecke algebras attached to different complexes.

At 5, the local comparison is [CN, Theorem 4.2.15]; its field, degree, residue-field, and decomposed-generic hypotheses hold by assumption. At R, the same torus-polynomial calculation gives the character-type condition. At Q, the distinct residual eigenvalues identify the two local characters and their diamond actions, as in [ACC, Proposition 6.5.11]. The fixed determinant and the resulting surjective deformation-ring map are obtained from these constructions as in [CN, Propositions 5.5.2 and 5.6.2–5.6.3]; the additional condition established here only takes a quotient of its target. Combining the finitely many nilpotent ideals above gives a uniformly nilpotent ideal $J_{\chi,Q}$. The resulting map factors through $R_{S_{\chi,Q}}$, remains surjective, and is $O[\Delta_Q]$-linear.

For compatibility, the coefficient systems for χ and 1 have the same reduction modulo ϖ. The derived change-of-coefficient isomorphism identifies the actions of the same abstract Hecke and diamond operators, and hence their common reduced image $\overline{\mathbb{T}}_Q$. The two maps agree after passage to

$$\overline{\mathbb{T}}_Q/(\overline{J}_{\chi,Q} + \overline{J}_{1,Q}).$$

Indeed, their Frobenius traces and determinants agree, so absolute residual irreducibility identifies the unframed deformations. The protected quotient is identical in the two problems. Under the augmentation isomorphism of [CN, Lemma 5.6.4], the same operators and Frobenius polynomials agree after killing the diamond variables. Quotienting by the image of the finite-level nilpotent ideal and the original-level ideal gives equality of the corresponding unframed deformations. These are exactly the reduction and augmentation compatibilities required in [ACC, §6.4.1, (5) and the subsequent augmentation conditions]. □

The lifting theorem

Theorem 4.2. *In the prepared setting above, let $\rho : G_F \to \mathrm{GL}_2(\overline{\mathbb{Q}}_5)$ be continuous, with residual representation $\bar{\rho}$ and determinant ε_5^{-1}. Suppose that:*

1. *ρ is unramified outside S and unipotently ramified at $R \cup P$;*
2. *at S_5^{cr}, it is Barsotti–Tate, and is ordinary exactly where $r_\iota(\pi)$ is ordinary;*
3. *at S_5^{st}, it is a noncrystalline extension of ε_5^{-1} by the trivial character.*

Then ρ is automorphic of weight zero.

Proof. Put $T = S \setminus P = S_5 \cup R$, $t = |T|$, and $M = \mathrm{ad}^0\,\bar{\rho}$. Thus T is nonempty. We frame at T, keeping the Steinberg condition at P in every global deformation functor. In the notation of [AKT, §A.2], the relative dual Selmer group for an augmented problem is

$$\ker\left(H^1(G_{F,S\cup Q}, M(1)) \longrightarrow \prod_{v\in P\cup Q} H^1(F_v, M(1))/\mathcal{L}_v^{\perp}\right).$$

Omitting a local factor from the framing base thus retains its dual condition in this kernel. Lemma 3.2 applies at P, while $\mathcal{L}_w^{\perp} = 0$ at every Taylor–Wiles prime w.

Lemma 3.4 supplies sets Q_N of a fixed sufficiently large cardinality r for which this group vanishes. The generator and Euler formulas of [AKT, Proposition A.4(1)–(2)] give a presentation over the relative local base with

$$g = -h^0(F, M(1)) - 1 + t - \sum_{v|\infty} h^0(F_v, M) + \sum_{v \in P \cup Q_N} \big(\dim_k \mathcal{L}_v - h^0(F_v, M)\big),$$

and hence

$$g = r - 3d - 1 + t.$$

Here the global invariant term vanishes, the d complex places each contribute 3, a Taylor–Wiles place has tangent dimension 2 and invariant dimension 1, and a protected place has both dimensions equal to 1. Increase r if necessary to make g nonnegative. The generator-surjection assertion in Proposition A.4(1) needs no local smoothness hypothesis. We do not invoke its stronger assertion about the number of relations, and do not assume unrestricted local H^2-vanishing at P.

For each χ, form

$$R_{\mathrm{loc},\chi} = \widehat{\bigotimes}_{v \in T} R_{v,\chi}.$$

The protected conditions remain imposed in R_{S_χ}; their rings are absent only from the relative local base. At R choose a second tuple χ with $\chi_v^2 \neq 1$ and $\chi_v \equiv 1$ modulo ϖ, enlarging O as needed. Such characters exist because $q_v \equiv 1$ modulo 5. The two local problems have identified reductions modulo ϖ.

Use the localized complexes and the finite-level maps of Lemma 4.1. Perfectness at the original, possibly non-neat, level follows from [CN, Proposition 5.5.3]. Perfectness over $O[\Delta_{Q_N}]$ and the identification of the Taylor–Wiles augmentation with the original complex are supplied by [CN, Lemma 5.6.4], using [AKT, Theorem 5.11].

Partial framing adds $4t - 1$ variables [ACC, Lemma 6.2.4]. Let $\mathcal{A}$ be that power-series ring over O and set

$$\begin{aligned} S_\infty &= \mathcal{A}[[\mathbb{Z}_5^r]], \\ R_\infty &= R_{\mathrm{loc},1}[[x_1, \ldots, x_g]], \\ R'_\infty &= R_{\mathrm{loc},\chi}[[x_1, \ldots, x_g]]. \end{aligned}$$

For each N, the presentations above give surjections from these power-series rings onto the corresponding T-framed global rings. Choose a common relative cotangent-space basis at the original level and lifts giving generators for the two original-level rings, compatible modulo ϖ. The augmentation from each framed augmented ring to its original-level ring is surjective, and so is its map on relative cotangent spaces. Extend lifts of the fixed original basis to a basis of the augmented relative cotangent space, choosing the remaining vectors in the augmentation kernel. The preceding presentation bound gives at most g basis elements, so pad the lists, including the original-level list, with zeros to obtain g generators. The actual lifts can be chosen with both the fixed augmentation values and common reductions modulo ϖ: for a surjection $B_N \to B_0$, the map

$$B_N \longrightarrow B_0 \times_{B_0/\varpi} (B_N/\varpi)$$

is surjective. Apply this to the framed rings and their primed counterparts. The resulting presentations are compatible both modulo ϖ and with augmentation; the identification between framed and unframed rings also respects the $4t-1$ framing variables. This supplies the presentation data in [ACC, §6.4.1, (3)–(4) and its augmentation conditions]. Lemma 4.1 supplies the compatible Hecke maps, so the abstract patching construction of [ACC, §6.4] applies with this framing algebra. Its augmentation maps give

$$R_\infty/\mathfrak{a}_\infty \twoheadrightarrow R_{S_1}, \qquad R'_\infty/\mathfrak{a}_\infty \twoheadrightarrow R_{S_\chi}$$

[ACC, Lemma 6.4.15]. In particular, the target defines a point of the patched spectrum.

The local dimensions give

$$\dim R_{\mathrm{loc},1} = 1 + 3t + 2d,$$
$$\dim R_\infty = r + 4t - d, \qquad \dim S_\infty = r + 4t.$$

The same formulas hold for the primed problem. Thus the dimension deficit is d. No unrestricted local H^2-vanishing is required at P: its imposed quotient is smooth and its relative Euler contribution is zero.

We apply the support argument of [CN, §5.4]. The local component and specialization properties at 5 are [CN, Proposition 5.3.2 and Lemmas 5.3.3–5.3.4]. The character-type properties at R are [AKT, Lemma A.2]; in the primed base those factors are irreducible. After enlarging coefficients, the relevant components may be taken geometrically irreducible. These facts verify the component hypotheses of [CN, Assumption 5.4.1].

The source π gives a nonzero automorphic point. Characteristic-zero cohomology is concentrated in the usual interval of length d. Source and target lie on matching components at 5: ordinary or nonordinary at the crystalline places, and noncrystalline ordinary at the others. There are no protected-place component labels in the local base.

A source component and a component containing the target consequently specialize to the same primed component. Proposition 5.4.2 of [CN] transfers support to the target component, and its Corollary 5.4.3 preserves support at augmentation. Hence the target occurs in characteristic-zero automorphic cohomology. The identification of Hecke eigenvalues with Frobenius traces gives a cuspidal weight-zero Π with $r_\iota(\Pi) \simeq \rho$. □

5 Constructing the modular auxiliary curve

Throughout this section, assume $\mu_5 \subset K$. We produce the protected place on A_t and then apply Theorem 4.2.

A multiplicative degeneration of width three

Lemma 5.1. *The family A_t admits a split multiplicative local specialization with minimal discriminant valuation prime to 5. It can be imposed at one place v over a rational prime $q \geq 7$, while A_t has good reduction at every other place over q. These conditions are compatible with Proposition 2.3.*

Proof. In the covering family choose a boundary point R_0 of exact order $6p$, so that $[3p]R_0$ is nonzero 2-torsion. The branch sections S and $-S$ collide. After blowing up the collision and normalizing the degree-three cover, the main component is a connected étale elliptic cover. Its defining torsion class is $-2pR_0$, of exact order 3. A rational tail meets this component in three nodes, each of thickness one.

The involution exchanges two nodes and fixes the third. On both branches of the fixed node it acts as -1. Its local quotient changes $xy = s$ to $uv = s^2$. The exchanged pair contributes one node of thickness one. The quotient components are rational; resolving the thickness-two node produces a cycle of three rational components. Thus the degeneration has type I_3.

The normalization $f(O) = 1$ gives the section represented by $(O, 1)$, so the genus-one quotient and its Jacobian have the same degeneration. Since R_0 is not 2-torsion, the x-coordinate is étale there; passage to the t-parameter preserves the width.

Choose a finite number field over which R_0, the components, the nodes, and their individual branches are defined. Spread the semistable model and its smoothing parameter s out over a ring of integers with finitely many primes inverted. Enlarge this finite set so that the node and branch data and the stated thicknesses persist, E has good reduction, and the nonzero leading coefficient of $j(A_t)^{-1}$ at $s = 0$ is a unit. Choose $q \geq 7$, different from 5 and p, outside this set and split completely in the chosen field and the normal closure of K. The resulting I_3 degeneration is split at the selected

place v. Its inverse j-invariant has order three in s; the inverse Tate j-series, or equivalently Tate uniformization [Si, Chapter V], therefore gives, on a sufficiently small v-adic neighborhood,

$$q_{A_t} = cs^3 h(s), \qquad c \in O_v^\times, \qquad h(0) = 1.$$

Shrink the neighborhood so that $h(s)$ is a unit. Choose $v(s) = k > 0$ sufficiently large with $5 \nmid k$. Then A_t is split multiplicative and $v(\Delta_{\min}) = 3k$ is prime to 5. These are open local conditions. At all other q-places, including v^c, use the near-O good-reduction open; the choice of q ensures that E has good reduction there and $v \neq v^c$. Proposition 2.3 permits all these conditions together. □

In a split Tate curve, residual inertia is nontrivial unipotent precisely when its discriminant valuation is nonzero modulo 5. Since $\mu_5 \subset K$, residual Frobenius has diagonal entries 1; changing its tame lift by a power of inertia makes it the identity. Thus the protected place has exactly the form used in Lemma 3.1.

A modular source and the prepared field

Proposition 5.2. *Let A/K be an elliptic curve with full $\mathrm{SL}_2(\mathbb{F}_5)$ residual image, decomposed generic residual representation, and a protected place as in Lemma 5.1. There is a finite solvable CM extension F/K over which $H^1(A)$ and a modular elliptic curve satisfy the hypotheses of Theorem 4.2. Every completion above the selected rational prime q can be kept unchanged.*

Proof. We use the proof of the modular-source construction [CN, Proposition 6.1.5], which allows an arbitrary finite Galois avoid field and finitely many nonempty local opens. Let $M = K(A[5])$, and choose a finite Galois extension D/K^+ containing M and any further avoid data. Choose a finite set V of places witnessing the residual image and decomposed genericity, away from $2, 3, 5, q$ and the rational primes of bad reduction of A. Include all places above the rational primes so chosen. In particular, V contains every place above a rational decomposed-generic witness. The image witnesses may be chosen to detect every proper intermediate field of M/K, as in [ACC, §6.5.12].

We first make the preliminary base change in the cited modular-source proof explicitly over K^+. At each relevant finite place u of K^+, prescribe a finite Galois local extension containing the local fields and the required auxiliary extensions for every place of K above u. Thus, at a split pair v, v^c, take a compositum of the prescriptions for both places and its local Galois closure; the two residual representations need not coincide. Above $2, 3, 5$, these prescriptions make $A[5]$ trivial and satisfy the preliminary local conditions of [CN, Proposition 6.1.5]. Above 5, they also give A good reduction or split multiplicative reduction according to its category, and residue fields larger than $\mathbb{F}_5$. At every place above q or below V, prescribe the trivial extension, and at every real place prescribe $\mathbb{R}$.

The exact local prescription lemma [CHT, Lemma 4.1.2] gives a totally real, solvable Galois extension E_0^+/K^+, linearly disjoint from D, with these completions. Put $L_0 = KE_0^+$. Then L_0/K is solvable Galois and CM. Disjointness gives $L_0 \cap M = K$, so the residual image remains full. Every q-completion is unchanged, and every place of V splits completely. At q we impose no condition trivializing the residual representation: its nontrivial inertia at the distinguished place is retained.

Now apply the weak approximation and Hilbert irreducibility step on the rational twisted modular curve $X(5)$, with the prescribed reduction categories at 5. Add the split-Tate open at every place above the distinguished v, good-reduction opens at the other q-places, and good-reduction opens at the places above V. The added opens are nonempty because A supplies local points, and their residue characteristics are disjoint from the mandatory 2-, 3-, and 5-adic conditions. The proof of [CN, Proposition 6.1.5] therefore gives a modular elliptic curve B/L_0, with $B[5] \simeq A[5]$, satisfying these conditions. Its residual image is full and remains decomposed generic.

Prepare both A and B at their remaining bad places away from the protected set. Choose local extensions making their common residual representation trivial and both reductions semistable. They may be enlarged to finite Galois extensions. Apply the same construction again over K^+: for each place of K^+, combine the required local extensions for all places of L_0 above it before taking the local Galois closure. Keep trivial prescriptions above q and below V, and real infinite completions.

Apply [CHT, Lemma 4.1.2] with an avoid field containing a Galois closure of L_0M/K^+. If E_1^+ is the resulting totally real extension, replace L_0 by $L_0E_1^+$. These composita are Galois and solvable over K; disjointness preserves the full residual image, and splitting at V preserves decomposed genericity. The source stays modular by solvable base change [ACC, Proposition 6.5.13]; its residual full image ensures that the base change remains cuspidal.

Next adjoin a totally real multiquadratic extension of $\mathbb{Q}$ of degree eight, disjoint from the fields already used and split at 5, q, and the rational primes below V. Such extensions follow from weak approximation for square classes, with new auxiliary ramification ensuring independence. Each 5-adic local degree is then at most one eighth of the global real degree.

This ratio cannot increase under further extension: a local degree grows by at most the global degree. Omitting any two 5-adic places therefore leaves at least $3d/4$, giving the strict inequality required in §4.

Finally, use the imaginary-quadratic helper construction of [ACC, §6.5.12], incorporating q and the rational primes below V into its finite splitting prescriptions. It arranges that every rational prime below the final deformation set, or ramifying in F, splits in an imaginary quadratic subfield of F. In particular, the relevant places split over F^+. The helpers are quadratic over $\mathbb{Q}$, so the final extension F/K remains Galois and solvable, and all its q-completions remain unchanged. The full residual image survives these final solvable Galois extensions as well: a nontrivial intersection with the residual field would give a nontrivial solvable quotient of the perfect group $\mathrm{SL}_2(\mathbb{F}_5)$.

Let P be the places above the original distinguished v; their conjugates remain good for both curves and lie outside S. Let R be the union of the remaining bad places of A and B, away from 5 and P. Then $S = S_5 \sqcup R \sqcup P$, the residual representation is trivial at $S_5 \cup R$, and the source is unramified outside $R \cup P \cup S_5^{\mathrm{st}}$ with Iwahori invariants on that set. At P both curves remain split Tate curves with nontrivial residual inertia. At 5 their ordinary, nonordinary, and noncrystalline ordinary categories match. For the modular source, the good and multiplicative reduction conditions give the required unramified and ordinary automorphic local conditions by [CN, Lemmas 6.1.3 and 6.1.7]. These are all the hypotheses of Theorem 4.2. □

Corollary 5.3. *The parameter in Proposition 2.3 can be chosen so that A_t is modular over K.*

Proof. Add the local opens of Lemma 5.1 to the specialization. Proposition 2.3 gives the full residual image and decomposed genericity. Proposition 5.2 supplies a modular source and the prepared field. Theorem 4.2 applied to $H^1(A_t)$ proves automorphy over that field. Solvable descent [ACC, Proposition 6.5.13] gives modularity over K. The residual full image also excludes geometric CM, so the resulting representation is cuspidal. □

6 Completion of the proof

We first record how automorphy moves between coefficient primes.

Lemma 6.1. *Suppose that a system V_θ from Proposition 2.1 is realized by a weight-zero cuspidal automorphic representation π with trivial central character at one coefficient prime. Then π realizes the entire compatible system. At every place covered by Proposition 2.2, its local parameter has zero monodromy in the potentially good case and Steinberg type in the potentially multiplicative case.*

Proof. The good-place characteristic polynomials of V_θ determine those of every coefficient realization of π. Semisimplicity, Chebotarev, and Brauer–Nesbitt identify the representations at all coefficient primes. Existence of these Galois representations is the CM Galois-representation theorem [HLTT].

For the local assertion, compare at an auxiliary coefficient prime different from the residue characteristic. Semisimplified local–global compatibility identifies the underlying semisimple Weil representation [V]. In the potentially good case, after a finite extension the two geometric Frobenius eigenvalues have the same complex absolute value. Nonzero rank-two monodromy would force their ratio to be a nontrivial norm character, contradicting that equality. Thus the automorphic monodromy is zero.

In the potentially multiplicative case, the geometric monodromy has rank one and the two Weil characters differ by the norm. The zero-monodromy parameter with these characters corresponds to a nongeneric one-dimensional representation. Since a cuspidal GL_2 representation is locally generic, π_v is a Steinberg twist. Its trivial central character makes the twisting character quadratic or trivial. The local calculation [O, Lemma 9.2(ii)] gives weight-zero ordinariness, including for ramified quadratic twists; compare [CN, Lemma 6.1.3(4)]. □

Proposition 6.2. *If $\mu_5 \subset K$ and E/K has no geometric complex multiplication, then E is modular.*

Proof. Choose p and t as in Proposition 2.3 and Corollary 5.3. Put $F = K(\mu_{3p})$, $A = A_t$, and $B = B_t$. This is an abelian CM extension. Base-change the cuspidal representation of A to F. Its Galois representation remains absolutely irreducible, so the base change is cuspidal. It realizes V_η.

We apply [CN, Theorem 5.2] three times:

$$V_\eta \xrightarrow{p} V_{\eta\psi} \xrightarrow{3} V_\psi \xrightarrow{p} V_1.$$

Proposition 2.1 supplies determinant ε_ℓ^{-1}, labelled weights $\{0, 1\}$, finite ramification, and potential semistability. It also supplies the residual match. For the first step, the common residual representation comes from C_3 modulo p; for the second, from C_p modulo 3; for the last, from B modulo p. Proposition 2.3 gives absolute irreducibility over the cyclotomic fields and decomposed genericity in each case. At a change of coefficient prime, Lemma 6.1 supplies the automorphic source at the new prime.

At a potentially ordinary place, source and target are potentially crystalline and potentially ordinary. At a supersingular place, both are potentially crystalline and neither is potentially ordinary. In both cases the source's automorphic monodromy is zero. At a potentially multiplicative place, both are noncrystalline and the source is ordinary of Steinberg type. Thus Proposition 2.2 and Lemma 6.1 give all local hypotheses of the lifting theorem. The coefficient primes are 3 and $p > 100$, so its exceptional mod-5 clause never occurs.

The last step makes $H^1(B/F)$ automorphic. Undoing the quadratic twist gives automorphy of $H^1(E/F)$. Descend through a cyclic tower for F/K. At each step, the prescribed Galois representation makes the automorphic representation invariant by strong multiplicity one, so cyclic base change supplies a cuspidal descent [AC]; [ACC, Proposition 6.5.13].

For clarity, the finite-character correction is determined by the Galois representation. Two extensions of the same absolutely irreducible representation of a finite-index normal subgroup differ by a character of the finite quotient: after identifying their restrictions, Schur's lemma makes their pointwise discrepancy scalar. Twisting the descended automorphic representation by the inverse corresponding Hecke character removes that discrepancy. Every intermediate field is CM, and the required Galois representations exist. This gives precisely $H^1(E/K)$. □

Proof of Theorem 1.1. First suppose E has no geometric CM. The field $K' = K(\mu_5)$ is CM and abelian over K. Proposition 6.2 applies to E/K', and the same solvable descent gives its modularity over K.

Agreement of the compatible systems identifies the elliptic local parameters, including monodromy, by [CN, Lemma 6.1.3]. This gives the stated local factors and conductor. The determinant ε_ℓ^{-1} identifies the unitary central character as trivial.

Now suppose that E has geometric complex multiplication by an imaginary quadratic field M. Its endomorphisms are defined over KM. The main theorem of complex multiplication supplies the corresponding algebraic Hecke characters [Si, Chapter II, §§9–10]. If $M \subset K$, $H^1(E)$ is a sum of two characters and the associated automorphic representation is their isobaric sum. Otherwise KM/K is quadratic and $H^1(E)$ is induced from one character of KM. Quadratic automorphic induction gives a cuspidal representation over K [AC]. The two infinity characters are distinct, so the character is not invariant under the nontrivial automorphism of KM/K. The CM theorem and compatibility of induction identify the local parameters. □

Acknowledgments

We thank the GitHub user `toby-gee` for pointing out the relevant precedents in the work of Taylor and Khare–Thorne in issue 1.

References

[O] OpenAI. *Modularity of elliptic curves over imaginary quadratic fields*. Preprint, 4 October 2026, 30 pp. Cited version and source repository. Cited source version: repository commit `adc7f1241b42e322a6451854ab7e4b4c146bf78a`.

[CN] Ana Caraiani and James Newton. *On the modularity of elliptic curves over imaginary quadratic fields*. arXiv:2301.10509v3, 27 March 2025. Version used for the theorem numbering.

[ACC] Patrick B. Allen, Frank Calegari, Ana Caraiani, Toby Gee, David Helm, Bao V. Le Hung, James Newton, Peter Scholze, Richard Taylor, and Jack A. Thorne. *Potential automorphy over CM fields*. Annals of Mathematics (2) **197** (2023), no. 3, 897–1113. doi:10.4007/annals.2023.197.3.2. Full text.

[AKT] Patrick B. Allen, Chandrashekhar Khare, and Jack A. Thorne. *Modularity of* $\mathrm{GL}_2(\mathbb{F}_p)$*-representations over CM fields*. Cambridge Journal of Mathematics **11** (2023), no. 1, 1–158. doi:10.4310/cjm.2023.v11.n1.a1. Preprint.

[Tay03] Richard Taylor. *On icosahedral Artin representations, II*. American Journal of Mathematics **125** (2003), no. 3, 549–566. doi:10.1353/ajm.2003.0021. Author's manuscript.

[KT] Chandrashekhar B. Khare and Jack A. Thorne. *Automorphy of some residually* S_5 *Galois representations*. Mathematische Zeitschrift **286** (2017), nos. 1–2, 399–429. doi:10.1007/s00209-016-1766-y. Preprint.

[CHT] Laurent Clozel, Michael Harris, and Richard Taylor. *Automorphy for some l-adic lifts of automorphic mod l Galois representations*. Publications Mathématiques de l'IHÉS **108** (2008), 1–181. doi:10.1007/s10240-008-0016-1.

[AC] James Arthur and Laurent Clozel. *Simple Algebras, Base Change, and the Advanced Theory of the Trace Formula*. Annals of Mathematics Studies **120**, Princeton University Press, 1989. Publisher record.

[S72] Jean-Pierre Serre. *Propriétés galoisiennes des points d'ordre fini des courbes elliptiques*. Inventiones Mathematicae **15** (1972), no. 4, 259–331. doi:10.1007/BF01405086.

[SG] Jean-Pierre Serre. *Topics in Galois Theory*. Second edition, with notes by Henri Darmon. Research Notes in Mathematics **1**, A K Peters, 2008.

[F] Gerd Faltings. *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*. Inventiones Mathematicae **73** (1983), no. 3, 349–366; erratum, **75** (1984), 381. Article; erratum.

[HLTT] Michael Harris, Kai-Wen Lan, Richard Taylor, and Jack Thorne. *On the rigid cohomology of certain Shimura varieties*. Research in the Mathematical Sciences **3** (2016), Paper No. 37, 308 pp. doi:10.1186/s40687-016-0078-5.

[V] Ila Varma. *Local-global compatibility for regular algebraic cuspidal automorphic representations when* $\ell \neq p$. Forum of Mathematics, Sigma **12** (2024), Paper No. e21, 32 pp. doi:10.1017/fms.2024.7.

[Si] Joseph H. Silverman. *Advanced Topics in the Arithmetic of Elliptic Curves*. Graduate Texts in Mathematics **151**, Springer-Verlag, 1994. doi:10.1007/978-1-4612-0851-8.