$$\begin{array}{c} \dot{V}_{0} = \frac{\partial V_{1}(\mathbf{x})}{\partial x_{1}} + \gamma_{2} z_{2} + \frac{1}{\gamma_{1}} \hat{M}_{0} \hat{M}_{1} & (2) \\ & = \frac{\partial V_{1}(\mathbf{x})}{\partial x_{1}} \left(\mathbf{x}_{1}(\mathbf{x}_{1}) + \mathbf{y}_{1}(\mathbf{x}_{2}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(z_{2} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{2}} \right) \left(z_{1}(\mathbf{y}_{1}) + \mathbf{y}_{1}(\mathbf{y}_{1}) z_{2} \right) + \frac{1}{\gamma_{0}} \hat{M}_{2} \hat{M}_{1} \\ & = \frac{\partial V_{1}(\mathbf{x})}{\partial x_{1}} \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(\mathbf{f}_{1}(\mathbf{x}_{1}) + \mathbf{g}_{1}(\mathbf{x}_{1}) z_{2} \right) + \gamma(z_{2} - m_{1}(\mathbf{x}_{1})) \right) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) \left(m_{1} - \frac{\partial m_{1}(\mathbf{x})}{\partial x_{1}} \right) + \frac{\gamma}{h_{0}} \hat{\mathbf{g}}_{1} \hat{\mathbf{g}}_{1$$

(1)

 $V_3 = V_1 + \frac{\gamma}{2} z_2^2 + \frac{\gamma}{2h} \tilde{M}_L^2$