

Alec Mabhiza Chirawu - 亚历克上

Student ID: M202161029

Information Theory and Network Coding

FINAL EXAM

ALEC MABHIZA CHIRAWU (亚历克上)
M202161029
INFORMATION THEORY AND NETWORK CODING
FINAL EXAMINATION

1) The different goals of source coding, channel coding and network coding

- ⇒ Source coding
 - It represent information as accurately as possible using as few bits as possible.
 - The main aim is to convert information waveforms for example text, audio images and videos.
 - It deals with data compression, and error control.
- ⇒ Channel coding
 - Channel coding provides the GSM receiver with the ability to detect transmission errors.
 - It correct errors from a bit point of view.
 - The main aim for channel coding is to convert bits to signal waveform (encoder) and decoder converts waveform back to bits.
- ⇒ Network coding
 - Network coding is just a networking technique in which transmitted data is encoded and decoded.
 - The main aim is to increase throughput and reduce delay.

M202161029

2) The Source Coding Theorem

→ Source coding theorem is known as Shannon's source coding theorem.
 → It is also called noiseless coding theorem.

→ Source coding theorem states that we can compress N independent and identically distributed (i.i.d) random variables each with entropy (H) down to (NH) bits with negligible loss of information as $N \rightarrow \infty$.

→ The theorem is:

Let X be an ensemble with $H(X) = H$ bits.

→ given $\epsilon > 0$ and $0 < E < 1$

→ N is positive N_0 for $N > N_0$.

$$\left| \frac{1}{N} H_Y(X^N) - H \right| < \epsilon$$

The channel coding theorem


→ Shannon's noisy channel

→ It states that all rates below capacity C are achievable,

→ For every rate $R \leq C$, there exists a sequence of $(2^{nR}, n)$ codes with maximum probability of error $P_E \rightarrow 0$.

M202161029

3)



Information measures corresponding to:

(i) Area $A = H(X|Y, Z)$

(ii) Area $D = I(X, Y|Z)$

(iii) Area $A+B+D = H(X, Y|Z)$

(iv) Area $E+F+G = I(X, Y; Z)$

(v) G is possible to take a negative value.

4) 4 binary codes:

M202161029

$C_1 =$

code	$L(x)$
000	3
11	2
001	3
010	3
101	3

$\Rightarrow 4 \cdot \frac{1}{2^3} + \frac{1}{2^2} \Rightarrow 0.75$

prefix

$C_2 \Rightarrow$

code	$L(x)$
0	1
10	2
0010	4
0011	4

$\Rightarrow \frac{1}{2} + \frac{1}{2} + 2 \cdot \frac{1}{2^4} \Rightarrow 0.875$

not prefix

C_3

code	$L(x)$
0	1
10	2
11	2
0100	4

$\Rightarrow \frac{1}{2} + 2 \cdot \frac{1}{2^2} + \frac{1}{2^4} \Rightarrow 1 \frac{1}{16}$

not prefix

$C_4 \Rightarrow$

code	$L(x)$
01	2
00	2
011	3
001	3

$\Rightarrow 2 \cdot \frac{1}{2} + 2 \cdot \frac{1}{2^3} \Rightarrow 0.75$

not prefix

a) C_3 does not satisfy Kraft's inequality

b) C_2 includes (0, 10 and 0010, 0011); C_3 includes (0, 10 and 0100)

C_4 includes (01, 00 and 011, 001) are not prefix codes and are not uniquely decodable.

$$\text{II} \quad \left[\begin{array}{l} P(X=0, Y=a) = \frac{1}{4} \\ P(X=0, Y=b) = 0 \\ P(X=0, Y=c) = \frac{1}{4} \\ P(X=1, Y=a) = 0 \\ P(X=1, Y=b) = \frac{1}{4} \\ P(X=1, Y=c) = \frac{1}{4} \end{array} \right]$$

$$P_X(x) = \sum_y P(X,Y) \quad \text{margin 6/02/24} \\ P_X(a) = \sum_y P_{XY}(X=0, y) = \frac{1}{4} + 0 + \frac{1}{4} = \frac{1}{2}$$

$$\begin{aligned} H(X,Y) &= \sum_x \sum_y P_{XY}(x,y) \log P_{XY}(x,y) \\ &\Rightarrow \frac{1}{4} \log_2 4 + 2 \cdot 0 \cdot \frac{1}{4} \log_2 4 + \frac{1}{4} \log_2 4 \\ &\Rightarrow \underline{\underline{2}} \end{aligned}$$

$$\begin{aligned} H(X) &\Rightarrow \sum P(x) \cdot \log_2 P(x) \\ &\Rightarrow \frac{1}{2} \log_2 2 + \frac{1}{2} \log_2 2 \\ &\Rightarrow \underline{\underline{1}} \end{aligned}$$

$$\begin{aligned} H(Y|X) &\Rightarrow \sum P(x) H(Y|X=x) \\ &\Rightarrow H(X,Y) - H(X) \\ &\Rightarrow 2 - 1 \\ &= \underline{\underline{1}} \end{aligned}$$


$$\begin{aligned} H(Y|X=0) &\Rightarrow P_X(a) + P_X(c) \\ &\Rightarrow \frac{1}{2} + \frac{1}{2} \\ &\Rightarrow \underline{\underline{1}} \end{aligned}$$

$$\begin{aligned} I(X;Y) &\Rightarrow H(X) - H(X|Y) \\ &\Rightarrow H(X) + H(Y) - H(X,Y) \\ &\Rightarrow 1 + 1 - 2 \\ &\Rightarrow \underline{\underline{0}} \end{aligned}$$

III. $X = \{x_1, x_2, x_3\} = \{0,6; 0,3; 0,1\}$

M202161029

1) Huffman code X :




$H_{code X} \Rightarrow \{0, 01, 11\}$

$$\bar{L}_1 \Rightarrow (1 \cdot 0,6) + (2 \cdot 0,3) + (2 \cdot 0,1) \Rightarrow 1,4$$

$$\begin{aligned} \eta_1 &= H(X) / \bar{L}_1 \Rightarrow H(0,6; 0,3; 0,1) / 1,4 \\ &\Rightarrow \frac{(3/5 \log_2 5/3) + (3/10 \log_2 9/3) + (1/10 \log_2 10)}{1,4} \\ &\Rightarrow 0,92532988 \\ &\Rightarrow 0,925 \end{aligned}$$

2) Huffman code for $X^2 = (0,6; 0,3; 0,1)^2 \Rightarrow (0,6; 0,3; 0,1)(0,6; 0,3; 0,1) \Rightarrow (0,36; 0,18; 0,06; 0,18; 0,09; 0,03; 0,06; 0,03; 0,01)$



$$\begin{aligned} \bar{L}_2 &\Rightarrow (1 \cdot 0,36) + (3 \cdot 0,18) + (3 \cdot 0,18) + (4 \cdot 0,09) + (4 \cdot 0,06) + (4 \cdot 0,06) + (5 \cdot 0,03) + (6 \cdot 0,03) + (6 \cdot 0,01) \\ &\Rightarrow 2,67 \end{aligned}$$

$$\begin{aligned} \eta_2 &= H(X^2) / \bar{L}_2 \Rightarrow 2H(X) / \bar{L}_2 \\ &\Rightarrow \frac{36}{100} \log_2 \frac{100}{36} + 2 \cdot \frac{18}{100} \log_2 \frac{100}{18} + \frac{9}{100} \log_2 \frac{100}{9} + 2 \cdot \frac{6}{100} \log_2 \frac{100}{6} + 2 \cdot \frac{3}{100} \log_2 \frac{100}{3} + \frac{1}{100} \log_2 100 \\ &\Rightarrow 2,59092368 / 2,67 \\ &\Rightarrow 0,9703834 \\ &\Rightarrow 0,970 \end{aligned}$$

IV) 1. $H(X_j) \Rightarrow H\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right) = 4 \cdot \frac{1}{4} \log_2 4 \Rightarrow \underline{2}$

$$H(X_{j+1}|X_j) = \underbrace{p(X_j=a)}_{\frac{1}{4}} H(X_2|X_j=a) + \underbrace{p(X_j=b)}_{\frac{1}{4}} H(X_2|X_j=b) + \underbrace{p(X_j=c)}_{\frac{1}{4}} H(X_2|X_j=c) + \underbrace{p(X_j=d)}_{\frac{1}{4}} H(X_2|X_j=d)$$

~~$H(X_{j+1}|X_j)$~~

$$H(X_2|X_j=a) = (Y_2, Y_4, 0, Y_4) = \frac{1}{2} \log_2 2 + 2 \cdot \frac{1}{4} \log_2 4 + 0 \log_2 0 \Rightarrow 1,5$$

$$H(X_2|X_j=b) \Rightarrow (0, Y_2, \frac{1}{4}, \frac{1}{4}) \Rightarrow 0 \log_2 0 + \frac{1}{2} \log_2 2 + 2 \cdot \frac{1}{4} \log_2 4 \Rightarrow 1,5$$

$$H(X_2|X_j=c) \Rightarrow (Y_4, \frac{1}{4}, \frac{1}{4}, 0) \Rightarrow \frac{1}{4} \log_2 4 + \frac{1}{4} \log_2 4 + \frac{1}{2} \log_2 2 + 0 \Rightarrow 1,5$$

$$H(X_2|X_j=d) \Rightarrow (Y_4, 0, Y_4, \frac{1}{2}) \Rightarrow 2 \cdot \frac{1}{4} \log_2 4 + 0 + \frac{1}{2} \log_2 2 \Rightarrow 1,5$$

$$\therefore H(X_j|X_{j+1}) \Rightarrow \frac{1}{4}(1,5) + \frac{1}{4}(1,5) + \frac{1}{4}(1,5) + \frac{1}{4}(1,5) \Rightarrow 1,5$$

$$\Rightarrow H(X_j, X_{j+1})/2 \Rightarrow \frac{H(X_j) + H(X_{j+1}|X_j)}{2} = \frac{2 + 1,5}{2} \Rightarrow \underline{1,75}$$

$$\Rightarrow H_x \Rightarrow \lim_{j \rightarrow \infty} H(X_{j+1}|X_j, X_{j-1}, \dots, X_1) = H(X_{j+1}|X_j) \Rightarrow \underline{1,5}$$

$\Rightarrow H_x$ in descending order:

$$H(X_j) > \frac{H(X_j) + H(X_{j+1}|X_j)}{2} > H_x$$

$$\underline{2 > 1,75 > 1,5}$$

2) $I(X_{j-1}, X_{j+1}|X_j) \Rightarrow \underline{1,5}$

$$V) X \in \{0, 1\}$$

$$Y \in \{0, e, 1\}$$

M202161024

1) transition matrix :

$$P(Y/X) = \begin{matrix} & \begin{matrix} 0 & e & 1 \end{matrix} \\ \begin{matrix} 0 \\ 1 \end{matrix} & \begin{bmatrix} P(Y=0/X=0) & P(Y=e/X=0) & P(Y=1/X=0) \\ P(Y=0/X=1) & P(Y=e/X=1) & P(Y=1/X=1) \end{bmatrix} \end{matrix}$$

$$\Rightarrow \begin{bmatrix} \frac{2}{3} & \frac{1}{3} & 0 \\ 0 & \frac{1}{3} & \frac{2}{3} \end{bmatrix}$$

$$H(Y/X) \Rightarrow H(\frac{2}{3}, \frac{1}{3}, 0) = \frac{2}{3} \log_2 \frac{3}{2} + \frac{1}{3} \log_2 3 \Rightarrow 0,91829583 \Rightarrow \underline{0,918}$$

$$P(X=0)=0, P(X=1)=1$$

$$P(X) = (0, 1)$$

$$P(Y/X) = \frac{P(X,Y)}{P(X)}$$

$$P(X,Y) = P(Y/X) \cdot P(X)$$

$$P(X,Y) \Rightarrow \begin{matrix} 0 \times [\frac{2}{3}, \frac{1}{3}, 0] \\ 1 \times [0, \frac{1}{3}, \frac{2}{3}] \end{matrix}$$

$$\Rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{3} & \frac{2}{3} \end{bmatrix}$$

$$P(X/Y) = \frac{P(X,Y)}{P(Y)}$$

$$P(Y) \Rightarrow [0, \frac{1}{3}, \frac{2}{3}]$$

$$P(X/Y) = \frac{\begin{bmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{3} & \frac{2}{3} \end{bmatrix}}{\begin{bmatrix} 0 & \frac{1}{3} & \frac{2}{3} \end{bmatrix}}$$

$$\Rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

$$I(X;Y) = H(X) - H(X/Y)$$

$$\Rightarrow H(X) - P(H(X))$$

$$\Rightarrow \frac{2}{3} H(X)$$

$$\Rightarrow \frac{2}{3} P(X)$$

$$\Rightarrow \frac{2}{3} \cdot 0 + \frac{2}{3} \cdot 1$$

$$\Rightarrow \frac{2}{3} \text{ bits/symbol}$$

$$2) H(Y/X) = \underline{0,918}$$

$$H(Y) \Rightarrow \underline{0}$$

$$I(X;Y) \Rightarrow \underline{\frac{2}{3}} \text{ bits/symbol}$$

$$3) P(X) = (\frac{1}{2}, \frac{1}{2})$$

$$P(X,Y) \Rightarrow \frac{1}{2} \cdot \begin{bmatrix} \frac{2}{3} & \frac{1}{3} & 0 \\ 0 & \frac{1}{3} & \frac{2}{3} \end{bmatrix} + \frac{1}{2} \cdot \begin{bmatrix} \frac{1}{3} & \frac{2}{3} & 0 \\ 0 & \frac{1}{3} & \frac{2}{3} \end{bmatrix}$$

$$P(X,Y) = \frac{P(X,Y)}{P(Y)} \because P(Y) \Rightarrow [\frac{1}{3}, \frac{1}{3}, \frac{2}{3}]$$

$$\Rightarrow \frac{\begin{bmatrix} \frac{1}{6} & \frac{1}{6} & 0 \\ 0 & \frac{1}{6} & \frac{1}{3} \end{bmatrix}}{\begin{bmatrix} \frac{1}{3} & \frac{1}{3} & \frac{2}{3} \end{bmatrix}} \Rightarrow \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

$$I(X;Y) \Rightarrow \frac{2}{3} H(X)$$

$$\Rightarrow (\frac{2}{3} \cdot \frac{1}{2}) H(\frac{1}{2}, \frac{1}{2})$$

$$\Rightarrow \frac{2}{3} \text{ bits/symbol}$$

$$H(Y) = [1 \ 1 \ \frac{1}{2}]$$

$$= 2 \cdot 1 \log 1 + \frac{1}{2} \log 2$$

$$\Rightarrow 1$$

$$3) H(Y/X) \Rightarrow 0,918$$

$$H(Y) \Rightarrow 1$$

$$I(X;Y) \Rightarrow \frac{2}{3} \text{ bits/symbol}$$

$$4) C = \max [I(X;Y)]$$

$$\Rightarrow \max [\frac{2}{3} H(X)]$$

$\therefore \max H(X) \Rightarrow \frac{2}{3} \text{ bits/symbol}$
 \Rightarrow It becomes maximum only if the transmitted symbol is 1.