

1 Probability and Combinatorics

$$\binom{n}{r}=\binom{n-1}{r}+\binom{n-1}{r-1}$$

$$\binom{-x}{r}=(-1)^r\binom{x+r-1}{r}$$

$$\binom{n}{r_1,r_2,\ldots,r_k}=\binom{n}{r_1}\cdots\binom{n-r_1-\cdots-r_{k-1}}{r_k}$$

$$\binom{x}{r}=\frac{x(x-1)\ldots(x-r+1)}{r!}$$

$$\binom{a}{0}\binom{b}{n}+\binom{a}{1}\binom{b}{n-1}+\cdots+\binom{a}{n}\binom{b}{0}=\binom{a+b}{n}$$

$$\text{Bernoulli}(p)=p^x(1-p)^{1-x}$$

$$p\mid p(1-p)\mid 1-p+pe^t$$

$$\text{Binomial}(n,p)=\binom{n}{x}p^x(1-p)^{n-x}$$

$$np\mid np(1-p)\mid (1-p+pe^t)^n$$

$$\text{Poisson}(\lambda)=\frac{\lambda^xe^{-\lambda}}{x!}\mid \lambda\mid \lambda\mid e^{\lambda(e^t-1)}$$

$$\text{Geometric}(p)=(1-p)^{x-1}p$$

$$\frac{1}{p}\mid \frac{1-p}{p^2}\mid \frac{pe^t}{1-(1-p)e^t}$$

$$\text{Uniform}(a,b)=\frac{1}{b-a}$$

$$\frac{a+b}{2}\mid \frac{(b-a)^2}{12}\mid \frac{e^{tb}-e^{ta}}{t(b-a)}$$

$$\text{Normal}(\mu,\sigma^2)=\frac{1}{\sqrt{2\pi}\sigma}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

$$\text{E}(Z^{2r+1})=0\mid \text{E}(Z^{2r})=\frac{(2r)!}{(2^rr!)}$$

$$\mu\mid \sigma^2\mid e^{\mu t+\frac{1}{2}\sigma^2t^2}$$

$$\text{Gamma}(\alpha,\beta)=\frac{\beta^\alpha}{\Gamma(\alpha)}x^{\alpha-1}e^{-\beta x}$$

$$\frac{\alpha}{\beta}\mid \frac{\alpha}{\beta^2}\mid \left(\frac{\beta}{\beta-t}\right)^\alpha$$

$$\text{E}(\ln(X))=\frac{\Gamma'(\alpha)}{\Gamma(\alpha)}-\ln(\beta)$$

$$\theta=\frac{1}{\beta}\mid \sum\Gamma(\alpha_i,\beta)=\Gamma(\sum\alpha_i,\beta)$$

$$\Gamma(1,\beta)=\text{Exponential}(\beta)\mid \Gamma(\frac{n}{2},\frac{1}{2})=\chi_n^2$$

$$\text{Beta}(\alpha,\beta)=\frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)}$$

$$\frac{\alpha}{\alpha+\beta}\mid \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}\mid \varnothing$$

$$\text{B}(a,b)=\frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$$

$$\text{Beta}(1,1)=\text{U}[0,1]\mid \mu_r'=\text{E}(X^r)$$

$$\text{ZIP}(\phi,\lambda):=\text{ZX}|(1-\phi)\lambda|(1-\phi)\lambda(1+\phi\lambda)$$

$$Z=\text{Bernoulli}(1-\phi),X=\text{Poisson}(\lambda)$$

$$\text{M}_{\text{ZIP}}(t)=\phi+(1-\phi)e^{\lambda(e^t-1)}$$

$$\frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}}e^{-\frac{1}{2(1-\rho^2)}(\frac{(x-\mu_1)^2}{\sigma_1^2}-2\rho\frac{(x-\mu_1)(y-\mu_2)}{\sigma_1\sigma_2}+\frac{(y-\mu_2)^2}{\sigma_2^2})}$$

$$X_1|X_2\mid \mu_1+\rho\frac{\sigma_1}{\sigma_2}(x_2-\mu_2)\mid \sigma_1^2(1-\rho^2)$$

$$\text{Skewness}=\frac{\mu_3}{\sigma^3}\mid \text{Kurtosis}=\frac{\mu_4}{\sigma^4}$$

$$\text{P}(g(X)\geq c)\leq \frac{\text{E}(g(x))}{C}$$

$$\text{P}(|X-\mu|\geq c\sigma)\leq \frac{1}{c^2}$$

$$\text{P}(|X-\mu|\geq c)\leq \frac{\text{E}(|X|^r)}{c^r}$$

$$\text{E}(XY)^2\leq \text{E}(X^2)\text{E}(Y^2)$$

$$\text{E}(g(X))\geq g(\text{E}(X)),\forall \text{凸函数 }g(x^2)$$

$$\text{E}(X)=\text{E}[\text{E}(X|Y)]$$

$$\text{Var}(X)=\text{E}[\text{Var}(X|Y)]+\text{Var}[\text{E}(X|Y)]$$

$$\text{q分位数}\xi_q:=\text{F}(\xi_q)=\text{P}(X\leq \xi_q)\geq q$$

$$\text{pgf}:=\text{G}_X(z)=\text{E}(z^X)=\sum_{x\in\mathbb{S}}z^x\text{P}_X(X=x)$$

$$\text{Additivity}:\text{N, Poisson, NB, Binomial}$$

$$IBF:=f_X(x)\propto \frac{f_{X|Y}(x|y_0)}{f_{Y|X}(y_0|x)}$$