# WARING–GOLDBACH PROBLEM: ONE SQUARE, FOUR FIFTH POWERS AND FIFTEEN SIXTH POWERS

RUNBO LI

Abstract. We prove that for every sufficiently large even integer $n$, the equation
$$n = p_1^2 + p_2^5 + p_3^5 + p_4^5 + p_5^5 + p_6^6 + \cdots + p_{20}^6$$
has a solution with all $p_j$ prime.

Contents

1. Introduction 1
2. Preliminaries 1
3. The major arcs 2
4. The minor arcs 5
5. Proof of Theorem 1.1 6
References 6

## 1. Introduction

Let $n$ be a sufficiently large positive integer that satisfies some necessary congruence conditions. Using the Hardy–Littlewood circle method, many mathematicians studied the number of representations of $n$ as the sum of mixed powers of prime numbers. Examples of recent works on this topic include [1], [2], [3], [4], [5], [9] and [11]. In this paper, we prove the following result.

**Theorem 1.1.** *For every sufficiently large even integer $n$, the equation*
$$n = p_1^2 + p_2^5 + p_3^5 + p_4^5 + p_5^5 + p_6^6 + \cdots + p_{20}^6 \tag{1}$$
*has a solution with all $p_j$ prime.*

## 2. Preliminaries

Let $n$ be a sufficiently large positive integer, $P_k = n^{\frac{1}{k}}$ and $L = \log n$. Let
$$\lambda_1 = 1, \quad \lambda_2 = \frac{27}{32}, \quad \lambda_3 = \frac{729}{1024}, \quad \lambda_4 = \frac{19683}{32768}, \quad \lambda_5 = \frac{531441}{1048576}, \quad \lambda_6 = \frac{14348907}{33554432}, \quad \lambda_7 = \frac{387420489}{1073741824},$$
$$\lambda_8 = \frac{10460353203}{34359738368}, \quad \lambda_9 = \frac{9365374360779057135}{36385377104158523392}, \quad \lambda_{10} = \frac{248229150174961425}{1137043034504953856},$$
$$\lambda_{11} = \frac{13208549498258175}{71065189656559616}, \quad \lambda_{12} = \frac{1418162385496725}{8883148707069952}, \quad \lambda_{13} = \frac{1234060169123925}{8883148707069952},$$
$$\lambda_{14} = \lambda_{15} = \frac{560936440510875}{4441574353534976},$$
and write
$$\Lambda = \sum_{j=1}^{15} \lambda_j = \frac{169503075438540046537}{28398343105684701184}, \qquad \Theta = \frac{3}{10} + \frac{\Lambda}{6} = \frac{1103100465143862543341}{851950293170541035520} \approx 1.29479.$$

*2020 Mathematics Subject Classification.* 11P32, 11P55.
*Key words and phrases.* Waring–Goldbach problem, circle method.

We define

$$f_k(\alpha)=\sum_{\frac{1}{2}P_k<p\leqslant P_k} e(\alpha p^k)\log p$$

and, for $1\leqslant j\leqslant 15$,

$$f_{k,j}(\alpha)=\sum_{\frac{1}{2}P_k^{\lambda_j}<p\leqslant P_k^{\lambda_j}} e(\alpha p^k)\log p.$$

We also define

$$F(\alpha)=f_2(\alpha)f_5(\alpha)^4\prod_{j=1}^{15} f_{6,j}(\alpha).$$

Then by the orthogonality of additive characters, the integral

$$\int_0^1 F(\alpha)e(-\alpha n)d\alpha \tag{2}$$

counts the number of solutions of (1) with prime variables lying in certain ranges weighted by a log factor. Our goal is to show the lower bound

$$\int_0^1 F(\alpha)e(-\alpha n)d\alpha \gg n^{\Theta} \tag{3}$$

holds for all even $n$.

Fix $B\geqslant 1$. Put

$$\mathfrak{M}(q,a)=\left\{\alpha\in[0,\ 1]:\left|\alpha-\frac{a}{q}\right|\leqslant\frac{L^B}{n}\right\},\qquad \mathfrak{M}=\bigcup_{1\leqslant q\leqslant L^B}\bigcup_{\substack{0\leqslant a\leqslant q\\(a,q)=1}}\mathfrak{M}(q,a),\qquad \mathfrak{m}=[0,\ 1]\backslash\mathfrak{M}.$$

Then (3) can be obtained if we get

$$\int_{\mathfrak{M}} F(\alpha)e(-\alpha n)d\alpha \gg n^{\Theta} \tag{4}$$

and

$$\int_{\mathfrak{m}} F(\alpha)e(-\alpha n)d\alpha \ll n^{\Theta}L^{-1}. \tag{5}$$

## 3. The major arcs

Define

$$S_k(q,a)=\sum_{\substack{x=1\\(x,q)=1}}^{q} e\left(\frac{ax^k}{q}\right),\qquad v_{k,r}(\beta)=\frac{1}{k}\sum_{2^{-k}n^r<m\leqslant n^r} m^{1/k-1}e(\beta m),\qquad v_k(\beta)=v_{k,1}(\beta),$$

$$U(q,a)=S_2(q,a)S_5(q,a)^4S_6(q,a)^{15},\qquad w(\beta)=v_2(\beta)v_5(\beta)^4\prod_{j=1}^{15}v_{6,\lambda_j}(\beta).$$

**Lemma 3.1.** *Let $q,a$ be integers such as $1\leqslant q\leqslant L^B$, $0\leqslant a\leqslant q$ and $(a,q)=1$. Then for $\alpha\in\mathfrak{M}(q,a)$, we have*

$$\sum_{\frac{1}{2}P_k^r<p\leqslant P_k^r} e(\alpha p^k)\log p=\frac{1}{\varphi(q)}S_k(q,a)v_{k,r}(\alpha-a/q)+O\left(P_k^r\exp(-c\sqrt{L})\right)$$

*for some constant $c>0$.*

*Proof.* This lemma follows from [[6], Lemma 6] and partial summation. □

We shall use this lemma to give an approximation of $F(\alpha)$ using functions $S_k(q,a)$ and $v_{k,r}$ defined above. By Lemma 3.1, we have

$$F(\alpha)=\frac{1}{\varphi(q)^{20}}U(q,a)w(\alpha-a/q)+O\left(n^{1+\Theta}L^{-4B}\right). \tag{6}$$

An integration over $\mathfrak{M}$ gives

$$\int_{\mathfrak{M}}F(\alpha)e(-\alpha n)d\alpha=\sum_{q\leqslant L^B}\frac{A_n(q)}{\varphi(q)^{20}}\int_{-L^B/n}^{L^B/n}w(\beta)e(-\beta n)d\beta+O\left(n^{\Theta}L^{-B}\right), \tag{7}$$

where

$$A_n(q)=\sum_{\substack{a=1\\(a,q)=1}}^{q}U(q,a)e(-an/q).$$

We first show that

$$\int_{-L^B/n}^{L^B/n}w(\beta)e(-\beta n)d\beta\gg n^{\Theta}. \tag{8}$$

**Lemma 3.2.** *([[10], Lemma 6.2]). Suppose that $|\beta|\leqslant\frac{1}{2}$. Then,*

$$v_k(\beta)\ll n^{\frac{1}{k}}(1+n|\beta|)^{-1}.$$

By Lemma 3.2 we know that

$$w(\beta)\ll n^{1+\Theta}(1+n|\beta|)^{-2}. \tag{9}$$

By (9), we get

$$\begin{aligned}
&\int_{-L^B/n}^{L^B/n}w(\beta)e(-\beta n)d\beta\\
=&\int_{-1/2}^{1/2}w(\beta)e(-\beta n)d\beta+O\left(\int_{L^B/n}^{1/2}|w(\beta)|d\beta\right)\\
=&\int_{-1/2}^{1/2}w(\beta)e(-\beta n)d\beta+O\left(n^{\Theta}L^{-B}\right)\\
=&\frac{1}{2\cdot5^4\cdot6^{15}}\sum_{\substack{m_1+\cdots+m_{20}=n\\2^{-2}n<m_1\leqslant n\\2^{-5}n<m_2,m_3,m_4,m_5\leqslant n\\2^{-6}n^{\lambda_{j-5}}<m_j\leqslant n^{\lambda_{j-5}}(6\leqslant j\leqslant 20)}}m_1^{-\frac{1}{2}}(m_2m_3m_4m_5)^{-\frac{4}{5}}\prod_{j=6}^{20}m_j^{-\frac{5}{6}}+O\left(n^{\Theta}L^{-B}\right)\\
\gg&\sum_{\substack{m_1+\cdots+m_{20}=n\\\frac{1}{4}n<m_1\leqslant\frac{1}{2}n\\\frac{1}{32}n<m_2,m_3,m_4\leqslant\frac{3}{64}n\\\frac{1}{64}n^{\lambda_{j-5}}<m_j\leqslant\frac{3}{128}n^{\lambda_{j-5}}(6\leqslant j\leqslant 20)}}m_1^{-\frac{1}{2}}(m_2m_3m_4m_5)^{-\frac{4}{5}}\prod_{j=6}^{20}m_j^{-\frac{5}{6}}\gg n^{\Theta}.
\end{aligned} \tag{10}$$

Next, we want to deal with the sum

$$\sum_{q\leqslant L^B}\frac{A_n(q)}{\varphi(q)^{20}}. \tag{11}$$

Define the series

$$\mathfrak{S}(n)=\sum_{q=1}^{\infty}\frac{A_n(q)}{\varphi(q)^{20}}.$$

**Lemma 3.3.** *([[6], Lemma 5]). For $k\geqslant 2$, we have*

$$S_k(q,a)\ll q^{\frac{1}{2}+\varepsilon}.$$

**Lemma 3.4.** *([10], Lemma 4.3]). For $(a,p)=1$, we have*

$$|S_k(p,a)| \leqslant ((k,p-1)-1)\sqrt{p}+1.$$

**Lemma 3.5.** *([6], Lemma 4]). Let $p^n \mid k$, $p^{n+1} \nmid k$ and*

$$l = \begin{cases} n+2, & p=2, \\ n+1, & p \neq 2. \end{cases}$$

*If $t > l$, then*

$$S_k(p^t,a)=0.$$

By Lemma 3.3, we know that $U(q,a) \ll q^{10+\varepsilon}$ for $(a,q)=1$, and hence $A_n(q) \ll q^{11+\varepsilon}$ uniformly in $n$. Thus, $\mathfrak{S}(n)$ converges absolutely, and

$$\sum_{q \leqslant L^B} \frac{A_n(q)}{\varphi(q)^{20}} = \mathfrak{S}(n) + O\left(L^{-B}\right). \tag{12}$$

Since the complete exponential sums are multiplicative, for $(q_1,q_2)=1$ we have

$$A_n(q_1q_2) = A_n(q_1)A_n(q_2).$$

Moreover, By Lemma 3.5 and the fact that $A_n(4) = S_5(4,a) = 0$, we have $A_n(q)=0$ unless $q$ is square-free. Hence

$$\mathfrak{S}(n) = \prod_p \mathfrak{S}_p(n) = \prod_p \left(1 + \frac{A_n(p)}{(p-1)^{20}}\right). \tag{13}$$

By the orthogonality of additive characters modulo $p$, we have

$$\mathfrak{S}_p(n) = \frac{pM_n(p)}{(p-1)^{20}}, \tag{14}$$

where $M_n(p)$ denote the number of solutions $(x_1,\dots,x_{20}) \in \left(\mathbb{F}_p^{\times}\right)^{20}$ of

$$x_1^2 + x_2^5 + x_3^5 + x_4^5 + x_5^5 + \sum_{j=6}^{20} x_j^6 \equiv n (\mathrm{mod} p). \tag{15}$$

When $p=2$, every unit is 1, and

$$x_1^2 + x_2^5 + x_3^5 + x_4^5 + x_5^5 + \sum_{j=6}^{20} x_j^6 \equiv 0 (\mathrm{mod} 2).$$

Hence $n \equiv 0 (\mathrm{mod} 2)$ is necessary.

When $p=3$, every square and sixth power is 1, and every fifth power is $\pm 1$. It is easy to see that the sum of four fifth powers contains all residue classes modulo 3, and (15) has a solution.

When $p=5$, we have $x^5 = x$ and $x^6 = \pm 1$. Fix $x_1 = 1$. Since $\mathbb{F}_5^{\times} + \mathbb{F}_5^{\times} = \mathbb{F}_5$, the sum of two fifth powers contains all residue classes modulo 5, and (15) has a solution.

When $p=7$, we have $x^6 = 1$. Fix $x_1, x_2, x_3 = 1$. Since $\mathbb{F}_7^{\times} + \mathbb{F}_7^{\times} = \mathbb{F}_7$, the sum of two fifth powers contains all residue classes modulo 7, and (15) has a solution.

When $p \geqslant 11$, we define $\mathcal{A}_k = \mathcal{A}_k(p) = \{x^k : x \in \mathbb{F}_p\}$. It is known that $|\mathcal{A}_k(p)| = \frac{p-1}{(k,p-1)}$. Applying the Cauchy–Davenport theorem repeatedly, we have

$$\begin{aligned} |\mathcal{A}_2 + 4\mathcal{A}_5 + 15\mathcal{A}_6| &\geqslant \min(p, |\mathcal{A}_2| + 4|\mathcal{A}_5| + 15|\mathcal{A}_6| - 19) \\ &\geqslant \min\left(p, \frac{p-1}{2} + \frac{4(p-1)}{5} + \frac{15(p-1)}{6} - 19\right) \\ &\geqslant \min\left(p, \frac{19p-114}{5}\right). \end{aligned}$$

Since $p \geqslant 11$, we have $\frac{19p-114}{5} > p$, and thus $\mathcal{A}_2 + 4\mathcal{A}_5 + 15\mathcal{A}_6 = \mathbb{F}_p$. Now (15) has a solution for every prime $p \geqslant 11$.

Combining above cases, we know that $M_n(p) \geqslant 1$ if $n$ is even, and also $\mathfrak{S}(n) > 0$ by (13). Now, by Lemma 3.4 and trivial estimates, we have

$$\begin{aligned}\left|\frac{A_n(p)}{(p-1)^{20}}\right| &\leqslant \frac{p(\sqrt{p}+1)(4\sqrt{p}+1)^4(5\sqrt{p}+1)^{15}}{(p-1)^{20}}\\ &= \frac{p^{11}}{(p-1)^{20}}\left(1+\frac{1}{\sqrt{p}}\right)\left(4+\frac{1}{\sqrt{p}}\right)^4\left(5+\frac{1}{\sqrt{p}}\right)^{15}\\ &\leqslant \frac{p^{11}}{(p/2)^{20}}\left(1+\frac{1}{\sqrt{p}}\right)\left(4+\frac{1}{\sqrt{p}}\right)^4\left(5+\frac{1}{\sqrt{p}}\right)^{15}\\ &\leqslant \frac{2^{20}\cdot 2\cdot 5^4\cdot 6^{15}}{p^9}\end{aligned}$$

for $p > 10^{10}$. Thus, we get

$$\prod_{p>10^{10}}\left(1+\frac{A_n(p)}{(p-1)^{20}}\right) \geqslant \prod_{p>10^{10}}\left(1-\frac{2^{21}\cdot 5^4\cdot 6^{15}}{p^9}\right) > c \tag{16}$$

with some $c > 0$. By (13), (14), (16) and $M_n(p) \geqslant 1$, we have

$$\mathfrak{S}(n) > c\prod_{p\leqslant 10^{10}}\frac{pM_n(p)}{(p-1)^{20}} > c\prod_{p\leqslant 10^{10}} p^{-19} \gg 1. \tag{17}$$

Now, by (7), (10), (12) and (17), the proof of (4) is completed.

## 4. The minor arcs

We first recall some mean value estimates proved by Andrade [3] and Zhao [12]. Note that we can remove the logarithmic weights and the restriction to primes in $f_k(\alpha)$ and $f_{k,j}(\alpha)$ at the cost of a power of $L$, and deal with the classical Weyl sums.

**Lemma 4.1.** *We have*

$$\int_0^1 |f_2(\alpha)|^2\,|f_5(\alpha)|^6\,d\alpha \ll n^{\frac{6}{5}+\varepsilon}.$$

*Proof.* Taking $k_1 = k_2 = k_3 = 5$ in [[3], Lemma 4], we get Lemma 4.1. □

**Lemma 4.2.** *We have*

$$\int_0^1 \left|\prod_{j=1}^{15} f_{6,j}(\alpha)\right|^2 d\alpha \ll P_6^{\Lambda+\varepsilon}.$$

*Proof.* Taking $t = 1$ and $U = P_6$ in [[12], Lemma 6.1], we get Lemma 4.2. □

Write $\tau = 2^{-100}$. Define

$$\mathcal{E} = \{\alpha \in [0,\ 1] : |f_5(\alpha)| \leqslant P_5^{(\Lambda-4)/2-10\tau}\}.$$

Using Hölder's inequality, together with Lemma 4.1, Lemma 4.2 and the definition of $\mathcal{E}$, we get

$$\begin{aligned}\int_{\mathcal{E}} |F(\alpha)|d\alpha &\leqslant \left(\int_0^1 |f_2(\alpha)|^2\,|f_5(\alpha)|^6\,d\alpha\right)^{\frac{1}{2}}\left(\int_0^1\left|\prod_{j=1}^{15} f_{6,j}(\alpha)\right|^2 d\alpha\right)^{\frac{1}{2}}\sup_{\alpha\in\mathcal{E}}|f_5(\alpha)|\\ &\ll n^{\frac{3}{5}+\varepsilon}P_6^{\Lambda/2+\varepsilon}P_5^{(\Lambda-4)/2-10\tau} \ll n^{\theta-\tau},\end{aligned} \tag{18}$$

where

$$\theta = \frac{441062789418431385223}{340780117268216414208} \approx 1.29427.$$

Note that $\Theta - \theta > 0.0005$.

## 5. Proof of Theorem 1.1

**Lemma 5.1.** *Suppose that* $X^{\frac{1}{24}} \leqslant P \leqslant X^{\frac{9}{20}}$. *Then either we have*

$$\sum_{X<p\leqslant 2X} e(\alpha p^5) \ll X^{\frac{47}{48}+\varepsilon},$$

*or there exist integers* $q$ *and* $a$ *such that* $1 \leqslant q \leqslant P$, $(a,q)=1$ *and* $|q\alpha - a| \leqslant PX^{-5}$.

*Proof.* Taking $k=5$ in [[8], Lemma 2.2], we get Lemma 5.1. □

Now suppose that $\alpha \in [0,\ 1]\backslash\mathcal{E}$. We have $|f_5(\alpha)| > P_5^{(\Lambda-4)/2-10\tau}$ in this case. Taking $X = \frac{1}{2}P_5$ and $P = \frac{1}{32}n^{\frac{1}{119}}$ in Lemma 5.1, we find that the first case will never occur since $\frac{\Lambda-4}{2} - \frac{47}{48} > 0.005$. Hence, we must have

$$[0,\ 1]\backslash\mathcal{E} \subset \mathfrak{R} = \bigcup_{1\leqslant q\leqslant \frac{1}{32}n^{\frac{1}{119}}} \bigcup_{\substack{0\leqslant a\leqslant q\\(a,q)=1}} \mathfrak{R}(q,a), \tag{19}$$

where

$$\mathfrak{R}(q,a) = \{\alpha \in [0,\ 1] : |q\alpha - a| \leqslant n^{-\frac{118}{119}}\}. \tag{20}$$

Define a function $\Upsilon : \mathfrak{R} \to [0,\ 1]$ by

$$\Upsilon(\alpha) = (q + n|q\alpha - a|)^{-1}.$$

Taking $k = 2, 5$ in [[7], Theorem 2] and by partial summation, we have

$$f_2(\alpha) \ll P_2 L^C \Upsilon(\alpha)^{\frac{1}{2}-\varepsilon} \tag{21}$$

and

$$f_5(\alpha) \ll P_5 L^C \Upsilon(\alpha)^{\frac{1}{2}-\varepsilon} \tag{22}$$

for some constant $C > 0$. By (21), (22) and trivial upper bounds for $f_{6,j}(\alpha)$ $(1 \leqslant j \leqslant 15)$, we have

$$F(\alpha) = f_2(\alpha)f_5(\alpha)^4 \prod_{j=1}^{15} f_{6,j}(\alpha) \ll n^{1+\Theta} L^{20C} \Upsilon(\alpha)^{\frac{49}{20}}. \tag{23}$$

Integrating over $\mathfrak{R}\backslash\mathfrak{M}$ gives

$$\int_{\mathfrak{R}\backslash\mathfrak{M}} |F(\alpha)|d\alpha \ll n^{1+\Theta} L^{20C} \int_{\mathfrak{R}\backslash\mathfrak{M}} |\Upsilon(\alpha)|^{\frac{49}{20}} d\alpha \ll n^{\Theta} L^{20C-\frac{9}{20}B+1}. \tag{24}$$

Choosing $B = 200C + 100$ in (24) and by the fact that $\mathfrak{m} \subset \mathcal{E} \cup \mathfrak{R}\backslash\mathfrak{M}$, the proof of (5) is completed. From (4) and (5) we get (3), and the proof of Theorem 1.1 is completed.

## References


[1] G. M. L. Andrade. On some Waring–Goldbach problems. *Ramanujan J.*, 69(67), 2026.
[2] G. M. L. Andrade. On sums of two squares, two cubes, and higher powers of primes. *Bull. Braz. Math. Soc., New Series*, 57(35), 2026.
[3] G. M. L. Andrade. Waring–Goldbach problems for one square and higher powers. *Indag. Math., in press, corrected proof. arXiv e-prints*, page arXiv:2603.05550v1, 2026.
[4] G. M. L. Andrade and H. Godinho. Representing integers as sums of mixed powers of primes. *Indag. Math., in press, corrected proof. arXiv e-prints*, page arXiv:2512.05154v1, 2025.
[5] J. Brüdern. Another Waring–Goldbach problem. *Acta. Arith.*, 210:125–132, 2023.
[6] L.-K. Hua. Some results in the additive prime number theory. *Quart. J. Math. Oxford Ser.*, 9(1):68–80, 1938.
[7] A. V. Kumchev. On Weyl sums over primes and almost primes. *Michigan Math. J.*, 54(2):243–268, 2006.
[8] A. V. Kumchev and T. D. Wooley. On the Waring–Goldbach problem for seventh and higher powers. *Monatsh. Math.*, 183(2):303–310, 2017.
[9] L. Long, J. Li, and M. Zhang. On Waring–Goldbach problem for two squares, three biquadratics, one quintic and one septic. *arXiv e-prints*, page arXiv:2610.15142v1, 2026.
[10] R. C. Vaughan. *The Hardy–Littlewood method, 2nd Edition*, volume 125 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1997.
[11] H. Wang and S. Tian. On Waring–Goldbach problem with mixed powers. *arXiv e-prints*, page arXiv:2608.12397v1, 2026.
[12] L. Zhao. On the Waring–Goldbach problem for fourth and sixth powers. *Proc. London Math. Soc.*, 108(3):1593–1622, 2014.

Department of Mathematics, Imperial College London, London SW7 2AZ, UK
*Email address*: runbo.li26@imperial.ac.uk, runbo.li.carey@gmail.com