

CSci 243 Homework 2

Due: Wednesday, Sept 21

My name

1. Using logical identities and laws, show the logic equivalence of

- (a) (5 points) $p \leftrightarrow q$ and $(p \wedge q) \vee (\neg p \wedge \neg q)$
(b) (5 points) $p \leftrightarrow q$ and $\neg p \leftrightarrow \neg q$

$$\begin{aligned} p &\leftrightarrow q \\ &= (p \rightarrow q) \wedge (q \rightarrow p) \\ &= (\neg p \vee q) \wedge (\neg q \vee p) \end{aligned}$$

2. (10 points) Prove that if $3n + 2$ is even integer then n is even, using

- (a) a proof by contraposition.
(b) a proof by contradiction.

$$(p \wedge q) \vee \neg(p \vee q)$$

(Hint: see a similar set of proofs in the book).

3. (5 points) Disprove the statement that “for any integer $x > 1$, there are positive integers y, z , such that $x^2 = y^2 + z^2$ ”, by finding a counterexample.
4. (5 points) Prove that there are no positive perfect cubes (i.e., x^3) less than 1000 that are the sum of the cubes of two positive integers (i.e., $\exists x, y, z > 0 (x^3 < 1000 \wedge x^3 = y^3 + z^3)$).
5. (5 points) Prove that $3 \cdot 10^{100} + 11$ is not a perfect square **OR** $3 \cdot 10^{100} + 12$ is not a perfect square. Is your proof constructive or nonconstructive?
6. (5 points) Given three numbers, prove that at least one pair of them has nonnegative product.

1. Using logical identities and laws, show the logic equivalence of

(a) (5 points) $p \leftrightarrow q$ and $(p \wedge q) \vee (\neg p \wedge \neg q)$

(b) (5 points) $p \leftrightarrow q$ and $\neg p \leftrightarrow \neg q$

TABLE 8 Logical Equivalences Involving Biconditional Statements.

$$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$$

$$p \leftrightarrow q \equiv \neg p \leftrightarrow \neg q$$

$$p \leftrightarrow q \equiv (p \wedge q) \vee (\neg p \wedge \neg q)$$

$$\neg(p \leftrightarrow q) \equiv p \leftrightarrow \neg q$$

(a)

$$\begin{aligned} ① \quad p \leftrightarrow q &\equiv (p \rightarrow q) \wedge (q \rightarrow p) \\ ② \quad &\equiv (\neg p \vee q) \wedge (\neg q \vee p) \\ &\text{Some thing off} \\ &\equiv (\neg q \rightarrow \neg p) \wedge (\neg p \rightarrow \neg q) \\ &= (q \vee \neg p) \wedge (p \vee \neg q) \\ &= (\neg p \vee q) \wedge (p \vee \neg q) \end{aligned}$$

Distributive?

$$\begin{aligned} &\Rightarrow \neg(p \wedge \neg q) \wedge \neg(\neg p \wedge q) \\ &\equiv \neg(\neg(p \wedge \neg q)) \vee \neg(\neg p \wedge q) \\ &\equiv (p \wedge \neg q) \vee (\neg p \wedge q) \end{aligned}$$

$$\equiv (\neg p \vee q) \wedge (p \vee \neg q)$$

Distribute!

$$\equiv (\neg p \wedge p) \vee (\neg p \wedge \neg q) \vee (q \wedge p) \vee (q \wedge \neg q)$$

Identity

$$\equiv F \vee (\neg p \wedge \neg q) \vee (q \wedge p) \vee F$$

$$\equiv (\neg p \wedge \neg q) \vee (q \wedge p)$$

(b) $p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$

$$\neg p \vee q$$

$$\neg p \Rightarrow \neg q$$

(b)

TABLE 7 Logical Equivalences Involving Conditional Statements.

$$p \rightarrow q \equiv \neg p \vee q$$

$$p \rightarrow q \equiv \neg q \rightarrow \neg p$$

$$p \vee q \equiv \neg p \rightarrow q$$

$$p \wedge q \equiv \neg(p \rightarrow \neg q)$$

$$\neg(p \rightarrow q) \equiv p \wedge \neg q$$

$$(p \rightarrow q) \wedge (p \rightarrow r) \equiv p \rightarrow (q \wedge r)$$

$$(p \rightarrow r) \wedge (q \rightarrow r) \equiv (p \vee q) \rightarrow r$$

$$(p \rightarrow q) \vee (p \rightarrow r) \equiv p \rightarrow (q \vee r)$$

$$(p \rightarrow r) \vee (q \rightarrow r) \equiv (p \wedge q) \rightarrow r$$

TABLE 6 Logical Equivalences.

Equivalence	Name
$p \wedge T \equiv p$	Identity laws
$p \vee F \equiv p$	
$\begin{array}{l} p \vee T \equiv T \\ p \vee F \equiv F \end{array}$	Domination laws
$p \vee p \equiv p$	Idempotent laws
$\neg(\neg p) \equiv p$	Double negation law
$p \vee q \equiv q \vee p$	Commutative laws
$p \wedge q \equiv q \wedge p$	
$(p \vee q) \vee r \equiv p \vee (q \vee r)$	Associative laws
$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$	
$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$	Distributive laws
$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$	
$\neg(p \wedge q) \equiv \neg p \vee \neg q$	De Morgan's laws
$\neg(p \vee q) \equiv \neg p \wedge \neg q$	
$p \vee (p \wedge q) \equiv p$	Absorption laws
$p \wedge (p \vee q) \equiv p$	
$p \vee \neg p \equiv T$	Negation laws
$p \wedge \neg p \equiv F$	

$$\begin{aligned} &\neg(p \vee q) \\ &\equiv \neg(\neg p \wedge \neg q) \end{aligned}$$

=

1. Using logical identities and laws, show the logic equivalence of

(a) (5 points) $p \leftrightarrow q$ and $(p \wedge q) \vee (\neg p \wedge \neg q)$

(b) (5 points) $p \leftrightarrow q$ and $\neg p \leftrightarrow \neg q$

(a)

$$\begin{aligned} \textcircled{1} \quad p \leftrightarrow q &\equiv (p \rightarrow q) \wedge (q \rightarrow p) \\ &\equiv (\neg p \vee q) \wedge (\neg q \vee p) \\ &\equiv (\neg p \vee q) \wedge (p \vee \neg q) \end{aligned}$$

Distribute!

$$\begin{aligned} &\equiv (\neg p \wedge p) \vee (\neg p \wedge \neg q) \vee (q \wedge p) \vee (q \wedge \neg q) \\ &\equiv F \vee (\neg p \wedge \neg q) \vee (q \wedge p) \vee F \\ &\equiv (\neg p \wedge \neg q) \vee (q \wedge p) \end{aligned}$$

Identity

(b)

$$p \leftrightarrow q \cancel{=}$$

$$\begin{aligned} \neg p \leftrightarrow \neg q &\equiv (\neg p \rightarrow \neg q) \wedge (\neg q \rightarrow \neg p) \\ &\equiv (q \rightarrow r) \wedge (r \rightarrow q) \\ &\equiv p \leftrightarrow q \end{aligned}$$

TABLE 8 Logical Equivalences Involving Biconditional Statements.

$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$
$p \leftrightarrow q \equiv \neg p \leftrightarrow \neg q$
$p \leftrightarrow q \equiv (p \wedge q) \vee (\neg p \wedge \neg q)$
$\neg(p \leftrightarrow q) \equiv p \leftrightarrow \neg q$

TABLE 7 Logical Equivalences Involving Conditional Statements.

$p \rightarrow q \equiv \neg p \vee q$
$p \rightarrow q \equiv \neg q \rightarrow \neg p$
$p \vee q \equiv \neg p \rightarrow q$
$p \wedge q \equiv \neg(p \rightarrow \neg q)$
$\neg(p \rightarrow q) \equiv p \wedge \neg q$
$(p \rightarrow q) \wedge (p \rightarrow r) \equiv p \rightarrow (q \wedge r)$
$(p \rightarrow r) \wedge (q \rightarrow r) \equiv (p \vee q) \rightarrow r$
$(p \rightarrow q) \vee (p \rightarrow r) \equiv p \rightarrow (q \vee r)$
$(p \rightarrow r) \vee (q \rightarrow r) \equiv (p \wedge q) \rightarrow r$

TABLE 6 Logical Equivalences.

Equivalence	Name
$p \wedge T \equiv p$	Identity laws
$p \vee F \equiv p$	
$p \vee T \equiv T$	Domination laws
$p \wedge F \equiv F$	
$p \vee p \equiv p$	Idempotent laws
$p \wedge p \equiv p$	
$\neg(\neg p) \equiv p$	Double negation law
$p \vee q \equiv q \vee p$	Commutative laws
$p \wedge q \equiv q \wedge p$	
$(p \vee q) \vee r \equiv p \vee (q \vee r)$	Associative laws
$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$	
$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$	Distributive laws
$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$	
$\neg(p \wedge q) \equiv \neg p \vee \neg q$	De Morgan's laws
$\neg(p \vee q) \equiv \neg p \wedge \neg q$	
$p \vee (p \wedge q) \equiv p$	Absorption laws
$p \wedge (p \vee q) \equiv p$	
$p \vee \neg p \equiv T$	Negation laws
$p \wedge \neg p \equiv F$	

2. (10 points) Prove that if $3n+2$ is even integer then n is even, using

(a) a proof by contraposition.

(b) a proof by contradiction.

(Hint: see a similar set of proofs in the book).

(a)

contraposition

assume q is false \rightarrow show p is false



assume n is not even

↳ adding +2 to an odd number = still an odd #
every odd +2 = odd

so, $3n+2$ shares even/odd nature with $3n$

3 is odd, and assuming n is odd

odd times odd = odd (\downarrow I need to explain more!)

therefore $3n$ is odd and $3n+2$ is odd when n is odd

when n is odd, $3n+2$ is odd

$$\neg q \rightarrow \neg p$$

Therefore if $3n+2$ is even then n is even

$$\neg q \rightarrow \neg p \equiv p \rightarrow q$$

(b)

contradiction: assume p is false, show a contradiction ($\neg r \wedge \neg \neg r$)

assume $3n+2$ is odd

$3n$ is odd b/c same nature with $3n+2$

odd times odd makes an odd
 n has to be odd for $3n$ to be odd

this doesn't show a contradiction or anything

assume that both p and $\neg q$ are true

$\overbrace{3n+2 \text{ is even}}$ $\overbrace{n \text{ is odd}}$



because we wanna prove

$3n+2$ is even $\rightarrow n$ is even

so say that

$3n+2$ is even $\wedge n$ is odd

we show that it is not true

$3n$ is even, b/c same nature like $3n+2$

odd times even makes even



3 times n

so, n is even



$(\neg q \wedge q)$ contradiction

CSci 243 Homework 2

Due: Wednesday, Sept 21

My name

1. Using logical identities and laws, show the logic equivalence of

- (a) (5 points) $p \leftrightarrow q$ and $(p \wedge q) \vee (\neg p \wedge \neg q)$
- (b) (5 points) $p \leftrightarrow q$ and $\neg p \leftrightarrow \neg q$

2. (10 points) Prove that if $3n + 2$ is even integer then n is even, using

- (a) a proof by contraposition.
- (b) a proof by contradiction.

(Hint: see a similar set of proofs in the book).

3. (5 points) Disprove the statement that “for any integer $x > 1$, there are positive integers y, z , such that $x^2 = y^2 + z^2$ ”, by finding a counterexample.
4. (5 points) Prove that there are no positive perfect cubes (i.e., x^3) less than 1000 that are the sum of the cubes of two positive integers (i.e., $\exists x, y, z > 0 (x^3 < 1000 \wedge x^3 = y^3 + z^3)$).
5. (5 points) Prove that $3 \cdot 10^{100} + 11$ is not a perfect square **OR** $3 \cdot 10^{100} + 12$ is not a perfect square. Is your proof constructive or nonconstructive?
6. (5 points) Given three numbers, prove that at least one pair of them has nonnegative product.

counter example

③ $x = 2$

$x^2 = 4$

$y^2 + z^2$

smallest is $y = 1 \quad z = 1$

$y^2 + z^2 = 2$

next is $y = 1 \quad z = 2$

$y^2 + z^2 = 1 + 4 = 5$

already passed 4

$0, 1, 4, 9, 16$

↳ no combo of 2 of these makes 3

$y=0$	$z=0$	$y^2+z^2=0$	$y=2$	$z=0$
$y=0$	$z=1$	= 1	$y^2+z^2=4$	
$y=1$	$z=1$	= 2		
$y=0$	$z=2$	= 4		
$y=1$	$z=2$	= 5		

→ no way to make 5

so let's make $x=3$

1

CSci 243 Homework 2

Due: Wednesday, Sept 21

My name

- Using logical identities and laws, show the logic equivalence of

- (a) (5 points) $p \leftrightarrow q$ and $(p \wedge q) \vee (\neg p \wedge \neg q)$
- (b) (5 points) $p \leftrightarrow q$ and $\neg p \leftrightarrow \neg q$

- Prove that if $3n + 2$ is even integer then n is even, using

- (a) a proof by contraposition.
- (b) a proof by contradiction.

(Hint: see a similar set of proofs in the book).

- Disprove the statement that “for any integer $x > 1$, there are positive integers y, z , such that $x^2 = y^2 + z^2$ ”, by finding a counterexample.
- (5 points) Prove that there are no positive perfect cubes (i.e., x^3) less than 1000 that are the sum of the cubes of two positive integers (i.e., $\exists x, y, z > 0 (x^3 < 1000 \wedge x^3 = y^3 + z^3)$).
- (5 points) Prove that $3 \cdot 10^{100} + 11$ is not a perfect square **OR** $3 \cdot 10^{100} + 12$ is not a perfect square. Is your proof constructive or nonconstructive?
- (5 points) Given three numbers, prove that at least one pair of them has nonnegative product.

④

perfect cubes less than 1000

$x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9$

goal: combine two of these to make another one

$x_n - x_{n-1}$	m_n
$729 - 512 = 217$	
$512 - 343 = 169$	
$343 - 216 = 127$	
$216 - 125 = 91$	
$125 - 64 = 61$	
$64 - 27 = 37$	
$27 - 8 = 19$	
$8 - 1 = 7$	

$$m_n > x_{n-2} + x_{n-3}$$

one of the two cubes we pick,
one has to be x_{n-1} , because
 x_{n-2} combined with anything smaller
is always less than x_n

why?

because $x_n > x_{n-2} + x_{n-3}$ already.
which mean any smaller counter won't $< x_n$ as well.

$x_{n-2} + x_{n-3} = m_n < x_n$
$343 + 216 = 559 < 729$
$216 + 125 = 341 < 512$
$125 + 64 = 189 < 343$
$64 + 27 = 91 < 216$
$27 + 8 = 35 < 125$
$8 + 1 = 9 < 64$
N/A 25
N/A 1

(4)

$$1, 8, 27, 64, 125, 216, 343, 512, 729$$

$$x_1 \quad x_2 \quad x_3 \quad x_4 \quad x_5 \quad x_6 \quad x_7 \quad x_8 \quad x_9$$

goal: combine two of these to make another one

$x_{n-2} + x_{n-3} =$	$t_n < x_n$
$343 + 216 =$	$559 < 729$
$216 + 125 =$	$341 < 512$
$125 + 64 =$	$189 < 343$
$64 + 27 =$	$91 < 216$
$27 + 8 =$	$35 < 125$
$8 + 1 =$	$9 < 64$
N/A	25
N/A	1



$x_n - x_{n-1}$	m_n
$729 - 512 =$	217
$512 - 343 =$	169
$343 - 216 =$	127
$216 - 125 =$	91
$125 - 64 =$	61
$64 - 27 =$	37
$27 - 8 =$	19
$8 - 1 =$	7

here we take the second and third square below the square we want to sum to (x_{n-2}, x_{n-1}) and see that their sum (t_n) is always less than the square we need for (x_n)

this shows that any combination of only the squares below x_n will be smaller than x_n .

therefore, x_{n-1} has to be one of the two squares used in the sum.

here we take the square we want to sum to (x_n) and subtract the next square of a lower value than x_n (x_{n-1})

we see that the differences (m_n) are not equal to any of the other squares of positive integers smaller than x_n .

Because we know that the perfect square x_n has to be one of the numbers used in the sum, and none of the numbers m_n seen here are also perfect squares, there are no possible sums of two perfect squares such that the sum is a perfect square less than 1000.

CSci 243 Homework 2

Due: Wednesday, Sept 21

My name

1. Using logical identities and laws, show the logic equivalence of
 - (a) (5 points) $p \leftrightarrow q$ and $(p \wedge q) \vee (\neg p \wedge \neg q)$
 - (b) (5 points) $p \leftrightarrow q$ and $\neg p \leftrightarrow \neg q$
2. (10 points) Prove that if $3n + 2$ is even integer then n is even, using
 - (a) a proof by contraposition.
 - (b) a proof by contradiction.

(Hint: see a similar set of proofs in the book).

3. (5 points) Disprove the statement that “for any integer $x > 1$, there are positive integers y, z , such that $x^2 = y^2 + z^2$ ”, by finding a counterexample.
4. (5 points) Prove that there are no positive perfect cubes (i.e., x^3) less than 1000 that are the sum of the cubes of two positive integers (i.e., $\exists x, y, z > 0 (x^3 < 1000 \wedge x^3 = y^3 + z^3)$).
5. (5 points) Prove that $3 \cdot 10^{100} + 11$ is not a perfect square **OR** $3 \cdot 10^{100} + 12$ is not a perfect square. Is your proof constructive or nonconstructive?
*existence proof
probably nonconstructive*
6. (5 points) Given three numbers, prove that at least one pair of them has nonnegative product.

(5) $3 \cdot 10^{100} + 11$

if it was a perfect square

$$\sqrt{3 \cdot 10^{100} + 11} = \text{an integer}$$

$$\begin{aligned}\sqrt{3(10^{100} + 4)} &= \text{an integer} \\ \sqrt{3} \sqrt{10^{100} + 4} &= \text{an integer} \\ \sqrt{3} \text{ is irrational} \\ \text{whether } \sqrt{10^{100} + 4} \text{ is rational or irrational, doesn't matter} \\ \text{an irrational} \times \text{rational} &= \text{irrational} \\ \text{an irrational} \times \text{irrational} &= \text{irrational} \\ \text{integers can't be irrational}\end{aligned}$$

CSci 243 Homework 2

Due: Wednesday, Sept 21

My name

1. Using logical identities and laws, show the logic equivalence of
 - (a) (5 points) $p \leftrightarrow q$ and $(p \wedge q) \vee (\neg p \wedge \neg q)$
 - (b) (5 points) $p \leftrightarrow q$ and $\neg p \leftrightarrow \neg q$
2. (10 points) Prove that if $3n + 2$ is even integer then n is even, using
 - (a) a proof by contraposition.
 - (b) a proof by contradiction.
- (Hint: see a similar set of proofs in the book).
3. (5 points) Disprove the statement that “for any integer $x > 1$, there are positive integers y, z , such that $x^2 = y^2 + z^2$ ”, by finding a counterexample.
4. (5 points) Prove that there are no positive perfect cubes (i.e., x^3) less than 1000 that are the sum of the cubes of two positive integers (i.e., $\exists x, y, z > 0 (x^3 < 1000 \wedge x^3 = y^3 + z^3)$).
5. (5 points) Prove that $3 \cdot 10^{100} + 11$ is not a perfect square **OR** $3 \cdot 10^{100} + 12$ is not a perfect square. Is your proof constructive or nonconstructive?
6. (5 points) Given three numbers, prove that at least one pair of them has nonnegative product.

⑥ $x \ y \ z$
prove $xy > 0$ or $yz > 0$ or $xz > 0$

if x, y, z all > 0
all products of pairs are > 0

if one is < 0
the product of the other 2 is > 0

if two are < 0
their product is > 0

if all three are < 0
all products are $>$