

CSci 243 Homework 8

Due: Wednesday, November 9, end of day

****My name****

1. (5 points) Use bubble sort to sort 1, 3, 8, 2, 9, 4, showing the lists obtained at each outer step. *gross but ok*
2. For each of the following program fragments, give an analysis of the running time. You may use summations to evaluate the running times of nested loops. Use Θ notation (i.e., ignore constants) and show your work. *gross like*

(a) (6 points)

```
sum = 0
for i = 0 to n-1
  for j = 0 to i
    for k = 1 to n/(2^j)          (2^j) means 2 to the power j
      sum ++
```

(b) (6 points)

```
for i = 1 to n
  j=1
  while (j <= i) {
    k=1
    while (k <= n) {
      k=k*2
      sum ++
    }
    j=2*j
  }
```

3. (8 points) Suppose that a store offers gift certificates in denominations of \$25 and \$40. Determine the possible total amounts you can form using these gift certificates. Prove your answer using strong induction. *hitch ok I'll watch a TT vid or smth*
4. (10 points) Consider the dot game we saw in class. Again there are two rows of dots, with n_1 and n_2 dots respectively, and players can remove any number of dots during their turn, but only from one row. However, now, the player who removes the last dot loses.

If $n_1 = n_2 > 1$, prove that there is a winning strategy for the second player. What happens if $n_1 \neq n_2$?

I wasn't paying attention in class but ok possibly simpler code and ez

CSci 243 Homework 8

Due: Wednesday, November 9, end of day

****My name****

- (5 points) Use bubble sort to sort 1, 3, 8, 2, 9, 4, showing the lists obtained at each outer step.
- For each of the following program fragments, give an analysis of the running time. You may use summations to evaluate the running times of nested loops. Use Θ notation (i.e., ignore constants) and show your work.

(a) (6 points)

```
sum = 0
for i = 0 to n-1
  for j = 0 to i
    for k = 1 to n/(2^j)
      sum ++
```

1 3 8 2 9 4
 1 3 8 2 9 4

(2^j) means 2 to the power j

(b) (6 points)

```
for i = 1 to n
  j=1
  while (j <= i) {
    k=1
    while (k <= n) {
      k=k*2
      sum ++
    }
    j=2*j
  }
```

1 3 8 2 9 4

1 3 2 8 9 4

1 3 2 8 9 4

1 3 2 8 4 9

1 3 8 2 9 4
 1 3 2 8 9 4
 1 3 2 8 4 9
 1 2 3 8 4 9
 1 2 3 4 8 9

- (8 points) Suppose that a store offers gift certificates in denominations of \$25 and \$40. Determine the possible total amounts you can form using these gift certificates. Prove your answer using strong induction.
- (10 points) Consider the dot game we saw in class. Again there are two rows of dots, with n_1 and n_2 dots respectively, and players can remove any number of dots during their turn, but only from one row. However, now, the player who removes the last dot loses.

If $n_1 = n_2 > 1$, prove that there is a winning strategy for the second player. What happens if $n_1 \neq n_2$?

CSci 243 Homework 8

Due: Wednesday, November 9, end of day

```
**My name**
```

1. (5 points) Use bubble sort to sort 1, 3, 8, 2, 9, 4, showing the lists obtained at each outer step.
2. For each of the following program fragments, give an analysis of the running time. You may use summations to evaluate the running times of nested loops. Use Θ notation (i.e., ignore constants) and show your work.

(a) (6 points)

```
sum = 0
for i = 0 to n-1
  for j = 0 to i
    for k = 1 to n/(2^j)      (2^j) means 2 to the power j
      sum ++
```

(b) (6 points)

```

for i = 1 to n
    j=1
    while (j <= i) {
        k=1
        while (k <= n) {
            k=k*2
            sum ++
        }
        j=2*j
    }
}

```

To prove $P(n)$ true for integer $n \geq 1$

$$(p(1) \wedge (\forall k \geq 1, p(1) \wedge p(2) \wedge \dots \wedge p(k)) \rightarrow p(k+1)) \rightarrow \forall n p(n)$$

Recursion step: verify that $n(1)$ is true

3. (8 points) Suppose that a store offers gift certificates in denominations of \$25 and \$40. Determine the possible total amounts you can form using these gift certificates. Prove your answer using strong induction.

what can I suppose to prove???

4. (10 points) Consider the dot game we saw in class. Again there are two rows of dots, with n_1 and n_2 dots respectively, and players can remove any number of dots during their turn, but only from one row. However, now, the player who removes the last dot loses.

If $n_1 = n_2 > 1$, prove that there is a winning strategy for the second player. What happens if $n_1 \neq n_2$?

well

$$n_1 = n_2 = 2$$

player 1 will remove 1 or 2 dots from a row

if removes 1 dot, p2 can remove 2 dots from other row + win

If remove 2 left, place on remove 1 left from remaining row + with

It for all $2 \leq j \leq k$ p_2 can write

25	75
40	90
50	15 15
65	70 70
75	25 15
80	40
90	5 20
105	25 40 40
125	40 40
100	25 25 25
115	15 25 40
130	65 70 10
145	25 40 40
160	40 40

$u = k+1$
 p1 removes $1 \leq r \leq k+1$ dots from a row
 In if $r > k+1$, p2 can remove k dots from other row, leaving 1 \rightarrow win
 In if $r = k$, p2 can remove all dots from other row, leaving 1 \rightarrow win
~~In if $r < k$, p2 can remove r dots from other row, leaving $k-r$ dots \rightarrow win~~
 In if $r < k$, p2 can remove r dots from other row, leaving $k-r$ dots \rightarrow win

CSci 243 Homework 8

Due: Wednesday, November 9, end of day

****My name****

- (5 points) Use bubble sort to sort 1, 3, 8, 2, 9, 4, showing the lists obtained at each outer step.
- For each of the following program fragments, give an analysis of the running time. You may use summations to evaluate the running times of nested loops. Use Θ notation (i.e., ignore constants) and show your work.

(a) (6 points)

```

sum = 0
for i = 0 to n-1
  for j = 0 to i
    for k = 1 to n/(2^j)          (2^j) means 2 to the power j
      sum ++
  
```

(b) (6 points)

```

for i = 1 to n
  j=1
  while (j <= i) {
    k=1
    while (k <= n) {
      k=k*2
      sum ++
    }
    j=2*j
  }

```

- (8 points) Suppose that a store offers gift certificates in denominations of \$25 and \$40. Determine the possible total amounts you can form using these gift certificates. Prove your answer using strong induction.
- (10 points) Consider the dot game we saw in class. Again there are two rows of dots, with n_1 and n_2 dots respectively, and players can remove any number of dots during their turn, but only from one row. However, now, the player who removes the last dot loses.

If $n_1 = n_2 > 1$, prove that there is a winning strategy for the second player. What happens if $n_1 \neq n_2$?