

Theorems:

Week 1: Theorem 1.1 Let A and B be statements. Then $A \Rightarrow B$ is equivalent to $\overline{A} \vee B$. That is, $(A \Rightarrow B) \Leftrightarrow (\overline{A} \vee B)$.

Theorem 1.2 Let A and B be statements. Then $A \Rightarrow B$ is equivalent to $\overline{B} \Rightarrow \overline{A}$. That is, $(A \Rightarrow B) \Leftrightarrow (\overline{A} \vee B)$.

Theorem 1.3 Let A and B be statements. Then $\overline{A \vee B}$ is equivalent to $\overline{A} \wedge \overline{B}$.

Theorem 2.4 Given any two real numbers $a < b$, there is some $r \in \mathbb{Q}$ such that $a < r < b$.

Theorem 2.5 The number $\sqrt{2}$ is irrational.

Week 2:

Theorem 3.7 -AM-GM Inequality Let $n \in \mathbb{N}$ and $x_1, \dots, x_n \geq 0$. Then $(x_1 \cdot x_2 \cdot \dots \cdot x_n)^{1/n} \leq \frac{(x_1 + \dots + x_n)}{n}$

Theorem 3.8 -Cauchy-Schwarz Inequality Let $n \in \mathbb{N}$, $x_1, \dots, x_n, y_1, \dots, y_n \in \mathbb{R}$. Then $x_1 y_1 + \dots + x_n y_n \leq$

$$\sqrt{x_1^2 + \dots + x_n^2} \sqrt{y_1^2 + \dots + y_n^2}$$

Theorem 3.8 Triangle Inequality For $x, y \in \mathbb{R}$, $|x + y| \leq |x| + |y|$.

Corollary 3.12 Reverse Triangle Inequality $||x| - |y|| \leq |x + y|$. in particular: $|x| - |y| \leq |x + y|$

Theorem 4.1 -Geometric Series Let $x \neq 1$ be a real number. Then for $N \in \mathbb{N}$, $\sum_{n=1}^N x^n = \frac{x-x^{N+1}}{1-x}$ and $\sum_{n=0}^N x^n = \frac{1-x^{N+1}}{1-x}$

Theorem 4.2 Every Integer $n \geq 2$ can be written as a product of primes $n = p_1 \dots p_k$

Week 3: Completeness Axiom for the real numbers: Every nonempty subset of $\mathbb{R}$ that is bounded above has a least upper bound.

Theorem 5.1 the Archimedean property holds. That is, for any $x, y \in \mathbb{R}$ with $x, y > 0$, there is some $n \in \mathbb{N}$ such that $ny > x$.

Theorem 6.6 Let $x > 0$ and $k \in \mathbb{N}$. Then there is a unique $y > 0$ such that $y^k = x$.

Week 4: Theorem 7.1 Monotone Convergence Theorem Let (a_n) be an increasing sequence of real numbers that is bounded above (i.e. there is M so that $a_n \leq M$ for all n). Then (a_n) converges to some limit. If (a_n) is a decreasing sequence and is bounded below, then (a_n) converges to some limit.

Theorem 8.1 Given a sequence (a_n) with $a_n \in 0, 1, \dots, 9$ for all n , the decimal expansion $x = 0.a_1 a_2 \dots a_n \dots$ given by the limit of the sequence $x_n = \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n}$ defines a real number x satisfying $0 \leq x \leq 1$

Theorem 8.2 Let $x \in \mathbb{R}$ satisfy $0 \leq x < 1$. Then there is a sequence of integers (a_n) , with $a_n \in 0, 1, \dots, 9$ for all n , so that if we let $x_n = \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n}$ then $x_n \rightarrow x$ as $n \rightarrow \infty$

Theorem 8.3 Let $0 \leq x < 1$ and suppose that $x = 0.a_1 a_2 \dots = 0.b_1 b_2 \dots$. Let ℓ be the smallest integer for which $a_\ell \neq b_\ell$ and suppose that $a_\ell < b_\ell$. Then $b_\ell = a_\ell + 1$, and for all $k > \ell$ we have $b_k = 0$ and $a_k = 9$

Lemma 8.4 if $x \geq 0$ is rational, then it has a periodic decimal expansion, that is, $x = a_0.a_1 a_2 \dots \overline{a_k b_1 b_2 \dots b_\ell}$

Corollary 8.5 If p and q are positive integers, the rational number p/q has a decimal expansion with period at most q .

Lemma 8.6 If $x \geq 0$ has a periodic decimal expansion, then x is rational.

Theorem 8.7 A number $x \geq 0$ is rational $\Leftrightarrow$ it has a periodic decimal expansion.

Theorem 8.11 -Comparison Test If $b_j \geq 0$ for all j and $\sum_{j=1}^{\infty} b_j$ is convergent, and if $|a_j| \leq b_j$ for all j , then $\sum_{j=1}^{\infty} a_j$ is also convergent, and moreover $|\sum_{j=1}^{\infty} a_j| \leq \sum_{j=1}^{\infty} b_j$

Week 5: Lemma 9.1 For all $u, v \in \mathbb{C}$: (1) $\overline{u+v} = \overline{u} + \overline{v}$ (2) $\overline{uv} = \overline{u} * \overline{v}$ (3) $|u+v| = |u| + |v|$

Theorem 9.2 -De Moivre's If we let $z = r(\cos \theta + i \sin \theta)$, and $n \in \mathbb{N}$ then $z^n = r^n (\cos n\theta + i \sin n\theta)$ $z^{-n} = r^{-n} (\cos -n\theta + i \sin -n\theta) = \frac{1}{r^{-n}} (\cos(n\theta) - i \sin(n\theta))$

Theorem 9.3 -Eulers Formula $e^{i\theta} = \cos(\theta) + i \sin(\theta)$.

Theorem 9.4 -Roots of Unity The solutions to $z^n = 1$ are $1, w, \dots, w^{n-1}$ where $w = e^{\frac{2\pi i}{n}}$ That is, they are $e^{\frac{2\pi k i}{n}}$ for $k = 0, 1, \dots, n-1$.

Theorem 10.1 -Abel-Ruffini There is no formula (like the quadratic formula) for the roots of a polynomial of degree ≥ 5

Theorem 10.2 -Fundamental Theorem of Algebra Any complex polynomial has at least one root in $\mathbb{C}$.

Theorem 10.3 -Factorization Theorem If p is a degree n polynomial, then there are n roots $r_1, \dots, r_n \in \mathbb{C}$ and a number $a \in \mathbb{C}$ so that $p(z) = a(z - r_1)(z - r_2) \dots (z - r_n)$.

Theorem 10.4 -Real Polynomials have conjugate roots If $p(x)$ has *real* coefficients and r is a root, so is $\bar{r}$.

Theorem 10.5 -Root-Coefficient Theorem $p(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$, has roots $r_1, \dots, r_n$ (counting multiplicities), then $r_1 + \dots + r_n = -a_{n-1}$ $r_1 \dots r_n = (-1)^n a_0$. In general, if s_j denotes the sum of all products of j -tuples of the roots (e.g. $s_2 = r_1 r_2 + r_1 r_3 + r_2 r_3 + \dots$), then $s_j = (-1)^j a_{n-j}$.

Week 6: Theorem 11.1 - The Remainder Theorem Let $a \in \mathbb{N}$ and $b \in \mathbb{Z}$. There are unique integers $q \in \mathbb{Z}$ and $0 \leq r < a$ such that $b = qa + r$

Lemma 11.2 If a divides b and b divides a then $a = \pm b$

Lemma 11.3 -Linear Combination Lemma Let a, b , and d be integers. If d divides both a and b , then d divides every integer linear combination of a and b . In other words, for every pair of integers $m, n \in \mathbb{Z}$, the integer d also divides $ma + nb$.

Theorem 11.4 -Bezout's Identity Let a and b be integers. Then $\gcd(a, b)$ is an integer linear combination of a and b . That is, there are integers s and t such that $\gcd(a, b) = sa + tb$.

Corollary 11.5 Let $a, b, c \in \mathbb{Z}$. Suppose $c \neq 0$, c divides ab and $\gcd(a, c) = 1$. Then c divides b .

Corollary 11.6 Let $a, b \in \mathbb{Z}$ and let $d \in \mathbb{Z}$ be such that d divides a and d divides b . Then d divides $\gcd(a, b)$.

Corollary 11.7 Let d be a common divisor of a and b , which is divisible by all divisors of a and b . Then $d = \pm \gcd(a, b)$.

Corollary 11.8 If $a, b \in \mathbb{Z}$, p is prime, and p divides ab , then either p divides a or p divides b (or both).

Corollary 11.9 If $n = p_1^{m_1} \dots p_k^{m_k}$, where each of $p_1, \dots, p_k$ is prime, and if p is a prime number that divides n , then $p = p_i$ for some $i = 1, \dots, k$.

Theorem 11.10 -Fundamental Theorem of Arithmetic Let $n \geq 2$ be an integer. *Existence* Then n is equal to a product $n = p_1^{r_1} \dots p_k^{r_k}$ of powers of prime numbers, where $p_1 < \dots < p_k$ and $r_i \geq 0$ for all i . (Uniqueness) The factorization is unique. That is, if we also have $p_1^{r_1} \dots p_k^{r_k} = n = q_1^{s_1} \dots q_\ell^{s_\ell}$ where $q_1 < \dots < q_\ell$ are primes and $s_i \geq 0$, then $k = \ell$, $p_i = q_i$, and $r_i = s_i$ for all i .

Lemma 11.11 -Existence of prime decomposition, Part 1. Every integer $n \geq 2$ can be written as a product of primes $n = p_1 \dots p_k$.

Lemma 11.12 -Existence of prime decomposition, Part 2. Every integer $n \geq 2$ can be written as a product of powers of primes $n = p_1^{r_1} \dots p_k^{r_k}$ where $p_1 < \dots < p_k$ and $r_i \geq 0$.

Lemma 11.13 -Uniqueness of Prime decomposition. Every integer n can be written as a *unique* product of powers of primes $n = p_1^{r_1} \dots p_k^{r_k}$

Theorem 11.14 Let $n = p_1^{a_1} \dots p_k^{a_k}$ be a prime decomposition. Then m divides n if and only if $m = p_1^{b_1} \dots p_k^{b_k}$ with each $0 \leq b_i \leq a_i$

Theorem 11.15 Let a and b have prime factorizations, $a = p_1^{r_1} \dots p_m^{r_m}$, $b = p_1^{s_1} \dots p_m^{s_m}$. Here p_i 's are distinct, but r_i and s_i are allowed to be zero. Then: (1) $\gcd(a,b) = p_1^{\min(r_1,s_1)} \dots p_m^{\min(r_m,s_m)}$ (2) $\text{lcm}(a,b) = p_1^{\max(r_1,s_1)} \dots p_m^{\max(r_m,s_m)}$ (3) $\text{lcm}(a,b) = ab/\gcd(a,b)$.

Lemma 11.16 Let $r \in \mathbb{Q}$. Then there exist unique coprime integers a and b such that $r = \frac{a}{b}$ and $b \geq 1$.

Theorem 11.17 Let n be a positive integer. Then $\sqrt{n} \in \mathbb{Q}$ if and only if n is a perfect square, that is, $n = m^2$ for some integer m .

Theorem 11.18 If $a, b \in \mathbb{N}$ are coprime, and ab is an n th power, then so are a and b .

Week 7: Theorem 12.1 Let $m \in \mathbb{N}$ and $a, b \in \mathbb{Z}$. The following are equivalent: (1) $a \equiv b \pmod{m}$; (2) m divides $a - b$; (3) a and b have same remainder when divided by m ; (4) $b = qm + a$ for some $q \in \mathbb{Z}$.

Theorem 12.2 We have the following properties for integers $a, b, c \in \mathbb{Z}$ and $m \in \mathbb{N}$: (1) $a \equiv a \pmod{m}$; (2) If $a \equiv b \pmod{m}$, then $b \equiv a \pmod{m}$. (3) If $a \equiv b \pmod{m}$ and $b \equiv c \pmod{m}$, then $a \equiv c \pmod{m}$.

Theorem 12.3 Suppose $a \equiv x \pmod{m}$ and $b \equiv y \pmod{m}$. Then (1) $-a \equiv -x \pmod{m}$; (2) $a + b \equiv x + y \pmod{m}$; (3) $ab \equiv xy \pmod{m}$; (4) for any natural number k , one has $a^k \equiv x^k \pmod{m}$.

Theorem 12.4 Let a and m be coprime. If $x, y \in \mathbb{Z}$ and $xa \equiv ya \pmod{m}$, then $x \equiv y \pmod{m}$. In particular, if p is prime and p does not divide a , then $xa \equiv ya \pmod{p}$ implies that $x \equiv y \pmod{p}$.

Theorem 12.5 The equation $ax \equiv b \pmod{m}$ has a solution if and only if $\gcd(a, m)$ divides b .

Theorem 12.6 Let $m \geq 2$ be a natural number. Then $\mathbb{Z}/m$ is a field if and only if m is prime.

Theorem 12.7 -Fermat's Little Theorem Let a be an integer, and let p be a prime number. If p does not divide a , then $a^{p-1} \equiv 1 \pmod{p}$

Theorem 12.8 -Fermat's Little Theorem Let $n \in \mathbb{N}$ and p be a prime number. Assume n and $p - 1$ are coprime and p does not divide b . Then the equation $x^n \equiv b \pmod{p}$ has exactly one solution in $x \in 1, \dots, p - 1$