

Instrumental Variables

UNDERSTANDING IV



Roadmap

Where do (Good) Instruments Come From?

- True Lotteries

- Natural Experiments

- Panel Data

2SLS Mechanics

- Just-Identified IV

- Overidentification

Weak and Many Instruments

- Weak IVs

- Many IVs

Subtlties of the Validity Condition

To apply IV, we need to make a good case for instrument validity
(note we can always check relevance!)

Subtleties of the Validity Condition

To apply IV, we need to make a good case for instrument validity (note we can always check relevance!)

Consider our simple causal model, $Y_i = \alpha + \beta D_i + \varepsilon_i$. Validity, $Cov(Z_i, \varepsilon_i) = 0$, intuitively requires two distinct assumptions:

Subtleties of the Validity Condition

To apply IV, we need to make a good case for instrument validity (note we can always check relevance!)

Consider our simple causal model, $Y_i = \alpha + \beta D_i + \varepsilon_i$. Validity, $Cov(Z_i, \varepsilon_i) = 0$, intuitively requires two distinct assumptions:

- **As-good-as-random assignment**: individuals with higher/lower potential earnings face the same distribution of Z_i
- **Exclusion**: the “assignment” of Z_i only affects Y_i through D_i

Subtlties of the Validity Condition

To apply IV, we need to make a good case for instrument validity (note we can always check relevance!)

Consider our simple causal model, $Y_i = \alpha + \beta D_i + \varepsilon_i$. Validity, $Cov(Z_i, \varepsilon_i) = 0$, intuitively requires two distinct assumptions:

- **As-good-as-random assignment**: individuals with higher/lower potential earnings face the same distribution of Z_i
- **Exclusion**: the “assignment” of Z_i only affects Y_i through D_i

Confusingly, old-school econometrics texts sometimes refer to $Cov(Z_i, \varepsilon_i) = 0$ as the “exclusion restriction”

Subtleties of the Validity Condition

To apply IV, we need to make a good case for instrument validity (note we can always check relevance!)


Consider our simple causal model, $Y_i = \alpha + \beta D_i + \varepsilon_i$. Validity, $Cov(Z_i, \varepsilon_i) = 0$, intuitively requires two distinct assumptions:

- **As-good-as-random assignment**: individuals with higher/lower potential earnings face the same distribution of Z_i
- **Exclusion**: the “assignment” of Z_i only affects Y_i through D_i


Confusingly, old-school econometrics texts sometimes refer to $Cov(Z_i, \varepsilon_i) = 0$ as the “exclusion restriction”

- More modern IV texts take care to distinguish between these two conceptually distinct requirements...


A Valid Instrument



A Violation of As-Good-As-Random Assignment



A Violation of Exclusion



Where do IVs Come From?

1. *True Lotteries*

One sure-fire way to ensure that Z_i is as-good-as-randomly assigned is...

Where do IVs Come From?

1. *True Lotteries*

One sure-fire way to ensure that Z_i is as-good-as-randomly assigned is... to randomly assign it!

Where do IVs Come From?

1. True Lotteries

One sure-fire way to ensure that Z_i is as-good-as-randomly assigned is... to randomly assign it!

- Some of the best IVs come from lotteries, either run by the researcher (e.g. an RCT) or so-called “natural experiments”
- We still need to worry about violations of the exclusion restriction
- Relevance holds when Z_i has some effect on D_i

Where do IVs Come From?

1. True Lotteries

One sure-fire way to ensure that Z_i is as-good-as-randomly assigned is... to randomly assign it!

- Some of the best IVs come from lotteries, either run by the researcher (e.g. an RCT) or so-called “natural experiments”
- We still need to worry about violations of the exclusion restriction
- Relevance holds when Z_i has some effect on D_i

“Gold standard” IV: a randomized offer to participate in a program, with D_i recording program participation

- Exclusion restriction likely to hold for any Y_i , by construction
- Relevance almost guaranteed (provided people want the program!)

Example

Charter School Lotteries

Abdulkadiroglu et al. (2016) are interested in whether going to a “charter” middle school increases standardized test scores

- Charter students tend to score better, but we worry about selection
- History of doubting educational inputs, since Coleman (1966)

Example

Charter School Lotteries

We leverage an institutional feature of charters: *admission lotteries*

- When more kids want to enroll than there are seats, admission offers $Z_i \in \{0, 1\}$ are effectively drawn from a hat

Example

Charter School Lotteries

We leverage an institutional feature of charters: *admission lotteries*

- When more kids want to enroll than there are seats, admission offers $Z_i \in \{0, 1\}$ are effectively drawn from a hat
- Offers plausibly only affect later test scores Y_i by changing charter enrollment $D_i \in \{0, 1\}$, so are plausibly valid instruments

Example

Charter School Lotteries

We leverage an institutional feature of charters: *admission lotteries*

- When more kids want to enroll than there are seats, admission offers $Z_i \in \{0, 1\}$ are effectively drawn from a hat
- Offers plausibly only affect later test scores Y_i by changing charter enrollment $D_i \in \{0, 1\}$, so are plausibly valid instruments
- We need to control for lottery fixed effects (“risk sets”) to make Z_i as-good-as-randomly assigned – more on this soon

Example

Charter School Lotteries

We leverage an institutional feature of charters: *admission lotteries*

- When more kids want to enroll than there are seats, admission offers $Z_i \in \{0, 1\}$ are effectively drawn from a hat
- Offers plausibly only affect later test scores Y_i by changing charter enrollment $D_i \in \{0, 1\}$, so are plausibly valid instruments
- We need to control for lottery fixed effects (“risk sets”) to make Z_i as-good-as-randomly assigned – more on this soon

We study a particular charter (UP Academy), which is “takeover”

- Two offer IVs: “immediate” (on lottery night) and from a waitlist

Lottery IV Estimates of UP Test Score Effects

TABLE 8—LOTTERY IV ESTIMATES OF UP EFFECTS

| | | 2SLS | | | | |
|----------------------------|------------------|---------------------------------|------------------|---------------------------|-----------------------------|--------------------------|
| | | First stage | | | Enrollment effect (5) | |
| | | Comparison group mean (1) | OLS (2) | Immediate offer (3) | | Waitlist offer (4) |
| <i>Panel A. All grades</i> | | | | | | |
| (Sixth through eighth) | Math (N = 2,202) | 0.059 | 0.301 (0.022) | 0.760 (0.063) | 0.562 (0.067) | 0.270 (0.056) |
| | ELA (N = 2,205) | 0.103 | 0.148 (0.020) | 0.759 (0.063) | 0.562 (0.067) | 0.118 (0.051) |

Where do IVs Come From?

2. *Natural Experiments*

Without appealing to literal randomization, we may credibly argue Z_i is as-good-as-randomly assigned conditional on some \mathbf{W}_i

- Such “natural experiments” rely on a selection-on-observables argument (for Z_i , instead D_i)
- Still worry about exclusion: Z_i cannot affect Y_i except through D_i

Example

Quarter-of-Birth

Angrist and Krueger (1991) famously estimate labor market returns to schooling with a creative IV: student quarter-of-birth

- Compulsory schooling requirements prevent students from dropping before the day they turn 16 (used to be more binding)
- Fixed school start dates mean students who drop out at 16 get more or less schooling depending on their birth date

Example


Quarter-of-Birth

Angrist and Krueger (1991) famously estimate labor market returns to schooling with a creative IV: student quarter-of-birth


- Compulsory schooling requirements prevent students from dropping before the day they turn 16 (used to be more binding)
- Fixed school start dates mean students who drop out at 16 get more or less schooling depending on their birth date
- Quarter-of-birth seems quasi-randomly assigned — is it excludable?
See Buckles and Hungerman (2013)...

The Quarter-of-Birth Natural Experiment: Visualized

A. Average Education by Quarter of Birth (first stage)



B. Average Weekly Wage by Quarter of Birth (reduced form)



Quarter-of-Birth IV Estimates of Returns to Schooling

Table 4.1.1: 2SLS estimates of the economic returns to schooling

| | OLS | | 2SLS | | | |
|---------------------------|-------------------|-------------------|-----------------------|--------------------------------------|-----------------------|-------------------------------|
| | (1) | (2) | (3) | (4) | (5) | (6) |
| Years of education | 0.075 (0.0004) | 0.072 (0.0004) | 0.103 (0.024) | 0.112 (0.021) | 0.106 (0.026) | 0.108 (0.019) |
| <i>Covariates:</i> | | | | | | |
| 9 year of birth dummies | | ✓ | | | ✓ | ✓ |
| 50 state of birth dummies | | ✓ | | | ✓ | ✓ |
| <i>Instruments:</i> | | | | | | |
| | | | dummy for QOB=1 | dummy for QOB=1 or QOB=2 | dummy for QOB=1 | full set of QOB dummies |

Where do IVs Come From?

3. Panel Data

We might also combine IV + difference-in-differences identification

- E.g. instrument with $Z_i \times Post_t$, controlling for Z_i and $Post_t$ FEs
- This requires two parallel trends assumptions, for the RF and FS
- Still need to worry about the exclusion restriction, as always

Example


Charter School Takeovers

In Abdulkadiroglu et al. (2016), we complement the lottery analysis of takeover charters with an instrumented diff-in-diff analysis


- Students enrolled in the “legacy” public school were eligible for being “grandfathered” into UP, without having to apply to the charter
- We compare their trends in test scores & enrollment to a matched comparison group of observably-similar students at other schools

Grandfathering IV: Visualized

Panel A. Score levels



Panel B. Score DD



Grandfathering IV Estimates of UP Test Score Effects

TABLE 7—GRANDFATHERING IV ESTIMATES OF UP EFFECTS

| | | Comparison group mean (1) | OLS (2) | 2SLS | |
|----------------------------|------------------|---------------------------------|------------------|-----------------------|-----------------------------|
| | | | | First stage (3) | Enrollment effect (4) |
| <i>Panel A. All grades</i> | | | | | |
| (Seventh through eighth) | Math (N = 1,543) | −0.233 | 0.400 (0.032) | 1.051 (0.040) | 0.321 (0.039) |
| | ELA (N = 1,539) | −0.214 | 0.296 (0.035) | 1.040 (0.041) | 0.394 (0.044) |

Roadmap

Where do (Good) Instruments Come From?

True Lotteries

Natural Experiments

Panel Data

2SLS Mechanics

Just-Identified IV

Overidentification

Weak and Many Instruments

Weak IVs

Many IVs

Just-Identified IV

The Stata `ivregress`/`ivreg2` commands (or `fixes::feols` in R) allows for controls and multiple treatments / instruments

- When # treatment = # instruments, we say the IV is “just-identified”:

Just-Identified IV

The Stata `ivregress`/`ivreg2` commands (or `fixes::feols` in R) allows for controls and multiple treatments / instruments

- When # treatment = # instruments, we say the IV is “just-identified”:

$$Y_i = \beta D_i + \mathbf{W}_i' \boldsymbol{\gamma} + \varepsilon_i \quad (\text{second stage})$$

$$D_i = \pi Z_i + \mathbf{W}_i' \boldsymbol{\mu} + \eta_i \quad (\text{first stage})$$

where \mathbf{W}_i includes a constant.

Just-Identified IV

The reduced form is:

$$Y_i = \rho Z_i + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$$

Just-Identified IV

The reduced form is:

$$Y_i = \rho Z_i + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$$

Same identification logic as before:

- Validity: $Cov(Z_i, \varepsilon_i) = 0$, allowing $Cov(Z_i, \mathbf{W}_i) \neq 0$

Just-Identified IV

The reduced form is:

$$Y_i = \rho Z_i + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$$

Same identification logic as before:

- Validity: $Cov(Z_i, \varepsilon_i) = 0$, allowing $Cov(Z_i, \mathbf{W}_i) \neq 0$
- Relevance: $\pi \neq 0$, so Z_i and D_i are correlated controlling for \mathbf{W}_i

Just-Identified IV

The reduced form is:

$$Y_i = \rho Z_i + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$$

Same identification logic as before:

- Validity: $Cov(Z_i, \varepsilon_i) = 0$, allowing $Cov(Z_i, \mathbf{W}_i) \neq 0$
- Relevance: $\pi \neq 0$, so Z_i and D_i are correlated controlling for \mathbf{W}_i

IV is still “reduced form over first stage”: $(\beta^{IV} = \rho^{OLS} / \pi^{OLS})$

Just-Identified IV

The reduced form is:

$$Y_i = \rho Z_i + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$$

Same identification logic as before:

- Validity: $Cov(Z_i, \varepsilon_i) = 0$, allowing $Cov(Z_i, \mathbf{W}_i) \neq 0$
- Relevance: $\pi \neq 0$, so Z_i and D_i are correlated controlling for \mathbf{W}_i

IV is still “reduced form over first stage”: $(\beta^{IV} = \rho^{OLS} / \pi^{OLS})$

- Can use Frisch-Waugh-Lovell to “partial out” \mathbf{W}_i from Y_i , X_i , D_i , and so get back to an IV regression without controls

Overidentification

Sometimes we have more than one instrument $Z_{i\ell}$, for $\ell = 1, \dots, L$.

This leads to an “overidentified” IV regression:

$$Y_i = \beta D_i + \mathbf{W}_i' \boldsymbol{\gamma} + \varepsilon_i \quad (\text{second stage})$$

$$D_i = \mathbf{Z}_i' \boldsymbol{\pi} + \mathbf{W}_i' \boldsymbol{\mu} + \eta_i \quad (\text{first stage})$$

where $\mathbf{Z}_i = [Z_{i1}, \dots, Z_{iL}]'$. Reduced form: $Y_i = \mathbf{Z}_i' \boldsymbol{\rho} + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$

Overidentification

Sometimes we have more than one instrument $Z_{i\ell}$, for $\ell = 1, \dots, L$.

This leads to an “overidentified” IV regression:

$$Y_i = \beta D_i + \mathbf{W}_i' \boldsymbol{\gamma} + \varepsilon_i \quad (\text{second stage})$$

$$D_i = \mathbf{Z}_i' \boldsymbol{\pi} + \mathbf{W}_i' \boldsymbol{\mu} + \eta_i \quad (\text{first stage})$$

where $\mathbf{Z}_i = [Z_{i1}, \dots, Z_{iL}]'$. Reduced form: $Y_i = \mathbf{Z}_i' \boldsymbol{\rho} + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$

Validity: $Cov(Z_{i\ell}, \varepsilon_i) = 0$ for all ℓ

- “Overidentified” b/c we could use any $Z_{i\ell}$ to identify $\beta = \boldsymbol{\rho}_\ell / \boldsymbol{\pi}_\ell$

Overidentification

Sometimes we have more than one instrument $Z_{i\ell}$, for $\ell = 1, \dots, L$.

This leads to an “overidentified” IV regression:

$$Y_i = \beta D_i + \mathbf{W}_i' \boldsymbol{\gamma} + \varepsilon_i \quad (\text{second stage})$$

$$D_i = \mathbf{Z}_i' \boldsymbol{\pi} + \mathbf{W}_i' \boldsymbol{\mu} + \eta_i \quad (\text{first stage})$$

where $\mathbf{Z}_i = [Z_{i1}, \dots, Z_{iL}]'$. Reduced form: $Y_i = \mathbf{Z}_i' \boldsymbol{\rho} + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$

Validity: $Cov(Z_{i\ell}, \varepsilon_i) = 0$ for all ℓ

- “Overidentified” b/c we could use any $Z_{i\ell}$ to identify $\beta = \boldsymbol{\rho}_\ell / \boldsymbol{\pi}_\ell$
- Relevance: $\boldsymbol{\pi}_\ell \neq 0$ for at least some ℓ

Overidentification

Sometimes we have more than one instrument $Z_{i\ell}$, for $\ell = 1, \dots, L$.

This leads to an “overidentified” IV regression:

$$Y_i = \beta D_i + \mathbf{W}_i' \boldsymbol{\gamma} + \varepsilon_i \quad (\text{second stage})$$

$$D_i = \mathbf{Z}_i' \boldsymbol{\pi} + \mathbf{W}_i' \boldsymbol{\mu} + \eta_i \quad (\text{first stage})$$

where $\mathbf{Z}_i = [Z_{i1}, \dots, Z_{iL}]'$. Reduced form: $Y_i = \mathbf{Z}_i' \boldsymbol{\rho} + \mathbf{W}_i' \boldsymbol{\kappa} + \nu_i$

Validity: $\text{Cov}(Z_{i\ell}, \varepsilon_i) = 0$ for all ℓ

- “Overidentified” b/c we could use any $Z_{i\ell}$ to identify $\beta = \boldsymbol{\rho}_\ell / \boldsymbol{\pi}_\ell$
- Relevance: $\boldsymbol{\pi}_\ell \neq 0$ for at least some ℓ

Overidentification can yield tests of IV validity

- Intuitively, 2SLS checks whether all the $Z_{i\ell}$ yields the same IV estimate, which is sensible in a constant-effects model...

Putting the “2S” in “2SLS”

You'll notice I haven't actually defined 2SLS beyond the simple case

- Before we had $\beta^{IV} = \frac{\text{Cov}(Z_i, Y_i)}{\text{Cov}(Z_i, D_i)}$ leading to $\hat{\beta}^{IV} = \frac{\widehat{\text{Cov}}(Z_i, Y_i)}{\widehat{\text{Cov}}(Z_i, D_i)}$
- General form follows similarly (as a sample analog) but is notation-heavy, so we won't go into it here

Putting the “2S” in “2SLS”

You'll notice I haven't actually defined 2SLS beyond the simple case

- Before we had $\beta^{IV} = \frac{\text{Cov}(Z_i, Y_i)}{\text{Cov}(Z_i, D_i)}$ leading to $\hat{\beta}^{IV} = \frac{\widehat{\text{Cov}}(Z_i, Y_i)}{\widehat{\text{Cov}}(Z_i, D_i)}$
- General form follows similarly (as a sample analog) but is notation-heavy, so we won't go into it here

A more useful way to define 2SLS is by a two-step procedure:

- First regress D_i on all instruments $Z_{i\ell}$ and controls W_{ik}
- Then regress Y_i on the “fitted values” \hat{D}_i and controls W_{ik}

Putting the “2S” in “2SLS”

You'll notice I haven't actually defined 2SLS beyond the simple case

- Before we had $\beta^{IV} = \frac{\text{Cov}(Z_i, Y_i)}{\text{Cov}(Z_i, D_i)}$ leading to $\hat{\beta}^{IV} = \frac{\widehat{\text{Cov}}(Z_i, Y_i)}{\widehat{\text{Cov}}(Z_i, D_i)}$
- General form follows similarly (as a sample analog) but is notation-heavy, so we won't go into it here

A more useful way to define 2SLS is by a two-step procedure:

- First regress D_i on all instruments $Z_{i\ell}$ and controls W_{ik}
- Then regress Y_i on the “fitted values” \hat{D}_i and controls W_{ik}

The proof of this follows from some (simple) linear algebra

- Intuitively, regressing Y_i on $\hat{\pi}^{OLS} Z_i$ gives a scaled RF:

$$\hat{\beta}^{IV} = \frac{\hat{\rho}^{OLS}}{\hat{\pi}^{OLS}}$$

Avoid Manual 2SLS!

Although easy, you should never do such “manual 2SLS” yourself!

- Your point estimates will be right, but your SEs won't be!
- Also might forget to include some controls in the second stage, etc

Just let Stata/R do everything for you...

2SLS Done Right

IV (2SLS) estimation

Estimates efficient for homoskedasticity only
Statistics robust to heteroskedasticity

| | | | |
|-----------------------|---|-----------------|-----------|
| | | Number of obs = | 69 |
| | | F(2, 66) = | 5.16 |
| | | Prob > F = | 0.0083 |
| Total (centered) SS | = | Centered R2 | = -2.5922 |
| Total (uncentered) SS | = | Uncentered R2 | = 0.3491 |
| Residual SS | = | Root MSE | = 5480 |

```
clear all  
sysuse auto
```

```
ivreg2 price (mpg=rep78) weight, r
```

| price | Coef. | Robust Std. Err. | z | P> z | [95% Conf. Interval] | |
|--------|-----------|---------------------|-------|-------|----------------------|----------|
| mpg | 1404.283 | 1499.569 | 0.94 | 0.349 | -1534.819 | 4343.384 |
| weight | 10.38214 | 8.57869 | 1.21 | 0.226 | -6.431778 | 27.19607 |
| _cons | -55229.89 | 57542.19 | -0.96 | 0.337 | -168010.5 | 57550.73 |

Underidentification test (Kleibergen-Paap rk LM statistic): 1.200
Chi-sq(1) P-val = 0.2734

Weak identification test (Cragg-Donald Wald F statistic): 1.459
(Kleibergen-Paap rk Wald F statistic): 1.083
Stock-Yogo weak ID test critical values: 10% maximal IV size 16.38
15% maximal IV size 8.96
20% maximal IV size 6.66
25% maximal IV size 5.53

Source: Stock-Yogo (2005). Reproduced by permission.

NB: Critical values are for Cragg-Donald F statistic and i.i.d. errors.

Hansen J statistic (overidentification test of all instruments): 0.000
(equation exactly identified)

Instrumented: mpg
Included instruments: weight
Excluded instruments: rep78

Roadmap

Where do (Good) Instruments Come From?

- True Lotteries

- Natural Experiments

- Panel Data

2SLS Mechanics

- Just-Identified IV

- Overidentification

Weak and Many Instruments

- Weak IVs

- Many IVs

Weak Instruments

When running just-identified IV, you should always worry about the “strength” of your instrument

- Specifically the first stage F-statistic, which tests $\pi^{OLS} = 0$

Weak Instruments

When running just-identified IV, you should always worry about the “strength” of your instrument

- Specifically the first stage F-statistic, which tests $\pi^{OLS} = 0$

If π^{OLS} is small relative to its standard error, the IV is “weak”

- Typically use the rule-of-thumb of $F < 10$ (Staiger and Stock 1997)
- In this case the second-stage SEs will be large and the 2SLS estimate will tend to be biased towards the corresponding OLS

Weak Instruments

When running just-identified IV, you should always worry about the “strength” of your instrument

- Specifically the first stage F-statistic, which tests $\pi^{OLS} = 0$

If π^{OLS} is small relative to its standard error, the IV is “weak”


- Typically use the rule-of-thumb of $F < 10$ (Staiger and Stock 1997)
- In this case the second-stage SEs will be large and the 2SLS estimate will tend to be biased towards the corresponding OLS

Much made of this over the years, but Angrist and Kolesár (2022) argue recently that we shouldn’t worry too much

- The SE increase tends to be large enough to “cover up” the bias
- Just-id. 2SLS is “approximately median-unbiased”


Weak Instruments: Visualized

Monte Carlo: $Y_i = \varepsilon_i$, $D_i = \Pi Z_i + \eta_i$: $\Pi = \text{Var}(\varepsilon_i) = \text{Var}(\eta_i) = 1$




Weak Instruments: Visualized

Monte Carlo: $Y_i = \varepsilon_i$, $D_i = \Pi Z_i + \eta_i$: $\Pi = 0.1$ (Weaker)



Weak Instruments: Visualized

Monte Carlo: $Y_i = \varepsilon_i$, $D_i = \Pi Z_i + \eta_i$: $\Pi = 0.01$ (Very Weak)



Many IVs

A more pernicious problem is many-instrument bias, when overid

- Also tends to manifest in low first-stage F 's, so also good to check

Many IVs

A more pernicious problem is many-instrument bias, when overid

- Also tends to manifest in low first-stage F 's, so also good to check

Many-IV bias is also towards OLS. But unlike before, the SEs go *down*

- Intuitively, a more flexible FS tends to fit D_i better \rightarrow more power
- But we can have *overfitting* with lots of $Z_i \rightarrow$ essentially recreate D_i

Many IVs

A more pernicious problem is many-instrument bias, when overid

- Also tends to manifest in low first-stage F 's, so also good to check

Many-IV bias is also towards OLS. But unlike before, the SEs go *down*


- Intuitively, a more flexible FS tends to fit D_i better \rightarrow more power
- But we can have *overfitting* with lots of $Z_i \rightarrow$ essentially recreate D_i

As we'll see, this bias is especially relevant in judge IV designs

- Potentially many judge assignment indicators as the instrument
- Leave-out corrections (e.g. Angrist et al. 1999) have been adapted to this setting in recent years (e.g. Kolesár 2013)


Many Instruments: Visualized

Monte Carlo: $Y_i = \varepsilon_i$, $D_i = \Pi Z_{i1} + \eta_i$: IV with one Z_{i1}




Many Instruments: Visualized

Monte Carlo: $Y_i = \varepsilon_i$, $D_i = \Pi Z_{i1} + \eta_i$: IV with ten Z_{ij}



Many Instruments: Visualized

Monte Carlo: $Y_i = \varepsilon_i$, $D_i = \Pi Z_{i1} + \eta_i$: IV with 100 Z_{ij}



What to Do?

Check your F's after every IV regression

- State of the art: Montiel Olea and Pflueger '15; `weakivtest` in Stata
- Staiger-Stock rule-of-thumb ($F > 10$) still seems widely held
- See Lee et al. (2020) for a recent alternative approach

If your F is small, some things to consider:

- Is there a different instrument that's stronger?
- Is there a better functional form for the instrument you have?
- Do interactions with covariates help? (note: beware many-weak!)
- Does changing the covariate set help? (note: beware invalidity!)