

Engineering topologically protected edge states

Signals in the LDOS: CdGM analogs

Full 2D simulation: band bending and the solid-core model

Disorder-induced mode-mixing: a new mechanism for topology

Conclusions

# Full-shell Majorana nanowires

## A theoretical description

Carlos Payá

Instituto de Ciencia de Materiales de Madrid (ICMM), CSIC

January 10, 2024




# Outline

- ① Engeniering topologically protected edge states**
- ② Signals in the LDOS: CdGM analogs**
- ③ Full 2D simulation: band bending and the solid-core model**
- ④ Disorder-induced mode-mixing: a new mechanism for topology**
- ⑤ Conclusions**

# The Kitaev chain

- ▶ Chain of  $N$  spin-less fermions ( $p$ -wave superconductivity):

$$H = -\mu \sum_{j=1}^N \left( c_j^\dagger c_j - \frac{1}{2} \right) + \sum_{j=1}^{N-1} \left[ -t (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger) \right]$$



R. Aguado 2017, *Rivista del Nuovo Cimento*.  
E. Prada *et al.* 2020, *Nature Reviews Physics*.  
A. Y. Kitaev 2001, *Physics-Uspekhi*.


# The Kitaev chain

- ▶ Chain of  $N$  spin-less fermions ( $p$ -wave superconductivity):

$$H = -\mu \sum_{j=1}^N \left( c_j^\dagger c_j - \frac{1}{2} \right) + \sum_{j=1}^{N-1} \left[ -t (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger) \right]$$

- ▶ Majorana representation:

$$c_j = \frac{1}{2} (\gamma_j^A + i\gamma_j^B), \quad c_j^\dagger = \frac{1}{2} (\gamma_j^A - i\gamma_j^B)$$



R. Aguado 2017, *Rivista del Nuovo Cimento*.  
E. Prada *et al.* 2020, *Nature Reviews Physics*.  
A. Y. Kitaev 2001, *Physics-Uspekhi*.

# The Kitaev chain

- Chain of  $N$  spin-less fermions ( $p$ -wave superconductivity):


$$H = -\mu \sum_{j=1}^N \left( c_j^\dagger c_j - \frac{1}{2} \right) + \sum_{j=1}^{N-1} \left[ -t (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger) \right]$$

- Majorana representation:

$$c_j = \frac{1}{2} (\gamma_j^A + i\gamma_j^B), \quad c_j^\dagger = \frac{1}{2} (\gamma_j^A - i\gamma_j^B)$$


- Hamiltonian in terms of Majorana operators:

$$H = -\frac{i\mu}{2} \sum_{j=1}^N \gamma_j^A \gamma_j^B + \frac{i}{2} \sum_{j=1}^{N-1} [(\Delta + t)\gamma_j^B \gamma_{j+1}^A + (\Delta - t)\gamma_j^A \gamma_{j+1}^B]$$



# Kitaev chain energy dispersion

Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:




c

d

## Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



d

# Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



R. Aguado 2017, *Rivista del Nuovo Cimento*.  
 E. Prada *et al.* 2020, *Nature Reviews Physics*.  
 A. Y. Kitaev 2001, *Physics-Uspekhi*.

# Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



R. Aguado 2017, *Rivista del Nuovo Cimento*.  
E. Prada *et al.* 2020, *Nature Reviews Physics*.  
A. Y. Kitaev 2001, *Physics-Uspekhi*.

# Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



- ▶ Two distinct phases characterized by a  $\mathbb{Z}_2$  invariant,  $M = (-1)^\nu$ .

# Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



- ▶ Two distinct phases characterized by a  $\mathbb{Z}_2$  invariant,  $M = (-1)^\nu$ .
- ▶  $\nu$  is the number of times the energy gap closes in the Brillouin zone.

# Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



- ▶ Two distinct phases characterized by a  $\mathbb{Z}_2$  invariant,  $M = (-1)^\nu$ .
- ▶  $\nu$  is the number of times the energy gap closes in the Brillouin zone.
- ▶  $M = 1 \Rightarrow$  no unpaired MZM.  $M = -1 \Rightarrow$  unpaired MZM (bulk-boundary correspondance).

R. Aguado 2017, *Rivista del Nuovo Cimento*.  
E. Prada et al. 2020, *Nature Reviews Physics*.  
A. Y. Kitaev 2001, *Physics-Uspekhi*.

# Majoranas for qubits



- ▶ MZM are non-Abelian anyons.
- ▶ Gap closing/reopening  $\Rightarrow$  topological protection.

R. Aguado 2017, *Rivista del Nuovo Cimento*.  
 E. Prada *et al.* 2020, *Nature Reviews Physics*.  
 A. Y. Kitaev 2001, *Physics-Uspekhi*.

# We need a $p$ -wave superconductor!

- ▶ The superconducting pairing term in the Kitaev chain is spinless:  
$$\Delta \left( c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger \right).$$

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

# We need a $p$ -wave superconductor!

- ▶ The superconducting pairing term in the Kitaev chain is spinless:  
 $\Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger)$ .
- ▶  $p$ -wave is very rare in nature. We need to engineer it.

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

# We need a $p$ -wave superconductor!

- ▶ The superconducting pairing term in the Kitaev chain is spinless:  
 $\Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger)$ .
- ▶  $p$ -wave is very rare in nature. We need to engineer it.
- ▶ Fu and Kane:  $s$ -wave pairing behaves as  $p$ -wave when projected onto the basis of helical electrons.

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

# We need a $p$ -wave superconductor!


- ▶ The superconducting pairing term in the Kitaev chain is spinless:  
 $\Delta \left( c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger \right)$ .
- ▶  $p$ -wave is very rare in nature. We need to engineer it.
- ▶ Fu and Kane:  $s$ -wave pairing behaves as  $p$ -wave when projected onto the basis of helical electrons.
- ▶ Lutchyn and Oreg: proximitize semiconductors with strong spin-orbit coupling.

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*


Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

# Rashba, Zeeman and helical bands







R. Aguado 2017, *Rivista del Nuovo Cimento*.

# Rashba, Zeeman and helical bands





# Rashba, Zeeman and helical bands




R. Aguado 2017, Rivista del Nuovo Cimento.

# Rashba, Zeeman and helical bands



R. Aguado 2017, Rivista del Nuovo Cimento.


# Topological phase transition



$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*  
R. Aguado 2017, *Rivista del Nuovo Cimento*.

# Topological phase transition




$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

- $V_Z$  comes from  $g$ -factor.

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*  
R. Aguado 2017, *Rivista del Nuovo Cimento*.

# Topological phase transition




$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

- $V_Z$  comes from  $g$ -factor.
- $V_Z \gtrsim \Delta$ .

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

R. Aguado 2017, *Rivista del Nuovo Cimento*.

# Topological phase transition




$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

- ▶  $V_Z$  comes from  $g$ -factor.
- ▶  $V_Z \gtrsim \Delta$ .
- ▶ Need high  $g$ -factor materials and high magnetic fields.

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*  
R. Aguado 2017, *Rivista del Nuovo Cimento*.

# Searching for Majoranas

- ▶ Strong experimental interest.




Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.

Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.



Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.  
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

## ► Drawbacks:

- Strong experimental interest.
- Zero-bias anomalies detected with non-topological explanations.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.

Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

- ▶ Drawbacks:
  - ▶ Multimode effects.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.  
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

- ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.  
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

## ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.  
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

## ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.  
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

## ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.  
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

## ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.
- ▶ Charge and pairing inhomogeneities.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.

Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

## ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.
- ▶ Charge and pairing inhomogeneities.
- ▶ Disorder.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.

Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.


## ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.
- ▶ Charge and pairing inhomogeneities.
- ▶ Disorder.
- ▶ QD physics.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.


Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

# The full-shell nanowire



S. Vaitiekėnas et al. 2020, *Science*.  
 P. San-Jose et al. 2023, *Phys. Rev. B*.  
 C. Payá et al. 2023, *arXiv*.

# The Little-Parks effect



- ▶ Cylinder  $\Leftrightarrow$  vortex.
- ▶ Too thin for full Meissner.
- ▶ Quantized winding of the order parameter:  $\Delta = |\Delta| e^{in\varphi}$ .
- ▶  $n \in \mathbb{Z}$  and jumps every flux quantum  $\Phi_0$ .
- ▶ Quasi-quantization of flux  $\Rightarrow$  pairing presents LP lobes.
- ▶ Depends on  $R$ , SC thickness  $d$  and  $\xi_d$ , the SC coherence length.

W. A. Little and R. D. Parks 1962, *Phys. Rev. Lett.*  
 R. D. Parks and W. A. Little 1964, *Phys. Rev.*

# The full-shell nanowire: analytical hollow-core model




► Effective Zeeman field:

$$V_Z = \phi \left( \frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- $\phi = n - \frac{\Phi}{\Phi_0}$ , magnetic flux.
- $n$  number of fluxoids.
- No need for  $g$ -factor.  $\Phi \sim \Phi_0$ .

S. Vaitiekėnas et al. 2020, *Science*.  
 P. San-Jose et al. 2023, *Phys. Rev. B*.  
 C. Payá et al. 2023, *arXiv*.

# The full-shell nanowire: analytical hollow-core model



- ▶ Good generalized angular momentum  
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$ :

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$

- ▶ Effective Zeeman field:

$$V_Z = \phi \left( \frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶  $\phi = n - \frac{\Phi}{\Phi_0}$ , magnetic flux.
- ▶  $n$  number of fluxoids.
- ▶ No need for  $g$ -factor.  $\Phi \sim \Phi_0$ .

S. Vaitiekėnas et al. 2020, *Science*.  
 P. San-Jose et al. 2023, *Phys. Rev. B*.  
 C. Payá et al. 2023, *arXiv*.

# The full-shell nanowire: analytical hollow-core model



- ▶ Good generalized angular momentum  
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$ :

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$

- ▶ ⇒ Computationally affordable.


- ▶ Effective Zeeman field:

$$V_Z = \phi \left( \frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶  $\phi = n - \frac{\Phi}{\Phi_0}$ , magnetic flux.
- ▶  $n$  number of fluxoids.
- ▶ No need for  $g$ -factor.  $\Phi \sim \Phi_0$ .

S. Vaitiekėnas et al. 2020, *Science*.  
P. San-Jose et al. 2023, *Phys. Rev. B*.  
C. Payá et al. 2023, *arXiv*.

# The full-shell nanowire: analytical hollow-core model



- ▶ Good generalized angular momentum  
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$ :

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$

- ▶ ⇒ Computationally affordable.
- ▶ Easier to understand physics.


- ▶ Effective Zeeman field:

$$V_Z = \phi \left( \frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶  $\phi = n - \frac{\Phi}{\Phi_0}$ , magnetic flux.
- ▶  $n$  number of fluxoids.
- ▶ No need for  $g$ -factor.  $\Phi \sim \Phi_0$ .

S. Vaitiekėnas et al. 2020, *Science*.  
 P. San-Jose et al. 2023, *Phys. Rev. B*.  
 C. Payá et al. 2023, *arXiv*.

# The full-shell nanowire: analytical hollow-core model



- ▶ Effective Zeeman field:

$$V_Z = \phi \left( \frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶  $\phi = n - \frac{\Phi}{\Phi_0}$ , magnetic flux.
- ▶  $n$  number of fluxoids.
- ▶ No need for  $g$ -factor.  $\Phi \sim \Phi_0$ .



- ▶ Good generalized angular momentum  
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$ :

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$

- ▶ ⇒ Computationally affordable.
- ▶ Easier to understand physics.
- ▶ SOC and chemical potential non-tunable.



S. Vaitiekėnas et al. 2020, *Science*.  
P. San-Jose et al. 2023, *Phys. Rev. B*.  
C. Payá et al. 2023, *arXiv*.

# The CdGM analog states





C. Payá et al. 2023, arXiv.  
P. San-Jose et al. 2023, Phys. Rev. B.

# The CdGM analog states



C. Payá et al. 2023, arXiv.  
P. San-Jose et al. 2023, Phys. Rev. B.



# The CdGM analog states



C. Payá et al. 2023, arXiv.



P. San-Jose et al. 2023, Phys. Rev. B.

# The CdGM analog states




C. Payá et al. 2023, arXiv.  
P. San-Jose et al. 2023, Phys. Rev. B.

# The CdGM analog states



C. Payá et al. 2023, arXiv.  
P. San-Jose et al. 2023, Phys. Rev. B.


# LDOS vs. flux



C. Payá et al. 2023, arXiv.

P. San-Jose et al. 2023, Phys. Rev. B.


# LDOS vs. flux



C. Payá et al. 2023, arXiv.

P. San-Jose et al. 2023, Phys. Rev. B.


# The tubular-core model



►  $\frac{\Phi_{dp}}{\Phi_0} = \left( \frac{R_{LP}}{R_{av}} \right)^2$ , CdGM's slope:  $\frac{1}{2mR_{LP}^2} \frac{\Phi}{\Phi_0} m_J \tau_z$  at degeneracy points.



C. Payá et al. 2023, arXiv.

# The tubular-core model




- Phase Diagram just shifts when increasing  $w$ .
- True topological protection only for small islands.

## The modified hollow-core model




- $w \leq 0.5R \Rightarrow$  all physics can be recuperated just with  $R_{av}$

# A solid core simulation: first radial mode




# A solid core simulation: first radial mode




► Conduction band **bends** close to the interface.

# A solid core simulation: first radial mode



- ▶ Conduction band **bends** close to the interface.
- ▶ Different boundary conditions: WF can extend to  $r = 0$ .


# A solid core simulation: first radial mode




- ▶ Conduction band **bends** close to the interface.
- ▶ Different boundary conditions: WF can extend to  $r = 0$ .
- ▶ If all WF are in first radial mode, physics similar to the tubular-core.

C. Payá *et al.* 2023, arXiv.

# A solid core simulation: second radial mode




# A solid core simulation: second radial mode




- When the second radial mode is occupied, the ZEP expands over the full lobe, but CdGMs cover it.

# A solid core simulation: second radial mode




- When the second radial mode is occupied, the ZEP expands over the full lobe, but CdGMs cover it.
- The tubular-core model is not a good approximation anymore.

# More radial modes in the Phase Diagram



- ▶ Notice axis are mean  $\alpha$  and  $U_{min}$ , the minimum of the dome-profile.
- ▶ One wedge per radial mode. No islands outside the first radial mode.

# Topological invariant



- ▶  $N_M$  is the number of MBS.

# Where is topology in the Hamiltonian?

## Hamiltonian

$$\langle m_J | H | m_J \rangle = H_{K,m_J} \tau_z + V_Z \sigma_z + A_{m_J} + C_{m_J} \sigma_z \tau_z + \alpha k_z \sigma_y \tau_z$$

- ▶  $\sigma_i, \tau_i$  Pauli matrices in spin and electron-hole space.
- ▶  $H_{K,m_J}$  is the kinetic term (+ effective chemical potential).
- ▶  $V_Z$  is the effective Zeeman term.
- ▶  $A_{m_J}$  and  $C_{m_J}$  is the coupling of  $J_z$  with the magnetic field and the spin.
- ▶  $\alpha k_z \sigma_y \tau_z$  allows topological transitions when  $m_J = 0$ .


# Topology through mode-mixing

- ▶ A  $\pm m_J$  crossing is parabolic  $\epsilon \sim k_z^2$ .
- ▶ It can be shown that any mode-mixing term  $M \sim \mathbb{I}, \sigma_z, \tau_z$ :

$$\langle m_J | M | -m_J \rangle \sim \alpha k_z.$$

- ▶ ⇒ mode-mixing acts as  $p$ -wave pairing between  $m_J \leftrightarrow -m_J$  states.

# Shaping the wave-function with radial harmonics




$$R(\varphi) = R_0$$

$$H = \begin{pmatrix} \ddots & & & \\ & h_{i,i} & & \\ & & \ddots & \\ & & & h_{i+\ell,i+\ell} \\ & & & & \ddots \end{pmatrix}$$

# Shaping the wave-function with radial harmonics


$$R(\varphi) = R_0 + \operatorname{Re} \sum_{\ell=1}^{\ell_{\max}} \delta R_\ell e^{i\ell\varphi}$$

# Shaping the wave-function with radial harmonics




$$H = \begin{pmatrix} \ddots & & & \\ & h_{i,i} & \dots & h_{i,i+\ell} \\ \vdots & \ddots & \ddots & \vdots \\ h_{i+\ell,i} & \dots & h_{i+\ell,i+\ell} \\ & & \ddots & \ddots \end{pmatrix}$$

# Shaping the wave-function with radial harmonics



$$H = \begin{pmatrix} \ddots & & & \\ & h_{i,i} & \dots & h_{i,i+\ell} \\ \vdots & \ddots & \ddots & \vdots \\ h_{i+\ell,i} & \dots & h_{i+\ell,i+\ell} \\ & & \ddots & \ddots \end{pmatrix}$$

# Shaping the wave-function with radial harmonics




$$R(\varphi) = R_0 + \operatorname{Re} \sum_{\ell=1}^{\ell_{\max}} \delta R_\ell e^{i\ell\varphi}$$

$$\langle m_J | H | m_J + \ell \rangle = h_{m_J, m_J + \ell}(\ell, \delta R_\ell)$$

$$\delta R_\ell \in \mathbb{C}$$


$$\ell \in \mathbb{N}$$

# Effects on the LDOS




- Unperturbed cylinder.
- No topological protection.

# Effects on the LDOS




- ▶ Smooth distortion  $\sim$  defects in the nanowire profile.
- ▶ All  $m_J$  modes interact with each other, opening gaps at 0 energy or creating new MZM.
- ▶ Topology is now possible in all lobes, as it can origin from any  $m_J$  mode.

# Effects on the LDOS




- ▶ Non-smooth distortion  $\sim$  defects in the nanowire profile + atomic size defects.
- ▶ Topological minigaps are larger because harmonic pre-factors can be larger.

# Effects on the LDOS



C. Payá *et al.* 2023, arXiv.

# Tubular-core




- Follows a simple equation:

$$(\mu_{m_J} - C_{m_J})^2 - (A_{m_J} + V_Z)^2 + \Delta^2 = 0 \xrightarrow[m_J=0]{} V_Z = \sqrt{\Delta^2 + \mu_0^2}$$

- Valid for any disorder model.

# Solid-core



► Independent of the disorder model.

C. Payá et al. 2023, arXiv.

Engineering topologically protected edge states

Signals in the LDOS: CdGM analogs

Full 2D simulation: band bending and the solid-core model

Disorder-induced mode-mixing: a new mechanism for topology

Conclusions


Summary

Messages



# Summary




# Summary






# Summary




# Summary





# Summary




# Summary








# Summary



# Summary




# Summary



+ mode-mixing  
disorder

# Summary



# Conclusions

- ▶ In pristine full-shell hybrid nanowires:
  1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.

# Conclusions

- ▶ In pristine full-shell hybrid nanowires:
  1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
  2. MZMs are generally topologically unprotected except for small islands in parameter space.

# Conclusions

- ▶ In pristine full-shell hybrid nanowires:
  1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
  2. MZMs are generally topologically unprotected except for small islands in parameter space.
  3. Tubular-core nanowires are a good experimental candidate for protected MZMs.

# Conclusions

- ▶ In pristine full-shell hybrid nanowires:
  1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
  2. MZMs are generally topologically unprotected except for small islands in parameter space.
  3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
  4. The solid-core phenomenology is more complex and depends on the radial modes.

# Conclusions

- ▶ In pristine full-shell hybrid nanowires:
  1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
  2. MZMs are generally topologically unprotected except for small islands in parameter space.
  3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
  4. The solid-core phenomenology is more complex and depends on the radial modes.
- ▶ Adding mode-mixing:

# Conclusions

- ▶ In pristine full-shell hybrid nanowires:
  1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
  2. MZMs are generally topologically unprotected except for small islands in parameter space.
  3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
  4. The solid-core phenomenology is more complex and depends on the radial modes.
- ▶ Adding mode-mixing:
  5. Mode-mixing induced by disorder behaves as an effective  $p$ -wave pairing.

# Conclusions

- ▶ In pristine full-shell hybrid nanowires:
  1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
  2. MZMs are generally topologically unprotected except for small islands in parameter space.
  3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
  4. The solid-core phenomenology is more complex and depends on the radial modes.
- ▶ Adding mode-mixing:
  5. Mode-mixing induced by disorder behaves as an effective  $p$ -wave pairing.
  6. Generic disorder generates new MZMs and opens topological minigaps.

# Conclusions

## Take home message

*Majorana physics of full-shell nanowires is very rich. For pristine configurations, the tubular-core model is the optimal candidate but, in the presence of mode-mixing, half of the parameter space is suitable for topologically protected Majorana bound states.*

# Full-shell Majorana nanowires

## A theoretical description

Carlos Payá

Instituto de Ciencia de Materiales de Madrid (ICMM), CSIC


January 10, 2024



Cylindrical nanowire  
Mode-mixing

Hollow-core  
Modified hollow-core  
Tubular-core



# Hollow-core results




C. Payá *et al.* 2023, arXiv.

# Modified hollow-core results


a




c




d




e



# Destructive Little-Parks




# Hexagonal wave-function



- ▶ New red stripes. Hexagon has  $\ell = 6$ .
- ▶ Upper stripe:  $m_J = 0$  mixes with  $m_J = \pm 6$ .
- ▶ Lower stripe:  $m_J = 3$  mixes with  $m_J = -3$ .
- ▶ The MZM coming from  $m_J = \pm 3$  **cannot** interact with  $m_J = 0 \Rightarrow$  they overlap.
- ▶ The  $m_J = \pm 6$  MZM annihilates the  $m_J = 0$  MZM.

# Hexagonal wave-function



- Except for the new topological stripes and a region where the MZM splits, the system is equivalent to the cylinder.

C. Payá *et al.* 2023, arXiv.

# Full-shell Majorana nanowires

## A theoretical description

Carlos Payá

Instituto de Ciencia de Materiales de Madrid (ICMM), CSIC

January 10, 2024

