

Engineering topologically protected edge states

Signals in the LDOS: CdGM analogs

Full 2D simulation: band bending and the solid-core model

Disorder-induced mode-mixing: a new mechanism for topology

Conclusions

Full-shell Majorana nanowires

A theoretical description

Carlos Payá

Instituto de Ciencia de Materiales de Madrid (ICMM), CSIC

January 10, 2024




Outline

- ① Engeniering topologically protected edge states**
- ② Signals in the LDOS: CdGM analogs**
- ③ Full 2D simulation: band bending and the solid-core model**
- ④ Disorder-induced mode-mixing: a new mechanism for topology**
- ⑤ Conclusions**

The Kitaev chain

- ▶ Chain of N spin-less fermions (p -wave superconductivity):

$$H = -\mu \sum_{j=1}^N \left(c_j^\dagger c_j - \frac{1}{2} \right) + \sum_{j=1}^{N-1} \left[-t (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger) \right]$$



R. Aguado 2017, *Rivista del Nuovo Cimento*.
E. Prada *et al.* 2020, *Nature Reviews Physics*.
A. Y. Kitaev 2001, *Physics-Uspekhi*.


The Kitaev chain

- ▶ Chain of N spin-less fermions (p -wave superconductivity):

$$H = -\mu \sum_{j=1}^N \left(c_j^\dagger c_j - \frac{1}{2} \right) + \sum_{j=1}^{N-1} \left[-t (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger) \right]$$

- ▶ Majorana representation:

$$c_j = \frac{1}{2} (\gamma_j^A + i\gamma_j^B), \quad c_j^\dagger = \frac{1}{2} (\gamma_j^A - i\gamma_j^B)$$



R. Aguado 2017, *Rivista del Nuovo Cimento*.
E. Prada *et al.* 2020, *Nature Reviews Physics*.
A. Y. Kitaev 2001, *Physics-Uspekhi*.

The Kitaev chain

- Chain of N spin-less fermions (p -wave superconductivity):


$$H = -\mu \sum_{j=1}^N \left(c_j^\dagger c_j - \frac{1}{2} \right) + \sum_{j=1}^{N-1} \left[-t (c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j) + \Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger) \right]$$

- Majorana representation:

$$c_j = \frac{1}{2} (\gamma_j^A + i\gamma_j^B), \quad c_j^\dagger = \frac{1}{2} (\gamma_j^A - i\gamma_j^B)$$


- Hamiltonian in terms of Majorana operators:

$$H = -\frac{i\mu}{2} \sum_{j=1}^N \gamma_j^A \gamma_j^B + \frac{i}{2} \sum_{j=1}^{N-1} [(\Delta + t)\gamma_j^B \gamma_{j+1}^A + (\Delta - t)\gamma_j^A \gamma_{j+1}^B]$$



Kitaev chain energy dispersion

Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:




c

d

Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



d

Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



R. Aguado 2017, *Rivista del Nuovo Cimento*.
 E. Prada *et al.* 2020, *Nature Reviews Physics*.
 A. Y. Kitaev 2001, *Physics-Uspekhi*.

Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



R. Aguado 2017, *Rivista del Nuovo Cimento*.
E. Prada *et al.* 2020, *Nature Reviews Physics*.
A. Y. Kitaev 2001, *Physics-Uspekhi*.

Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



- ▶ Two distinct phases characterized by a \mathbb{Z}_2 invariant, $M = (-1)^\nu$.

Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



- ▶ Two distinct phases characterized by a \mathbb{Z}_2 invariant, $M = (-1)^\nu$.
- ▶ ν is the number of times the energy gap closes in the Brillouin zone.

Kitaev chain energy dispersion


Let's consider periodic boundary conditions and solve the eigenvalue problem in momentum space:



- ▶ Two distinct phases characterized by a \mathbb{Z}_2 invariant, $M = (-1)^\nu$.
- ▶ ν is the number of times the energy gap closes in the Brillouin zone.
- ▶ $M = 1 \Rightarrow$ no unpaired MZM. $M = -1 \Rightarrow$ unpaired MZM (bulk-boundary correspondance).

R. Aguado 2017, *Rivista del Nuovo Cimento*.
E. Prada et al. 2020, *Nature Reviews Physics*.
A. Y. Kitaev 2001, *Physics-Uspekhi*.

Majoranas for qubits



- ▶ MZM are non-Abelian anyons.
- ▶ Gap closing/reopening \Rightarrow topological protection.

R. Aguado 2017, *Rivista del Nuovo Cimento*.
 E. Prada *et al.* 2020, *Nature Reviews Physics*.
 A. Y. Kitaev 2001, *Physics-Uspekhi*.

We need a p -wave superconductor!

- ▶ The superconducting pairing term in the Kitaev chain is spinless:
$$\Delta \left(c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger \right).$$

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

We need a p -wave superconductor!

- ▶ The superconducting pairing term in the Kitaev chain is spinless:
 $\Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger)$.
- ▶ p -wave is very rare in nature. We need to engineer it.

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

We need a p -wave superconductor!

- ▶ The superconducting pairing term in the Kitaev chain is spinless:
 $\Delta (c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger)$.
- ▶ p -wave is very rare in nature. We need to engineer it.
- ▶ Fu and Kane: s -wave pairing behaves as p -wave when projected onto the basis of helical electrons.

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

We need a p -wave superconductor!


- ▶ The superconducting pairing term in the Kitaev chain is spinless:
 $\Delta \left(c_j c_{j+1} + c_{j+1}^\dagger c_j^\dagger \right)$.
- ▶ p -wave is very rare in nature. We need to engineer it.
- ▶ Fu and Kane: s -wave pairing behaves as p -wave when projected onto the basis of helical electrons.
- ▶ Lutchyn and Oreg: proximitize semiconductors with strong spin-orbit coupling.

L. Fu and C. L. Kane 2008, *Phys. Rev. Lett.*

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*


Y. Oreg, G. Refael, and F. von Oppen 2010, *Phys. Rev. Lett.*

Rashba, Zeeman and helical bands







R. Aguado 2017, *Rivista del Nuovo Cimento*.

Rashba, Zeeman and helical bands





Rashba, Zeeman and helical bands




R. Aguado 2017, Rivista del Nuovo Cimento.

Rashba, Zeeman and helical bands



R. Aguado 2017, Rivista del Nuovo Cimento.


Topological phase transition



$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*
R. Aguado 2017, *Rivista del Nuovo Cimento*.

Topological phase transition




$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

- V_Z comes from g -factor.

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*
R. Aguado 2017, *Rivista del Nuovo Cimento*.

Topological phase transition




$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

- V_Z comes from g -factor.
- $V_Z \gtrsim \Delta$.

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*

R. Aguado 2017, *Rivista del Nuovo Cimento*.

Topological phase transition




$$V_{Zc} = \sqrt{\Delta^2 + \mu^2}$$

- ▶ V_Z comes from g -factor.
- ▶ $V_Z \gtrsim \Delta$.
- ▶ Need high g -factor materials and high magnetic fields.

R. M. Lutchyn, J. D. Sau, and S. Das Sarma 2010, *Phys. Rev. Lett.*
R. Aguado 2017, *Rivista del Nuovo Cimento*.

Searching for Majoranas


- ▶ Strong experimental interest.



Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.



Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

► Drawbacks:

- Strong experimental interest.
- Zero-bias anomalies detected with non-topological explanations.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.

Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

- ▶ Drawbacks:
 - ▶ Multimode effects.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

- ▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.
Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.
- ▶ Charge and pairing inhomogeneities.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.

Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.

▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.
- ▶ Charge and pairing inhomogeneities.
- ▶ Disorder.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.

Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

Searching for Majoranas

- ▶ Strong experimental interest.
- ▶ Zero-bias anomalies detected with non-topological explanations.


▶ Drawbacks:

- ▶ Multimode effects.
- ▶ Electrostatic environment.
- ▶ Renormalized parameters.
- ▶ High magnetic fields.
- ▶ Orbital effects.
- ▶ Charge and pairing inhomogeneities.
- ▶ Disorder.
- ▶ QD physics.

Claims: V. Mourik *et al.* 2012, *Science*. S. M. Albrecht *et al.* 2016, *Nature*. M. T. Deng *et al.* 2016, *Science*.


Trivial explanations: E. J. H. Lee *et al.* 2012, *Phys. Rev. Lett.* M. Valentini, F. Peñaranda, *et al.* 2021, *Science*. M. Valentini, M. Borovkov, *et al.* 2022, *Nature*.

The full-shell nanowire



S. Vaitiekėnas et al. 2020, *Science*.
 P. San-Jose et al. 2023, *Phys. Rev. B*.
 C. Payá et al. 2023, *arXiv*.

The Little-Parks effect



- ▶ Cylinder \Leftrightarrow vortex.
- ▶ Too thin for full Meissner.
- ▶ Quantized winding of the order parameter: $\Delta = |\Delta| e^{in\varphi}$.
- ▶ $n \in \mathbb{Z}$ and jumps every flux quantum Φ_0 .
- ▶ Quasi-quantization of flux \Rightarrow pairing presents LP lobes.
- ▶ Depends on R , SC thickness d and ξ_d , the SC coherence length.

W. A. Little and R. D. Parks 1962, *Phys. Rev. Lett.*
 R. D. Parks and W. A. Little 1964, *Phys. Rev.*

The full-shell nanowire: analytical hollow-core model




- ▶ **Effective Zeeman field:**

$$V_Z = \phi \left(\frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶ $\phi = n - \frac{\Phi}{\Phi_0}$, magnetic flux.
- ▶ n number of fluxoids.
- ▶ No need for g -factor. $\Phi \sim \Phi_0$.

S. Vaitiekėnas et al. 2020, *Science*.
 P. San-Jose et al. 2023, *Phys. Rev. B*.
 C. Payá et al. 2023, *arXiv*.

The full-shell nanowire: analytical hollow-core model



- ▶ Good generalized angular momentum
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$:

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$

- ▶ Effective Zeeman field:

$$V_Z = \phi \left(\frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶ $\phi = n - \frac{\Phi}{\Phi_0}$, magnetic flux.
- ▶ n number of fluxoids.
- ▶ No need for g -factor. $\Phi \sim \Phi_0$.

S. Vaitiekėnas et al. 2020, *Science*.
 P. San-Jose et al. 2023, *Phys. Rev. B*.
 C. Payá et al. 2023, *arXiv*.

The full-shell nanowire: analytical hollow-core model



- ▶ Good generalized angular momentum
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$:

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$

- ▶ ⇒ Computationally affordable.


- ▶ Effective Zeeman field:

$$V_Z = \phi \left(\frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶ $\phi = n - \frac{\Phi}{\Phi_0}$, magnetic flux.
- ▶ n number of fluxoids.
- ▶ No need for g -factor. $\Phi \sim \Phi_0$.

S. Vaitiekėnas et al. 2020, *Science*.
P. San-Jose et al. 2023, *Phys. Rev. B*.
C. Payá et al. 2023, *arXiv*.

The full-shell nanowire: analytical hollow-core model



- ▶ Good generalized angular momentum
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$:

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$

- ▶ ⇒ Computationally affordable.
- ▶ Easier to understand physics.

- ▶ Effective Zeeman field:

$$V_Z = \phi \left(\frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶ $\phi = n - \frac{\Phi}{\Phi_0}$, magnetic flux.
- ▶ n number of fluxoids.
- ▶ No need for g -factor. $\Phi \sim \Phi_0$.

S. Vaitiekėnas et al. 2020, *Science*.
 P. San-Jose et al. 2023, *Phys. Rev. B*.
 C. Payá et al. 2023, *arXiv*.

The full-shell nanowire: analytical hollow-core model



- ▶ Effective Zeeman field:

$$V_Z = \phi \left(\frac{1}{4mR^2} + \frac{\alpha}{2R} \right)$$

- ▶ $\phi = n - \frac{\Phi}{\Phi_0}$, magnetic flux.
- ▶ n number of fluxoids.
- ▶ No need for g -factor. $\Phi \sim \Phi_0$.



- ▶ Good generalized angular momentum
 $J_z = -i\partial_\varphi + \frac{1}{2}\sigma_z + \frac{1}{2}n\tau_z$:

$$m_J = \begin{cases} \mathbb{Z} + 1/2, & \text{if } n \text{ even} \\ \mathbb{Z}, & \text{if } n \text{ odd} \end{cases}$$



- ▶ ⇒ Computationally affordable.
- ▶ Easier to understand physics.
- ▶ SOC and chemical potential non-tunable.

S. Vaitiekėnas et al. 2020, *Science*.
P. San-Jose et al. 2023, *Phys. Rev. B*.
C. Payá et al. 2023, *arXiv*.

The CdGM analog states





The CdGM analog states



C. Payá et al. 2023, arXiv.
P. San-Jose et al. 2023, Phys. Rev. B.



The CdGM analog states



C. Payá et al. 2023, arXiv.



P. San-Jose et al. 2023, Phys. Rev. B.

The CdGM analog states



C. Payá et al. 2023, arXiv.
P. San-Jose et al. 2023, Phys. Rev. B.


The CdGM analog states



C. Payá et al. 2023, arXiv.

P. San-Jose et al. 2023, Phys. Rev. B.


LDOS vs. flux



C. Payá et al. 2023, arXiv.

P. San-Jose et al. 2023, Phys. Rev. B.


LDOS vs. flux



C. Payá et al. 2023, arXiv.

P. San-Jose et al. 2023, Phys. Rev. B.


The tubular-core model



► $\frac{\Phi_{dp}}{\Phi_0} = \left(\frac{R_{LP}}{R_{av}} \right)^2$, CdGM's slope: $\frac{1}{2mR_{LP}^2} \frac{\Phi}{\Phi_0} m_J \tau_z$ at degeneracy points.



C. Payá *et al.* 2023, arXiv.

The tubular-core model




- Phase Diagram just shifts when increasing w .
- True topological protection only for small islands.

The modified hollow-core model




- $w \leq 0.5R \Rightarrow$ all physics can be recuperated just with R_{av}

A solid core simulation: first radial mode




A solid core simulation: first radial mode




- ▶ Conduction band **bends** close to the interface.

A solid core simulation: first radial mode



- ▶ Conduction band **bends** close to the interface.
- ▶ Different boundary conditions: WF can extend to $r = 0$.


A solid core simulation: first radial mode




- ▶ Conduction band **bends** close to the interface.
- ▶ Different boundary conditions: WF can extend to $r = 0$.
- ▶ If all WF are in first radial mode, physics similar to the tubular-core.

C. Payá *et al.* 2023, arXiv.

A solid core simulation: second radial mode




A solid core simulation: second radial mode




- When the second radial mode is occupied, the ZEP expands over the full lobe, but CdGMs cover it.

A solid core simulation: second radial mode




- When the second radial mode is occupied, the ZEP expands over the full lobe, but CdGMs cover it.
- The tubular-core model is not a good approximation anymore.

More radial modes in the Phase Diagram



- ▶ Notice axis are mean α and U_{min} , the minimum of the dome-profile.
- ▶ One wedge per radial mode. No islands outside the first radial mode.

Topological invariant



- ▶ N_M is the number of MBS.

Where is topology in the Hamiltonian?

Hamiltonian

$$\langle m_J | H | m_J \rangle = H_{K,m_J} \tau_z + V_Z \sigma_z + A_{m_J} + C_{m_J} \sigma_z \tau_z + \alpha k_z \sigma_y \tau_z$$

- ▶ σ_i, τ_i Pauli matrices in spin and electron-hole space.
- ▶ H_{K,m_J} is the kinetic term (+ effective chemical potential).
- ▶ V_Z is the effective Zeeman term.
- ▶ A_{m_J} and C_{m_J} is the coupling of J_z with the magnetic field and the spin.
- ▶ $\alpha k_z \sigma_y \tau_z$ allows topological transitions when $m_J = 0$.


Topology through mode-mixing

- ▶ A $\pm m_J$ crossing is parabolic $\epsilon \sim k_z^2$.
- ▶ It can be shown that any mode-mixing term $M \sim \mathbb{I}, \sigma_z, \tau_z$:

$$\langle m_J | M | -m_J \rangle \sim \alpha k_z.$$

- ▶ ⇒ mode-mixing acts as **p -wave pairing between $m_J \leftrightarrow -m_J$ states.**

Shaping the wave-function with radial harmonics



$$R(\varphi) = R_0$$


$$H = \begin{pmatrix} \ddots & & & \\ & h_{i,i} & & \\ & & \ddots & \\ & & & h_{i+\ell,i+\ell} \\ & & & & \ddots \end{pmatrix}$$

C. Payá *et al.* 2023, arXiv.

Shaping the wave-function with radial harmonics


$$R(\varphi) = R_0 + \operatorname{Re} \sum_{\ell=1}^{\ell_{\max}} \delta R_\ell e^{i\ell\varphi}$$

Shaping the wave-function with radial harmonics




$$H = \begin{pmatrix} \ddots & & & \\ & h_{i,i} & \dots & h_{i,i+\ell} \\ \vdots & \ddots & \ddots & \vdots \\ h_{i+\ell,i} & \dots & h_{i+\ell,i+\ell} \\ & & \ddots & \ddots \end{pmatrix}$$

Shaping the wave-function with radial harmonics



$$H = \begin{pmatrix} \ddots & & & \\ & h_{i,i} & \dots & h_{i,i+\ell} \\ \vdots & \ddots & \ddots & \vdots \\ h_{i+\ell,i} & \dots & h_{i+\ell,i+\ell} \\ & & \ddots & \ddots \end{pmatrix}$$

Shaping the wave-function with radial harmonics




$$R(\varphi) = R_0 + \operatorname{Re} \sum_{\ell=1}^{\ell_{\max}} \delta R_\ell e^{i\ell\varphi}$$

$$\langle m_J | H | m_J + \ell \rangle = h_{m_J, m_J + \ell}(\ell, \delta R_\ell)$$

$$\delta R_\ell \in \mathbb{C}$$


$$\ell \in \mathbb{N}$$

Effects on the LDOS




- Unperturbed cylinder.
- No topological protection.

Effects on the LDOS




- ▶ Smooth distortion \sim defects in the nanowire profile.
- ▶ All m_J modes interact with each other, opening gaps at 0 energy or creating new MZM.
- ▶ Topology is now possible in all lobes, as it can origin from any m_J mode.

Effects on the LDOS




- ▶ Non-smooth distortion \sim defects in the nanowire profile + atomic size defects.
- ▶ Topological minigaps are larger because harmonic pre-factors can be larger.

Effects on the LDOS



C. Payá *et al.* 2023, arXiv.

Tubular-core




- Follows a simple equation:

$$(\mu_{m_J} - C_{m_J})^2 - (A_{m_J} + V_Z)^2 + \Delta^2 = 0 \xrightarrow[m_J=0]{} V_Z = \sqrt{\Delta^2 + \mu_0^2}$$

- Valid for any disorder model.

Solid-core



► Independent of the disorder model.

C. Payá et al. 2023, arXiv.

Engineering topologically protected edge states

Signals in the LDOS: CdGM analogs

Full 2D simulation: band bending and the solid-core model

Disorder-induced mode-mixing: a new mechanism for topology

Conclusions


Summary

Messages



Summary





Summary






Summary




Summary





Summary




Summary








Summary



Summary




Summary



+ mode-mixing
disorder

Summary



Conclusions

- ▶ In pristine full-shell hybrid nanowires:
 1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.

Conclusions

- ▶ In pristine full-shell hybrid nanowires:
 1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
 2. MZMs are generally topologically unprotected except for small islands in parameter space.

Conclusions

- ▶ In pristine full-shell hybrid nanowires:
 1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
 2. MZMs are generally topologically unprotected except for small islands in parameter space.
 3. Tubular-core nanowires are a good experimental candidate for protected MZMs.

Conclusions

- ▶ In pristine full-shell hybrid nanowires:
 1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
 2. MZMs are generally topologically unprotected except for small islands in parameter space.
 3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
 4. The solid-core phenomenology is more complex and depends on the radial modes.

Conclusions

- ▶ In pristine full-shell hybrid nanowires:
 1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
 2. MZMs are generally topologically unprotected except for small islands in parameter space.
 3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
 4. The solid-core phenomenology is more complex and depends on the radial modes.
- ▶ Adding mode-mixing:

Conclusions

- ▶ In pristine full-shell hybrid nanowires:
 1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
 2. MZMs are generally topologically unprotected except for small islands in parameter space.
 3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
 4. The solid-core phenomenology is more complex and depends on the radial modes.
- ▶ Adding mode-mixing:
 5. Mode-mixing induced by disorder behaves as an effective p -wave pairing.

Conclusions

- ▶ In pristine full-shell hybrid nanowires:
 1. Majorana zero modes appear at odd LP lobes coexist with CdGM analog states.
 2. MZMs are generally topologically unprotected except for small islands in parameter space.
 3. Tubular-core nanowires are a good experimental candidate for protected MZMs.
 4. The solid-core phenomenology is more complex and depends on the radial modes.
- ▶ Adding mode-mixing:
 5. Mode-mixing induced by disorder behaves as an effective p -wave pairing.
 6. Generic disorder generates new MZMs and opens topological minigaps.

Conclusions

Take home message

Majorana physics of full-shell nanowires is very rich. For pristine configurations, the tubular-core model is the optimal candidate but, in the presence of mode-mixing, half of the parameter space is suitable for topologically protected Majorana bound states.

Full-shell Majorana nanowires

A theoretical description

Carlos Payá

Instituto de Ciencia de Materiales de Madrid (ICMM), CSIC


January 10, 2024



Cylindrical nanowire
Mode-mixing


Hollow-core
Modified hollow-core
Tubular-core

Hollow-core results




C. Payá et al. 2023, arXiv.

Modified hollow-core results




C. Payá et al. 2023, arXiv.

Destructive Little-Parks




Hexagonal wave-function



- ▶ New red stripes. Hexagon has $\ell = 6$.
- ▶ Upper stripe: $m_J = 0$ mixes with $m_J = \pm 6$.
- ▶ Lower stripe: $m_J = 3$ mixes with $m_J = -3$.
- ▶ The MZM coming from $m_J = \pm 3$ **cannot** interact with $m_J = 0 \Rightarrow$ they overlap.
- ▶ The $m_J = \pm 6$ MZM annihilates the $m_J = 0$ MZM.

Hexagonal wave-function



- Except for the new topological stripes and a region where the MZM splits, the system is equivalent to the cylinder.

C. Payá *et al.* 2023, arXiv.

Full-shell Majorana nanowires

A theoretical description

Carlos Payá

Instituto de Ciencia de Materiales de Madrid (ICMM), CSIC

January 10, 2024

