

Applied Data Mining Homework 04


Xun Zhao, xz2827

Problem 1:

The function f can be interpreted as:

$$f(x, y) = I\{g(x, y) > 0\}$$
$$g(x, y) = \sigma(x) - \sigma(y)$$

So the network is:



Here, the three ϕ nodes are:


$$\phi_1(\cdot) = \sigma(\cdot)$$
$$\phi_2(\cdot) = \sigma(\cdot)$$
$$\phi_3(\cdot) = I\{\cdot > 0\}$$

Problem 2:

According to the definition of sigmoid function, $\sigma(x) = \frac{1}{1+e^{-x}}$, we can get:

$$\begin{aligned}
-\sigma(-x) &= -\frac{1}{1+e^x} \\
&= \frac{e^x}{1+e^x} - 1 \\
&= \frac{1}{e^{-x}+1} - 1 \\
&= \sigma(x) - 1
\end{aligned}$$

So, the network is:



Here, the three ϕ nodes are:

$$\begin{aligned}
\phi_1(\cdot) &= \sigma(\cdot) \\
\phi_2(\cdot) &= c = 1 \\
\phi_3(\cdot) &= \cdot
\end{aligned}$$

Problem 3:


According basic logical relationship:

$$\begin{aligned}
x \oplus y &= (x = y = 1) \vee (x = y = 0) \\
&= (x \wedge y) \vee (\neg x \wedge \neg y)
\end{aligned}$$

So, the network is the **or** of two **and**.

The single unit of **and** can be presented as $f(x, y) = I\{x + y > 1\}$. And the single unit of **or** can be presented as $f(x, y) = I\{x + y > 0\}$

So the network is:



Here, the bias nodes are:

$$c_1 = 1$$

$$c_2 = 1$$

And other three ϕ nodes are:

$$\phi_1(\cdot) = I\{\cdot > 0\}$$

$$\phi_2(\cdot) = I\{\cdot > 0\}$$

$$\phi_3(\cdot) = I\{\cdot > 0\}$$

Problem 4:

Classification function is:

$$\begin{aligned}f(\vec{x}) &= \phi(\vec{x} \cdot \vec{w} + w_c \cdot c) \\&= I\{\vec{x} \cdot \vec{w} \geq c\}\end{aligned}$$

So, the classification result of \vec{x} is:

$$\begin{aligned}f(\vec{x}) &= I\left\{\begin{pmatrix} -3 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} - \frac{1}{2\sqrt{2}} > 0\right\} \\&= I\left\{-\frac{3}{\sqrt{2}} - \frac{1}{2\sqrt{2}} > 0\right\} \\&= I\left\{-\frac{7}{2\sqrt{2}} > 0\right\} \\&= 0\end{aligned}$$

And the classification result of \vec{x}' is:

$$\begin{aligned}f(\vec{x}') &= I\left\{\begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \end{pmatrix} \cdot \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} - \frac{1}{2\sqrt{2}} > 0\right\} \\&= I\left\{\frac{1}{\sqrt{2}} - \frac{1}{2\sqrt{2}} > 0\right\} \\&= I\left\{\frac{1}{2\sqrt{2}} > 0\right\} \\&= 1\end{aligned}$$

Problem 5:

$$\vec{x} \cdot \vec{y} = e^{\log(x_1) + \log(y_1)} + e^{\log(x_2) + \log(y_2)}$$

The network is:

