
YM
nyears = 67 proyears = 50 3 sampled iterations
M
Msd
Mgrad
M by Year
M/K
M−at−length
M−at−age
Mexp
Linf
Linfsd
Linfgrad
Linf by Year
t0
K
Ksd
Kgrad
K by Year
Steepness (h)
procsd
AC
Rec Devs by Year
Stock−Recruit
Depletion
L50
L95
turity−at−length (current yea
aturity−at−age (current yea
Length
Age
year


Sampled length−at−age curves
First historical year
Last historical year
Last projected year
Length
80
60
40
20
0
0
5
10
15
20
25
Age


Fleet: YM Rec comb Stock: YM
nyears = 67 proyears = 50 3 sampled iterations
Esd
qinc
qcv
Spat_targ
Fishing effort (historical)
Change in future catchability
Selectivity−at−length
First historical year
Last historical year
Last projected year
L5
LFS
Vmaxle
26
34
0.90
Selectivity−at−age
First historical year
Last historical year
Last projected year

Observation biases and sample sizes for observation object Generic_Obs



Observation time series plots for observation object Generic_Obs

























Implementation error time series plots for implementation object Perfect_Imp




















Spawning biomass
0
20000
0
10
20
30
40
50
60
Historical year


Stock depletion (SSB)
0
0.5
1
1.5
2
0
10
20
30
40
50
60
Historical year


Fishing mortality rate (apical)
0
0.2
0.4
0.6
0
10
20
30
40
50
60
Historical year


Annual catches
0
4000
8000
0
10
20
30
40
50
60
Historical year


Recruitment
0
4000
8000
0
10
20
30
40
50
60
Historical year


Recruitment
0
4000
8000
0
5000
15000
25000
35000
Spawning biomass


2.1%
16.7%
4.2%
F/F_MSY
B/B_MSY
0.0
0.5
1.0
1.5
2.0
2.5
3.0
1
67


Age vulnerability (0.25, 0.75)
Simulation 1
Simulation 2
Simulation 3
Age
0
5
15
25
0
10
20
30
40
50
60
Historical year