

# Stochastic Differential Equations

## Introduction

Amanda Groccia, Tatyana Moorehead, Carrie Rider

University of Connecticut, Norfolk State University, Clarkson University

June 20, 2014

# Outline

- 1 Background on Shrimpy
- 2 Brownian Motion
- 3 Modeling
- 4 Nondimensionalization

# Outline

- 1 Background on Shrimpy
  - The Problem
- 2 Brownian Motion
  - Fundamental Concepts
  - Integration
  - Itô's Formula
- 3 Modeling
- 4 Nondimensionalization

# The Problem

Southern German water routes have had several drastic population changes concerning gammarids[2].

Much of this is due to canal construction.

Native Species:

*Gammarus pulex* (Gp)

Invasive Species:

*Dikerogammarus villosus* (Dv)

*Dikerogammarus haemobaphes* (Dh)

*Dikerogammarus bispinosus* (Db)

*Echinogammarus berilloni* (Eb)

# Killer Shrimp




URL?

# Killer Shrimp



URL?





(Kinzler, 2008)



# Time Line

The time to prominence.

Years	Species
<1976	<i>Gp</i> native
1976-1994	<i>Dh</i> invades
1992-1995	<i>Dv</i> invades, <i>Dh</i> declines
>1995	All but <i>Dv</i> coexist separate from <i>Dv</i>



# Kinzler, 2008 Study

Isolated pairs of specimens in a controlled environment

- One freshly moulted (prey)
- One predator
- Total of 279 experiments
- Grouped by age, sex, and species

Results

- Found  $D_v$  to be clear strongest predator
- Found  $D_h$  to have highest cannibalism rate

# Basic Goals

**Determine long term population trends!**

Does  $Dv$  totally dominate in the end?

Which species survive?

Is there an equilibrium?

# Outline

- 1 Background on Shrimpy
  - The Problem
- 2 Brownian Motion
  - Fundamental Concepts
  - Integration
  - Itô's Formula
- 3 Modeling
- 4 Nondimensionalization

# Markov Process

(**Markov Process**) is a stochastic process with the following properties:

- ① The number of possible outcomes or states is finite
- ② The outcome at any stage depends only on the outcome of the previous stage.
- ③ The probabilities are constant over time.

# Brownian Motion

## Definition

(**Brownian Motion**) is a stochastic process that models random continuous motion. The stochastic process  $B = \{B(t), t \geq 0\}$  is standard Brownian Motion if the following holds:

- ①  $B$  has independent increments.
- ② For  $0 \leq s < t$ ,

$$B(t) - B(s) \sim N(0, t - s).$$

- ③ The paths of  $B$  are continuous with probability 1.
- ④  $B(0) = 0$

Random Walk  
Random Walk [1]



# Scaling a Brownian Motion

If  $x(t)$  is a Brownian Motion then  $\frac{x(\lambda t)}{\sqrt{\lambda}}$  is a Brownian motion.

$$\begin{aligned} & P\left(a \leq \frac{x(\lambda t)}{\sqrt{\lambda}} - \frac{x(\lambda s)}{\sqrt{\lambda}} \leq b\right) \\ &= P\left(a\sqrt{\lambda} \leq x(\lambda t) - x(\lambda s) \leq b\sqrt{\lambda}\right), \\ &= \frac{1}{\sqrt{2\pi(\lambda t - \lambda s)}} \int_{a\sqrt{\lambda}}^{b\sqrt{\lambda}} e^{\frac{-x^2}{2}(\lambda t - \lambda s)} dx, \\ &= \frac{1}{\sqrt{2\pi(t - s)}} \int_a^b e^{\frac{-u^2}{2}(t - s)} du. \end{aligned}$$

# Riemann-Stieltjes

**(Riemann-Stieltjes Integral):**

$$\int_a^b f(g)dg = \lim_{n \rightarrow \infty} \sum_{i=1}^N f(g(t_i)) \cdot (g(t_{i+1})) - g(t_i))$$

The main motivation for the Riemann-Stieltjes Integral comes from the concept of Cumulative Distribution Function (CDF) of a random variable.

# Wiener Process

A standard Wiener process (also called Brownian Motion) is a stochastic process  $\{W_t\}_{t \geq 0}$ , which has properties mutually consistent with those of Brownian motion.

**Wiener Integral** For a pair  $(W_t, f(t))$  of a Wiener Process  $W_t$ , a random process  $f(t)$ , we define the Itô integral

$$I(f) = \int_0^\infty f(t) dW_t$$

# Itô's Formula

Itô's Formula is used in Itô Calculus to find the differential of a time-dependent function of a stochastic process.

## Differential Form

$$\partial x_t = \left( \frac{\partial x}{\partial t} + \frac{1}{2} \left( \frac{\partial^2 x}{\partial B^2} \right) \right) \cdot dt + \frac{\partial x}{\partial B} \cdot dB$$

## Integral Form

$$F(t, B(t)) - F(a, B(a)) = \int_a^t \frac{\partial F}{\partial s} ds + \frac{1}{2} \frac{\partial^2 F}{\partial B^2} ds + \int_a^t \frac{\partial F}{\partial B} dB$$

# Itô's Formula

Let  $F = tB^2$

$$\frac{dF}{dt} = B^2$$

$$\frac{dF}{dB} = 2t + B$$

$$\frac{d^2F}{dB^2} = 2t$$

$$\begin{aligned} tB^2(t) - aB^2(a) &= \int_a^t B^s ds + \int_a^t 2sBdB + \int_a^t \frac{1}{2} 2s ds \\ &= \int_a^b B^2 + sds + \int_a^t 2sBdB \end{aligned}$$

OR

$$tB^2(t) - aB^2(a) = \int_a^t B^s ds + \int_a^t 2sBdBt + \frac{1}{2}t^2 - \frac{1}{2}a^2$$

# Outline

- 1 Background on Shrimpy
  - The Problem
- 2 Brownian Motion
  - Fundamental Concepts
  - Integration
  - Itô's Formula
- 3 Modeling
- 4 Nondimensionalization

# Outline

- 1 Background on Shrimpy
  - The Problem
- 2 Brownian Motion
  - Fundamental Concepts
  - Integration
  - Itô's Formula
- 3 Modeling
- 4 Nondimensionalization

# Nondimensionalization

The initial model is:

$$\begin{aligned}\frac{dx}{ds} &= rx \left(1 - \frac{x}{k}\right) - \alpha xy, \\ \frac{dy}{ds} &= \rho y \left(1 - \frac{y}{l}\right) - \beta xy.\end{aligned}$$

Let

$$x \rightarrow A\hat{x}(s)$$

$$y \rightarrow B\hat{y}(s)$$

$$t \rightarrow \tau \cdot s$$

When you substitute and group terms the system becomes nondimensionalized.



The nondimensionalized system is:

$$\begin{aligned}\frac{dx}{ds} &= rx(1-x) - \alpha xy, \\ \frac{dy}{ds} &= y(1-y) - \beta xy.\end{aligned}$$

# Heun's Method

- Heun's method is a numerical procedure for approximating ordinary differential equations with a given initial value.
- First you calculate the intermediate value  $\tilde{y}_{i+1}$  and then the final approximation  $y_{i+1}$  at the next generation point.

$$\tilde{y}_{i+1} = y_i + \Delta t f(t_i, y_i)$$

$$y_{i+1} = y_i + \frac{\Delta t}{2} [f(y_i, t_i) + f(\tilde{y}_{i+1}, t_{i+1})]$$

# Acknowledgments

We would like to thank the following for their support and funding:

- Dr. Joel Foisy, SUNY Potsdam
- Dr. Kelly Black, Clarkson University
- National Security Agency (number?)
- National Science Foundation (DSM-1262737)

# Bibliography



D. Higham.

An algorithmic introduction to numerical simulation of stochastic differential equations.

*SIAM Review*, 43(3):525–546, 2001.



Werner Kinzler, Axel Kley, Gerd Mayer, Dieter Waloszek, and Gerhard Maier.

Mutual predation between and cannibalism within several freshwater gammarids: *Dikergammarus villosus* versus one native and three invasives.

*Aquatic Ecology*, 43(2):457–464, 2009.



G A Polis, C A Myers, and R D Holt.

The ecology and evolution of intraguild predation: Potential competitors that eat each other.

*Annual Review of Ecology and Systematics*, 20(1):297–330, 1989.