FIITJEE

ALL INDIA TEST SERIES

FULL TEST – VII

JEE (Main)-2024

TEST DATE: 22-03-2024

ANSWERS, HINTS & SOLUTIONS

Physics

PART – A

SECTION – A


1. B

Sol. $\int_0^R \frac{k\lambda}{r} 2\pi r dr = \frac{q}{4\varepsilon_0}$

$k\lambda 2\pi R = \frac{q}{4\varepsilon_0}$

$\frac{\lambda R}{2\varepsilon_0} = \frac{q}{4\varepsilon_0}$


$\Rightarrow \quad \lambda = \frac{q}{2R}$



2. B

Sol. $F - \mu mg = mv\frac{dv}{dx}$

$10x - 10x^2 = K.E.$





3. C

For More Material Join: @JEEAdvanced_2024

Sol. $\frac{4x}{3H} = \frac{4R}{3\pi H}$

$\Rightarrow x = \frac{R}{\pi}$






4. C

Sol. $I = \frac{20}{500} A$

$I' = \frac{5}{1000} = \frac{1}{200} A$


$I_1 = \frac{1}{25} - \frac{1}{200} = \frac{7}{200}$

$\frac{I}{I_1} = \frac{1/25}{7/200} = \frac{8}{7}$



5. C

Sol. Average velocity = $\frac{\ell/\sqrt{3}}{\text{time}} = \frac{\ell/\sqrt{3}}{2\ell/3v} = \frac{\sqrt{3}v}{2}$



6. C

Sol. $I_R = I_T - I_{cut} = \frac{(\sigma\pi R^2)R^2}{4} - \left[\frac{\left(\frac{\sigma\pi R^2}{4}\right)\frac{R^2}{4}}{4} + \left(\frac{\sigma\pi R^2}{4}\right)\frac{R^2}{8}\right]$

$= \frac{\sigma\pi R^4}{4}\left[1 - \frac{1}{16} - \frac{1}{8}\right] == \frac{13\sigma\pi R^4}{64}$

7. A

Sol. $P = \frac{E_{rms}^2 \cos\phi}{z} \Rightarrow 400 = \frac{(250)^2}{z} \times 0.8$

$\cos\phi = \frac{R}{Z} = 0.8$

$z = \sqrt{X_L^2 + R^2}$

$R = 100\ \Omega$

$Z = 125\ \Omega$

$X_L = 75\ \Omega$

8. B

Sol. $m_1v_1 = m_2v_2$

For More Material Join: @JEEAdvanced_2024

$m_1 = m_2 8$

$\frac{m_1}{m_2} \Rightarrow \frac{R_1}{R_2} = (8)^{1/3} = 2:1$

9. B

Sol. $\frac{dQ}{dt} = \frac{k_1 40A}{L} = \frac{k_2 10A}{L} = \frac{k_3 50A}{L}$

Under steady state $\Rightarrow 4k_1 = k_2 = 5k_3$

10. C

Sol. induced emf $e = (-)L\left(\frac{dI}{dt}\right)$

11. C

Sol. Gravitational potential at point P

$V_P = -\frac{GM}{R}$


Work done, $W = m_0\left[V_\infty - V_P\right] = \frac{GMm_0}{R}$

12. C

Sol. $\delta_1 = 180° - 2 \times 53° = 74°$

$\delta_2 = 53°$

$\delta_{net} = \delta_1 + \delta_2 = 127°$



13. B

Sol. $15 \times \frac{2\pi m}{Bq} v\cos 37° = d$

14. C

Sol. No. of photons per second = $\frac{I}{\left(\frac{hC}{\lambda}\right)} \times \text{Area} = \frac{8 \times 10 \times 10^{-4}}{2 \times 1.6 \times 10^{-19}}$

No. of photoelectrons per second = $\frac{1}{5} \times \frac{8 \times 10 \times 10^{-4}}{2 \times 1.6 \times 10^{-19}} = 5 \times 10^{15}$

15. A

Sol. $\left[\beta x^2\right] = \left[k\theta\right]$

$\beta L^2 = ML^2T^{-2}$

$\beta = MT^{-2}$

Also,

For More Material Join: @JEEAdvanced_2024

$[W] = [\alpha\beta]$

$\left[ML^2T^{-2}\right] = \left[\alpha MT^{-2}\right]$

$\alpha = L^2$

16. B

Sol. $P_0V_0^2 = P4V_0^2$

$P = \frac{P_0}{4}$

$P_0V_0 = nRT_1$

$\frac{P_0}{4}(2V_0) = nRT_2$

$\Delta U = \frac{3}{2}nR(T_2 - T_1) = \frac{3}{2}(-)\frac{P_0V_0}{2} = -\frac{3}{4}P_0V_0$

17. B

Sol.

A	B	Y
1	0	1
1	1	1
0	0	0
0	1	1

18. C

Sol. $T_1 = 2\pi\sqrt{\frac{\ell}{g}},\ T_2 = 2\pi\sqrt{\frac{\ell}{g+a}}$

$T_2 = 2\pi\sqrt{\frac{\ell}{(g+4)}}$

$y = 2t^2,\ a = \frac{d^2y}{dt^2} = 4\ m/s^2$

$\left(\frac{T_2}{T_1}\right)^2 = \left(\frac{g}{g+4}\right) = \frac{10}{14} = \frac{5}{7}$

19. B

Sol. Potential energy

$U_i = -\vec{P}_2 \cdot \vec{E}_1 = -P\left(\frac{2kP}{d^3}\right)\cos 180°$

$U_i = \frac{2kP^2}{d^3} = \frac{P^2}{2\pi\varepsilon_0 d^3}$

$U_f = \frac{P^2}{16\pi\varepsilon_0 d^3}$

$\frac{P^2}{2\pi\varepsilon_0 d^3} - \frac{P^2}{16\pi\varepsilon_0 d^3} = \frac{1}{2}mv^2 \times 2$

$v = \sqrt{\frac{7P^2}{16\pi m\varepsilon_0 d^3}} = \frac{P}{4d}\sqrt{\frac{7}{\pi m\varepsilon_0 d}}$


For More Material Join: @JEEAdvanced_2024

20. B

Sol. $de = (B \sin\theta)\, dx \times \sin\theta\, \omega$

$$e = B\omega \sin^2\theta \int_0^L x\,dx$$

$$e = \frac{1}{2} B\omega L^2 \sin^2\theta$$



SECTION – B


21. 2

Sol. IAOR at (0, 1.5)

$v = r\omega$

$3 = 1.5\omega$

$\omega = 2$ rad/s



22. 2

Sol. $$\phi = \frac{2\pi}{\lambda} = \frac{2\pi}{\lambda}\frac{\lambda}{4}$$

$$\phi = \frac{\pi}{2}$$

$$I_R = I_0 \cos^2\left(\frac{\phi}{2}\right) = I_0 \cos^2\frac{\pi}{4}$$

$$I_R = \frac{I_0}{2}$$

23. 7

Sol. $$Av_1 = \frac{A}{4} v_2 \Rightarrow v_2 = 4v_1$$

$$P_1 + \rho gh + \frac{1}{2}Pv_1^2 + P_2 + \frac{1}{2}Pv_2^2$$

$$(P_1 - P_2) + \rho gh = \frac{1}{2}P\left(v_2^2 - \frac{v_2^2}{16}\right)$$

$$(P_1 - P_2) + \rho gh = \frac{15}{32}\rho v_2^2$$


$\Rightarrow v_2 = 7$ m/s

For More Material Join: @JEEAdvanced_2024

24. 6

Sol. $R_{eq} = 6\Omega \Rightarrow \dfrac{V}{R_{eq}} = I$

$I = 6A$



25. 4


Sol. At slipping

$\tan\theta = \mu_S$

$\tan\theta = \dfrac{1}{\sqrt{3}}$

$\theta = \dfrac{\pi}{6}$

$\ell = R\theta = \dfrac{24}{\pi} \times \dfrac{\pi}{6} = 4$ cm



26. 885

Sol. $I = \dfrac{1}{2}\varepsilon_0 E_0^2 c$

Radiation pressure $= \dfrac{2I}{c} = \varepsilon_0 E_0^2 = 8.85 \times 10^{-12} \times 10^4 = 885 \times 10^{-10}$

27. 20

Sol. $T_1 + T_2 = 20$

$T_2 L = 20\,x$

$T_2 = \dfrac{20x}{L} = 20x$

$T_1 = 20(1-x)$

Also,

$\dfrac{1}{2\ell}\sqrt{\dfrac{T_1}{\mu}} = \dfrac{1}{\ell}\sqrt{\dfrac{T_2}{\mu}}$


$T_1 = 4T_2$

$20(1-x) = 80\,x$

$20 - 20x = 80x$

$\Rightarrow \dfrac{20}{100} = x$

$\Rightarrow x = 20$ cm



28. 4

For More Material Join: @JEEAdvanced_2024


Sol. $\tau = 2mg \times \frac{\ell}{\sqrt{2}}$

$$\left[\frac{m\ell^2}{3} + \frac{m\ell^2}{12} + m\left(\frac{\sqrt{5}\ell}{2}\right)^2\right]\alpha = 2mg\frac{\ell}{\sqrt{2}}$$

$$\Rightarrow \frac{5}{3}\ell\alpha = \frac{2g}{\sqrt{2}}$$

$$\frac{5}{3}\frac{3}{\sqrt{2}}\alpha = \frac{20}{\sqrt{2}}$$

$\Rightarrow \alpha = 4$ rad/s^2



29. 4

Sol. $\frac{dy}{dx} = \tan\theta = 8x$


$u\cos\theta = v\sin\theta$

$eu\sin\theta = v\cos\theta$

$\tan^2\theta = 1/e$

$\tan\theta = 2$

$x = 1/4$ and $y = 1/4$



30. 4

Sol. $\text{Charge} = \frac{|\Delta\phi|}{R} = \frac{B\pi r^2}{4R}$

For More Material Join: @JEEAdvanced_2024

Chemistry

PART – B

SECTION – A

31. C

Sol. Ionic product of $MOH = \frac{0.2}{2}\left[\frac{10^{-5}}{2}\right]$

$= \frac{0.1}{2} \times 10^{-6}$

$= 5 \times 10^{-7}\ M^2$


K_{ip} for $M(OH)_2 = \frac{0.2}{2}\left[\frac{10^{-5}}{2}\right]^2 = \frac{0.1}{4} \times 10^{-10}$

$= 2.5 \times 10^{-12}\ M^3$

Only for $Pb(OH)_2$, $K_{ip} > K_{sp}$, hence $Pb(OH)_2$ for precipitate.

32. D

Sol.



33. B

Sol. $L_1 = 179$ pm, $L_2 = 163$ pm

34. A

Sol. $\Delta W = \frac{\pi(5V_1 - V_1)(7P_1 - P_1)}{4} = \frac{\pi 4V_1 \times 6P_1}{4}$

$\Delta W = 6\pi P_1 V_1$

For More Material Join: @JEEAdvanced_2024

35. A

Sol. $t_{1/2} = \frac{A_o}{2k} \Rightarrow k = \frac{A_o}{200}$

For zeroth $A_o - A = +kt$

$A_o - \frac{A_o}{8} = 8kt$

$\frac{7A_o}{8} = \frac{A_o}{200}t$

$t = \frac{700}{4} = 175$ min

36. D

Sol. $K_c = \frac{[D]^3}{[A][B]^2}$ unitless

$= M^o$

37. D

Sol. Due to relative lowering of vapour pressure.

38. B

Sol. pK_a, 2-fluorobenzoic acid = 3.27
2-bromobenzoic acid = 2.88
2-chlorobenzoic acid = 2.94
Due to SIR effect.


39. B

Sol. Vitamin B_1 – Beri-Beri
Rest are corrects.

40. A

Sol. Zn > Co > Cu > Ni
906 758 745 736 (kJ/mole)

41. C

Sol. 

329.3 K
350.8 K
310.5 K
300.8 K

42. D

Sol. $\Delta T_f \propto$ im

Higher value of im, lower the freezing point.

(i) im $= 2 \times 2 = 4$ (ii) im $= 5 \times 3 = 15$

(iii) im $= 3 \times 4 = 12$ (iv) im $= 2 \times 5 = 10$


For More Material Join: @JEEAdvanced_2024

43. A
Sol. Potassium has lilac colour on flame test.


44. C
Sol. $\left[Co(en)_2(NH_3)_2\right]^{+3}$ has 2 geometrical isomerism and it's cis form will have optical isomerism.

45. A
Sol. AgBr is a partially soluble in NH_3 and in $Na_2S_2O_3$, NaCN, Ag^+ form complexes.

46. B
Sol.



Non-planar structure



Planar and trans forms

47. C
Sol. Maltose on hydrolysis give two molecular of glucose only.

48. B
Sol. $CrO_4^{-2} + H_2O_2 + H^+ \longrightarrow CrO_5 + 3H_2O$

$4CrO_5 + 12H^+ \longrightarrow \underset{\text{green soln}}{4Cr^{+3}} + 7O_2 + 6H_2O$

49. A
Sol. $M^{+3} \xrightarrow{xV} M^{+2} \xrightarrow{z} M^{+1} \xrightarrow{?} M$

$\underbrace{\qquad\qquad}_{y}$

$x \times 1 + z \times 1 + Q \times 1 = y \times 3$

$Q = 3y - x - z$

50. C
Sol. $E = \left(\frac{hc}{\lambda}\right)$, λ should be reverse order of strength of ligands.

SECTION – B

51. 6
Sol. Elements follow $4d^{1\text{ to }10}\ 5s^1$ are Nb, Mo, Tc, Ru, Rh, Ag.

52. 5
Sol. Mischmetal is alloy of lanthanoid, which contain nearly 5% of Fe.

53. 12
Sol. By the ratio find % of C, H and O then formula

$C_3H_6O + 4O_2 \longrightarrow 3CO_2 + 3H_2O$


Y

3 mole compound will require 12 mole O_2.

For More Material Join: @JEEAdvanced_2024

54. 10

Sol.



55. 675

Sol. Balmer series $\frac{1}{\lambda_1} = R_H Z^2 \left(\frac{1}{2^2} - \frac{1}{4^2}\right)$ .... (1)

Paschen series $\frac{1}{\lambda_2} = R_H Z^2 \left(\frac{1}{3^2} - \frac{1}{4^2}\right)$ .... (2)

$$\frac{\frac{1}{\lambda_1}}{\frac{1}{\lambda_2}} = \left(\frac{\frac{1}{2^2} - \frac{1}{4^2}}{\frac{1}{3^2} - \frac{1}{4^2}}\right) = \left(\frac{\frac{4-1}{16}}{\frac{16-9}{9\times16}}\right) = \frac{3}{16} \times \frac{9\times16}{7}$$

$$\frac{\lambda_2}{\lambda_1} = \frac{27}{7} \Rightarrow \frac{\lambda_2}{350} = \frac{27}{7}$$

$$\frac{\lambda_2}{175} = \frac{27}{7} \Rightarrow \lambda_2 = \frac{175\times27}{7}$$

$\lambda_2 = 675$ nm

56. 200

Sol. $M = \frac{20}{40} \times \frac{1000}{500} = 1\,M$

$M_1V = M_2V_2$

$1\times V_1 = 0.2\times1000$

$V_1 = 200$ ml

57. 18

Sol. $n = 5$

$\ell = 4$

g(sub shell)

$m = -4, -3, -2, -1, 0, +1, +2, +3, +4$

9 orbitals.

Maximum electron = 18.

58. 3

Sol. $NO(g) + \frac{1}{2}O_2(g) \rightleftharpoons NO_2(g)$

$\Delta G_r^o = \Delta G_{NO_2}^o - \Delta G_{NO}^o = 5.1454 - 1 = 4.1454$ Kcal

$\Delta G^o = -2.303RT \log_{10} K_{eq}$

$4.1454\times10^3 = -2.303RT \log_{10} K_{eq}$

$4.1454\times10^3 = -2.303\times2\times300 \log_{10} K_{eq}$

For More Material Join: @JEEAdvanced_2024

$$\log_{10} K_{eq} = -\frac{4.1454}{2.303 \times 300 \times 2}$$

$$= -\frac{4145.4}{2.303 \times 600} = -3$$

$$K_{eq} = 10^{-3}$$

59. 70

Sol. x = 7, y = 10

60. 29

Sol. $E^{o}_{Cu^{2+}/Cu} = 0.34$ V

For More Material Join: @JEEAdvanced_2024

Mathematics

PART – C

SECTION – A

61. C

Sol. f(x) is discontinuous at $x = 1, \sqrt{2}, \sqrt{3}, 2, \sqrt{5}, \sqrt{6}, \sqrt{7}, \sqrt{8}, 3$, g(x) is non differentiable at $x = 1, \sqrt{2}, \sqrt{3}, 2, \sqrt{5}, \sqrt{6}, \sqrt{7}, \sqrt{8}$ and $\frac{3}{2}$.

62. B

Sol. $(x^2 - 4)(x - 1)(x^5 + 2x - 3) = 0$

$x^5 + 2x - 3 = 0$ has only real root.

63. B

Sol. $\begin{Bmatrix} x^2 - 12x + 37 & ; & x \le 2 \\ 2x - x^2 + 17 & ; & 2 < x < 5 \\ x^2 - 12x + 37 & ; & x \ge 5 \end{Bmatrix}$

64. C

Sol. $\lim\limits_{x \to 0} \dfrac{(A-B)x + \left(A - \frac{B}{2} + C\right)x^2 + \left(\frac{A}{2} - \frac{B}{3} - C\right)x^3 .....}{\frac{x^3}{2}} = 2024$

$\Rightarrow \quad A = B,\ A - \frac{B}{2} + C = 0, \frac{A}{2} - \frac{B}{3} - C = 1012$

$A = B = 1518,\ C = -759$

65. B

Sol. $J(x) = \int e^{\tan^2 x} \left(2\tan x\left(\sec^2 x \cos^2 x - \frac{1}{1+\tan^2 x}\right)\right)dx$

$= \int e^{\tan^2 x}(2\tan x \sec^2 x \cdot \cos^2 x - \sin 2x)dx$

$J(x) = e^{\tan^2 x}\cos^2 x + c$

$\because\ J(0) = 1 \ \Rightarrow c = 0$

$J\left(\frac{\pi}{4}\right) = \frac{e}{2}$

66. B

Sol. $\tan 45° = \left|\frac{m-2}{1+2m}\right| \Rightarrow m = -3,\ \frac{1}{3}$

$a\left|\frac{1}{m_1} - \frac{1}{m_2}\right| = 12.5$

$\Rightarrow \frac{10a}{3} = 12.5 \ \Rightarrow a = 3.75$

67. B

Sol. α and β lie on the chord $y = 1$ intersecting the circle $x^2 + y^2 - 8x - 6y + 17 = 0$

$\therefore \quad x^2 - 8x + 12 = 0 \ \Rightarrow x = 2, 6$

$\therefore \quad \alpha = 2 + i,\ \beta = 6 + i$

For More Material Join: @JEEAdvanced_2024

68. D

Sol. $\tan 45° = \left|\frac{m-3}{1+3m}\right| \Rightarrow m = -2, \frac{1}{2}$

69. B

Sol. $37^{2024} + 43^{2024} = 16\lambda + 2$

70. C

Sol. $13 + (N-1)\,20 \le 8093$

$\Rightarrow N \le 405$

71. C


Sol. $a + b = 5$

72. C

Sol. $a = 7$

73. C

Sol. Required area $= \int_0^2 4\sqrt{x}\,dx + \frac{1}{2}(2023 + 2022)$



74. D

Sol. $\sqrt{2}\sum_{k=1}^{9}\left[\tan\left(\theta + (2k+1)\frac{\pi}{8}\right) - \tan\left(\theta + (2k-1)\frac{\pi}{8}\right)\right]$

$\Rightarrow \sqrt{2}\left(\tan\left(\theta + \frac{19\pi}{8}\right) - \tan\left(\theta + \frac{\pi}{8}\right)\right) = \left(2 + \frac{2}{\sqrt{3}}\right)\sqrt{2}$

$\Rightarrow \frac{1}{\cos\left(\theta + \frac{\pi}{8}\right)\cos\left(\theta + \frac{3\pi}{8}\right)} = 2\left(1 + \frac{1}{\sqrt{3}}\right)\sqrt{2}$

$\Rightarrow \cos\frac{\pi}{4} + \cos\left(2\theta + \frac{\pi}{2}\right) = \frac{\sqrt{3}}{(\sqrt{3}+1)\sqrt{2}}$

$\Rightarrow \sin 2\theta = \frac{\sqrt{3}-1}{2\sqrt{2}} \Rightarrow \theta = \frac{\pi}{24}, \frac{11\pi}{24}$

75. C

Sol. $x > 0,\ \tan^{-1}\left(\frac{10x}{25 - x^2}\right) = \frac{\pi}{6} \Rightarrow x = 10 - 5\sqrt{3}$

$x < 0,\ \tan^{-1}\left(\frac{10x}{25 - x^2}\right) = -\frac{\pi}{3} \Rightarrow x = -\frac{5}{\sqrt{3}}$

76. C

For More Material Join: @JEEAdvanced_2024

Sol. $a = 5, \angle B = \frac{5\pi}{12}$

$\sin A = \frac{a}{2R} = \frac{5}{5\sqrt{2}} = \frac{1}{\sqrt{2}} \Rightarrow A = \frac{\pi}{4}$

Also $b = 2R \sin B = 5\sqrt{2}\left(\frac{\sqrt{3}+1}{2\sqrt{3}}\right) = \frac{5(\sqrt{3}+1)}{2}$

$\angle C = \frac{\pi}{3}$

Hence $\Delta = \frac{1}{2}ab\sin C = \frac{1}{2}\times 5\times 5\frac{(\sqrt{3}+1)}{2}\times\frac{\sqrt{3}}{2} = \frac{25\sqrt{3}(\sqrt{3}+1)}{8}$

77. B

Sol. $32\sin^5\alpha + 243\cos^5\beta + 1 + 1 + 1 \geq 5(2^5 3^5 \sin^5\alpha \cos^5\beta)^{1/5} \geq 30\sin\alpha\cos\beta$

$\Rightarrow \quad 32\sin^5\alpha = 243\cos^5\beta = 1$

$\Rightarrow \quad \sin\alpha = \frac{1}{2}, \cos\beta = \frac{1}{3}$

$\cos 2\alpha = \frac{1}{2}, \cos 2\beta = -\frac{7}{9}$

78. C

Sol. $(a, a) \in R$: R is reflexive

Let $a = \frac{q}{p}$ and b is rational then $q - pb + q$ is irrational but $pb - p\frac{q}{p} + q$ is rational

$\therefore$ R is not symmetric

Let $pa - pb + q$ and $pb - pc + q$ both are irrational but $p(a - c) + 2q$ may be rational if $a - c = -\frac{2q}{p}$

$\therefore$ non transitive.

79. C

Sol. $\int \frac{x\cos x - \sin x}{x\sqrt{x^2 - \sin^2 x}}dx = \int \frac{\frac{x\cos x - \sin x}{\sin^2 x}}{\frac{x}{\sin x}\sqrt{\left(\frac{x}{\sin x}\right)^2 - 1}}dx = \operatorname{cosec}^{-1}\left(\frac{x}{\sin x}\right) + c$

80. A

Sol. $\frac{1}{(y+2024)^2}\frac{dy}{dx} + \frac{1}{y+2024}x^2 = e^{x^3/3}$

$\Rightarrow \quad \frac{dt}{dx} - tx^2 = e^{x^3/3}; t = -\frac{1}{y+2024}$

$-\frac{1}{y+2024}e^{-x^3/3} = x + c$

$\because \ y(0) = -\frac{6071}{3} \ \therefore \ c = -3$

$y + 2024 = -\frac{1}{(x-3)e^{x^3/3}}$

For More Material Join: @JEEAdvanced_2024

SECTION – B

81. 1728

Sol. $n(C) = (2^{506} - 1)2^{1518}$

$m - p = 1012.$

82. 16

Sol. Equation of perpendicular bisector of the line segment joining the points (3, 5) and (7, 9) is $x + y - 12 = 0$. Solving it with the equation of diameter $3x - y - 12 = 0$ we get centre as (6, 6)


$\therefore$ Area of rectangle = $2\sqrt{2} \times 4\sqrt{2} = 16$

83. 6

Sol. $\frac{z_3}{z_1} = \cos 3\theta e^{i3\theta}$

$\frac{z_2}{z_1} = \cos\theta e^{i\theta}$

$\left(\frac{z_2}{z_1}\right)^3 = \frac{\cos^3\theta}{\cos 3\theta} \times \frac{z_3}{z_1} \Rightarrow \frac{\cos^3\theta}{\cos 3\theta} = \frac{5+3\sqrt{3}}{8} \Rightarrow \theta = \frac{\pi}{12}$



84. 171

Sol. Let $P_n = \alpha^n + \beta^n$ then $P_{n+1} - P_{n-1} = 13P_n$

85. 300

Sol. 111138, 111146, 113222, 111226, 111432

$\frac{6!}{4!} \times 2 + \frac{6!}{3!2!} \times 2 + \frac{6!}{3!}$

$= 60 + 120 + 120 = 300$

86. 14

Sol. $5^{n(n-1)} 3^{(n-1)^3} = 5^{42} 3^{216} \Rightarrow n = 7$

$$\pi \int_{n-14}^{n} x \sin \pi x \, dx = 2\pi \left[\left(-\frac{x\cos\pi x}{\pi} \right)_0^7 + \frac{1}{\pi} \int_0^7 \cos \pi x \, dx \right]$$

87. 280

Sol. $L_1 : \frac{x-8}{8} = \frac{y-0}{-6} = \frac{z-0}{0}$

$L_2 : \frac{x}{0} = \frac{y+8}{8} = \frac{z}{-5}$

$$d = \frac{\begin{vmatrix} 8 & 8 & 0 \\ 8 & -6 & 0 \\ 0 & 8 & -5 \end{vmatrix}}{\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 8 & -6 & 0 \\ 0 & 8 & -5 \end{vmatrix}} = \frac{280}{\sqrt{1649}}$$

For More Material Join: @JEEAdvanced_2024

88. 10118

Sol. $(2023\vec{a}+2024\vec{b})\cdot\left\{\left(2|\vec{b}|^2\,\vec{a}+3|\vec{a}|^2\,\vec{b}\right)\right\}$

$=(2023\vec{a}+2024\vec{b})\cdot(2\vec{a}+3\vec{b})$

89. 180

Sol. Possible values of f(1), f(2), f(3) are

(1, 2, 3), (2, 3, 4), (3, 4, 5), (4, 5, 6), (5, 6, 7)

(1, 3, 5), (2, 4, 6), (3, 5, 7), (1, 4, 7)

$P(E)=\dfrac{n(E)}{n(S)}=\dfrac{18.4!}{7!}=\dfrac{18.4!}{7.6.5.4!}=\dfrac{3}{35}$

90. 200

Sol. $x + y = 21$

$x^2 + y^2 = 225$

$\Rightarrow x = 9, y = 12$

$\therefore$ M. D $=\dfrac{\sum_{i=1}^{6}|x_i-\bar{x}|}{6}=\dfrac{10}{3}$

For More Material Join: @JEEAdvanced_2024