

Variational Calculus

Morten Hjorth-Jensen^{1,2}

Scott Pratt¹

Carl Schmidt³

¹Department of Physics and Astronomy and National Superconducting Cyclotron Laboratory, Michigan State University, USA

²Department of Physics, University of Oslo, Norway

³Department of Physics and Astronomy, Michigan State University, USA

Jun 6, 2019

Introduction

The calculus of variations involves problems where the quantity to be minimized or maximized is an integral.

In the general case we have an integral of the type

$$E[\Phi] = \int_a^b f(\Phi(x), \frac{\partial \Phi}{\partial x}, x) dx,$$

where E is the quantity which is sought minimized or maximized. The problem is that although f is a function of the variables Φ , $\partial\Phi/\partial x$ and x , the exact dependence of Φ on x is not known. This means again that even though the integral has fixed limits a and b , the path of integration is not known. In our case the unknown quantities are the single-particle wave functions and we wish to choose an integration path which makes the functional $E[\Phi]$ stationary. This means that we want to find minima, or maxima or saddle points. In physics we search normally for minima. Our task is therefore to find the minimum of $E[\Phi]$ so that its variation δE is zero subject to specific constraints. In our case the constraints appear as the integral which expresses the orthogonality of the single-particle wave functions. The constraints can be treated via the technique of Lagrangian multipliers