

Why Measure Theory

Carter Deslauriers

September 19, 2026

1 Introduction: Motivation

1.1 Definitions

I will begin with the definition of Riemann Integration.

Definition 1 (Partition). Let $[a, b]$ be the domain of a function then P a partition is of the form

$$P = \{x_0, x_1, \dots, x_n\}$$

Such that

$$a = x_0 < x_1 < \dots < x_n = b$$

Definition 2 (Upper sums). Let f be a bounded function $f: [a, b] \rightarrow \mathbb{R}$, the upper sums for f

$$U(f, P_\epsilon) = \sum_{k=1}^n (x_k - x_{k-1}) \sup_{[x_{k-1}, x_k]} f$$

Where $U(f)$ is defined as

$$U(f) = \inf_P U(f, P)$$

Definition 3 (Lower sums). Let f be a bounded function $f: [a, b] \rightarrow \mathbb{R}$, the lower sums for f

$$L(f, P_\epsilon) = \sum_{k=1}^n (x_k - x_{k-1}) \inf_{[x_{k-1}, x_k]} f$$

Where $L(f)$ is defined as

$$L(f) = \sup_P L(f, P)$$

Definition 4 (Reimann Integration). Let f be a bounded function $f: [a, b] \rightarrow \mathbb{R}$.

If for every $\epsilon > 0$ there exists a partition P_ϵ such that

$$U(f, P_\epsilon) - L(f, P_\epsilon) < \epsilon \tag{1}$$

then $f(x)$ is Riemann Integrable with

$$\int f = U(f) = L(f) \tag{2}$$

1.2 Riemann Failure

1.2.1 Function

Let the sequence of rational numbers $\{r_1, r_2, \dots\}$ enumerate the rationals on $[0, 1]$.

Let $f_n(x)$ be a series of functions of the form

$$f_n(x) = \begin{cases} 1 : & x \in \{r_1, r_2, \dots, r_n\} \\ 0 : & x \notin \{r_1, r_2, \dots, r_n\} \end{cases} \quad (3)$$

1.2.2 Upper and Lower Sums

Fix $n \in \mathbb{N}$.

Take the partition P_n such that $(x_k - x_{k-1}) = \frac{\epsilon}{4n}$ for all consecutive pairs of points in P_n .

The total number of intervals in which there are rational numbers is $2n$; since there are n rationals where each can be in at most 2 intervals at once.

Note. if $r_k = x_k$ for instance then $[x_{k-1}, x_k]$ and $[x_k, x_{k+1}]$ both contains r_k

This gives

$$U(f_n, P_n) = \left(\frac{\epsilon}{4n}\right)(2n)(1) + (1)(0) = \frac{\epsilon}{2}$$

Where each length $(x_k - x_{k-1}) = \left(\frac{\epsilon}{4n}\right)$, there exists at most $2n$ of these intervals, each with $\sup_{[x_k - x_{k-1}]} f = 1$. All other possible intervals have a max combined length of 1, each with $\sup_{[x_k - x_{k-1}]} f = 0$.

Since $\mathbb{I}$ is dense in $\mathbb{R}$, between any two real numbers there exists an irrational. Meaning on any interval $[x_{k-1}, x_k]$, $\inf_{[x_{k-1}, x_k]} f = 0$ giving

$$L(f_n, P_n) = 0$$

This gives for any $\epsilon > 0$ for some $N \in \mathbb{N}$ whenever $n \geq N$ it follows

$$U(f_n, P_n) - L(f_n, P_n) = \frac{\epsilon}{2} < \epsilon$$

meaning the function is Riemann integrable with

$$\int_0^1 f_n = L(f_n) = 0$$

1.2.3 Limit

Since the sequence of values r_k are countable we have

$$\lim_{n \rightarrow \infty} f_n = \begin{cases} 1 : & x \in \mathbb{Q} \cap [0, 1] \\ 0 : & x \notin \mathbb{Q} \cap [0, 1] \end{cases} \quad (4)$$

Call this function f

Now for any partition P , since both $\mathbb{Q}$ is dense in $\mathbb{R}$ as well as $\mathbb{I}$ dense in $\mathbb{R}$, on any interval $[x_{k-1}, x_k] \in P$ there exists both a rational and irrational.

This gives $\inf_{[x_{k-1}, x_k]} f = 0$ as well $\sup_{[x_k - x_{k-1}]} f = 1$ on every interval. Which forces

$$U(f, P) = 1$$

and

$$L(f, P) = 0$$

for every partition P implying $U(f) = 1$ and $L(f) = 0$ which says that f is not Riemann integrable.

1.3 Lebesgue Explanation

1.3.1 Measure 0

Definition 5 (Measure 0). A is a measure zero set if for every $\epsilon > 0$ there exists a countable covering $\bigcup_n O_n$ where each O_n is an open interval such that

$$A \subset \bigcup_n O_n \quad (5)$$

Where

$$\sum_n l(O_n) \leq \epsilon \quad (6)$$

and $l()$ denotes the length of a open interval defined as $b - a$ for any open interval (a, b) .

1.3.2 Lebesgue Criterion

By the Lebesgue criterion

Theorem 1 (Lebesgue Criterion). *A bounded function $f: [a, b] \rightarrow \mathbb{R}$ is Riemann integrable iff the set of discounties is Measure 0*

1.3.3 f discontinuous everywhere

Recall that

$$\lim_{n \rightarrow \infty} f_n = \begin{cases} 1 : & x \in \mathbb{Q} \cap [0, 1] \\ 0 : & x \notin \mathbb{Q} \cap [0, 1] \end{cases}$$

I want to show that f is discontinuous on all of $[0, 1]$ First observe the definition of continuity.

Definition 6. A function f is continuous at a point c if for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $|x - c| < \delta$ it follows that $|f(x) - f(c)| < \epsilon$

Now fix $\epsilon_0 \leq 1$ take $\epsilon_0 = \frac{1}{2}$. Choose any point $c \in [0, 1]$. Again both $\mathbb{Q}$ and

$\mathbb{I}$ dense in $\mathbb{R}$.

First consider $c \in \mathbb{Q}$ then for any δ chosen there exists some irrational point

$$x \in (c - \delta, c + \delta) \implies |f(x) - f(c)| = 1 \geq \epsilon_0 \quad (7)$$

Consider the case $c \in \mathbb{I}$ then for any δ chosen there exists some rational point

$$x \in (c - \delta, c + \delta) \implies |f(x) - f(c)| = 1 \geq \epsilon_0 \quad (8)$$

Meaning the set of discontinuous points of f is all of $[0, 1]$ where $|[0, 1]| = 1$.

Note. I do not prove it here but it is proved in section 2.2

To show that the set of discontinuities for every f_n is measure 0 first consider the set of points at which f_n is discontinuous, call it D_n .

For any points $c \in \{r_1, r_2, \dots, r_n\}$ since $\mathbb{I}$ dense in $\mathbb{R}$ we have for a fixed $\epsilon < 1$ and any $\delta > 0$

$$|x - c| < \delta \implies |f(x) - f(c)| = 1 \geq \epsilon$$

Making

$$D_n = \{r_1, r_2, \dots, r_n\}$$

For any points $c \notin \{r_1, r_2, \dots, r_n\}$ we can take $\delta = \frac{\min(|x-r_1|, |x-r_2|, \dots, |x-r_n|)}{2}$ since there are a fixed amount of rationals for any fixed $n \in \mathbb{N}$.

This would give us for

$$|x - c| < \delta \implies |f(x) - f(c)| = 0 < \epsilon$$

making f_n continuous on any points $x \notin \{r_1, r_2, \dots, r_n\}$.

Now consider the open covering $\bigcup_k O_k$ such that each

$$O_k = (r_k - \frac{\epsilon}{2^{k+1}}, r_k + \frac{\epsilon}{2^{k+1}})$$

Then

$$\begin{aligned} \sum_k l(O_k) &= \sum_k \frac{\epsilon}{2^k} \\ &= \frac{\frac{\epsilon}{2}}{1 - \frac{1}{2}} \\ &= \epsilon \end{aligned}$$

giving

$$\sum_k l(O_k) \leq \epsilon$$

making D_n measure 0 by definition.

We now understand why each f_n integrates but f itself fails to integrate. Our understanding lays not in the language of the Riemann integral itself but instead in the language of measure.

We also see that the method by which he partition the domain fails for functions that oscillate too quickly and will fail to be defined in spaces outside of $\mathbb{R}$

1.4 Lebesgue Savior

2 Outer Measure

Definition 7 (Outer Measure). $|A| = \inf \sum_k l(O_k)$ Where inf is taken over all open coverings such that $A \subset \bigcup_k O_k$

2.1 Agrees everyrwhere

Outer measure agrees with length on open sets and $[a, b] = (a, b)$ in terms of measure. Translation invariance, subadditivity, monotonicity

Theorem 2. 1. If $A \subset B$ then $|A| \leq |B|$

2. $|[a, b]| = b - a$

3. If $A_1, A_2, \dots$ are sets such that $\bigcup_k A_k = B$, then $|B| \leq \sum_k |A_k|$

4. $|A + t| = |A|$

I will now prove each in order.

Proof: Monotonicity. For monotonicity (1) first observe that since

$$A \subset B \subset \bigcup_k O_k$$

any open covering of B is an open covering of A .

Now outer measure is defined as inf over all open coverings. Therefore there exists an open covering $\bigcup_k O_k$, such that

$$\bigcup_k l(O_k) < |B| + \epsilon$$

Since any open covering of B is an open covering of A

$$|A| \leq \sum_k l(O_k) < |B| + \epsilon$$

Where ϵ is arbitrary > 0 . Meaning $|A| \leq |B|$

□

Proof: $|[a, b]| = b - a$. For the second property of outer measure consider the open cover for $[a, b]$

$$(a - \epsilon, a + \epsilon) \cup (a, b) \cup (b - \epsilon, b + \epsilon)$$

Since $|[a, b]|$ is inf over all coverings we have

$$|[a, b]| \leq 2\epsilon + b - a + 2\epsilon$$

Where ϵ is arbitrary giving

$$|[a, b]| \leq b - a$$

For the other direction consider any open covering of $[a, b]$. I want to show that

$$\sum_k l(O_k) \geq b - a$$

hence giving

$$\inf \sum_k l(O_k) \geq b - a$$

Now $[a, b]$ is a compact interval, by Heine-Borel for every open cover

$$[a, b] \subset \bigcup_k O_k$$

there exists a finite subcollection such that

$$[a, b] \subset \bigcup_{k=1}^n O_k$$

Where

$$\sum_{k=1}^n l(O_k) \leq \sum_{k=1}^{\infty} l(O_k)$$

Since length is non negative.

Suppose the claim holds for n intervals.

Begin with $n = 1$ intervals. Any interval containing $[a, b]$ ie $[a, b] \subset O_k$ has length $l(O_k) \geq b - a$, since

$$(a, b) \subset [a, b] \subset O_k$$

Now consider $n + 1$ intervals covering $[a, b]$.

b must be contained in at least one of these intervals call it (c, d) . If $c < a$ then $l(c, d) \geq b - a$ so consider $a < c < b < d$; since (c, d) covers none of $[a, c]$ we can invoke the inductive hypothesis

$$\sum_{k=1}^n l(O_k) \geq c - a$$

for O_k covering $[a, c]$. Adding this to $l(c, d)$ gives

$$\begin{aligned} \sum_{k=1}^{n+1} l(O_k) &\geq (c - a) + (d - c) \\ &= d - a \\ &\geq b - a \end{aligned}$$

Completing the proof. □

Proof: Countable Subadditivity.

□

Proof: Translation Invariance. I would like to show $|A + t| = |A|$.

First observe that

$$l((a, b)) = l((a + t, b + t))$$

since

$$l((a + t, b + t)) = (b + t) - (a + t) = b - a$$

Now observe that for any open cover of A call it O_A , O_{A+t} covers A .

Where O_{A+t} is defined as

$$\bigcup_k (O_k + t)$$

for $O_k \in O_A$.

Take any point in $A + t$ call it y . Then $y = x + t$ for some $x \in A$. Where $x \in O_k$ for some $O_k \in O_A$. Then $y \in O_k + t$. Where

$$l(O_k) = l(O_k + t)$$

Making the set of covers of A , call them $S \subset S_t$, where S_t is the set of open covers of $A + t$.

Meaning since $|A|$ inf over all covers ie inf S and $S \subset S_t$, it follows $|A + t| \leq |A|$.

Note. Say $B \subset C$ then $\inf C \leq \inf B$; since taking inf over larger set is smaller.

Now consider the sets $A+t+(-t)$ and $A+t$ where the left side is a translations

of the right side. Giving

$$|A + t + (-t)| \leq |A + t| \implies |A| \leq |A + t|$$

Which follows from the above. Completing the proof $|A + t| = |A|$.

□

2.2 Failure of Outer Measure as a Measure

Actually doesn't give us countable additivity of disjoint sets though if we consider the Vitali set. This comes from axiom of choice and we want sigma algebras to be closed under complementation, countable union, countable intersection from de Morgan, and contain null and the set itself, because then we can use all of our operations and write proof that are completely consistent

3 Integration on Measure

We can now generalize to any space $(X, \mathcal{S}, \mu)$ and we have measure which splits based on output and reverse maps to inputs where we get the measure of those sets times the inf of the function on the sets, which we can then use for probability where we know for certain our methods are coherent.