


[Draft] Weathering Potential Explorer User Guide

The Weathering Potential Explorer includes several user-adjustable inputs and filters that shape how suitability is visualized. Use this map as a starting point to narrow down promising areas for ERW and not as a substitute for site-level planning, MRV modeling, or carbon flux estimation.


For the best experience, please use desktop fullscreen mode by clicking this button  on the top right corner.

Controls Tab

Percentile Ranking Legend (set as default)



The Explorer visualizes relative weathering potential using a color scale based on percentile rankings across global agricultural soils. Each pixel is ranked against other cropland pixels worldwide based on its environmental conditions for rock weathering.



Here's how to interpret the color categories:

- 80-100th percentile (Top 20%) (**Red**): Among the highest weathering potential areas on cropland globally
- 60-80th percentile (**Red Orange**)
- 40-60th percentile (**Orange**)
- 20-40th percentile (**Light Orange**)
- 0-20th percentile (Bottom 20%) (**Light Tan**): Among the least favorable croplands for weathering based on current model inputs. This doesn't mean weathering is infeasible, just that conditions are less likely to be favorable for fast reactions, and local factors may matter more.

These rankings are based only on land currently classified as cropland according to the dataset selected (defaulted to GFSAD1000). Non-agricultural lands are not included in the percentile calculation. This scale is most useful for comparing regions or prioritizing areas for further investigation. Just because a site ranks lower does not mean it is a bad site, especially if other factors (like rock type, supply chains, or agronomic fit) make it work in practice. We explore more of these limitations in the [\[to link\] full explainer article](#).

Weathering Multiplier

Show Weathering Multiplier (~1 min to load, zoom helps)

The Weathering Multiplier toggle offers an alternative way to interpret the model output. Instead of showing percentile rankings, this view tells you how much potentially faster weathering is expected in a given location compared to the global median across all croplands.

Weathering Potential Magnitude vs Agricultural Median)

- 5.0x+ (5+ times agricultural median)
- 4.0x (4 times agricultural median)
- 3.0x (3 times agricultural median)
- 2.0x (2 times agricultural median)
- 1.0x (equals agricultural median)

Here's how to interpret the color categories:

- A value of 1.0x indicates the global median weathering rate.
- Light pink areas are near the median (~1.0x).
- Pink areas are around 3x faster than the median.
- Darker purple areas are 5x or more faster than the median weathering rate.

This view is especially useful for understanding the magnitude of difference between locations. While the percentile map shows how a location ranks, it doesn't tell you how much faster weathering might be from one region to another. For instance, the 80th and 90th percentile might seem close, but if one is only 1.3x the median and the other is 10x, that's a major difference!

The Weathering Multiplier highlights these kinds of gaps, helping users see where weathering could happen substantially faster than average agricultural soils worldwide. This view also helps describe the shape of the distribution underlying the percentile rankings. It answers not just "how high is this ranked?" but also "how much *more* weathering is potentially likely to occur here?"

Show Individual Layers


Show Individual Layers

Clicking "Show Individual Layers" will load all of the map's underlying environmental layers into the display. You'll briefly see the layers "flash" as they load - this is normal.




To view layers individually, click the “Layers” icon in the top-right corner of the map. This opens a control panel where:

- Each layer is listed separately
- Layers are stacked top-to-bottom, meaning the top-most selected layer is the one that’s visible
- You can use the opacity slider next to each layer to blend or compare them




For each environmental variable (pH, temperature, moisture), you’ll see 2 types of layers:

1. **Raw Data Values:** showing the original environmental conditions



2. **Effect on Weathering Potential:** showing how each variable influences weathering rate in the model (scaled from 0 to 1, where **blue** is slower and **red** is faster)



These individual layers are useful for understanding what’s driving the map output in different regions, and for exploring how each variable contributes to the model’s estimation of relative weathering potential.

Masks Tab

Cropland Masking

Users can optionally restrict the analysis to agricultural land by applying one of two cropland filters:

- **ESA WorldCover (10 m):** Offers higher spatial resolution and general land cover classification, ideal for detecting finer-scale agricultural patterns.
- **GFSAD1000 (1km) (set as default):** Includes more detailed cropland type distinctions (e.g., irrigated vs. rainfed) but at coarser resolution, which may be better suited for large-scale regional planning.

Cropland Options

Cropland Dataset:

GFSAD Cropland (Custom)

- ☒ Class 1: Irrigated
- ☒ Class 2: Rainfed
- ☒ Class 3: Cropland/Natural Veg
- ☒ Class 4: Cropland/Pasture
- ☒ Mask to Cropland Only

Both layers help highlight areas where ERW is operationally feasible (i.e. croplands that are already managed, routinely amended, and where ERW has been most commonly piloted), but may miss smallholder systems or mosaic landscapes in tropical regions.

pH Depth and Masking

The Weathering Potential Explorer uses soil pH in H₂O from the SoilGrids dataset. You can customize two aspects of how pH is handled in the model:

1. **Select pH Depth:** Choose the soil depth from which pH values are drawn. Available depths range from 0–5 cm to 100–200 cm.
2. **Set pH Filter Range:** Optionally exclude soils with pH values outside a specified range. The default filter excludes soils with pH below 4.5 or above 8.0.

pH Depth Selection

SoilGrids pH Depth:

0-5cm (Surface)

pH Range Options

☒ Mask out pH

Minimum pH (keep areas >= this value):

4.5

Maximum pH (keep areas <= this value):

8

SOC masking

The Weathering Potential Explorer includes an optional filter for soil organic carbon (SOC), allowing you to exclude areas with very low or very high SOC content. SOC is measured as a percentage by mass from the OpenLandMap dataset at 0-5 cm depth. You can set both a minimum and maximum SOC threshold. The default setting keeps areas with SOC between 0% and 5%.

SOC (Soil Organic Carbon) Options

☒ Mask out SOC

Minimum SOC % (keep areas >= this value):

0

Maximum SOC % (keep areas <= this value):

5

Soil texture masking

The Weathering Potential Explorer includes an optional filter for soil texture, using USDA classifications from the OpenLandMap dataset (0–5 cm depth). When you enable "Mask by Soil Texture", all 12 texture classes are selected by default, so no filtering is applied initially. All textures remain visible and included in the analysis.

When you deselect classes, unchecking texture types will:

- Hide those classes from the map
- Exclude them from the weathering potential analysis
- Limit results to areas with the remaining selected textures

Soil Texture Options

☒ Mask by Soil Texture

- ☒ Clay (Heavy clay soils)
- ☒ Silty Clay (Fine-textured soils)
- ☒ Sandy Clay (Clay with sand content)
- ☒ Clay Loam (Balanced clay-rich soils)
- ☒ Silty Clay Loam (Fine-textured loams)
- ☒ Sandy Clay Loam (Coarser clay loams)
- ☒ Loam (Well-balanced soils)
- ☒ Silty Loam (Smooth-textured soils)
- ☒ Sandy Loam (Light, well-draining soils)
- ☒ Silt (Very fine particles)
- ☒ Loamy Sand (Sandy soils with structure)
- ☒ Sand (Coarse, fast-draining soils)

Regional Analysis Tab

On-click regional statistics

Clicking on any region returns a pop-up summary that shows:

- Total land area and cropland area (if a cropland mask is applied)
- The distribution of pixels across relative weathering potential categories (from low to high)

These summaries reflect all active filters, including cropland, pH, SOC, and soil texture masks. That means the reported land areas represent only the portion of the region that meets your selected criteria. This helps you understand not just environmental favorability, but also how much land corresponds to your masks.

These summaries are based on administrative boundaries from the FAO GAUL Level 1 dataset (e.g., provinces, counties, or equivalent).

Weathering Potential Explorer Minimize ×

Controls

Masks

Regional Analysis

Global Areas

Click on Map for Regional Analysis
Click anywhere on the map to see state and country level weathering rate statistics.

State/Province: Kaduna
Total Region Area: 44,494.66 km²
Land Area: 44,164.42 km² (99.3% of region)
Cropland Area (GFSAD): 29,688.27 km² (66.7% of region, 67.2% of land)
Weathering potential coverage: 99.7% of cropland area

- 0-20th percentile (Bottom 20%): 1,116.85 km² (3.1% of cropland)
- 20-40th percentile: 0.00 km² (0.0% of cropland, 0.0% of land)
- 40-60th percentile: 32.22 km² (0.1% of cropland, 0.1% of land)
- 60-80th percentile: 27,427.68 km² (92.4% of cropland, 92.4% of land)
- 80-100th percentile (Top 20%): 1,016.77 km² (3.4% of cropland)

Load Country Data for Nigeria

Below the regional summary, you'll also see a "Load country data for [Country Name]" button. Clicking this runs the same analysis for the entire country that contains the selected region, allowing you to quickly compare local and national suitability profiles.

Global Areas Tab

The Global Areas tab provides a summary of how land is distributed across relative weathering potential categories, aggregated globally.

Clicking the "Calculate Global Areas" button runs the same analysis used in the Regional tab, but across the entire globe. It returns total land area and cropland area (if masked), broken down by percentile categories, reflecting all active filters, including pH, SOC, and soil texture.

Use this view to get a quick sense of how much land exists globally under your filtering scenario.

Weathering Potential Explorer Minimize ×

Controls

Masks

Regional Analysis

Global Areas

Global Weathering Potential by Percentile
Calculate Global Percentile Areas
Global percentile calculation complete!

- 0-20th percentile (Bottom 20%)**
Area: 1.86 million km²
- 20-40th percentile**
Area: 3.18 million km²
- 40-60th percentile**
Area: 5.14 million km²
- 60-80th percentile**
Area: 4.31 million km²
- 80-100th percentile (Top 20%)**
Area: 1.76 million km²

WALKTHROUGH EXAMPLE: NYANZA, KENYA

Let's walk through how you might use the Weathering Potential Explorer to assess relative weathering potential in a region of interest: in this example, Nyanza, Kenya.

Step 1: Apply Cropland Masking

To focus on regions where ERW is likely to be operationally feasible, you apply the GFSAD cropland mask. This restricts your analysis to agricultural lands and filters out urban, forested, and non-arable areas. You keep all of the subclasses checked for this initial analysis.

Cropland Options
Cropland Dataset:

GFSAD Cropland (Custom)

☒ Class 1: Irrigated
☒ Class 2: Rainfed
☒ Class 3: Cropland/Natural Veg
☒ Class 4: Cropland/Pasture
☒ Mask to Cropland Only

Step 2: Select a pH Depth

You select the 5–15 cm pH depth layer to be more representative of the root zone, where water movement, root activity, and mineral interactions are more stable and relevant for longer-term weathering.

pH Depth Selection
SoilGrids pH Depth:

5-15cm (Shallow)

Step 3: Enable SOC + pH Filtering

To avoid areas where bicarbonate generation may be reduced due to harsher soil conditions, you enable the filter that excludes land with soil organic carbon greater than 5% and pH less than 5 and greater than 8.5. Your reasoning is that while weathering can still occur in these environments, carbon may not be stored as effectively, or there may be a higher risk of releasing CO₂ from soil organic matter.

pH Range Options	SOC (Soil Organic Carbon) Options
<input checked="" type="checkbox"/> Mask out pH	<input checked="" type="checkbox"/> Mask out SOC
Minimum pH (keep areas >= this value): <div>5</div>	Minimum SOC % (keep areas >= this value): <div>0</div>
Maximum pH (keep areas <= this value): <div>8.5</div>	Maximum SOC % (keep areas <= this value): <div>5</div>

Step 4: Enable Soil Texture Filtering


To refine your analysis based on physical soil properties, you enable the soil texture filter and only uncheck the “clay” category to focus on areas with better drainage and more dynamic water movement, which are more favorable for sustained weathering reactions.

Soil Texture Options

- ☒ Mask by Soil Texture
 - ☐ Clay (Heavy clay soils)
 - ☒ Silty Clay (Fine-textured soils)
 - ☒ Sandy Clay (Clay with sand content)
 - ☒ Clay Loam (Balanced clay-rich soils)
 - ☒ Silty Clay Loam (Fine-textured loams)
 - ☒ Sandy Clay Loam (Coarser clay loams)
 - ☒ Loam (Well-balanced soils)
 - ☒ Silty Loam (Smooth-textured soils)
 - ☒ Sandy Loam (Light, well-draining soils)
 - ☒ Silt (Very fine particles)
 - ☒ Loamy Sand (Sandy soils with structure)
 - ☒ Sand (Coarse, fast-draining soils)

Step 5: Zoom In and Interpret the Map

You zoom into Kenya and observe clusters of orange and red cells. These correspond to a relative weathering potential ranking in the 40-100th percentile.



Step 6: Click for Regional Statistics

You click on the Nyanza region. A pop-up displays a breakdown of land area by weathering potential rankings, both for total land and for cropland (based on your selected filters).

State/Province: Nyanza	State/Province: Nyanza
Total Region Area: 16,350.20 km ²	Total Region Area: 16,350.20 km ²
Land Area: 12,425.50 km ² (76.0% of region)	Land Area: 12,425.50 km ² (76.0% of region)
Cropland Area (GFSAD): 6,054.29 km ² (37.0% of region, 48.7% of land)	Cropland Area (GFSAD): 6,054.29 km ² (37.0% of region, 48.7% of land)
Weathering potential coverage: 51.4% of cropland area	Weathering potential coverage: 51.4% of cropland area
0-20th percentile (Bottom 20%): 460.90 km ² (7.6% of cropland, 0.3% of land)	0-20th percentile (Bottom 20%): 460.90 km ² (7.6% of cropland, 0.3% of land)
20-40th percentile: 0.00 km ² (0.0% of cropland, 0.0% of land)	20-40th percentile: 0.00 km ² (0.0% of cropland, 0.0% of land)
40-60th percentile: 0.99 km ² (0.0% of cropland, 0.0% of land)	40-60th percentile: 0.99 km ² (0.0% of cropland, 0.0% of land)
60-80th percentile: 2,600.85 km ² (43.0% of cropland, 20.9% of land)	60-80th percentile: 2,600.85 km ² (43.0% of cropland, 20.9% of land)
80-100th percentile (Top 20%): 50.96 km ² (0.8% of cropland, 0.4% of land)	80-100th percentile (Top 20%): 50.96 km ² (0.8% of cropland, 0.4% of land)

You repeat the process for the North Eastern region, which shows a slightly different profile with more extensive land area, but less cropland coverage.

State/Province: North Eastern	State/Province: North Eastern
Total Region Area: 128,086.67 km ²	Total Region Area: 128,086.67 km ²
Land Area: 127,503.41 km ² (99.5% of region)	Land Area: 127,503.41 km ² (99.5% of region)
Cropland Area (GFSAD): 3,500.97 km ² (2.7% of region, 2.7% of land)	Cropland Area (GFSAD): 3,500.97 km ² (2.7% of region, 2.7% of land)
Weathering potential coverage: 99.9% of cropland area	Weathering potential coverage: 99.9% of cropland area
0-20th percentile (Bottom 20%): 207.24 km ² (5.9% of cropland, 0.2% of land)	0-20th percentile (Bottom 20%): 207.24 km ² (5.9% of cropland, 0.2% of land)
20-40th percentile: 1,093.20 km ² (31.2% of cropland, 0.9% of land)	20-40th percentile: 1,093.20 km ² (31.2% of cropland, 0.9% of land)
40-60th percentile: 499.21 km ² (14.3% of cropland, 0.4% of land)	40-60th percentile: 499.21 km ² (14.3% of cropland, 0.4% of land)
60-80th percentile: 1,515.34 km ² (43.3% of cropland, 1.2% of land)	60-80th percentile: 1,515.34 km ² (43.3% of cropland, 1.2% of land)
80-100th percentile (Top 20%): 183.00 km ² (5.2% of cropland, 0.1% of land)	80-100th percentile (Top 20%): 183.00 km ² (5.2% of cropland, 0.1% of land)

These breakdowns help you assess not just how favorable the region is environmentally, but also how much land actually meets your filtering criteria.

While the Weathering Potential Explorer is useful for highlighting promising areas, it is ultimately a first-pass screening tool. For local policymakers, researchers, or project developers, additional layers of information — such as feedstock availability, land tenure, infrastructure access, existing cropping systems, and soil amendment practices — will be essential for assessing where ERW can be viably and responsibly scaled.