

# RPA Accumulation during Class Switch Recombination Represents 5'–3' DNA-End Resection during the S–G2/M Phase of the Cell Cycle

Arito Yamane,<sup>1</sup> Davide F. Robbiani,<sup>3</sup> Wolfgang Resch,<sup>1</sup> Anne Bothmer,<sup>3,7</sup> Hirotaka Nakahashi,<sup>1</sup> Thiago Oliveira,<sup>3</sup> Philipp C. Rommel,<sup>3</sup> Eric J. Brown,<sup>5</sup> Andre Nussenzweig,<sup>6</sup> Michel C. Nussenzweig,<sup>3,4,\*</sup> and Rafael Casellas<sup>1,2,\*</sup>

<sup>1</sup>Genomics & Immunity, NIAMS

<sup>2</sup>Center of Cancer Research, NCI

National Institutes of Health, Bethesda, MD 20892, USA

<sup>3</sup>Laboratory of Molecular Immunology

<sup>4</sup>Howard Hughes Medical Institute

The Rockefeller University, New York, NY 10065, USA

<sup>5</sup>Abramson Family Cancer Research Institute and Department of Cancer Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

<sup>6</sup>Laboratory of Genome Integrity, NCI, National Institutes of Health, Bethesda, MD 20892, USA

<sup>7</sup>Present address: Cancer Genetics Program, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215, USA

\*Correspondence: [nussen@rockefeller.edu](mailto:nussen@rockefeller.edu) (M.C.N.), [casellar@mail.nih.gov](mailto:casellar@mail.nih.gov) (R.C.)

<http://dx.doi.org/10.1016/j.celrep.2012.12.006>

## SUMMARY

Activation-induced cytidine deaminase (AID) promotes chromosomal translocations by inducing DNA double-strand breaks (DSBs) at immunoglobulin (*Ig*) genes and oncogenes in the G1 phase. RPA is a single-stranded DNA (ssDNA)-binding protein that associates with resected DSBs in the S phase and facilitates the assembly of factors involved in homologous repair (HR), such as Rad51. Notably, RPA deposition also marks sites of AID-mediated damage, but its role in *Ig* gene recombination remains unclear. Here, we demonstrate that RPA associates asymmetrically with resected ssDNA in response to lesions created by AID, recombination-activating genes (RAG), or other nucleases. Small amounts of RPA are deposited at AID targets in G1 in an ATM-dependent manner. In contrast, recruitment in the S–G2/M phase is extensive, ATM independent, and associated with Rad51 accumulation. In the S–G2/M phase, RPA increases in nonhomologous-end-joining-deficient lymphocytes, where there is more extensive DNA-end resection. Thus, most RPA recruitment during class switch recombination represents salvage of unrepaired breaks by homology-based pathways during the S–G2/M phase of the cell cycle.

## INTRODUCTION


Antigen receptor genes are assembled from variable (V), diversity (D), and joining (J) gene segments by the V(D)J recombinase

Rag1 and Rag2 (Fugmann et al., 2000). This process requires generation of DNA DSBs in lymphocyte precursors in the G1 phase of the cell cycle. These targeted DNA double-strand breaks (DSBs) are joined by nonhomologous end joining (NHEJ) as revealed by a complete block in *Ig* and T cell receptor (TCR) gene assembly in the absence of NHEJ (Jung et al., 2006; Rooney et al., 2004).

Upon antigenic stimulation the *Ig* genes in B lymphocytes undergo additional diversification by somatic hypermutation (SHM) and class switch recombination (CSR). CSR is a deletional recombination reaction between highly repetitive switch regions that replaces the heavy chain constant domain *Igμ* with one of several downstream isotypes *Igγ*, *Igε*, or *Igα* (Honjo et al., 2002; Stavnezer et al., 2008). Similar to V(D)J recombination, NHEJ plays a key role in the resolution of DSBs incurred during CSR (Helmink and Sleckman, 2012).

Both SHM and CSR are initiated by activation-induced cytidine deaminase (AID), an enzyme that converts cytidines to uracils at *Ig* variable genes and switch (S) regions (Maul et al., 2011). dU:dG mismatches are recognized by base excision and mismatch repair proteins leading to formation of DNA nicks and DSBs that are obligate intermediates in CSR (Stavnezer et al., 2008). In addition to *Ig* loci, AID can also target a large number of non-*Ig* genes (Liu et al., 2008; Pasqualucci et al., 1998; Shen et al., 1998; Yamane et al., 2011), including oncogenes that are frequently translocated to *Ig* in human and mouse B cell tumors (Chiarle et al., 2011; Klein et al., 2011; Küppers and Dalla-Favera, 2001).

How AID is targeted to *Ig* and non-*Ig* loci is still unknown, but transcriptional pausing has been implicated (Peters and Storb, 1996; Rajagopal et al., 2009; Wang et al., 2009). In support of this idea, the RNA exosome and RNA polymerase II stalling factor Spt5 appear to be required for AID to access its target genes (Basu et al., 2011; Pavri et al., 2010). Another potential AID cofactor, the single-stranded DNA (ssDNA) binding



**Figure 1. RPA Recruitment to AID, RAG, and I-SceI-Induced Breaks**

(A) RPA occupancy at the *Igh* locus of 53BP1<sup>-/-</sup> (lane I), H2AX<sup>-/-</sup> (lane II), and H2AX<sup>-/-</sup>UNG<sup>-/-</sup>Msh2<sup>-/-</sup> (lane III) B cells activated ex vivo in the presence of LPS+IL-4. Deep-sequencing read densities were normalized to adjust for library size and bin width (reads per million per kb; RPKM). Numbers in parentheses represent average read density in RPKM for the displayed genomic window.

(B) TCRα locus showing recruitment of RPA in 53BP1<sup>-/-</sup> (lane I) and 53BP1<sup>+/+</sup> (lane III) control thymocytes. Rag2 occupancy is also included in lane II.

(C) Kinetics of RPA recruitment at an I-SceI site engineered at Myc intron 1. Activated B cells were transduced with retroviruses expressing ER-I-SceI and cells were harvested at 0, 0.5, 3, or 24 hr after tamoxifen treatment. See also Figure S1.

Replication Protein A (RPA), was isolated as part of a biochemical screen for activities that enhance AID hypermutation of in-vitro-transcribed substrates (Chaudhuri et al., 2004). Consistent with this idea, RPA associates with *Ig* and non-*Ig* AID target genes (Vuong et al., 2009; Yamane et al., 2011) in a manner that is directly proportional to the extent of AID activity (Hakim et al., 2012). However, the precise role of RPA in CSR remains unknown.

In eukaryotic cells, RPA forms a complex with ssDNA that is essential for DNA replication, telomere maintenance, DNA recombination, DNA repair, and DNA damage checkpoint activation (Oakley and Patrick, 2010; Wold, 1997). There are at least three ways whereby RPA might impact CSR. First, RPA could help stabilize AID's ssDNA targets during *Ig* gene transcription (Chaudhuri et al., 2004). Second, RPA might stimulate long patch base-excision repair (DeMott et al., 1998; Ranalli et al., 2002) or mismatch repair (Genschel and Modrich, 2003; Lin et al., 1998), which play critical roles in the processing of *Ig* gene deamination (Stavnezer et al., 2008). Third, RPA might associate with and stabilize resected ssDNA that cannot easily be repaired by classical NHEJ, thereby facilitating salvage by homology-mediated repair pathways (Bothmer et al., 2010; Hasham et al., 2010; Zhang et al., 2010).

## RESULTS

### RPA Accumulates in Response to DNA Breaks


Deletion of 53BP1 markedly increases RPA recruitment to *Ig* genes, particularly in the presence of the *Igκ*AID transgene (Figure S1; Hakim et al., 2012). To clarify the nature of RPA recruitment at AID targets, we monitored RPA by chromatin immuno-

precipitation (ChIP)-Seq in the absence of H2AX, a factor that like 53BP1 limits DNA-end resection during intrachromosomal recombination (Bothmer et al., 2010; Helmink et al., 2011). B cells were stimulated with lipopolysaccharide (LPS) and interleukin 4 (IL-4) to promote CSR from *Igμ* to *Igγ1* and *Igε*. Similar to 53BP1<sup>-/-</sup> B cells, we found broad RPA islands centered at recombining S<sub>μ</sub>, S<sub>γ1</sub>, and to a lesser extent S<sub>ε</sub> (Figure 1A, lanes I and II). Thus, absence of H2AX enhances RPA deposition at *Igh*.

To ascertain whether RPA recruitment requires DSBs, as opposed to AID recruitment, we performed the same experiment in H2AX<sup>-/-</sup>UNG<sup>-/-</sup>Msh2<sup>-/-</sup> B cells. In this genetic background, AID is targeted to and deaminates *Ig* switch regions normally, but the resulting uracils are not processed to produce DSBs (Maul et al., 2011; Rada et al., 2004; Xue et al., 2006). Notably, RPA signals fell to background levels in the absence of DSBs (Figure 1A, lane III). This finding is consistent with the notion that RPA deposition requires the formation of DSBs, but is independent of DNA deamination and AID targeting.

To determine whether RPA is also associated with other DSBs produced in G1, we examined CD4<sup>+</sup>CD8<sup>+</sup> double-positive thymocytes, which actively undergo recombination-activating gene (RAG)-mediated TCR-α recombination. Consistent with extensive nucleolytic processing of coding ends in 53BP1<sup>-/-</sup> T cells (Difilippantonio et al., 2008), we found prominent RPA signals in Jα (Figure 1B, lane I), a profile that is reminiscent of RAG2 recruitment to this locus (Figure 1B, lane II; Ji et al., 2010). In contrast to wild-type B cells, which accumulate RPA at the *Ig* locus during CSR, RPA accumulation in T cells required 53BP1 deletion, because ChIP signals were indistinguishable from background in wild-type thymocytes (Figure 1B, lane III).

To verify that DSBs are sufficient to stimulate RPA accumulation and to examine the kinetics of recruitment, we induced DNA breaks in 53BP1<sup>-/-</sup>AID<sup>-/-</sup>Myc<sup>+/+</sup> B cells by expression of an estrogen inducible I-SceI meganuclease (ER-I-SceI) (Robbiani



**Figure 2. RPA Interacts with Resected ssDNA Downstream of AID or I-Ppol Endonuclease**

(A) RPA association with + and - DNA strands at the *Grap* gene locus from *IgkAID*-53BP1<sup>-/-</sup>-activated B cells. RPKM values for the specified window (chr11:61,401,372-61,543,038) are provided in parenthesis.

(B) RPA bound to *Igh* locus. As in (A), ChIP-seq signals were resolved into upper and lower strands. Technical replicates were treated with *E. coli* Exol or RecJ exonucleases prior to deep-sequencing library preparation.

(C) PolII binding to + and - strands at the *Grap* locus.

(D) Technical IP replicates from (A) and (C) samples incubated in the presence of Exol.

(E) Composite diagram showing RPA profiles at 19 I-Ppol mouse genomic sites in 53BP1<sup>-/-</sup> MEFs transduced with retroviruses expressing the I-Ppol homing endonuclease.

See also Figure S2.


et al., 2008). RPA recruitment became detectable 3 hr following induction of ER-I-SceI and persisted for at least 24 hr (Figure 1C). Taken together, the data demonstrate that in 53BP1<sup>-/-</sup> lymphocytes RPA accumulates rapidly in response to physiological DSBs introduced in G1 (RAGs, AID) as well as nuclease-induced (I-SceI) damage.

#### RPA Associates with Resected ssDNA

During DNA repair RPA associates with ssDNA created by end resection (Symington and Gautier, 2011). Resected DNA is typically asymmetric and immunoprecipitated RPA associated with such DNA structures would be expected to reflect that asymmetry. Indeed, when RPA ChIP-seq libraries from 53BP1<sup>-/-</sup>*IgkAID* B cells were resolved into upper (+) and lower (-) DNA strands RPA was asymmetrically distributed around AID target sites. As exemplified by *Grap*, a non-*Ig* AID target gene (Klein et al.,

2011; Yamane et al., 2011), RPA was enriched on the + strand upstream, and on the - strand downstream of intron 1 (Figure 2A). This distribution is consistent with 5'-3' DNA-end resection (Figure 2A, schematics). Directionality in RPA recruitment was also observed at *Igh* and other AID target genes in 53BP1<sup>-/-</sup> B cells (Figure 2B and not shown). Similar results were also obtained with Myc<sup>+</sup> cells transduced with I-SceI (Figure S2A). In contrast to RPA, PolII ChIP-seq libraries did not display any obvious strand biases at *Grap* or other AID target genes (Figure 2B and not shown).

To confirm that RPA binds to resected DNA at AID target genes, we incubated RPA and control PolII immunoprecipitates with *Escherichia coli* single-strand exonuclease Exol, which digests ssDNA from 3'-5' before library preparation. While PolII signals were largely unchanged, RPA enrichment at *Grap* or other AID targets could not be detected after nuclease treatment



**Figure 3. Rad51 Recruitment to AID On- and Off-Targets**

(A) Rad51 accumulation on the upper (+) and lower (–) strand at the *Igh* locus of *IgkAID-53BP1*<sup>–/–</sup>-activated B cells.

(B) Rad51 (blue), RPA (red), and translocation profiles at *Mir155* and *Cd83* genes in chromosomes 16 and 13, respectively, from *IgkAID-53BP1*<sup>–/–</sup> B cells.

(C) Chromosomal translocations (TSS  $\pm$  2 kb) involving *Igh* or *Myc* at Rad51-recruiting genes. Chromosomes 12 and 15 carrying the I-SceI sites were excluded from the analysis. Correlation between the two data sets is calculated using Spearman's  $\rho$ .

(D) Rad51 levels (TSS  $\pm$  2 kb) at RPA+ and RPA– genes in *IgkAID-53BP1*<sup>–/–</sup> B cells.

See also Figure S3.

(Figures 2C and 2D; Figure S2B). Although to a lesser extent, treatment of IP DNA with RecJ, a 5'–3' exonuclease, also diminished RPA signals substantially (Figures 2B and 2D; Figure S2B). This observation is consistent with the notion that resected, 200 bp sonicated DNA is susceptible to both 5' and 3' nucleolytic attack (Figure S2C). On the basis of these findings, we conclude that RPA associates with resected DNA in response to AID or nuclease-mediated DNA damage.

In yeast, DNA-end resection can extend up to 10 kb in the absence of a homologous donor or Rad51 (Bishop et al., 1992;

Sugawara et al., 1995; Zhu et al., 2008). The extent and dynamics of DNA-end resection in mammalian cells however are unknown. Because AID-mediated damage is not site specific but occurs over a relatively large genomic domain ( $\sim$ 1–3 kb; Klein et al., 2011; Yamane et al., 2011), we monitored DNA-end resection in 53BP1<sup>–/–</sup> MEFs transduced with the I-Ppol endonuclease, which recognizes 19 sites in the C57BL/6 genome (San Filippo et al., 2008). RPA recruitment to all 19 I-Ppol sites was strand biased, centered on the DSB, and extended over an area of slightly more than 20 kb (Figure 2E). Analogous results were also obtained at *Myc* upon I-SceI expression in *Myc*<sup>+</sup> B cells (Figure S2A). We conclude that in mammalian cells deficient in 53BP1, resection occurs for up to 10–15 kb on either side of a DSB.


### Rad51 Recruitment to AID-Mediated Breaks

RPA has been studied primarily in the context of HR where it recruits Rad51, a recombinase that forms the presynaptic nucleoprotein filament required for strand invasion (Ogawa et al., 1993; Sung and Roberson, 1995). To determine whether Rad51 also associates with resected DNA damaged by AID, we performed Rad51 ChIP-seq on 53BP1<sup>–/–</sup>*IgkAID* B cells. We found extensive Rad51 recruitment at *Igh* (Figure 3A). Analogous to RPA, the Rad51 ChIP signal was centered at recombining switch domains and displayed a marked DNA strand bias (Figure 3A). In addition, Rad51 was also found at AID off-target genes, such as *Mir155* and *Cd83* (Figure 3B). Additional examples are provided in Figure S3. At these sites, the domain separating the strand-specific Rad51 islands overlapped with hotspots of AID-induced chromosomal translocations (Figure 3B), as determined by TC-Seq (Klein et al., 2011). Moreover, the relative amount of Rad51 deposition per off-target gene (TSS  $\pm$  2 kb) was directly proportional to its translocation frequency (Spearman  $\rho$  = 0.65, Figure 3C). This result is consistent with the notion that, as described for RPA (Hakim et al., 2012), the extent of Rad51 occupancy is a function of AID activity. In contrast, genes not associated with RPA displayed background levels of Rad51 (Figure 3D and not shown). We conclude that in the absence of 53BP1, RPA and Rad51 associate with sites of AID-mediated damage at *Ig* and non-*Ig* genes.

### RPA Recruitment Occurs Mostly in S–G2/M

Recruitment of RPA and Rad51 to resected DNA might occur in G1 during NHEJ repair of V(D)J or CSR DSBs. Alternatively, RPA could be recruited in S and G2/M phases of the cell cycle as part of a salvage mechanism for unrepaired DSBs. To address this question, we purified H2AX<sup>–/–</sup> lymphocytes in the G1, S, and G2/M phases of the cell cycle and performed RPA ChIP-seq. Consistent with limited DNA-end resection during NHEJ (Bothmer et al., 2010; Helmink et al., 2011), G1 cells displayed localized RPA deposition at the *Igh* locus (Figure 4A) in a manner that overlapped with sites of AID activity (Hakim et al., 2012; Klein et al., 2011). In contrast, lymphocytes from S and G2/M phases recruited far greater amounts of RPA (Figure 4A). Analogous results were obtained using 53BP1<sup>–/–</sup> B cells (see below). Thus, RPA recruitment occurs in both G1 and S–G2/M, with the majority of the signal accumulating in cells that progress to S–G2/M.





**Figure 4. RPA and  $\gamma$ H2AX Recruitment during Cell Cycle**

(A) RPA accumulation in H2AX<sup>-/-</sup>-activated B cells at the *Igh* locus during the cell cycle. Samples were stained with Hoechst dye and sorted into G1, S, and G2/M-phased cells. RPKM values for the specified genomic windows are provided in parentheses.

(B) Extent of H2AX phosphorylation ( $\gamma$ H2AX) at the *Igh* locus in wild-type (G1, lane I; S, lane II), H2AX<sup>-/-</sup> (G1, lane III), or 53BP1<sup>-/-</sup> (G1, lane IV; S, lane V)-activated B cells.

(C) RPA (G1, G2/M) and  $\gamma$ H2AX (G2/M) deposition at *Igh* in cells from H2AX<sup>-/-</sup> or 53BP1<sup>-/-</sup> respectively. ChIP-seq data were split into + and - strands.

Our results suggested that in end-joining-deficient cells, a significant fraction of CSR lesions produced in G1 persist into S and G2/M, where they are processed for HR. To explore this possibility, we monitored *Igh* breaks in cycling cells by ChIP for the phosphorylated histone H2AX ( $\gamma$ H2AX), which accumulates at sites of DSBs (Rogakou et al., 1999). In wild-type B cells in G1,  $\gamma$ H2AX was primarily associated with *Igh* (1.40 reads per million per kb [RPKM], Figure 4B, lane I), and these signals decreased in S phase cells to levels found in H2AX<sup>-/-</sup> controls (Figure 4B, lanes II and III). These results support the idea that most CSR breaks are resolved before DNA replication (Hasham et al., 2012; Petersen et al., 2001; Schrader et al., 2007; Sharbeen et al., 2012). In contrast, 53BP1<sup>-/-</sup> B cells showed similar levels of  $\gamma$ H2AX accumulation in G1 and in the S phase (Figure 4B, lane V), indicating that in end-joining-deficient cells a significant fraction of CSR breaks persist beyond G1, even in the presence of intact ATM and p53 (Callén et al., 2007). These data indicate that unrepaired AID-mediated DNA damage does not efficiently activate cell-cycle checkpoints in the absence of 53BP1 or H2AX.

As expected based on results with unsorted samples (Figure S2A), RPA displayed strand biases at *Igh* in all cell-cycle stages (Figure 4C; data not shown). In contrast,  $\gamma$ H2AX signals

were not asymmetric around S domains (Figure 4C), indicating that this chromatin mark, unlike RPA, is mostly associated with unresected double-strand lesions. The persistence of  $\gamma$ H2AX signals in G2/M-phased cells is consistent with the presence of frequent chromosome 12 breaks in NHEJ-deficient metaphases (Bothmer et al., 2011; Bunting et al., 2010; Gostissa et al., 2011).

#### ATM Is Required for Resection in G1 but Not S-G2/M


ATM is required for DNA-end resection during CSR, V(D)J recombination, and in cells treated with clastogenic agents (Bothmer et al., 2010, 2011; Helmink et al., 2011; Jazayeri et al., 2006). To examine the role of ATM in RPA recruitment during CSR, we treated LPS+IL-4-activated B cells with the small molecule ATM inhibitor KU-55933 (ATMi) and performed RPA ChIP-seq on purified G1 and S-G2/M phase cells. ATMi-treated G1 phase cells showed only background RPA signals at non-*Ig* AID targets in the absence of 53BP1. In addition, RPA was 3-fold lower at *Igh* in ATMi-treated cells relative to controls ( $p = 0.02$ ; Figure 5B;  $n = 3$ ). In contrast, activated B cells treated with ATR inhibitor (ATR45) (Charrier et al., 2011; Schoppa et al., 2012) did not show this effect (Figure 5B), which is consistent with the observation that ATR is activated only after significant resection has occurred (Brown and Baltimore, 2003; Pellicioli et al., 2001; Shiotani and Zou, 2009; Zou and Elledge, 2003). Furthermore, ATR inhibitor treatment did not affect the extent of resection in S-G2/M (Morin et al., 2008).

To confirm a role for ATM in initiating resection in G1, we examined 53BP1<sup>-/-</sup> and 53BP1<sup>-/-</sup>ATM<sup>-/-</sup> thymocytes. The analysis showed a substantial reduction in RPA deposition at the TCR $\alpha$  locus in the absence of ATM (Figure 5C). Consistent with ATMi results, RPA signals at *Igh* in unfractionated B cells undergoing CSR from the same mice were not affected (Figure 5D). Taken together, these findings demonstrate that ATM is required to promote initiation of RPA recruitment to resected DNA in G1- but is not required for its recruitment in S-G2/M-phased lymphocytes.

We have shown that in NHEJ-deficient cells (H2AX<sup>-/-</sup> or 53BP1<sup>-/-</sup>), a fraction of DNA breaks that are normally repaired in G1 become substrates for homologous-mediated repair in S/G2/M. To investigate whether this also occurs under physiological conditions, we measured Rad51 and RPA occupancy in activated wild-type B cells undergoing CSR. We found strand-biased recruitment of Rad51 and RPA at *Ig* $\mu$  and *Ig* $\gamma$ 1 (Figure 5E), consistent with 5'-3' resection activity in wild-type cells. Resection required AID activity because RPA signals were increased in the presence of *Ig* $\kappa$ AID and were indistinguishable from background in AID<sup>-/-</sup> lymphocytes (Figure 5E). As expected, the extent of resection in wild-type cells was reduced relative to H2AX<sup>-/-</sup> or 53BP1<sup>-/-</sup> (compare RPKM values to Figures 2B and 5D, for instance). We thus conclude that recombining *Ig* genes are resected under physiological conditions, arguing that nonhomologous and homologous repair pathways contribute to repair during CSR.

#### DISCUSSION

Our results show that RPA is recruited to resected DNA DSBs produced by AID, RAGs, and site-specific endonucleases.



**Figure 5. ATM Is Required for G1 but Not S-G2/M Resection**

(A) RPA accumulation at *Pim1* and *IL4Rα* loci from 53BP1<sup>-/-</sup>-activated B cells that were either treated (lower two panels) or not treated (upper panels) with the ATM inhibitor KU-55933. Samples were sorted into G1 or S/G2/M-phased cells using the Hoechst dye 33342. Numbers in parentheses represent RPKM values within the specified genomic windows.

(B) RPA accumulation at *Igμ* and *Igγ1* loci in WT, ATMi<sup>-</sup>, or ATRi<sup>-</sup>-treated B cells. Values represent the RPKM ratio between G1 and S-G2/M-phased cells. Error bars represent the SD of four biological replicates (two for *Igμ* and two for *Igγ1*) for each experimental condition with the exception of ATRi. p value was calculated using the unpaired t test.

(C and D) RPA recruitment to the *TCRα* in thymocytes (C) or *IgH* in activated B cells (D) from 53BP1<sup>-/-</sup> (upper) or 53BP1<sup>-/-</sup> ATM<sup>-/-</sup> (lower) mice.

(E) RPA and Rad51 recruitment to activated B cells with an intact NHEJ: *IgκAID* transgenics, AID<sup>+/+</sup>, and AID<sup>-/-</sup>.


RPA accumulation begins within 3 hr in the G1 phase of the cell cycle and unrepaired lesions persist into the S and G2/M phases where HR or other salvage repair mechanisms resolve DSBs. While our findings do not exclude the possibility that RPA can enhance AID activity (Chaudhuri et al., 2004), we can conclude that RPA recruitment is not essential for *Ig* gene deamination by AID.

RPA recruitment to DNA damage is asymmetric around DSBs, displaying a strong bias for the + DNA strand 5', and the - strand 3' of the break. This profile is best explained by the 5'-3' directionality of DNA-end resection, which we find spreads 10–15 kb away from the site of the break in both directions as previously described in yeast (Bishop et al., 1992; Sugawara et al., 1995; Zhu et al., 2008). Consistent with this idea, exonuclease treatment of immunoprecipitated DNA, a procedure that specifically removes ssDNA 3' tracks, ablates RPA ChIP signals. In addition, Rad51, which is required for strand invasion during HR, mirrors RPA deposition. The striking overlap between RPA and Rad51 profiles is consistent with the idea that RPA accumulation on resected DNA in S-G2/M facilitates HR (New et al., 1998, Figure 6).

DNA-end resection has been primarily studied in yeast, where the Sae2 (CtIP) nuclease and the MRX (MRN) complex initiate the trimming phase, and the Exo1 and the RecQ helicase Sgs1 extend the resected tracks (Mimitou and Symington, 2008; Symington and Gautier, 2011; Zhu et al., 2008). The mechanistic details of end resection in mammalian cells are less clear, but both CtIP and ATM appear to play a role. During V(D)J recombination,

depletion of CtIP in H2AX<sup>-/-</sup> cells protects DSBs from end resection (Helmink et al., 2011). In like manner, inhibition of ATM in 53BP1<sup>-/-</sup> B cells protects CSR breaks from degradation (Bothmer et al., 2010). Our results confirm a role for ATM in the resection of V(D)J and CSR lesions, but notably, only in the G1 phase of the cell cycle. Thus, an ATM-independent pathway controls resection at post-replicative stages.

The accumulation of RPA at resected AID target genes in the S-G2/M phases of the cell cycle suggest that some AID-induced DSBs are carried over from G1 to S where they would be preferentially repaired by HR. Consistent with this model, B cells deficient in the HR factor XRCC2 accumulate unrepaired AID-mediated breaks at postreplicative stages of the cell cycle (Hasham et al., 2010, 2012). However, the mechanism that allows these lesions to persist beyond G1 was not resolved. ATM is an essential mediator of the G1-S phase checkpoint, and when it is deleted DSBs persist through cell-cycle phase transitions leading to chromosome breaks and genome instability (Bredemeyer et al., 2006; Callén et al., 2007; Derheimer and Kastan, 2010). While it is conceivable that the relative low number of AID-mediated DSBs per cell might be insufficient to activate the checkpoint, this scenario seems unlikely because ATM can be activated by only a few breaks (Bakkenist and Kastan, 2003; Chen et al., 2000; Petersen et al., 2001). Moreover, AID activity engages ATM as indicated by formation of γH2AX foci on recombining *IgH* loci (Petersen et al., 2001). Cells that resolve these breaks by NHEJ extinguish ATM and progress. In contrast, DSBs that fail to resolve would be expected to undergo further processing, including end resection and RPA recruitment, which can suppress ATM signaling and abort the G1-S checkpoint (Callén et al., 2007; Shiotani and Zou, 2009). Thus, RPA



**Figure 6. DNA-End Resection during CSR**

Model showing how AID-mediated breaks at *Igh* are either processed by C-NHEJ proteins into efficient switch recombination, or are resected by ATM-dependent A-NHEJ in G1 or by the HR in S-G2/M, leading to RPA recruitment and repair. Based on previous work (Shiotani and Zou, 2009), extensive resection in connection with HR is expected to inhibit ATM activity and potentially reduce the formation of chromosomal translocations via A-NHEJ.

accumulation in G1 may help avert the G1-S checkpoint and facilitate homology-mediated repair by enabling progression of cells carrying unrepaired breaks from G1 into the S phase.

What is the role of homology-mediated DNA repair during CSR? AID initiates CSR by introducing multiple DSBs in highly repetitive switch regions, which vary in length from 1 to ~10 kb (Honjo et al., 2002). NHEJ repair of paired DSBs in heterologous switch regions results in productive CSR. However, ~20% of the time productive recombination fails, and DSBs are rejoined by an alternative form of nonhomologous end joining (A-NHEJ) that promotes intraswitch deletions (Bottaro et al., 1998; Dudley et al., 2002; Gu et al., 1993; Hummel et al., 1987; Reina-San-Martin et al., 2003; Winter et al., 1987). We propose that this salvage mechanism is mediated by S region microhomology uncovered by DNA-end resection (Bothmer et al., 2010, 2011) and RPA recruitment (Figure 6). The presence of localized RPA at *Igμ* and *Igγ1* in the G1 phase of the cell cycle is consistent with this idea. Whether A-NHEJ is also active at *Igh* in S-G2/M remains to be determined. Our results also imply that DNA breaks not processed by the canonical or alternative NHEJ pathways would be further resected in S-G2/M and repaired by HR using the second, intact *Igh* allele (Figure 6). The physiologic advantage of engaging error-free HR during CSR is that switch regions thus repaired would be available for subsequent recombination attempts in the next G1 phase. In addition, HR should be able to counteract AID-induced mutations and limit the formation of chromosomal translocations.

## EXPERIMENTAL PROCEDURES

### Mice

*IgκAID* (Robbiani et al., 2009); *UNG*<sup>-/-</sup> (Endres et al., 2004); *ATM*<sup>-/-</sup> (Barlow et al., 1996); *AID*<sup>-/-</sup> (Muramatsu et al., 2000), *53BP1*<sup>-/-</sup> (Ward et al., 2003),

*Myc*<sup>l</sup> (Robbiani et al., 2008); *Msh2*<sup>-/-</sup> (Reitmair et al., 1995); and *H2AX*<sup>-/-</sup> (Bassing et al., 2003) mice were previously described. All experiments were in accordance with protocols approved by the NIAMS and Rockefeller Institutional Animal Care and Use Committee.

### ChIP-seq

Ex vivo cultured B cells or thymocytes were fixed with 1% formaldehyde (Sigma) for 10 min at 37°C. The fixation was quenched by addition of glycine (Sigma) at a final concentration of 125 mM. Twenty million fixed cells were washed with PBS and then resuspended in 1 ml of RIPA buffer (10 mM Tris [pH 7.6], 1 mM EDTA, 0.1% SDS, 0.1% sodium deoxycholate, 1% Triton X-100, 1× Complete Mini EDTA free proteinase inhibitor [Roche]) or stored at -80°C until further processing. The sonication was performed on S2 sonicator (Covaris) at duty cycle 20%, intensity 5, cycle/burst 200 for 30 min. Ten micrograms of anti-RPA32 (EMD, NA19L), anti-RAD51 (Santa Cruz, H-92), or anti-γH2AX (Epitomics, 2212-1) was incubated with 40 μl of Dynabeads Protein A for 40 min at room temperature. The antibody-bound beads were added to 500 μl of sonicated chromatin, incubated at 4°C overnight, and washed twice with RIPA buffer, twice with RIPA buffer containing 0.3M NaCl, twice with LiCl buffer (0.25 M LiCl, 0.5% Igepal-630, 0.5% sodium deoxycholate), once with TE (pH 8) plus 0.2% Triton X-100, and once with TE (pH 8). Crosslinking was reversed by incubating the beads at 65°C for 4 hr with 0.3% SDS and 1 mg/ml Proteinase K. ChIP DNA was purified by phenol-chloroform extraction followed by ethanol precipitation. The DNA was subsequently blunt-ended with End-It DNA end repair kit (Epicenter) and A-tailed with Taq DNA polymerase (Invitrogen) in the presence of 200mM of dATP for 40 min at 70°C. The sample was purified by phenol-chloroform extraction after each reaction. Illumina compatible adaptors (Illumina or Bioo Scientific) were then ligated with T4 DNA ligase (Enzymatics), and the reaction was purified once with AMPure XP magnetic beads (Beckman Coulter). Samples were PCR amplified for 18 cycles with KAPA HiFi DNA polymerase mix (KAPA Biosystems). The amplicon was run on a 2% agarose gel and size-selected at 200–300 bp. Thirty-six or 50 bp of sequencing data were acquired on GAll or HiSeq2000 (Illumina). For nuclease treatment, ChIP DNA was incubated for 2 hr with 20 units of *E. coli* Exol (New England Biolabs) at 37°C or 30 units of RecJf nuclease (New England Biolabs) at 37°C for 2 hr prior to the blunt-end ligation step. How ssDNA is amplified and detected during the Illumina library protocol has been recently addressed by two independent studies (Croucher et al., 2009; Khil et al., 2012). In brief, ssDNA is rendered double-stranded by intermolecular or intramolecular (hairpin) priming, facilitated by DNA microhomologies (Croucher et al., 2009; Khil et al., 2012). Table S1 contains a list of all ChIP-seq experiments as they appear in the manuscript figures. Biological replicates or comparable experiments that further support the results presented in each panel are also provided.

### Cell Culture and Retroviral Infection

For I-SceI assays, CD43<sup>+</sup> resting B cells were isolated from spleens, stimulated, and infected as previously described (Bothmer et al., 2011). The pMX-IRES-GFP based retrovirus encoding I-SceI was previously described (Robbiani et al., 2008). The new pMX-ER-I-SceI was generated by fusing in-frame the optimized ligand-binding domain of the human estrogen receptor (ERT2) with I-SceI. For I-Ppol assays, immortalized 53BP1<sup>-/-</sup> mouse embryonic fibroblasts (MEFs) were transduced with the pMY retroviral vector encoding BFP-I-Ppol fusion gene. Following spin-infection MEFs were reseeded until they reached ~70% confluency, at which time cells were trypsinized for 5 seconds at 37°C, and trypsin was quenched with complete media containing 1.25% formaldehyde for 10 seconds. Following incubation a final concentration of 125 mM glycine was added to stop crosslinking.

### Flow Cytometry and Cell Sorting

I-SceI-GFP B cells and BFP-I-Ppol MEFs were sorted on a FACSAriaIII (Becton Dickinson) or MoFlo (Beckman Coulter). For cell-cycle sorting, B cells were cultured for 72 hr with LPS, IL-4, and anti-Cd180 antibody and incubated for an additional 45 min in the presence of 10 μg/ml of Hoechst 33342 (Invitrogen). Cells were then fixed the same conditions used for ChIP-seq sample preparation. Samples were resuspended in HBSS buffer containing

Hoechst 33342 and G1, S, or G2/M phase cells were sorted with Influx or Arial machines (BD Biosciences) using 355 or 375 nm lasers, respectively.

## ACCESSION NUMBERS

Deep-sequencing data are available at GEO under accession number GSE42451.

## SUPPLEMENTAL INFORMATION

Supplemental Information includes three figures and one table and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2012.12.006>.

## LICENSING INFORMATION

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

## ACKNOWLEDGMENTS

We thank all members of the Casellas and Nussenzweig labs for helpful discussions, Matthias F. Muellenbeck, Jim Simone, Jeff Lay, and Gustavo Gutierrez for technical assistance. The work was supported in part by the intramural program of NIAMS at the NIH. This study utilized the high-performance computational capabilities of the Helix/BioWolf Systems at the National Institutes of Health, Bethesda, MD (<http://helix.nih.gov>).

Received: September 21, 2012

Revised: November 14, 2012

Accepted: December 12, 2012

Published: January 3, 2013

## REFERENCES

- Bakkenist, C.J., and Kastan, M.B. (2003). DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. *Nature* 421, 499–506.
- Barlow, C., Hirotsumi, S., Paylor, R., Liyanage, M., Eckhaus, M., Collins, F., Shiloh, Y., Crawley, J.N., Ried, T., Tagle, D., and Wynshaw-Boris, A. (1996). Atm-deficient mice: a paradigm of ataxia telangiectasia. *Cell* 86, 159–171.
- Bassing, C.H., Suh, H., Ferguson, D.O., Chua, K.F., Manis, J., Eckersdorff, M., Gleason, M., Bronson, R., Lee, C., and Alt, F.W. (2003). Histone H2AX: a dosage-dependent suppressor of oncogenic translocations and tumors. *Cell* 114, 359–370.
- Basu, U., Meng, F.L., Keim, C., Grinstein, V., Pefanis, E., Eccleston, J., Zhang, T., Myers, D., Wasserman, C.R., Wesemann, D.R., et al. (2011). The RNA exosome targets the AID cytidine deaminase to both strands of transcribed duplex DNA substrates. *Cell* 144, 353–363.
- Bishop, D.K., Park, D., Xu, L., and Kleckner, N. (1992). DMC1: a meiosis-specific yeast homolog of E. coli recA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell* 69, 439–456.
- Bothmer, A., Robbiani, D.F., Feldhahn, N., Gazumyan, A., Nussenzweig, A., and Nussenzweig, M.C. (2010). 53BP1 regulates DNA resection and the choice between classical and alternative end joining during class switch recombination. *J. Exp. Med.* 207, 855–865.
- Bothmer, A., Robbiani, D.F., Di Virgilio, M., Bunting, S.F., Klein, I.A., Feldhahn, N., Barlow, J., Chen, H.T., Bosque, D., Callen, E., et al. (2011). Regulation of DNA end joining, resection, and immunoglobulin class switch recombination by 53BP1. *Mol. Cell* 42, 319–329.
- Bottaro, A., Young, F., Chen, J., Serwe, M., Sablitzky, F., and Alt, F.W. (1998). Deletion of the IgH intronic enhancer and associated matrix-attachment regions decreases, but does not abolish, class switching at the mu locus. *Int. Immunol.* 10, 799–806.
- Bredemeyer, A.L., Sharma, G.G., Huang, C.Y., Helmink, B.A., Walker, L.M., Khor, K.C., Nuskey, B., Sullivan, K.E., Pandita, T.K., Bassing, C.H., and Sleckman, B.P. (2006). ATM stabilizes DNA double-strand-break complexes during V(D)J recombination. *Nature* 442, 466–470.
- Brown, E.J., and Baltimore, D. (2003). Essential and dispensable roles of ATR in cell cycle arrest and genome maintenance. *Genes Dev.* 17, 615–628.
- Bunting, S.F., Callén, E., Wong, N., Chen, H.T., Polato, F., Gunn, A., Bothmer, A., Feldhahn, N., Fernandez-Capetillo, O., Cao, L., et al. (2010). 53BP1 inhibits homologous recombination in Brca1-deficient cells by blocking resection of DNA breaks. *Cell* 141, 243–254.
- Callén, E., Jankovic, M., Difilippantonio, S., Daniel, J.A., Chen, H.T., Celeste, A., Pellegrini, M., McBride, K., Wangsa, D., Bredemeyer, A.L., et al. (2007). ATM prevents the persistence and propagation of chromosome breaks in lymphocytes. *Cell* 130, 63–75.
- Charrier, J.D., Durrant, S.J., Golec, J.M., Kay, D.P., Knechtel, R.M., MacGormick, S., Mortimore, M., O'Donnell, M.E., Pinder, J.L., Reaper, P.M., et al. (2011). Discovery of potent and selective inhibitors of ataxia telangiectasia mutated and Rad3 related (ATR) protein kinase as potential anticancer agents. *J. Med. Chem.* 54, 2320–2330.
- Chaudhuri, J., Khuong, C., and Alt, F.W. (2004). Replication protein A interacts with AID to promote deamination of somatic hypermutation targets. *Nature* 430, 992–998.
- Chen, H.T., Bhandoola, A., Difilippantonio, M.J., Zhu, J., Brown, M.J., Tai, X., Rogakou, E.P., Brotz, T.M., Bonner, W.M., Ried, T., and Nussenzweig, A. (2000). Response to RAG-mediated VDJ cleavage by NBS1 and gamma-H2AX. *Science* 290, 1962–1965.
- Chiarle, R., Zhang, Y., Frock, R.L., Lewis, S.M., Molinie, B., Ho, Y.J., Myers, D.R., Choi, V.W., Compagno, M., Malkin, D.J., et al. (2011). Genome-wide translocation sequencing reveals mechanisms of chromosome breaks and rearrangements in B cells. *Cell* 147, 107–119.
- Croucher, N.J., Fookes, M.C., Perkins, T.T., Turner, D.J., Marguerat, S.B., Keane, T., Quail, M.A., He, M., Assefa, S., Bähler, J., et al. (2009). A simple method for directional transcriptome sequencing using Illumina technology. *Nucleic Acids Res.* 37, e148.
- DeMott, M.S., Zigman, S., and Bambara, R.A. (1998). Replication protein A stimulates long patch DNA base excision repair. *J. Biol. Chem.* 273, 27492–27498.
- Derheimer, F.A., and Kastan, M.B. (2010). Multiple roles of ATM in monitoring and maintaining DNA integrity. *FEBS Lett.* 584, 3675–3681.
- Difilippantonio, S., Gapud, E., Wong, N., Huang, C.Y., Mahowald, G., Chen, H.T., Kruhlak, M.J., Callen, E., Livak, F., Nussenzweig, M.C., et al. (2008). 53BP1 facilitates long-range DNA end-joining during V(D)J recombination. *Nature* 456, 529–533.
- Dudley, D.D., Manis, J.P., Zarrin, A.A., Kaylor, L., Tian, M., and Alt, F.W. (2002). Internal IgH class switch region deletions are position-independent and enhanced by AID expression. *Proc. Natl. Acad. Sci. USA* 99, 9984–9989.
- Endres, M., Biniszkiwicz, D., Sobol, R.W., Harms, C., Ahmadi, M., Lipski, A., Katchanov, J., Mergenthaler, P., Dirnagl, U., Wilson, S.H., et al. (2004). Increased postischemic brain injury in mice deficient in uracil-DNA glycosylase. *J. Clin. Invest.* 113, 1711–1721.
- Fugmann, S.D., Lee, A.I., Shockett, P.E., Vile, J., and Schatz, D.G. (2000). The RAG proteins and V(D)J recombination: complexes, ends, and transposition. *Annu. Rev. Immunol.* 18, 495–527.
- Genschel, J., and Modrich, P. (2003). Mechanism of 5'-directed excision in human mismatch repair. *Mol. Cell* 12, 1077–1086.
- Gostissa, M., Alt, F.W., and Chiarle, R. (2011). Mechanisms that promote and suppress chromosomal translocations in lymphocytes. *Annu. Rev. Immunol.* 29, 319–350.
- Gu, H., Zou, Y.R., and Rajewsky, K. (1993). Independent control of immunoglobulin switch recombination at individual switch regions evidenced through Cre-loxP-mediated gene targeting. *Cell* 73, 1155–1164.
- Hakim, O., Resch, W., Yamane, A., Klein, I., Kieffer-Kwon, K.-R., Jankovic, M., Oliveira, T., Bothmer, A., Voss, T.C., Ansarah-Sobrinho, C., et al. (2012). DNA



- damage defines sites of recurrent chromosomal translocations in B lymphocytes. *Nature* 484, 69–74.
- Hasham, M.G., Donghia, N.M., Coffey, E., Maynard, J., Snow, K.J., Ames, J., Wilpan, R.Y., He, Y., King, B.L., and Mills, K.D. (2010). Widespread genomic breaks generated by activation-induced cytidine deaminase are prevented by homologous recombination. *Nat. Immunol.* 11, 820–826.
- Hasham, M.G., Snow, K.J., Donghia, N.M., Branca, J.A., Lessard, M.D., Stavnezer, J., Shopland, L.S., and Mills, K.D. (2012). Activation-induced cytidine deaminase-initiated off-target DNA breaks are detected and resolved during S phase. *J. Immunol.* 189, 2374–2382.
- Helmink, B.A., and Sleckman, B.P. (2012). The response to and repair of RAG-mediated DNA double-strand breaks. *Annu. Rev. Immunol.* 30, 175–202.
- Helmink, B.A., Tubbs, A.T., Dorsett, Y., Bednarski, J.J., Walker, L.M., Feng, Z., Sharma, G.G., McKinnon, P.J., Zhang, J., Bassing, C.H., and Sleckman, B.P. (2011). H2AX prevents CtIP-mediated DNA end resection and aberrant repair in G1-phase lymphocytes. *Nature* 469, 245–249.
- Honjo, T., Kinoshita, K., and Muramatsu, M. (2002). Molecular mechanism of class switch recombination: linkage with somatic hypermutation. *Annu. Rev. Immunol.* 20, 165–196.
- Hummel, M., Berry, J.K., and Dunnick, W. (1987). Switch region content of hybridomas: the two spleen cell IgH loci tend to rearrange to the same isotype. *J. Immunol.* 138, 3539–3548.
- Jazayeri, A., Falck, J., Lukas, C., Bartek, J., Smith, G.C., Lukas, J., and Jackson, S.P. (2006). ATM- and cell cycle-dependent regulation of ATR in response to DNA double-strand breaks. *Nat. Cell Biol.* 8, 37–45.
- Ji, Y., Resch, W., Corbett, E., Yamane, A., Casellas, R., and Schatz, D.G. (2010). The in vivo pattern of binding of RAG1 and RAG2 to antigen receptor loci. *Cell* 141, 419–431.
- Jung, D., Giallourakis, C., Mostoslavsky, R., and Alt, F.W. (2006). Mechanism and control of V(D)J recombination at the immunoglobulin heavy chain locus. *Annu. Rev. Immunol.* 24, 541–570.
- Khil, P.P., Smagulova, F., Brick, K.M., Camerini-Otero, R.D., and Petukhova, G.V. (2012). Sensitive mapping of recombination hotspots using sequencing-based detection of ssDNA. *Genome Res.* 22, 957–965.
- Klein, I.A., Resch, W., Jankovic, M., Oliveira, T., Yamane, A., Nakahashi, H., Di Virgilio, M., Bothmer, A., Nussenzweig, A., Robbiani, D.F., et al. (2011). Translocation-capture sequencing reveals the extent and nature of chromosomal rearrangements in B lymphocytes. *Cell* 147, 95–106.
- Küppers, R., and Dalla-Favera, R. (2001). Mechanisms of chromosomal translocations in B cell lymphomas. *Oncogene* 20, 5580–5594.
- Lin, Y.L., Shivji, M.K., Chen, C., Kolodner, R., Wood, R.D., and Dutta, A. (1998). The evolutionarily conserved zinc finger motif in the largest subunit of human replication protein A is required for DNA replication and mismatch repair but not for nucleotide excision repair. *J. Biol. Chem.* 273, 1453–1461.
- Liu, M., Duke, J.L., Richter, D.J., Vinuesa, C.G., Goodnow, C.C., Kleinstein, S.H., and Schatz, D.G. (2008). Two levels of protection for the B cell genome during somatic hypermutation. *Nature* 451, 841–845.
- Maul, R.W., Saribasak, H., Martomo, S.A., McClure, R.L., Yang, W., Vaisman, A., Gramlich, H.S., Schatz, D.G., Woodgate, R., Wilson, D.M., 3rd, and Gearhart, P.J. (2011). Uracil residues dependent on the deaminase AID in immunoglobulin gene variable and switch regions. *Nat. Immunol.* 12, 70–76.
- Mimitou, E.P., and Symington, L.S. (2008). Sae2, Exo1 and Sgs1 collaborate in DNA double-strand break processing. *Nature* 455, 770–774.
- Morin, I., Ngo, H.P., Greenall, A., Zubko, M.K., Morrice, N., and Lydall, D. (2008). Checkpoint-dependent phosphorylation of Exo1 modulates the DNA damage response. *EMBO J.* 27, 2400–2410.
- Muramatsu, M., Kinoshita, K., Fagarasan, S., Yamada, S., Shinkai, Y., and Honjo, T. (2000). Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* 102, 553–563.
- New, J.H., Sugiyama, T., Zaitseva, E., and Kowalczykowski, S.C. (1998). Rad52 protein stimulates DNA strand exchange by Rad51 and replication protein A. *Nature* 391, 407–410.
- Oakley, G.G., and Patrick, S.M. (2010). Replication protein A: directing traffic at the intersection of replication and repair. *Front. Biosci.* 15, 883–900.
- Ogawa, T., Yu, X., Shinohara, A., and Egelman, E.H. (1993). Similarity of the yeast RAD51 filament to the bacterial RecA filament. *Science* 259, 1896–1899.
- Pasqualucci, L., Migliazza, A., Fracchiolla, N., William, C., Neri, A., Baldini, L., Chaganti, R.S., Klein, U., Küppers, R., Rajewsky, K., and Dalla-Favera, R. (1998). BCL-6 mutations in normal germinal center B cells: evidence of somatic hypermutation acting outside Ig loci. *Proc. Natl. Acad. Sci. USA* 95, 11816–11821.
- Pavri, R., Gazumyan, A., Jankovic, M., Di Virgilio, M., Klein, I., Ansarah-Sobrinho, C., Resch, W., Yamane, A., Reina San-Martin, B., Barreto, V., et al. (2010). Activation-induced cytidine deaminase targets DNA at sites of RNA polymerase II stalling by interaction with Spt5. *Cell* 143, 122–133.
- Pelliccioli, A., Lee, S.E., Lucca, C., Foiani, M., and Haber, J.E. (2001). Regulation of Saccharomyces Rad53 checkpoint kinase during adaptation from DNA damage-induced G2/M arrest. *Mol. Cell* 7, 293–300.
- Peters, A., and Storb, U. (1996). Somatic hypermutation of immunoglobulin genes is linked to transcription initiation. *Immunity* 4, 57–65.
- Petersen, S., Casellas, R., Reina-San-Martin, B., Chen, H.T., Difilippantonio, M.J., Wilson, P.C., Hanitsch, L., Celeste, A., Muramatsu, M., Pilch, D.R., et al. (2001). AID is required to initiate Nbs1/gamma-H2AX focus formation and mutations at sites of class switching. *Nature* 414, 660–665.
- Rada, C., Di Noia, J.M., and Neuberger, M.S. (2004). Mismatch recognition and uracil excision provide complementary paths to both Ig switching and the A/T-focused phase of somatic mutation. *Mol. Cell* 16, 163–171.
- Rajagopal, D., Maul, R.W., Ghosh, A., Chakraborty, T., Khamlichi, A.A., Sen, R., and Gearhart, P.J. (2009). Immunoglobulin switch mu sequence causes RNA polymerase II accumulation and reduces dA hypermutation. *J. Exp. Med.* 206, 1237–1244.
- Ranalli, T.A., DeMott, M.S., and Bambara, R.A. (2002). Mechanism underlying replication protein a stimulation of DNA ligase I. *J. Biol. Chem.* 277, 1719–1727.
- Reina-San-Martin, B., Difilippantonio, S., Hanitsch, L., Masilamani, R.F., Nussenzweig, A., and Nussenzweig, M.C. (2003). H2AX is required for recombination between immunoglobulin switch regions but not for intra-switch region recombination or somatic hypermutation. *J. Exp. Med.* 197, 1767–1778.
- Reitmair, A.H., Schmits, R., Ewel, A., Bapat, B., Redston, M., Mitri, A., Waterhouse, P., Mittrücker, H.W., Wakeham, A., Liu, B., et al. (1995). MSH2 deficient mice are viable and susceptible to lymphoid tumours. *Nat. Genet.* 11, 64–70.
- Robbiani, D.F., Bothmer, A., Callen, E., Reina San-Martin, B., Dorsett, Y., Difilippantonio, S., Bolland, D.J., Chen, H.T., Corcoran, A.E., Nussenzweig, A., et al. (2008). Activation Induced Deaminase is Required for the Chromosomal Translocations in c-myc that lead to c-myc/IgH translocations. *Cell* 135, 1028–1038.
- Robbiani, D.F., Bunting, S., Feldhahn, N., Bothmer, A., Camps, J., Deroubaix, S., McBride, K.M., Klein, I.A., Stone, G., Eisenreich, T.R., et al. (2009). AID produces DNA double-strand breaks in non-Ig genes and mature B cell lymphomas with reciprocal chromosome translocations. *Mol. Cell* 36, 631–641.
- Rogakou, E.P., Boon, C., Redon, C., and Bonner, W.M. (1999). Megabase chromatin domains involved in DNA double-strand breaks in vivo. *J. Cell Biol.* 146, 905–916.
- Rooney, S., Chaudhuri, J., and Alt, F.W. (2004). The role of the non-homologous end-joining pathway in lymphocyte development. *Immunol. Rev.* 200, 115–131.
- San Filippo, J., Sung, P., and Klein, H. (2008). Mechanism of eukaryotic homologous recombination. *Annu. Rev. Biochem.* 77, 229–257.
- Schoppy, D.W., Ragland, R.L., Gilad, O., Shastri, N., Peters, A.A., Murga, M., Fernandez-Capetillo, O., Diehl, J.A., and Brown, E.J. (2012). Oncogenic stress sensitizes murine cancers to hypomorphic suppression of ATR. *J. Clin. Invest.* 122, 241–252.
- Schrader, C.E., Guikema, J.E., Linehan, E.K., Selsing, E., and Stavnezer, J. (2007). Activation-induced cytidine deaminase-dependent DNA breaks in

- class switch recombination occur during G1 phase of the cell cycle and depend upon mismatch repair. *J. Immunol.* 179, 6064–6071.
- Sharbeen, G., Yee, C.W., Smith, A.L., and Jolly, C.J. (2012). Ectopic restriction of DNA repair reveals that UNG2 excises AID-induced uracils predominantly or exclusively during G1 phase. *J. Exp. Med.* 209, 965–974.
- Shen, H.M., Peters, A., Baron, B., Zhu, X., and Storb, U. (1998). Mutation of BCL-6 gene in normal B cells by the process of somatic hypermutation of Ig genes. *Science* 280, 1750–1752.
- Shiotani, B., and Zou, L. (2009). Single-stranded DNA orchestrates an ATM-to-ATR switch at DNA breaks. *Mol. Cell* 33, 547–558.
- Stavnezer, J., Guikema, J.E., and Schrader, C.E. (2008). Mechanism and regulation of class switch recombination. *Annu. Rev. Immunol.* 26, 261–292.
- Sugawara, N., Ivanov, E.L., Fishman-Lobell, J., Ray, B.L., Wu, X., and Haber, J.E. (1995). DNA structure-dependent requirements for yeast RAD genes in gene conversion. *Nature* 373, 84–86.
- Sung, P., and Robberson, D.L. (1995). DNA strand exchange mediated by a RAD51-ssDNA nucleoprotein filament with polarity opposite to that of RecA. *Cell* 82, 453–461.
- Symington, L.S., and Gautier, J. (2011). Double-strand break end resection and repair pathway choice. *Annu. Rev. Genet.* 45, 247–271.
- Vuong, B.Q., Lee, M., Kabir, S., Irimia, C., Macchiarulo, S., McKnight, G.S., and Chaudhuri, J. (2009). Specific recruitment of protein kinase A to the immunoglobulin locus regulates class-switch recombination. *Nat. Immunol.* 10, 420–426.
- Wang, L., Wuerffel, R., Feldman, S., Khamlichi, A.A., and Kenter, A.L. (2009). S region sequence, RNA polymerase II, and histone modifications create chromatin accessibility during class switch recombination. *J. Exp. Med.* 206, 1817–1830.
- Ward, I.M., Minn, K., van Deursen, J., and Chen, J. (2003). p53 Binding protein 53BP1 is required for DNA damage responses and tumor suppression in mice. *Mol. Cell. Biol.* 23, 2556–2563.
- Winter, E., Krawinkel, U., and Radbruch, A. (1987). Directed Ig class switch recombination in activated murine B cells. *EMBO J.* 6, 1663–1671.
- Wold, M.S. (1997). Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu. Rev. Biochem.* 66, 61–92.
- Xue, K., Rada, C., and Neuberger, M.S. (2006). The in vivo pattern of AID targeting to immunoglobulin switch regions deduced from mutation spectra in *msh2*<sup>-/-</sup> *ung*<sup>-/-</sup> mice. *J. Exp. Med.* 203, 2085–2094.
- Yamane, A., Resch, W., Kuo, N., Kuchen, S., Li, Z., Sun, H.W., Robbani, D.F., McBride, K., Nussenzweig, M.C., and Casellas, R. (2011). Deep-sequencing identification of the genomic targets of the cytidine deaminase AID and its cofactor RPA in B lymphocytes. *Nat. Immunol.* 12, 62–69.
- Zhang, Y., Gostissa, M., Hildebrand, D.G., Becker, M.S., Boboila, C., Chiarle, R., Lewis, S., and Alt, F.W. (2010). The role of mechanistic factors in promoting chromosomal translocations found in lymphoid and other cancers. *Adv. Immunol.* 106, 93–133.
- Zhu, Z., Chung, W.H., Shim, E.Y., Lee, S.E., and Ira, G. (2008). Sgs1 helicase and two nucleases Dna2 and Exo1 resect DNA double-strand break ends. *Cell* 134, 981–994.
- Zou, L., and Elledge, S.J. (2003). Sensing DNA damage through ATRIP recognition of RPA-ssDNA complexes. *Science* 300, 1542–1548.