

DETECTION RECORDS

199

▲ 8% vs prev month

TOTAL DETECTIONS

210

99% disease incidence

LEADING DISEASE

Bud Root


123 occurrences

AVG CONFIDENCE

79.9%


98% GPS coverage

■ DISEASE DISTRIBUTION OVERVIEW




Disease	Detections	Avg Confidence	Severity
bud root	123	86.1%	Critical
leaf rot	31	81.5%	Moderate
Drying of Leaflets	18	68.3%	Moderate
pestalotiopsis	14	58.8%	Severe
cercospora	14	72.1%	Moderate
healthy	5	61.3%	Mild
Healthy	3	75.6%	Mild
Cercospora	1	37.8%	Moderate
Pestalotiopsis	1	27.4%	Severe

MONTHLY DETECTION TREND




The monthly trend shows detection activity across the reporting period. **bud root** remains the most frequent result. Disease conditions represent **98.6%** of all classified findings. ▲ **8% vs prev month** in submitted records.

DISEASED VS HEALTHY — MONTHLY BREAKDOWN




SEVERITY ANALYSIS BY MAP AREA





Area	Samples	Critical	Severe	Moderate	Mild
Area 1	0	0	0	0	0
Area 2	164	81	15	61	7
Area 3	0	0	0	0	0
Area 4	0	0	0	0	0

Priority alert: bud root is the leading classified disease with 123 detections at 86.1% avg confidence. Disease incidence stands at 98.6% of all detections across 199 submitted records.

DETECTION ACTIVITY BY DAY OF WEEK




■ VERIFICATION & SOURCE BREAKDOWN



196 of 199 records have GPS coordinates (98% coverage). Records from drone source: 0. Records from photo upload: 199.

■ CONFIDENCE SCORE DISTRIBUTION



■ TOP 5 HIGHEST-CONFIDENCE DETECTIONS

#	Disease	Confidence	Source	Timestamp
1	bud root	97.9%	Upload	2026-07-29 10:06:43
2	bud root	97.9%	Upload	2026-07-29 10:04:22
3	leaf rot	97.8%	Upload	2026-08-22 11:03:44
4	leaf rot	97.8%	Upload	2026-08-19 04:44:01
5	leaf rot	97.7%	Upload	2026-08-22 11:03:42

GPS Data Coverage: 196 of 199 records contain GPS coordinates (98.5%). Records without GPS are excluded from the map-area analysis above.

■ EXPERT TREATMENT RECOMMENDATIONS

Guidance below is sourced from the system's expert recommendation database and is aligned with Philippine Coconut Authority (PCA) guidelines. Always confirm diagnosis and pesticide registration with your local agricultural office before treatment.

Bud Rot Critical

Fertilizer: Do not use fertilizer as a cure. After removing active rot, restore palm vigor with organic matter and a soil-

Treatment: Treat as urgent Phytophthora bud rot: contact the Philippine Coconut Authority or local agricultural office. R

Prevention:

- Improve drainage and air flow; clean the crown before the rainy season.
- Remove and destroy infected nuts, fronds, and nearby alternate-host material.
- Inspect nearby palms, especially within about 20 m, and avoid wounding the crown.

PCA-aligned disease-management guidance. Confirm diagnosis and pesticide registration with the local

Leaf Rot Moderate

Fertilizer: sdsd

Treatment: dsdsfsdfe32

Prevention:

- Maintain good drainage, mulch, and water during dry periods.
- Avoid crown and leaf injury and remove diseased debris.
- Monitor palms regularly so leaf symptoms are diagnosed early.

Davao region recommendations (High confidence - follow recommendations closely). Location: davao def

Pestalotiopsis Severe

Fertilizer: Use a soil-test-based coconut fertilizer program. Keep potassium and chloride adequate; PCA reports KCl nutrit

Treatment: Practice clean culture: prune and remove severely infected leaves and debris. After confirmation, apply only a

Prevention:

- Use clean planting material and remove infected leaves promptly.
- Keep nursery and field sanitation high and improve airflow and drainage.
- Minimize leaf wounds and monitor frequently in wet weather.

PCA-aligned fungal leaf-spot guidance. Confirm the diagnosis locally before applying pesticide.

Cercospora

Moderate

- Fertilizer:** Fertilize from a soil or leaf analysis. Maintain adequate potassium and chloride nutrition—PCA notes that KCl
- Treatment:** Prune only heavily affected fronds and remove them from the farm. Have the diagnosis confirmed, then use only
- Prevention:**
- Remove diseased fallen leaves and maintain clean culture.
 - Improve spacing, airflow, and drainage; avoid prolonged leaf wetness where possible.
 - Disinfect pruning tools between palms.

PCA-aligned leaf-spot guidance. Use only products registered for the crop and disease in your area.

Healthy

Mild

- Fertilizer:** Apply balanced NPK 12:12:12 every 3 months. Use phosphate rich during flowering season.
- Treatment:** ddfdfg
- Prevention:**
- dsdfsdgdfg

Davao region recommendations (Good confidence - follow recommendations). Location: davao default