

Appendix A

FOURIER TRANSFORM

This appendix gives the definition of the one-, two-, and three-dimensional Fourier transforms as well as their properties.

A.1 One-Dimensional Fourier Transform

If we have a one-dimensional (1-D) function $f(x)$, its Fourier transform $F(m)$ is defined as

$$F(m) = \int_{-\infty}^{\infty} f(x) \exp(-2\pi i m x) dx, \quad (\text{A.1.1})$$

where m is a variable in the Fourier space. Usually it is termed the frequency in the Fourier domain. If x is a variable in the spatial domain, m is called the spatial frequency. If x represents time, then m is the temporal frequency which denotes, for example, the colour of light in optics or the tone of sound in acoustics.

In this book, we call Eq. (A.1.1) the inverse Fourier transform because a minus sign appears in the exponent. Eq. (A.1.1) can be symbolically expressed as

$$F(m) = \mathcal{F}^{-1}\{f(x)\}. \quad (\text{A.1.2})$$

where \mathcal{F}^{-1} denotes the inverse Fourier transform in Eq. (A.1.1).

Therefore, the direct Fourier transform in this case is given by

$$f(x) = \int_{-\infty}^{\infty} F(m) \exp(2\pi i m x) dm, \quad (\text{A.1.3})$$

which can be symbolically rewritten as

$$f(x) = \mathcal{F}\{F(m)\}. \quad (\text{A.1.4})$$

Substituting Eq. (A.1.2) into Eq. (A.1.4) results in

$$f(x) = \mathcal{F} \mathcal{F}^{-1}\{f(x)\}. \quad (\text{A.1.5})$$

Therefore we have the following unity relation:

$$\mathcal{F} \mathcal{F}^{-1} = 1. \quad (\text{A.1.6})$$

It means that performing a direct Fourier transform and an inverse Fourier transform of a function $f(x)$ successively leads to no change.

Using the identity $\exp(ix) = \cos x + i \sin x$ and Eq. (A.1.1), one has

$$F(m) = A(m) - iB(m), \quad (\text{A.1.7})$$

where

$$\begin{cases} A(m) = \int_{-\infty}^{\infty} f(x) \cos(2\pi m x) dx, \\ B(m) = \int_{-\infty}^{\infty} f(x) \sin(2\pi m x) dx. \end{cases} \quad (\text{A.1.8})$$

A.2 Two-Dimensional Fourier Transform

In a similar way, the direct and inverse Fourier transforms of a two-dimensional (2-D) function $f(x, y)$ can be expressed as

$$f(x, y) = \int_{-\infty}^{\infty} F(m, n) \exp[2\pi i(mx + ny)] dm dn, \quad (\text{A.2.1})$$

and

$$F(m, n) = \int_{-\infty}^{\infty} f(x, y) \exp[-2\pi i(mx + ny)] dx dy, \quad (\text{A.2.2})$$

respectively.

If x and y are spatial coordinates, the exponent in Eq. (A.2.2), $\exp[-2\pi i(mx + ny)]$, represents a plane wave if we recognize that $k_x = 2\pi m$ and $k_y = 2\pi n$. Here k_x and k_y are the components of the wave vector \mathbf{k} in the x and y directions, i.e.

$$k_x^2 + k_y^2 = k^2 = \frac{2\pi}{\lambda}, \quad (\text{A.2.3})$$

where λ is the illumination wavelength. In other words, the inverse Fourier transform of a spatial function is equivalent to resolving the function into a series of plane waves propagating in different directions and the direct Fourier transform means that the original function is represented by a superposition of these plane waves each of which has a particular weighting factor.

A.3 Three-Dimensional Fourier Transform

For a three-dimensional (3-D) function $f(x, y, z)$, we have the direct and inverse 3-D Fourier transforms as follows:

$$f(x, y, z) = \int_{-\infty}^{\infty} F(m, n, s) \exp[2\pi i(mx + ny + sz)] dm dn ds, \quad (\text{A.3.1})$$

and

$$F(m, n, s) = \int_{-\infty}^{\infty} f(x, y, z) \exp[-2\pi i(mx + ny + sz)] dx dy dz. \quad (\text{A.3.2})$$




Fig. A.3.1 A plane wave corresponds to an exponent in Eq. (A.3.1).

According to the discussion in Section A.2, the exponent in Eqs. (A.3.2) denotes a plane wave with the wave-vector components, k_x , k_y and k_z which are given by

$$\begin{cases} k_x = 2\pi m, \\ k_y = 2\pi n, \\ k_z = 2\pi s. \end{cases} \quad (\text{A.3.3})$$

If $2\pi(mx + ny + sz) = \text{constant} = A$, this equation gives a series of parallel planes. The phase on these planes is constant. If $A = 2\pi j$ ($j = 0, \pm 1, \pm 2, \dots$), the phase difference between two adjacent planes is 2π . The spatial periods along the x , y and z axes are

$$\begin{cases} \lambda_x = \frac{2\pi}{k_x} = \frac{1}{m}, \\ \lambda_y = \frac{2\pi}{k_y} = \frac{1}{n}, \\ \lambda_z = \frac{2\pi}{k_z} = \frac{1}{s}. \end{cases} \quad (\text{A.3.4})$$

Here Eq. (A.3.3) has been used. As we expect, m , n , and s correspond to the spatial frequencies in the x , y and z directions, respectively. A spatial frequency vector \mathbf{m} can be introduced with three components m , n , s . Thus the direct and inverse 3-D Fourier transforms can be expressed in a compact form:

$$f(\mathbf{r}) = \int_{-\infty}^{\infty} F(\mathbf{m}) \exp(2\pi i \mathbf{m} \bullet \mathbf{r}) d\mathbf{m}, \quad (\text{A.3.5})$$

and

$$F(\mathbf{m}) = \int_{-\infty}^{\infty} f(\mathbf{r}) \exp(-2\pi i \mathbf{m} \bullet \mathbf{r}) d\mathbf{r}, \quad (\text{A.3.6})$$

where the vector \mathbf{r} has three components x , y and z .

A.4 Fourier Transform Theorems

In this section, we give the Fourier transform theorems without giving the derivation procedure. The theorems are given for a 1-D case but the 2-D and 3-D forms of the theorems can be easily written down.

a) Similarity theorem

If $F(m) = \mathcal{F}\{f(x)\}$, then

$$\mathcal{F}\{f(ax)\} = \frac{1}{|a|} F\left(\frac{m}{a}\right). \quad (\text{A.4.1})$$

It means that a “stretching” of the coordinate in the x space leads to a contraction of the coordinates in the Fourier space and a change of the Fourier transform by a factor of $1/|a|$.

b) Shift theorem

If $F(m) = \mathcal{F}\{f(x)\}$, then

$$\mathcal{F}\{f(x - a)\} = F(m) \exp(-2\pi i m a). \quad (\text{A.4.2})$$

It means that a translation of a function in the x space leads to a linear phase shift in the Fourier space.

c) Parsval's theorem

If $F(m) = \mathcal{F}\{f(x)\}$, then

$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \int_{-\infty}^{\infty} |F(m)|^2 dm, \quad (\text{A.4.3})$$

which is a statement of the conservation of energy in physics.

d) Convolution theorem

If $F(m) = \mathcal{F}\{f(x)\}$ and $G(m) = \mathcal{F}\{g(x)\}$, then

$$\mathcal{F}\left\{\int_{-\infty}^{\infty} f(\xi) g(x - \xi) d\xi\right\} = F(m)G(m), \quad (\text{A.4.4})$$

or

$$\mathcal{F}\{f(x) \otimes g(x)\} = F(m)G(m). \quad (\text{A.4.5})$$

This theorem implies that the Fourier transform of a convolution operation of two functions in the x space is simply equivalent to the product of their corresponding Fourier transforms.

e) Autocorrelation theorem

If $F(m) = \mathcal{F}\{f(x)\}$, then

$$\mathcal{F}\left\{\int_{-\infty}^{\infty} f(\xi) f^*(\xi - x) d\xi\right\} = |F(m)|^2, \quad (\text{A.4.6})$$

or

$$\mathcal{F}\{|f(x)|^2\} = \int_{-\infty}^{\infty} F(\xi) F^*(\xi - m) d\xi. \quad (\text{A.4.7})$$

It is noted that the autocorrelation theorem is a special case of the convolution theorem if we let $g(x) = f^*(-x)$.

Appendix B

HANKEL TRANSFORM

In this appendix, a special form of the two-dimensional (2-D) Fourier transform, called the Hankel transform, is discussed.

Let us start with the 2-D Fourier transform in Cartesian coordinates, which is given by

$$F(m, n) = \int \int_{-\infty}^{\infty} f(x, y) \exp[2\pi i(xm + yn)] dx dy. \quad (\text{B.1.1})$$

The Hankel transform is the 2-D Fourier transform in a polar coordinate. The function $f(x, y)$ can be represented by a function $f(r, \phi)$ if the following coordinate transformation is adopted:

$$\begin{cases} x = r \cos \phi, \\ y = r \sin \phi, \end{cases} \quad (\text{B.1.2})$$

and

$$\begin{cases} m = l \cos \psi, \\ n = l \sin \psi, \end{cases} \quad (\text{B.1.3})$$

where r and ϕ are the polar coordinates in the $x - y$ plane, whereas l and ψ are the polar coordinates in the $m - n$ plane. Therefore Eq. (B.1.1) can be rewritten as

$$F(l, \theta) = \int_0^{2\pi} \int_0^{\infty} f(r, \phi) \exp[2\pi i r l \cos(\phi - \psi)] r dr d\phi. \quad (\text{B.1.4})$$

In the case of a circularly symmetric system, $f(r, \phi) = f(r)$. Thus the Fourier transform of $f(r)$ is also circularly symmetric, which may be denoted by $F(l)$. Finally, Eq. (B.1.4) can reduce to

$$F(l) = \int_0^\infty f(r) J_0(2\pi r l) 2\pi r dr, \quad (\text{B.1.5})$$

which is called the Hankel transform. In this expression,

$$J_0(x) = \frac{1}{2\pi} \int_0^{2\pi} \exp(\pm ix \cos \phi) d\phi \quad (\text{B.1.6})$$

is a Bessel function of the first kind of order zero.




Fig. B.1.1 Bessel functions of the first kind of the first five orders, $J_0(x)$, $J_1(x)$, $J_2(x)$, $J_3(x)$ and $J_4(x)$.

If $f(r)$ is a uniform function within a radius a :

$$f(r) = \begin{cases} 1 & , \quad r \leq a, \\ 0 & , \quad r > a, \end{cases} \quad (\text{B.1.7})$$

thus using the identity

$$\int_0^x x_0 J_0(x_0) dx_0 = x J_1(x), \quad (\text{B.1.8})$$

one can derive an analytical expression for Eq. (B.1.5):

$$F(l) = \pi a^2 \left[\frac{2J_1(2\pi al)}{2\pi al} \right], \quad (\text{B.1.9})$$

where $J_1(x)$ is a Bessel function of the first kind of order one. Fig. B.1.1 gives the Bessel functions of the first kind of the first four orders. The function $2J_1(x)/x$ is also termed the Airy function and is shown in Fig. B.1.2.




Fig. B.1.2 Airy function $2J_1(r)/r$: 2-D behaviour.

Appendix C

DELTA FUNCTIONS

This appendix summarizes the main properties of delta functions. For a one-dimensional (1-D) problem, the definition of a delta function $\delta(x)$ is given by

$$\delta(x) = \begin{cases} \infty & , \quad x = 0, \\ 0 & , \quad x \neq 0, \end{cases} \quad (\text{C.1.1})$$

and

$$\int_{-\infty}^{\infty} \delta(x) dx = 1. \quad (\text{C.1.2})$$

Eqs. (C.1.1) and (C.1.2) give a complete definition of a delta function. Mathematically, a delta function represents an infinity at the origin ($x = 0$). Physically, it represents an impulse response or action. For example, in optical imaging, a point illumination source or a point detector can be represented by a delta function.

In general, a delta function can be defined with respect to an arbitrary position x_0 . In this case, Eqs. (C. 1.1) and (C.1.2) should be rewritten as

$$\delta(x - x_0) = \begin{cases} \infty & , \quad x = x_0, \\ 0 & , \quad x \neq x_0, \end{cases} \quad (\text{C.1.3})$$

and

$$\int_{-\infty}^{\infty} \delta(x - x_0) dx = 1. \quad (\text{C.1.4})$$

One of the important properties of delta functions is mathematically expressed as

$$\int_{-\infty}^{\infty} \delta(x - x_0) f(x) dx = f(x_0), \quad (\text{C.1.5})$$

or

$$\int_{-\infty}^{\infty} \delta(x) f(x) dx = f(0). \quad (\text{C.1.6})$$

This property means that the integration of the product of a delta function and an arbitrary function $f(x)$ is equivalent to taking the value of the function $f(x)$ at $x = x_0$. If $f(x)$ denotes a physical quantity, Eq. (C.1.5) represents the selection of the value of the physical quantity at x_0 .

Now, let us assume $f(x)$ to be $\exp(-2\pi i m x)$. The inverse Fourier transform of a delta function can be easily derived, according to Eq. (A.1.5), as

$$F(m) = \int_{-\infty}^{\infty} \delta(x - x_0) \exp(-2\pi i m x) dx = \exp(-2\pi i m x_0). \quad (\text{C.1.7})$$

According to the definition of the direct Fourier transform in Eq. (A.1.3), a delta function can be expressed as a integration form given by

$$\delta(x - x_0) = \int_{-\infty}^{\infty} \exp[2\pi i m (x - x_0)] dm. \quad (\text{C.1.8})$$

A delta function can be represented by a series of normal functions in the limiting forms. Two examples of these limiting forms are

$$\delta_a(x) = \sqrt{\frac{a}{\pi}} \exp(-ax^2) \quad , \quad a \rightarrow 0, \quad (\text{C.1.9})$$

and

$$\delta_a(x) = \frac{a}{\pi} \sin c(ax) \quad , \quad a \rightarrow 0, \quad (\text{C.1.10})$$

where

$$\sin c(x) = \frac{\sin x}{x}. \quad (\text{C.1.11})$$

The above definitions and discussions also apply for the two- and three-dimensional cases.

SUBJECT INDEX

2

2-D amplitude point spread function 54, 57

3

3-D amplitude point spread function 57, 62, 66, 67
pulsed beam illumination 127
3-D convolution 65
3-D space-invariant 66

4

4f system 61, 97, 105

A

Abbe imaging theory 57
2-D Fourier transform 59
aberration 145
curvature of field 186
distortion 187
primary astigmatism 185
primary coma 184
primary spherical aberration 183
aberration function
Zernike's circle polynomials 182
Airy function 23
Airy pattern 23, 45, 207
amplitude point spread function 62
apodization 145
apodization function 151
Helmholtz condition 156
Herschel condition 155
sine condition 153
uniform projection condition 156
autocorrelation theorem 204
axial optical coordinate 46, 128

B

Babinet's principle 27
Bessel function 206
Born approximation 64

C

chromatic aberration 129
circular disk
pulsed beam illumination 119
coherent imaging 95
coherent transfer function 76, 77
aberration 78
annular pupil 78
high numerical aperture 157
pulsed beam illumination 133
confocal microscopy 1
two-photon fluorescence
microscopy 2
convolution theorem 203

D

Debye approximations 19, 145
Debye integral 147
aberration 177
Debye theory 19, 145
defocused amplitude point spread
function 57
delta function 209
depolarization 145
dielectric interface 169
aberration 173, 176
displacement theorem 181
doublet 130

E

effective object function 65
evanescent wave 2
Ewald sphere 75, 79, 160

F

far-field diffraction 9
femtosecond 2
Fourier integral 74
Fourier transform 74, 199
Fourier transform theorems 202
Fraunhofer diffraction 9, 22, 41
frequency chirp 129
Fresnel approximation 20
Fresnel diffraction 9, 21, 22, 41
 Airy pattern 23
 circular aperture 22
 circular disk 26
 doughnut aperture 32
 serrated aperture 28
Fresnel number 21, 44, 128

G

Gaussian-shaped pulse 112
 pulse width 113
 spectral width 113
Green's theorem 11
group-velocity dispersion 129

H

Hankel transform 205
harmonic grating 72
Helmholtz condition 156
Helmholtz equation 11
Herschel condition 155
Huygens-Fresnel principle 7, 8, 9
Huygens-Fresnel Principle
 spherical wavelet 10

I

in-focus transfer function 91
intensity point spread function 68

K

Kirchhoff scalar diffraction theory 11
 boundary conditions 13, 15
 Fresnel-Kirchhoff
 diffraction formula 15
 Kirchhoff diffraction integral 13

L

Lagrange condition 156
laser trapping 2
laser tweezers 2
lens law 53
line object 88

M

Marechal Criterion 191
mode-locking technique 110

N

near-field scanning optical
 microscopy 2
 evanescent wave 2
 laser trapping 2
nonlinear optical microscopy 2
numerical aperture 43

O

on-axis transfer function 93
optical data processing 103
optical singularity 31
optical transfer function 75, 79, 80, 84
 annular pupil 85
 high numerical aperture 157
 pulsed beam illumination 137

P

paraxial approximation 19, 20, 39, 150
 Fraunhofer approximation 21
 Fresnel approximation 19
 Fresnel diffraction 21
Parsval's theorem 203
phase contrast imaging 101
phase singularity 31

picosecond 2
Poisson's spot 27
primary aberrations 182, 188
primary astigmatism 185
primary coma 184
primary spherical aberration 183
propagation time difference 129
pulsed beam illumination 115
 circular aperture 115
 circular disk 119
 serrated aperture 121
pupil function
 annular lens 47
 circular lens 44
 doughnut lens 49

Q

Q-factor 109
Q-switching technique 109

R

radial optical coordinate 43, 46, 128
Rayleigh's quarter wavelength rule 191
Rayleigh-Sommerfeld
 diffraction theory 15
 first Rayleigh-Sommerfeld diffraction
 integral 17
 second Rayleigh-Sommerfeld
 diffraction integral 17

S

Sech-shaped pulse 114
serrated aperture
 pulsed beam illumination 121
shift theorem 203
similarity theorem 203
sine condition 153, 197
single lens
 annular lens 47
 axial magnification 66
 circular lens 42
 coherent imaging 51
 coherent transfer function 76, 77, 78
 defocused pupil function 46, 56
 doughnut lens 49

 focal length 39, 125
 focus 40
 Fresnel number 44
 incoherent imaging 67
 lens law 53
 optical transfer function 79, 80, 84
 pupil function 38, 42
 transmittance 38, 39
 transverse magnification 66
sinusoidal grating 99
spatial filter 99
spatial filtering 99, 105
spatial frequency 72
spatial frequency vector 202
spherical aberration 3
 balanced condition 197
 cover slip 195
 mismatching 3, 192
 tube length 196
Strehl intensity 180
superposition 8

T

tangent condition 156, 198
thick object 64
 layer 86
 planar 87
three-dimensional optical storage 3
transfer function
 1-D line 89
 1-D on-axis 89
 2-D defocused 88
 2-D in-focus 88, 92
 cut-off spatial frequency 74
 spatial frequency 72, 74
 spatial frequency vector 75
 spatial period 74
 thick object 74
transmission cross coefficient 76
transmittance 39
 pulsed beam illumination 124
tube length
 sine condition 197
 tangent condition 198
two-photon fluorescence microscopy 2

U

uniform projection condition 156

V

vectorial Debye integral 163

vectorial Debye theory 163

W

wave equation 11

Z

Zernike's circle polynomials 182

Springer Series in OPTICAL SCIENCES

New editions of volumes prior to volume 60

- 1 **Solid-State Laser Engineering**
By W. Koechner, 5th ed. 1999, 480 figs., 60 tabs., approx. 760 pages
- 14 **Laser Crystals**
Their Physics and Properties
By A. A. Kaminskii, 2nd ed. 1990, 89 figs., 56 tabs., XVI, 456 pages
- 15 **X-Ray Spectroscopy**
An Introduction
By B. K. Agarwal, 2nd ed. 1991, 239 figs., XV, 419 pages
- 36 **Transmission Electron Microscopy**
Physics of Image Formation and Microanalysis
By L. Reimer, 4th ed. 1997, 263 figs. XVI, 584 pages
- 45 **Scanning Electron Microscopy**
Physics of Image Formation and Microanalysis
By L. Reimer, 2nd ed. 1998, 260 figs., XIV, 527 pages

Published titles since volume 60

- 60 **Holographic Interferometry in Experimental Mechanics**
By Yu. I. Ostrovsky, V. P. Shchepinov, V. V. Yakovlev, 1991, 167 figs., IX, 248 pages
- 61 **Millimetre and Submillimetre Wavelength Lasers**
A Handbook of cw Measurements
By N. G. Douglas, 1989, 15 figs., IX, 278 pages
- 62 **Photoacoustic and Photothermal Phenomena II**
Proceedings of the 6th International Topical Meeting, Baltimore, Maryland, August, 1989
By J. C. Murphy, J. W. MacLachlan Spicer, L. C. Aamodt, B. S. H. Royce (Eds.), 1990, 389 figs., 23 tabs., XXI, 545 pages
- 63 **Electron Energy Loss Spectrometers**
The Technology of High Performance
By H. Ibach, 1991, 103 figs., VIII, 178 pages
- 64 **Handbook of Nonlinear Optical Crystals**
By V. G. Dmitriev, G. G. Gurzadyan, D. N. Nikogosyan, 3rd ed. 1999, 39 figs., XVI, 413 pages
- 65 **High-Power Dye Lasers**
By F. J. Duarte (Ed.), 1991, 93 figs., XIII, 252 pages
- 66 **Silver-Halide Recording Materials**
for Holography and Their Processing
By H. I. Bjelkhagen, 2nd ed. 1995, 64 figs., XX, 440 pages
- 67 **X-Ray Microscopy III**
Proceedings of the Third International Conference, London, September, 1990
By A. G. Michette, G. R. Morrison, C. J. Buckley (Eds.), 1992, 359 figs., XVI, 491 pages
- 68 **Holographic Interferometry**
Principles and Methods
By P. K. Rastogi (Ed.), 1994, 178 figs., 3 in color, XIII, 328 pages
- 69 **Photoacoustic and Photothermal Phenomena III**
Proceedings of the 7th International Topical Meeting, Doorwerth, The Netherlands, August, 1991
By D. Bicanic (Ed.), 1992, 501 figs., XXVIII, 731 pages

Springer Series in
OPTICAL SCIENCES

- 70 **Electron Holography**
By A. Tonomura, 2nd ed. 1999, 127 figs., XII, 162 pages
- 71 **Energy-Filtering Transmission Electron Microscopy**
By L. Reimer (Ed.), 1995, 199 figs., XIII, 424 pages
- 72 **Nonlinear Optical Effects and Materials**
By P. Günter (Ed.), 1999, 173 figs., 43 tabs., approx. 400 pages
- 73 **Evanescent Waves**
From Newtonian Optics to Atomic Optics
By F. de Fornel, 1999, 100 figs., approx. 300 pages
- 74 **International Trends in Optics and Photonics**
ICO IV
By T. Asakura (Ed.), 1999, 190 figs., 14 tabs., XX, 428 pages
- 75 **Advanced Optical Imaging Theory**
By M. Gu, 1999, 93 figs., XII, 216 pages