

Solutions to Exercises

1. Determine all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x - f(y)) = 1 - x - y$$

for all real numbers x, y . (Slovenia, 1999)

Solution. Setting $y = 0$, we get

$$f(x - f(0)) = 1 - x$$

for all x .

Let $c = f(0)$, and let $t = x - c$, so $x = t + c$, and

$$f(t) = 1 - (t + c) = 1 - t - c$$

for all t . Setting $t = 0$, we get $f(0) = 1 - c$. But $f(0) = c$, so $1 - c = c$, which means $c = \frac{1}{2}$.

Therefore, the solution is $f(x) = \frac{1}{2} - x$. It is easy to verify that this solution works.

Note. Another substitution that works nicely is setting $x = f(y)$.

2. Find all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$x^2 f(x) + f(1 - x) = 2x - x^4$$

for all $x \in \mathbb{R}$.

Solution. Substituting $1 - x$ for x , we get

$$(1 - x)^2 f(1 - x) + f(x) = 2(1 - x) - (1 - x)^4 = -x^4 + 4x^3 - 6x^2 + 2x + 1.$$

Thus, we have the following system of equations in $f(x)$ and $f(1 - x)$:


$$\begin{aligned} x^2 f(x) + f(1 - x) &= -x^4 + 2x, \\ f(x) + (1 - x)^2 f(1 - x) &= -x^4 + 4x^3 - 6x^2 + 2x + 1. \end{aligned}$$

Multiplying the first equation by $(1 - x)^2$, we get

$$\begin{aligned} x^2(x - 1)^2 f(x) + (1 - x)^2 f(1 - x) &= (1 - x)^2(-x^4 + 2x) \\ &= -x^6 + 2x^5 - x^4 + 2x^3 - 4x^2 + 2x. \end{aligned}$$

Subtracting the equation $f(x) + (1 - x)^2 f(1 - x) = -x^4 + 4x^3 - 6x^2 + 2x + 1$, we get

$$[x^2(x - 1)^2 - 1]f(x) = -x^6 + 2x^5 - 2x^3 + 2x^2 - 1,$$



Art of Problem Solving
WOOT 2013–14
Functional Equations



which factors as

$$(x^2 - x - 1)(x^2 - x + 1)f(x) = -(x^2 - x - 1)(x^2 - x + 1)(x^2 - 1).$$

Now, the quadratic equation $x^2 - x + 1 = 0$ has no real roots, so we may safely divide both sides by $x^2 - x + 1$. However, the quadratic equation $x^2 - x - 1 = 0$ does have real roots, namely $x = \frac{1 \pm \sqrt{5}}{2}$. So, if x is not a root of $x^2 - x - 1 = 0$, then we may divide both sides by $x^2 - x - 1$ to get

$$f(x) = -(x^2 - 1) = 1 - x^2.$$

We then check this solution: Let x be a real number that is not a root of $x^2 - x - 1 = 0$. Since the roots of $x^2 - x - 1 = 0$ add up to 1, $1 - x$ is also not a root of $x^2 - x - 1 = 0$. Therefore, $f(x) = 1 - x^2$ and $f(1 - x) = 1 - (1 - x)^2$, so

$$\begin{aligned} x^2 f(x) + f(1 - x) &= x^2(1 - x^2) + 1 - (1 - x)^2 \\ &= x^2 - x^4 + 1 - 1 + 2x - x^2 \\ &= 2x - x^4. \end{aligned}$$

Thus, the function $f(x) = 1 - x^2$ works, when x is not a root of $x^2 - x - 1 = 0$. But what if x is a root of $x^2 - x - 1 = 0$?

To take a closer look at this case, take $x = \frac{1 + \sqrt{5}}{2}$ and $x = \frac{1 - \sqrt{5}}{2}$, respectively, in the given functional equation. This gives us the system of equations


$$\begin{aligned} \frac{3 + \sqrt{5}}{2} f\left(\frac{1 + \sqrt{5}}{2}\right) + f\left(\frac{1 - \sqrt{5}}{2}\right) &= \frac{-5 - \sqrt{5}}{2}, \\ \frac{3 - \sqrt{5}}{2} f\left(\frac{1 - \sqrt{5}}{2}\right) + f\left(\frac{1 + \sqrt{5}}{2}\right) &= \frac{-5 + \sqrt{5}}{2}. \end{aligned}$$

To make these equations easier to work with, let $a = f\left(\frac{1 + \sqrt{5}}{2}\right)$ and $b = f\left(\frac{1 - \sqrt{5}}{2}\right)$, so these equations become

$$\begin{aligned} \frac{3 + \sqrt{5}}{2} a + b &= \frac{-5 - \sqrt{5}}{2}, \\ \frac{3 - \sqrt{5}}{2} b + a &= \frac{-5 + \sqrt{5}}{2}. \end{aligned}$$

From the first equation,

$$b = \frac{-5 - \sqrt{5}}{2} - \frac{3 + \sqrt{5}}{2} a.$$



Art of Problem Solving
WOOT 2013–14
Functional Equations



Substituting into the second equation, the left-hand side becomes

$$\begin{aligned}\frac{3 - \sqrt{5}}{2}b + a &= \frac{3 - \sqrt{5}}{2} \left(\frac{-5 - \sqrt{5}}{2} - \frac{3 + \sqrt{5}}{2}a \right) + a \\ &= \frac{(3 - \sqrt{5})(-5 - \sqrt{5})}{4} - \frac{(3 - \sqrt{5})(3 + \sqrt{5})}{4}a + a \\ &= \frac{-15 - 3\sqrt{5} + 5\sqrt{5} + 5}{4} - \frac{9 - 5}{4}a + a \\ &= \frac{-10 + 2\sqrt{5}}{4} \\ &= \frac{-5 + \sqrt{5}}{2},\end{aligned}$$

which is the right-hand side. Thus, the second equation is in fact equivalent to the first equation. There are no other conditions on the values a and b , so we are free to choose a , which then determines b . Hence, the complete solution is as follows:

$$f(x) = \begin{cases} 1 - x^2 & \text{if } x \neq \frac{1+\sqrt{5}}{2}, x \neq \frac{1-\sqrt{5}}{2}, \\ a & \text{if } x = \frac{1+\sqrt{5}}{2}, \\ \frac{-5-\sqrt{5}}{2} - \frac{3+\sqrt{5}}{2}a & \text{if } x = \frac{1-\sqrt{5}}{2}, \end{cases}$$

where a is any constant.

3. Let $F(x)$ be a real valued function defined for all real x except for $x = 0$ and $x = 1$ and satisfying the functional equation

$$F(x) + F\left(\frac{x-1}{x}\right) = 1 + x.$$

Find all functions $F(x)$ satisfying these conditions. (Putnam, 1971)

Solution. Substituting $(x-1)/x$ for x , we get


$$F\left(\frac{x-1}{x}\right) + F\left(\frac{(x-1)/x - 1}{(x-1)/x}\right) = 1 + \frac{x-1}{x},$$

which simplifies as

$$F\left(\frac{x-1}{x}\right) + F\left(\frac{1}{1-x}\right) = \frac{2x-1}{x}.$$

Substituting $1/(1-x)$ for x in the given functional equation, we get

$$F\left(\frac{1}{1-x}\right) + F\left(\frac{1/(1-x) - 1}{1/(1-x)}\right) = 1 + \frac{1}{1-x},$$



Art of Problem Solving
WOOT 2013–14
Functional Equations



which simplifies as

$$F\left(\frac{1}{1-x}\right) + F(x) = \frac{2-x}{1-x}.$$

Thus, we have the system of equations

$$\begin{aligned} F(x) + F\left(\frac{x-1}{x}\right) &= 1+x, \\ F\left(\frac{x-1}{x}\right) + F\left(\frac{1}{1-x}\right) &= \frac{2x-1}{x}, \\ F\left(\frac{1}{1-x}\right) + F(x) &= \frac{2-x}{1-x}. \end{aligned}$$

Adding the first and third equation and subtracting the second equation, we get

$$2F(x) = 1+x + \frac{2-x}{1-x} - \frac{2x-1}{x} = \frac{1+x^2-x^3}{x(1-x)},$$

so

$$F(x) = \frac{1+x^2-x^3}{2x(1-x)}.$$

We check that this solution works. Substituting into the given functional equation, we get

$$\begin{aligned} F(x) + F\left(\frac{x-1}{x}\right) &= \frac{1+x^2-x^3}{2x(1-x)} + \frac{1 + \left(\frac{x-1}{x}\right)^2 - \left(\frac{x-1}{x}\right)^3}{2 \cdot \frac{x-1}{x} \cdot \left(1 - \frac{x-1}{x}\right)} \\ &= \frac{1+x^2-x^3}{2x(1-x)} + \frac{x^3 + x(x-1)^2 - (x-1)^3}{2x(x-1)} \\ &= \frac{1+x^2-x^3}{2x(1-x)} + \frac{x^3 + x^2 - 2x + 1}{2x(x-1)} \\ &= \frac{1+x^2-x^3}{2x(1-x)} + \frac{-x^3 - x^2 + 2x - 1}{2x(1-x)} \\ &= \frac{2x - 2x^3}{2x(1-x)} \\ &= \frac{2x(1+x)(1-x)}{2x(1-x)} \\ &= 1+x, \end{aligned}$$

so our solution works.


4. Let \mathbb{R}^+ be the set of all positive real numbers. Find all functions $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that

$$x^2(f(x) + f(y)) = (x+y)f(f(x)y)$$

www.artofproblemsolving.com

Sponsored by The D. E. Shaw group, Two Sigma Investments LLC, Jane Street Capital, Citadel LLC, and Dropbox





Art of Problem Solving
WOOT 2013–14
Functional Equations



holds for any positive real numbers x and y . (Slovenia, 2005)

Solution. Let a be a fixed point of f (assuming such a fixed point exists). Setting $x = a$ and $y = 1$ into the given functional equation, we get

$$a^2(a + f(1)) = (a + 1)a.$$

Since a is positive, we can divide both sides by a , and the equation simplifies to

$$a^2 + [f(1) - 1]a - 1 = 0.$$

Since this is a quadratic, a can have at most two values. In fact, it can only have one positive value since the product of the two roots is -1 . So f has at most one fixed point.

Setting $y = x$ in the given functional equation, we get

$$2x^2 f(x) = 2xf(xf(x)).$$

Since x is positive, we can divide both sides by $2x$ to get

$$xf(x) = f(xf(x))$$

for all $x > 0$.

Hence, $xf(x)$ is a fixed point of f for all $x > 0$, which means that f has at least one fixed point. Furthermore, if a is a fixed point, then $af(a) = a^2$ is also a fixed point. So, a, a^2, a^4, \dots are all fixed points. However, we found earlier that f has at most one fixed point, so we must have $a = 1$. Therefore, the only possible fixed point f can have is 1, which means $xf(x) = 1$, or $f(x) = 1/x$ for all $x > 0$. It is easy to verify that this solution works.

5. Let $g : S \rightarrow S$ be a function such that g has exactly two fixed points, and $g \circ g$ has exactly four fixed points. Prove that there is no function $f : S \rightarrow S$ such that $g = f \circ f$.

Solution. Let a and b be the fixed points of g , so $g(a) = a$ and $g(b) = b$. Then $(g \circ g)(a) = g(g(a)) = g(a) = a$ and $(g \circ g)(b) = g(g(b)) = g(b) = b$, so a and b are also two of the fixed points of $g \circ g$. Let c and d be the other two fixed points of $g \circ g$, so $(g \circ g)(c) = c$ and $(g \circ g)(d) = d$.


We compute $(g \circ g \circ g)(c)$ in two different ways. First, $(g \circ g \circ g)(c) = g((g \circ g)(c)) = g(c)$. Also, $(g \circ g \circ g)(c) = (g \circ g)(g(c))$, so

$$(g \circ g)(g(c)) = g(c).$$


In other words, $g(c)$ is a fixed point of $g \circ g$. Therefore, $g(c) \in \{a, b, c, d\}$.

We claim that $g(c)$ cannot be a . If $g(c) = a$, then $g(a) = g(g(c)) = c$. But $g(a) = a$, contradiction, so $g(c)$ cannot be a . Similarly, $g(c)$ cannot be b . Also, $g(c)$ cannot be c (because then c would be a fixed point of g), so $g(c) = d$. Then $g(d) = g(g(c)) = c$.

For the sake of contradiction, suppose there exists a function $f : S \rightarrow S$ such that $g = f \circ f$.



Art of Problem Solving
WOOT 2013–14
Functional Equations



Then $f \circ g = g \circ f$, so in particular

$$(f \circ g)(a) = (g \circ f)(a).$$

But $(f \circ g)(a) = f(g(a)) = f(a)$, and $(g \circ f)(a) = g(f(a))$, so $g(f(a)) = f(a)$. Hence, $f(a)$ is a fixed point of g , which means $f(a)$ is a or b . Similarly, $f(b)$ is a or b .

Also,

$$(f \circ g)(c) = (g \circ f)(c).$$

But $(f \circ g)(c) = f(g(c)) = f(d)$, and $(g \circ f)(c) = g(f(c))$, so $g(f(c)) = f(d)$. Similarly, $g(f(d)) = f(c)$. Then $(g \circ g)(f(c)) = g(g(f(c))) = g(f(d)) = f(c)$, so $f(c)$ is a fixed point of $g \circ g$, which means $f(c) \in \{a, b, c, d\}$.

We consider the possible values of $f(c)$.

Case 1: $f(c) = a$.

If $f(c) = a$, then $g(c) = (f \circ f)(c) = f(f(c)) = f(a)$, which is a or b . But $g(c) = d$, contradiction.

Case 2: $f(c) = b$.

If $f(c) = b$, then $g(c) = (f \circ f)(c) = f(f(c)) = f(b)$, which is a or b . But $g(c) = d$, contradiction.

Case 3: $f(c) = c$.

If $f(c) = c$, then $g(c) = (f \circ f)(c) = f(f(c)) = f(c) = c$, so c is a fixed point of g . But the only fixed points of g are a and b , contradiction.

Case 4: $f(c) = d$.

If $f(c) = d$, then $g(c) = (f \circ f)(c) = f(f(c)) = f(d)$. But $g(c) = d$, so $f(d) = d$. Then $g(d) = (f \circ f)(d) = f(f(d)) = f(d) = d$, so d is a fixed point of g . But the only fixed points of g are a and b , contradiction.

Every possible case leads to a contradiction, so there is no function $f : S \rightarrow S$ such that $g = f \circ f$.

6. Let $S = \{0, 1, 2, \dots\}$. Find all functions defined on S taking their values in S such that

$$f(m + f(n)) = f(f(m)) + f(n)$$

for all m and n in S . (IMO, 1996)


Solution. Setting $m = n = 0$ in the given functional equation, we get

$$f(f(0)) = f(f(0)) + f(0),$$

so $f(0) = 0$.

Setting $m = 0$ in the given functional equation, we get

$$f(f(n)) = f(f(0)) + f(n),$$



Art of Problem Solving
WOOT 2013–14
Functional Equations



which simplifies to $f(f(n)) = f(n)$ for all n . Hence, the given functional equation implies that

$$f(m + f(n)) = f(m) + f(n) \quad (*)$$

for all m and n .

We see that the function $f(n) = 0$ satisfies the given functional equation, so assume that f is not identically 0.

Since $f(f(n)) = f(n)$ for all n , $f(n)$ is a fixed point of f for any n . Let a be the smallest nonzero fixed point of f , so $f(a) = a$. Setting $m = n = a$ in $(*)$, we get

$$f(a + f(a)) = 2f(a),$$

which simplifies to $f(2a) = 2a$.

Setting $m = 2a$ and $n = a$ in $(*)$, we get

$$f(2a + f(a)) = f(2a) + f(a),$$

which simplifies to $f(3a) = 3a$. By a straightforward induction argument, $f(ka) = ka$ for all positive integers k , i.e. ka is a fixed point of f for all $k \geq 1$. We claim that every fixed point of f is of the form ka , where $k \geq 1$.

Let b be a fixed point of f , and write

$$b = ka + r,$$

where $0 \leq r < a$. Then

$$f(b) = f(ka + r).$$

Setting $m = r$ and $n = ka$ in $(*)$, we get

$$f(r + f(ka)) = f(r) + f(ka).$$

But ka is a fixed point of f , i.e. $f(ka) = ka$, so this equation becomes $f(ka + r) = ka + f(r)$. Hence,

$$f(b) = f(ka + r) = ka + f(r).$$

Also, b is a fixed point of f , so $f(b) = b = ka + r$, and


$$ka + f(r) = ka + r,$$

which means $f(r) = r$. By definition, a is the smallest nonzero fixed point of f , and $r < a$. Therefore, $r = 0$, so $b = ka$. We conclude that every fixed point of f is a multiple of a .

Since $f(n)$ is a fixed point of f for any n , $f(n)$ is a multiple of a for all n . Let $f(n) = g(n)a$, so $g(n)$ is a positive integer for all n .

Let N be a nonnegative integer, and write

$$N = ka + r,$$



Art of Problem Solving
WOOT 2013–14
Functional Equations



where $0 \leq a < r$. Setting $m = r$ and $n = ka$ in $(*)$, we get

$$f(r + f(ka)) = f(r) + f(ka),$$

which simplifies to $f(ka + r) = ka + f(r)$. Since $N = ka + r$ and $f(r) = g(r)a$,

$$f(N) = ka + g(r)a, \quad (**)$$

where k and r are the quotient and remainder when N is divided by a , respectively. If $r = 0$, then $N = ka$ is divisible by a , which means N is a fixed point, so $f(N) = ka$. Then

$$ka = ka + g(0)a,$$

so $g(0) = 0$.

Thus, the function f is determined by:

- (i) A positive integer a , and
- (ii) the positive integers $g(1), g(2), \dots, g(a-1)$. (We set $g(0) = 0$.)

Conversely, we claim that any function so determined (by these values and $(**)$) satisfies the given functional equation.

To see this, let N and M be arbitrary nonnegative integers, and write $N = ka + r$ and $M = la + s$, where $0 \leq r, s < a$. Then

$$f(N) = ka + g(r)a,$$

$$f(M) = la + g(s)a,$$

so

$$\begin{aligned} f(M + f(N)) &= f(la + s + ka + g(r)a) \\ &= f((k + l + g(r))a + s) \\ &= [k + l + g(r)]a + g(s)a \\ &= [la + g(s)a] + [ka + g(r)a] \\ &= f(M) + f(N). \end{aligned}$$

Also, $f(M) = la + g(s)a$ is a multiple of a , and since $g(0) = 0$,

$$f(f(M)) = f(la + g(s)a) = la + g(s)a = f(M).$$


Therefore,

$$f(M + f(N)) = f(M) + f(N) = f(f(M)) + f(N),$$

so the given functional equation is always satisfied.

In summary, all the solutions can be described as follows: Either $f(n) = 0$ for all n , or there exists a positive integer a and positive integers $g(0) = 0, g(1), g(2), \dots, g(a-1)$ such that if $N = ka + r$, where k and r are the quotient and remainder when N is divided by a , respectively, then

$$f(N) = ka + g(r)a.$$



Art of Problem Solving
WOOT 2013–14
Functional Equations



7. A function f is *multiplicative* if $f(xy) = f(x)f(y)$ for all x and y . Find all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ that are both additive and multiplicative.

Solution. Since f is multiplicative,

$$f(xy) = f(x)f(y)$$

for all x and y . Setting $y = x$, we get

$$f(x^2) = f(x)^2$$

for all x .

This equation tells us that for any nonnegative real number x ,

$$f(x) = f(\sqrt{x})^2 \geq 0.$$

Since f is also additive, f is of the form $f(x) = cx$, where c is a constant.

By the multiplicative condition,

$$c^2xy = cxy$$

for all x and y , so $c^2 = c$, which means $c = 0$ or $c = 1$.

Therefore, the only functions $f : \mathbb{R} \rightarrow \mathbb{R}$ that are both additive and multiplicative are $f(x) = 0$ and $f(x) = x$. It is easy to verify that both solutions work.

8. Find all continuous functions $f : (0, \infty) \rightarrow (0, \infty)$ such that

$$f(xy) = f(x)f(y)$$

for all $x, y > 0$.

Solution. If we could turn the products in our functional equation into sums, then we could apply Cauchy. So, we seek a substitution that turns products into sums. This gets us thinking about logarithms and exponentials. Let's try the substitution $f(x) = e^{g(x)}$, which will produce a sum of functions on the right-hand side:

$$e^{g(xy)} = e^{g(x)}e^{g(y)} = e^{g(x)+g(y)},$$


so $g(xy) = g(x) + g(y)$. That takes care of the right side, but the left side is still a product. If we could replace xy with $\log xy$, then we could produce a sum. That suggests trying $f(x) = e^{h(\log x)}$. Let's give that a try in our original functional equation:

$$e^{h(\log xy)} = e^{h(\log x)}e^{h(\log y)} = e^{h(\log x)+h(\log y)},$$

so $h(\log xy) = h(\log x) + h(\log y)$, from which we have

$$h(\log x + \log y) = h(\log x) + h(\log y).$$

Success!



Art of Problem Solving
WOOT 2013–14
Functional Equations



Let $a = \log x$ and $b = \log y$, so

$$h(a+b) = h(a) + h(b)$$

for all a and b . Solving $f(x) = e^{h(\log x)}$ for $h(x)$ gives $h(x) = \log f(e^x)$. Since f is continuous, so is h . Finally, since h is continuous and $h(a+b) = h(a) + h(b)$, we have $h(x) = cx$ for some constant c . Then

$$f(t) = e^{h(\log t)} = e^{c \log t} = (e^{\log t})^c = t^c.$$

Therefore, the solutions are of the form $f(x) = x^c$, where c is a constant. It is easy to verify that any solution of this form works.

9. Find all continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x+y) = f(x) + f(y) + xy$$

for all $x, y \in \mathbb{R}$.

Solution. Setting $y = 1$, we get

$$f(x+1) = f(x) + f(1) + x$$

for all x . Let $a = f(1)$, so

$$f(x+1) = f(x) + x + a \tag{*}$$

for all x .

Setting $x = 1$ in $(*)$, we get

$$f(2) = f(1) + 1 + a = 2a + 1.$$

Setting $x = 2$ in $(*)$, we get

$$f(3) = f(2) + 2 + a = 3a + 3.$$

Setting $x = 3$ in $(*)$, we get

$$f(4) = f(3) + 3 + a = 4a + 6.$$

Setting $x = 4$ in $(*)$, we get

$$f(5) = f(4) + 4 + a = 5a + 10.$$

By a straightforward induction argument,


$$f(n) = an + \frac{n(n-1)}{2} = \frac{1}{2}n^2 + \left(a - \frac{1}{2}\right)n$$

for all positive integers n , so let

$$g(x) = f(x) - \frac{1}{2}x^2.$$

Then

$$f(x) = g(x) + \frac{1}{2}x^2.$$



Art of Problem Solving
WOOT 2013–14
Functional Equations



Substituting into the given functional equation, we get

$$g(x+y) + \frac{1}{2}(x+y)^2 = g(x) + \frac{1}{2}x^2 + g(y) + \frac{1}{2}y^2 + xy,$$

which simplifies as

$$g(x+y) = g(x) + g(y)$$

for all x and y . Since g is continuous, $g(x) = cx$ for some constant c . Then

$$f(x) = \frac{1}{2}x^2 + cx,$$

where c is a constant. It is easy to verify that any solution of this form works.

www.artofproblemsolving.com

Sponsored by The D. E. Shaw group, Two Sigma Investments LLC, Jane Street Capital, Citadel LLC, and Dropbox

