

# Synthetic Geometry

Matthew Brennan

## 1 Introduction

This handout includes a rough outline of the solutions presented in the talk and a list of some common theorems and results useful in many geometry problems. Many of the theorems and lemmas listed have interesting proofs and it is recommended that you try to prove some of them. The solutions to the examples are outlines which are intended to illustrate certain techniques. In some cases, calculations and dealing with special diagram cases are omitted.

## 2 Warmup

Here are two problems to start, both of which have short clean solutions. The first warmup applies a transformation after which the result becomes much clearer. The solution to the second warmup shifts the focus to a new triangle after which the problem becomes tractable.

**Warmup 1.** (*JBMO 2002*) An isosceles triangle  $ABC$  satisfies that  $CA = CB$ . A point  $P$  is on the circumcircle between  $A$  and  $B$  and on the opposite side of the line  $AB$  to  $C$ . If  $D$  is the foot of the perpendicular from  $C$  to  $PB$ , show that  $PA + PB = 2 \cdot PD$ .

**Solution:** Let the point  $Q$  be such that triangles  $QCB$  and  $PCA$  are congruent. Since  $PACB$  is cyclic,

$$\angle CBQ = \angle CAP = 180^\circ - \angle CBP$$

which implies that  $P$ ,  $B$  and  $Q$  are collinear. Since  $QCB$  and  $PCA$  are congruent,  $CPQ$  is isosceles and thus  $D$  is the midpoint of  $PQ$ . Therefore

$$PA + PB = PQ = 2 \cdot PD$$

**Warmup 2.** (*Russia 2005*) In an acute-angled triangle  $ABC$ ,  $AM$  and  $BN$  are altitudes. A point  $D$  is chosen on arc  $ACB$  of the circumcircle of the triangle. Let the lines  $AM$  and  $BD$  meet at  $P$  and the lines  $BN$  and  $AD$  meet at  $Q$ . Prove that  $MN$  bisects segment  $PQ$ .

**Solution:** Assume without the loss of generality that  $D$  is on arc  $AC$  not including  $B$ . Let  $H$  be the orthocenter of  $ABC$ . Since  $ADCB$  is cyclic,

$$\angle PAN = \angle DAC = \angle DBC = \angle QBM.$$

Also, it follows that

$$\angle NAH = 90^\circ - \angle ACB = \angle MBH.$$

Since  $HP \perp AN$  and  $HQ \perp BM$ ,  $PAN$  is similar to  $QBM$  and  $NAH$  is similar to  $MBH$ . Therefore

$$\frac{PM}{MH} = \frac{PM/BM}{MH/BM} = \frac{QN/AN}{NH/AN} = \frac{QN}{NH}$$

If  $X$  denotes the midpoint of  $PQ$ , then

$$\frac{PM}{MH} \cdot \frac{NH}{QN} \cdot \frac{QX}{XP} = 1$$

and by Menelaus' Theorem applied to triangle  $HPQ$ , points  $X$ ,  $M$  and  $N$  are collinear.

### 3 Redefining Points to be Easier

Often points in geometry problems are defined in ways that are difficult to deal with. For any points that seem difficult to understand or work with, it is often best to redefine them in a useful way. Specifically, if  $P$  is a point in the diagram that is difficult to deal with, it is often best to define  $P'$  in some other way using a property we think is true of  $P$  and which can be used to define  $P$ , and then prove that  $P' = P$ . One thing to note is that this method requires that we have a property of  $P$  in mind. Finding out what is true of  $P$  is usually the most difficult part of problems that can be solved using this method. There is no best way to look for properties of  $P$ . However, it is often useful to think about what properties of  $P$  would solve or yield significant progress on the problem and what properties seem as though they may be true based on the diagram.

Often the best conjectures are simple, such as  $P$  lies on a line in the diagram,  $P$  lies on a circle in the diagram or is concyclic with other points in the diagram, that two lines are parallel or perpendicular, or that two triangles are similar or congruent. It is important though to make sure that you are not only trying to conjecture about the diagram and also trying to make direct progress on the problem. The examples in this section are intended to illustrate how this method of redefining points can be applied to problems.

**Example 1.** *An acute-angled triangle  $ABC$  is inscribed in a circle  $\omega$ . A point  $P$  is chosen inside the triangle. Line  $AP$  intersects  $\omega$  at the point  $A_1$ . Line  $BP$  intersects  $\omega$  at the point  $B_1$ . A line  $\ell$  is drawn through  $P$  and intersects  $BC$  and  $AC$  at the points  $A_2$  and  $B_2$ . Prove that the circumcircles of triangles  $A_1A_2C$  and  $B_1B_2C$  intersect again on line  $\ell$ .*

We want to analyze the second intersection of the circumcircles of triangles  $A_1A_2C$  and  $B_1B_2C$ . How much we can prove about this intersection  $Q$  varies greatly with how we define  $Q$ . Consider the two different methods below:

**Method #1:** First let's try defining  $Q$  directly as the intersection of the circumcircles of triangles  $A_1A_2C$  and  $B_1B_2C$ . From this, we know that  $\angle CQB_2 = 180^\circ - \angle CB_1B_2$  and  $\angle CQA_2 = 180^\circ - \angle CA_1A_2$ . What we want is to show that  $\angle CQB_2 + \angle CQA_2 = 180^\circ$  which now is equivalent to  $\angle CB_1B_2 + \angle CA_1A_2 = 180^\circ$ . However, this is not immediately true given the conditions in the problem. Now consider a second method.

**Method #2:** From the diagram, it looks like  $B_1PQA_1$  is cyclic. From this information, consider defining  $Q'$  as the intersection of the circumcircle of  $B_1PA_1$  and  $\ell$ . From cyclic quadrilaterals, we have

$$\angle B_1Q'P = \angle B_1A_1P = \angle B_1CB_2$$

which implies that  $Q'$  is on the circumcircle of  $B_1B_2C$ . By a similar argument, we have that  $Q'$  is on the circumcircle of  $A_1A_2C$ . Together these imply that  $Q = Q'$ . Thus  $Q$  lies on  $\ell$ .

A solution can also be obtained by defining  $Q'$  as the intersection of the circumcircle of  $B_1B_2C$  and  $\ell$ . The way we define  $Q'$  above can be motivated by more than a conjecture based off of a conjecture from a diagram. We want to define  $Q'$  in some way and then use this way to show it lies on circles. The cleanest way to do this is to show the angle conditions for a cyclic quadrilateral. In order to get these angle conditions, one promising approach is to define  $Q'$  as the intersection of a circle with something, which in this case is  $\ell$ .

One note for completeness is that the condition  $\angle CB_1B_2 + \angle CA_1A_2 = 180^\circ$  in Method #1 is a direct implication of Pascal's Theorem. In this case, Method #2 has saved us having to cite a deep theorem such as Pascal's Theorem. In other cases, redefining points can avoid much more complicated applications of advanced theorems.

These next examples illustrate this same method applied in more situations. Particularly in Example 3, it is hard to find a clean solution without the observations used to define  $P'$ .

**Example 2.** (China 2012) In the triangle  $ABC$ ,  $\angle A$  is biggest. On the circumcircle of  $ABC$ , let  $D$  be the midpoint of arc  $ABC$  and  $E$  be the midpoint of arc  $ACB$ . The circle  $c_1$  passes through  $A, B$  and is tangent to  $AC$  at  $A$ , the circle  $c_2$  passes through  $A, E$  and is tangent  $AD$  at  $A$ . Circles  $c_1$  and  $c_2$  intersect at  $A$  and  $P$ . Prove that  $AP$  bisects  $\angle BAC$ .

If the result is true, then by the tangency conditions  $\angle APB = 180^\circ - \angle BAC$  and  $\angle PBA = 180^\circ - \angle APB - \angle PAB = \frac{1}{2}\angle BAC = \angle PAB$ . Therefore if the problem is true, then  $P$  lies on the perpendicular bisector of  $AB$ . This gives us the hint to try defining  $P$  based on this. The method below defines  $P'$  as the intersection of  $c_1$  and the perpendicular bisector of  $AB$ .

**Solution:** Let the center of  $c_1$  be  $O_1$  and let the center of  $c_2$  be  $O_2$ . Since  $c_1$  is tangent to  $AC$ , it follows that  $\angle BO_1A = 2\angle BAC$ . Since  $O_1$  and  $E$  both lie on the perpendicular bisector of  $AB$ , it follows that  $O_1E$  bisects angle  $\angle BO_1A$  which implies that  $\angle BO_1A = \angle BAC$  and hence that  $\angle BP'E = 90^\circ + \frac{1}{2}\angle BAC$ . However, since  $P'$  lies on the perpendicular bisector  $EO_1$  of  $AB$ ,  $A$  is the reflection of  $B$  about  $EO_1$  and  $\angle AP'E = \angle BP'E = 90^\circ + \angle BAC$ . Since  $c_2$

is tangent to  $AD$  and passes through  $E$ , it follows that  $\angle AO_2E = 2\angle DAE = 180^\circ - \angle BAC$ . Combining this with the angle relation above yields that  $P'$  lies on  $c_2$ . Hence  $P'$  lies on both  $c_1$  and  $c_2$  and  $P = P'$ . Therefore  $\angle BAP = \frac{1}{2}\angle BO_1P = \frac{1}{2}\angle BAC$  which implies the result.

**Example 3.** (IMO 2011) Let  $ABC$  be an acute triangle with circumcircle  $\Gamma$ . Let  $\ell$  be a tangent line to  $\Gamma$ , and let  $\ell_a, \ell_b$  and  $\ell_c$  be the lines obtained by reflecting  $\ell$  in the lines  $BC$ ,  $CA$  and  $AB$ , respectively. Show that the circumcircle of the triangle determined by the lines  $\ell_a, \ell_b$  and  $\ell_c$  is tangent to the circle  $\Gamma$ .

Experimenting with what we can prove yields that we can get almost no information that seems to lead to proving the desired result through standard techniques. The main issue is that we know almost nothing about the point of tangency if the problem is true. The key to the simplest solution to this problem is to find a way to define this supposed point of tangency. We try intersecting circumcircles in order to obtain angle information to prove that the point of intersection lies on  $\Gamma$ , the circumcircle of the triangle determined by the three lines and prove that the circles are tangent at this point.

**Solution Outline:** Let  $A', B'$  and  $C'$  be the intersections of  $\ell_b$  and  $\ell_c$ ,  $\ell_a$  and  $\ell_c$ , and  $\ell_a$  and  $\ell_b$ , respectively. Let  $P$  be the point of tangency between  $\Gamma$  and  $\ell$  and let  $Q$  be the reflection of  $P$  through  $BC$ . Now let  $T$  be the second intersection of the circumcircles of  $BB'Q$  and  $CC'Q$ . It can be shown that  $T$  lies on  $\Gamma$  and the circumcircle of  $A'B'C'$  by angle chasing. Similarly,  $T$  can be shown to be a point of tangency between the circles by angle chasing. The angle chasing is made easier by first showing that  $AA', BB'$  and  $CC'$  meet at the incenter  $I$  of  $A'B'C'$ .

**Example 4.** (CMO 2013) Let  $O$  denote the circumcenter of an acute-angled triangle  $ABC$ . Let point  $P$  on side  $AB$  be such that  $\angle BOP = \angle ABC$ , and let point  $Q$  on side  $AC$  be such that  $\angle COQ = \angle ACB$ . Prove that the reflection of  $BC$  in the line  $PQ$  is tangent to the circumcircle of triangle  $APQ$ .

Here, we use the method above to define the reflection  $R$  of the point of tangency in line  $PQ$  as the intersection of triangle  $OBP$  with side  $BC$ . This construction can be motivated either by noticing this pattern in the diagram, noting that this method of intersecting circles obtains angles in exactly the way needed to prove the result, or by trying to complete the Miquel configuration. The Miquel configuration is described in greater detail in Section 5.

**Solution:** Let the circumcircle of triangle  $OBP$  intersect side  $BC$  at the points  $R$  and  $B$  and let  $\angle A, \angle B$  and  $\angle C$  denote the angles at vertices  $A, B$  and  $C$ , respectively. Now note that since  $\angle BOP = \angle B$  and  $\angle COQ = \angle C$ , it follows that

$$\angle POQ = 360^\circ - \angle BOP - \angle COQ - \angle BOC = 360^\circ - (180^\circ - \angle A) - 2\angle A = 180^\circ - \angle A.$$

This implies that  $APOQ$  is a cyclic quadrilateral. Since  $BPOR$  is cyclic,

$$\angle QOR = 360^\circ - \angle POQ - \angle POR = 360^\circ - (180^\circ - \angle A) - (180^\circ - \angle B) = 180^\circ - \angle C.$$

This implies that  $CQOR$  is a cyclic quadrilateral. Since  $APOQ$  and  $BPOR$  are cyclic,

$$\angle QPR = \angle QPO + \angle OPR = \angle OAQ + \angle OBR = (90^\circ - \angle B) + (90^\circ - \angle A) = \angle C.$$

Since  $CQOR$  is cyclic,  $\angle QRC = \angle COQ = \angle C = \angle QPR$  which implies that the circumcircle of triangle  $PQR$  is tangent to  $BC$ . Further, since  $\angle PRB = \angle BOP = \angle B$ ,

$$\angle PRQ = 180^\circ - \angle PRB - \angle QRC = 180^\circ - \angle B - \angle C = \angle A = \angle PAQ.$$

This implies that the circumcircle of  $PQR$  is the reflection of  $\Gamma$  in line  $PQ$ . By symmetry in line  $PQ$ , this implies that the reflection of  $BC$  in line  $PQ$  is tangent to  $\Gamma$ .

## 4 Spiral Similarity and Applying Transformations

One of the most useful techniques in synthetic geometry problems is applying transformations to a diagram. Often solutions to difficult problems introduce a point to the diagram which allows for a clean quick solution. These points can often be viewed as completions of transformations already present in a diagram. For example, a diagram may contain a parallelogram  $ABCD$  in which cases there is a translation mapping  $AB$  to  $DC$ . A diagram may contain a trapezoid  $ABCD$  with  $AB \parallel CD$  in which case there is a homothety mapping  $AB$  to  $CD$ . The transformations that most commonly appear are spiral similarities, rotations, homotheties and translations. The first few examples illustrate different ways to apply spiral similarities. The first example is one direction of Ptolemy's Theorem.

**Example 5.** (Ptolemy) If  $ABCD$  is a cyclic quadrilateral, then

$$AB \cdot CD + AD \cdot BC = AC \cdot BD$$

Here we construct similar triangles by applying a spiral similarity with center  $A$  mapping the  $C$  to  $D$ . We let the point  $B$  be mapped to  $P$  under this transformation.

**Solution:** Let  $P$  be the point on  $BD$  such that  $\angle APD = \angle ABC$ . Note that since  $\angle ADP = \angle ACB$  which implies that triangles  $ABC$  and  $APD$  are similar. This implies that triangles  $ADC$  and  $APB$  are similar. Therefore  $\frac{AD}{AC} = \frac{PD}{BC}$  and  $\frac{AB}{AC} = \frac{BP}{CD}$ . Therefore

$$BD = BP + PD = \frac{AB \cdot CD}{AC} + \frac{AD \cdot BC}{AC}$$

which implies on multiplying up that  $AB \cdot CD + AD \cdot BC = AC \cdot BD$ .

**Example 6.** (IMO Shortlist 2000) Let  $ABCD$  be a convex quadrilateral. The perpendicular bisectors of its sides  $AB$  and  $CD$  meet at  $Y$ . Denote by  $X$  a point inside the quadrilateral

$ABCD$  such that  $\angle ADX = \angle BCX < 90^\circ$  and  $\angle DAX = \angle CBX < 90^\circ$ . Show that  $\angle AYB = 2 \cdot \angle ADX$ .

In this example we consider the spiral similarity with center  $B$  mapping line  $CX$  to the perpendicular bisector of  $AB$  in order to obtain the angle we want  $Y$  to have at the image  $Y'$  of  $C$ . We then show that  $Y = Y'$  in the same way as in the previous section.

**Solution:** Let  $X'$  and  $Y'$  be such that  $AX' = BX'$ ,  $AY' = BY'$ ,  $\angle AX'B = 2 \cdot \angle BXC$  and  $\angle AY'B = 2 \cdot \angle BCX$ . We have that  $AX'Y'$  and  $AXD$  are similar, and that  $BX'Y'$  and  $BXC$  are similar. These similarities imply that triangles  $AXX'$  and  $ADY'$  are similar and that triangles  $BXX'$  and  $BCY'$  are similar. The ratios of similarity give that

$$DY' = \frac{AY' \cdot XX'}{AX'} = \frac{BY' \cdot XX'}{BX'} = CY'$$

Thus  $Y'$  lies on the perpendicular bisector of  $CD$  and  $Y' = Y$ . Therefore  $\angle AYB = 2 \cdot \angle ADX$ .

**Example 7.** (IMO 1996) Let  $P$  be a point inside a triangle  $ABC$  such that

$$\angle APB - \angle ACB = \angle APC - \angle ABC.$$

Let  $D$ ,  $E$  be the incenters of triangles  $APB$ ,  $APC$ , respectively. Show that the lines  $AP$ ,  $BD$ ,  $CE$  meet at a point.

**Solution:** Here we use spiral similarity to construct exactly the given angle condition. By the angle bisector theorem, it suffices to show that  $\frac{AB}{BP} = \frac{AC}{CP}$ . Let  $Q$  be such that triangles  $APB$  and  $ACQ$  are similar. It follows that  $APC$  and  $ABQ$  are similar. It follows that

$$\angle CBQ = \angle APC - \angle ABC = \angle APB - \angle ACB = \angle BCQ$$

and thus  $BQ = CQ$ . Ratios of similarity finish the problem since

$$\frac{AB}{BP} = \frac{AQ}{CQ} = \frac{AQ}{BQ} = \frac{AC}{CP}$$

The next problem illustrates an often useful transformation when there is a midpoint of the side of a triangle. It is often useful to perform a  $180^\circ$  rotation about the midpoint to produce a parallelogram as in the example below which is from Challenging Problems in Geometry.

**Example 8.** Let  $ABC$  be a given triangle and  $M$  be the midpoint of  $BC$ . If  $\angle CAM = 2 \cdot \angle BAM$  and  $D$  is a point on line  $AM$  such that  $\angle DBA = 90^\circ$ , prove that  $AD = 2 \cdot AC$ .

**Solution:** There is a very short trigonometric solution to this problem, but we present a synthetic one to illustrate the transformation mentioned above. Let  $D$  be such that  $ABDC$  is a parallelogram. If  $N$  is the midpoint of  $AD$ , then  $M$  is the midpoint of  $AD$ . Now note

$$\angle BND = 2 \cdot \angle BAM = \angle CAM = \angle NDB$$

and thus  $BD = BN$ . This implies that  $AC = BD = BN = \frac{1}{2}AD$ .

The next example illustrates applying translations, which are particularly useful when there is a parallelogram in the diagram.

**Example 9.** (2013 British MO) *The point  $P$  lies inside triangle  $ABC$  so that  $\angle ABP = \angle PCA$ . The point  $Q$  is such that  $PBQC$  is a parallelogram. Prove that  $\angle QAB = \angle CAP$ .*

**Solution:** Let  $R$  be such that  $RACP$  is a parallelogram. It follows that  $\angle ARP = \angle PCA = \angle ABP$  which implies that  $RAPB$  is cyclic. It follows that  $BRP$  and  $QAC$  are congruent and thus  $\angle QAC = \angle BRP = \angle BAP$ . This implies that  $\angle QAB = \angle CAP$ .

## 5 Geometry Facts

The theorems and facts below are many that I find useful. This list is a work in progress so there are many useful ideas that I have omitted. I plan to add sections on collinearity, concurrency, and miscellaneous useful facts. If any come to mind, please let me know. For now, I have not included ideas from inversive and projective geometry. The theorems below all apply to points in a plane. Quadrilaterals are named  $ABCD$  such that the sides of the quadrilateral are  $AB, BC, CD$  and  $DA$ . The directed angle  $\angle ABC$  is the counter-clockwise angle between  $0$  and  $180^\circ$  needed to rotate line  $AB$  to line  $BC$ . A triangle  $ABC$  is taken to have angles  $a, b$  and  $c$ .

### Cyclic Quadrilaterals:

1. A convex quadrilateral  $ABCD$  is cyclic if and only if either:
  - (a)  $\angle ADB = \angle ACB$
  - (b)  $\angle DAB + \angle BCD = 180^\circ$
2. The above two conditions can be restated as a single condition in terms of directed angles: Four points  $A, B, C$  and  $D$  are concyclic if and only if  $\angle ABC = \angle ADC$ .
3. (Power of a Point) Let  $ABCD$  be a convex quadrilateral such that  $AB$  and  $CD$  intersect at  $P$  and diagonals  $AC$  and  $BD$  intersect at  $Q$ .  $ABCD$  is cyclic if and only if either:
  - (a)  $AQ \cdot QC = BQ \cdot QD$  or equivalently  $QAD$  and  $QBC$  are similar
  - (b)  $PA \cdot PB = PC \cdot PD$  or equivalently  $PAD$  and  $PCB$  are similar
4. Given a triangle  $ABC$ , the intersections of the internal and external bisectors of angle  $\angle BAC$  with the perpendicular bisector of  $BC$  both lie on the circumcircle of  $ABC$ .

5. (Ptolemy's Theorem) A quadrilateral  $ABCD$  is cyclic if and only if

$$AB \cdot CD + AD \cdot BC = AC \cdot BD$$

6. Let  $ABCD$  be a cyclic quadrilateral such that  $AB$  and  $CD$  intersect at  $P$  and diagonals  $AC$  and  $BD$  intersect at  $Q$ . Then:

$$\frac{BQ}{QD} = \frac{AB \cdot BC}{AD \cdot DC} \quad \text{and} \quad \frac{PB}{PA} = \frac{BC \cdot BD}{AC \cdot AD}$$

7. (Polars) Let  $ABCD$  be a cyclic quadrilateral inscribed in circle  $\Gamma$  such that  $AB$  and  $CD$  intersect at  $P$  and diagonals  $AC$  and  $BD$  intersect at  $Q$ . If the tangents drawn from  $P$  to  $\Gamma$  touch  $\Gamma$  at  $R$  and  $S$ , then  $R$ ,  $Q$  and  $S$  are collinear.

### Circles:

- (Power of a Point) Given a circle  $\Gamma$  with center  $O$  and a point  $P$  then for any line  $\ell$  through  $P$  that intersects  $\Gamma$  at  $A$  and  $B$ , the value  $PA \cdot PB$  is constant as  $\ell$  varies and is equal to the power of the point  $P$  with respect to  $\Gamma$ .
  - The power of  $P$  is equal to  $r^2 - PO^2$  if  $P$  is inside  $\Gamma$  and  $PO^2 - r^2$  otherwise.
  - If  $PA$  is tangent to  $\Gamma$ , then the power of  $P$  is equal to  $PA^2$ .
- (Radical Axis) Given two circles  $\Gamma_1$  and  $\Gamma_2$ , the set of all points  $P$  with equal powers with respect to  $\Gamma_1$  and  $\Gamma_2$  is a line which is the radical axis of the two circles.
  - The radical axis is perpendicular to the line through the centers of  $\Gamma_1$  and  $\Gamma_2$ .
  - If  $\Gamma_1$  and  $\Gamma_2$  intersect at  $A$  and  $B$ , then the radical axis passes through  $A$  and  $B$ .
  - If  $AB$  is a common tangent with  $A$  on  $\Gamma_1$  and  $B$  on  $\Gamma_2$ , then the radical axis passes through the midpoint of  $AB$ .
- (Radical Center) Given three circles  $\Gamma_1, \Gamma_2$  and  $\Gamma_3$ , the three radical axes between pairs of the three circles meet at a common point  $P$  which is the radical center of the circles.
- A point  $P$  is a circle of radius zero and the radical axis of  $P$  and a circle  $\Gamma$  is the line through the midpoints of  $PA$  and  $PB$  where  $A$  and  $B$  are points on  $\Gamma$  such that  $PA$  and  $PB$  are tangent to  $\Gamma$ .
- (Monge's Theorem) Given three circles  $\Gamma_1, \Gamma_2$  and  $\Gamma_3$ . If  $P$ ,  $Q$  and  $R$  are the external centers of homothety between pairs of the three circles, then  $P$ ,  $Q$  and  $R$  are collinear. If  $P$  and  $Q$  are internal centers of homothety, then  $P$ ,  $Q$  and  $R$  are also collinear.
- Two circles  $\Gamma_1$  and  $\Gamma_2$  intersect at  $R$  and have centers  $O_1$  and  $O_2$ . If  $P$  and  $Q$  are the internal and external centers of homothety between the two circles, then  $\angle PRQ = 90^\circ$ . The lines  $RP$  and  $RQ$  are the internal and external bisectors of  $\angle O_1RO_2$ .



**Triangle Geometry:**

1. (Angle Bisector Theorem) Let  $ABC$  be a given triangle and let  $P$  and  $Q$  be the intersections of the internal and external bisectors of angle  $\angle ABC$  with line  $AC$ . Then

$$\frac{AB}{BC} = \frac{AP}{PC} = \frac{AQ}{QC}$$

2. Angles around the centers of a triangle  $ABC$ :

- (a) If  $I$  is the incenter of  $ABC$  then  $\angle BIC = 90^\circ + \frac{a}{2}$ ,  $\angle IBC = \frac{b}{2}$  and  $\angle ICB = \frac{c}{2}$ .
- (b) If  $H$  is the orthocenter of  $ABC$  then  $\angle BHC = 180^\circ - a$ ,  $\angle HBC = 90^\circ - c$  and  $\angle HCB = 90^\circ - b$ .
- (c) If  $O$  is the circumcenter of  $ABC$  then  $\angle BOC = 2a$  and  $\angle OBC = \angle OCB = 90^\circ - a$ .
- (d) If  $I_a$  is the  $A$ -excenter of  $ABC$  then  $\angle AI_aB = \frac{c}{2}$ ,  $\angle AI_aC = \frac{b}{2}$  and  $\angle BI_aC = 90^\circ - \frac{a}{2}$ .

3. Pedal triangles of the centers of a triangle  $ABC$ :

- (a) If  $DEF$  is the triangle formed by projecting the incenter  $I$  onto sides  $BC$ ,  $AC$  and  $AB$ , then  $I$  is the circumcenter of  $DEF$  and  $\angle EDF = 90^\circ - \frac{a}{2}$ .
- (b) If  $DEF$  is the triangle formed by projecting the orthocenter  $H$  onto sides  $BC$ ,  $AC$  and  $AB$ , then  $H$  is the incenter of  $DEF$  and  $\angle EDF = 180^\circ - 2a$ .
- (c) The medial triangle of  $ABC$  is the pedal triangle of the circumcenter  $O$  of  $ABC$  and  $O$  is its orthocenter.

4. Alternate methods of defining the orthocenter and circumcenter:

- (a)  $O$  is the circumcenter of  $ABC$  if and only if  $\angle AOB = 2\angle ACB$  and  $OA = OB$ .
- (b)  $H$  is the orthocenter of  $ABC$  if and only if  $H$  lies on the altitude from  $A$  and satisfies that  $\angle BHC = 180^\circ - \angle BAC$ .

5. Facts related to the orthocenter  $H$  of a triangle  $ABC$  with circumcircle  $\Gamma$ :

- (a) If  $O$  is the circumcenter of  $ABC$ , then  $\angle BAH = \angle CAO$ .
- (b) If  $D$  is the point diametrically opposite to  $A$  on  $\Gamma$  and  $M$  is the midpoint of  $BC$ , then  $M$  is also the midpoint of  $HD$ .
- (c) If  $AH$ ,  $BH$  and  $CH$  intersect  $\Gamma$  again at  $D$ ,  $E$  and  $F$ , then there is a homothety centered at  $H$  sending the pedal triangle of  $H$  to  $DEF$  with ratio 2.
- (d) If  $D$  and  $E$  are the intersections of  $AH$  with  $BC$  and  $\Gamma$ , respectively, then  $D$  is the midpoint of  $HE$ .
- (e)  $H$  lies on the three circles formed by reflecting  $\Gamma$  about  $AB$ ,  $BC$  and  $AC$ .

- (f) If  $M$  is the midpoint of  $BC$  then  $AH = 2 \cdot OM$ .
- (g) If  $BH$  and  $CH$  intersect  $AC$  and  $AB$  at  $D$  and  $E$ , and  $M$  is the midpoint of  $BC$ , then  $M$  is the center of the circle through  $B, D, E$  and  $C$ , and  $MD$  and  $ME$  are tangent to the circumcircle of  $ADE$ .

6. Facts related to the incenter  $I$  and excenters  $I_a, I_b, I_c$  of  $ABC$  with circumcircle  $\Gamma$ :

- (a) If the incircle of  $ABC$  is tangent to  $AB$  and  $AC$  at points  $D$  and  $E$  and  $s$  is the semiperimeter of  $ABC$  then

$$AD = AE = \frac{AB + AC - BC}{2} = s - BC$$

- (b) If  $AI$  intersects  $\Gamma$  at  $D$  then  $DB = DI = DC$ ,  $D$  is the midpoint of  $II_a$ , and  $II_a$  is a diameter of the circle with center  $D$  which passes through  $B$  and  $C$ .
  - (c) If  $AI, BI$  and  $CI$  intersect  $\Gamma$  at  $D, E$  and  $F$ , then  $I_a I_b I_c$ ,  $DEF$  and the pedal triangle of  $I$  are similar and have parallel sides.
  - (d)  $I$  is the orthocenter of  $I_a I_b I_c$  and  $\Gamma$  is the nine-point circle of  $I_a I_b I_c$ .
  - (e) If  $BI$  and  $CI$  intersect  $\Gamma$  again at  $D$  and  $E$ , then  $I$  is the reflection of  $A$  in line  $DE$  and if  $M$  is the intersection of the external bisector of  $\angle BAC$  with  $\Gamma$ , then  $DMEI$  is a parallelogram.
  - (f) If the incircle and  $A$ -excircle of  $ABC$  are tangent to  $BC$  at  $D$  and  $E$ ,  $BD = CE$ .
  - (g) If the  $A$ -excircle of  $ABC$  is tangent to  $AB$ ,  $AC$  and  $BC$  at  $D$ ,  $E$  and  $F$  then  $AB + BF = AC + CF = AD = AE = s$  where  $s$  is the semi-perimeter of  $ABC$ .
  - (h) If  $M$  is the midpoint of arc  $BAC$  of  $\Gamma$ , then  $M$  is the midpoint of  $I_b I_c$  and the center of the circle through  $I_b, I_c, B$  and  $C$ .
7. (Nine-Point Circle) Given a triangle  $ABC$ , let  $\Gamma$  denote the circle passing through the midpoints of the sides of  $ABC$ . If  $H$  is the orthocenter of  $ABC$ , then  $\Gamma$  passes through the midpoints of  $AH, BH$  and  $CH$  and the projections of  $H$  onto the sides of  $ABC$ .
8. (Euler Line) If  $O, H$  and  $G$  are the circumcenter, orthocenter and centroid of a triangle  $ABC$ , then  $G$  lies on segment  $OH$  with  $HG = 2 \cdot OG$ .
9. (Symmedian) Given a triangle  $ABC$  such that  $M$  is the midpoint of  $BC$ , the symmedian from  $A$  is the line that is the reflection of  $AM$  in the bisector of angle  $\angle BAC$ .
- (a) If the tangents to the circumcircle  $\Gamma$  of  $ABC$  at  $B$  and  $C$  intersect at  $N$ , then  $N$  lies on the symmedian from  $A$  and  $\angle BAM = \angle CAN$ .
  - (b) If the symmedian from  $A$  intersects  $\Gamma$  at  $D$ , then  $AB/BD = AC/CD$ .
10. If the median from  $A$  in a triangle  $ABC$  intersects the circumcircle  $\Gamma$  of  $ABC$  at  $D$ , then  $AB \cdot BD = AC \cdot CD$ .

11. (Euler's Formula) Let  $O$ ,  $I$  and  $I_a$  be the circumcenter, incenter and  $A$ -excenter of a triangle  $ABC$  with circumradius  $R$ , inradius  $r$  and  $A$ -exradius  $r_a$ . Then:
  - (a)  $OI = \sqrt{R(R - 2r)}$ .
  - (b)  $OI_a = \sqrt{R(R - 2r_a)}$ .
12. (Poncelet's Porism) Let  $\Gamma$  and  $\omega$  be two circles with centers  $O$  and  $I$  and radii  $R$  and  $r$ , respectively, such that  $OI = \sqrt{R(R - 2r)}$ . Let  $A, B$  and  $C$  be any three points on  $\Gamma$  such that lines  $AB$  and  $AC$  are tangent to  $\omega$ . Then line  $BC$  is also tangent to  $\omega$ .
13. (Apollonius Circle) Let  $ABC$  be a given triangle and let  $P$  be a point such that  $AB/BC = AP/PC$ . If the internal and external bisectors of angle  $\angle ABC$  meet line  $AC$  at  $Q$  and  $R$ , then  $P$  lies on the circle with diameter  $QR$ .

### Trigonometry:

1. (Sine Law) Given a triangle  $ABC$  with circumradius  $R$

$$\frac{BC}{\sin \angle A} = \frac{AC}{\sin \angle B} = \frac{AB}{\sin \angle C} = 2R$$

2. (Cosine Law) Given a triangle  $ABC$

$$BC^2 = AB^2 + AC^2 - 2 \cdot AB \cdot AC \cdot \cos \angle A$$

3. (Pythagorean Theorem) If  $ABC$  is a triangle, then  $\angle ABC = 90^\circ$  if and only if

$$AB^2 + BC^2 = AC^2$$

4. Given a triangle  $ABC$  and a point  $D$  on line  $BC$ , then

$$\frac{\sin \angle BAD}{\sin \angle CAD} = \frac{BD \cdot AC}{CD \cdot AB}$$

### Miscellaneous Synthetic Facts:

1. (Spiral Similarity) Let  $OAB$  and  $OCD$  be directly similar triangles. Then  $OAC$  and  $OBD$  are also directly similar triangles.
2. The unique center of spiral similarity sending  $AB$  to  $CD$  is the second intersection of the circumcircles of  $QAB$  and  $QCD$  where  $AC$  and  $BD$  intersect at  $Q$ .
3. Lines  $AB$  and  $CD$  are perpendicular if and only if  $AC^2 - AD^2 = BC^2 - BD^2$ .
4. (Apollonius Circle) Given two points  $A$  and  $B$  and a fixed  $r > 0$ , then the locus of points  $Q$  such that  $AQ/BQ = r$  is a circle  $\Gamma$  with center at the midpoint of  $Q_1Q_2$  where  $Q_1$  and  $Q_2$  are the two points on line  $AB$  satisfying  $AQ_i/BQ_i = r$  for  $i = 1, 2$ .
5. Let  $ABCD$  be a convex quadrilateral. The four interior angle bisectors of  $ABCD$  are concurrent and there exists a circle  $\Gamma$  tangent to the four sides of  $ABCD$  if and only if  $AB + CD = AD + BC$ .

## 6 Problems

The problems below have been arranged roughly in order of difficulty. I divided the problems into three difficulty classes: A, B and C.

- A1. (CMO 1997) The point  $O$  is situated inside the parallelogram  $ABCD$  such that  $\angle AOB + \angle COD = 180^\circ$ . Prove that  $\angle OBC = \angle ODC$ .
- A2. (APMO 2007) Let  $ABC$  be an acute angled triangle with  $\angle BAC = 60^\circ$  and  $AB > AC$ . Let  $I$  be the incenter, and  $H$  the orthocenter of the triangle  $ABC$ . Prove that  $2\angle AHI = 3\angle ABC$ .
- A3. (IMO 2006) Let  $ABC$  be triangle with incenter  $I$ . A point  $P$  in the interior of the triangle satisfies  $\angle PBA + \angle PCA = \angle PBC + \angle PCB$ . Show that  $AP \geq AI$ , and that equality holds if and only if  $P = I$ .
- A4. (IMO 2008) Let  $H$  be the orthocenter of an acute-angled triangle  $ABC$ . The circle  $\Gamma_A$  centered at the midpoint of  $BC$  and passing through  $H$  intersects the sideline  $BC$  at points  $A_1$  and  $A_2$ . Similarly, define the points  $B_1, B_2, C_1$  and  $C_2$ . Prove that six points  $A_1, A_2, B_1, B_2, C_1$  and  $C_2$  are concyclic.
- A5. (Russia 2012) The points  $A_1, B_1$  and  $C_1$  lie on the sides  $BC, CA$  and  $AB$  of the triangle  $ABC$ , respectively. Suppose that  $AB_1 - AC_1 = CA_1 - CB_1 = BC_1 - BA_1$ . Let  $O_A, O_B$  and  $O_C$  be the circumcenters of triangles  $AB_1C_1, A_1BC_1$  and  $A_1B_1C$  respectively. Prove that the incenter of triangle  $O_AO_BO_C$  is the incenter of triangle  $ABC$ .
- A6. (IMO Shortlist 2006) Let  $ABCD$  be a trapezoid with parallel sides  $AB > CD$ . Points  $K$  and  $L$  lie on the line segments  $AB$  and  $CD$ , respectively, so that  $\frac{AK}{KB} = \frac{DL}{LC}$ . Suppose that there are points  $P$  and  $Q$  on the line segment  $KL$  satisfying  $\angle APB = \angle BCD$  and  $\angle CQD = \angle ABC$ . Prove that the points  $P, Q, B$  and  $C$  are concyclic.
- A7. (IMO Shortlist 2008) Let  $ABCD$  be a convex quadrilateral and let  $P$  and  $Q$  be points in  $ABCD$  such that  $PQDA$  and  $QPBC$  are cyclic quadrilaterals. Suppose that there exists a point  $E$  on the line segment  $PQ$  such that  $\angle PAE = \angle QDE$  and  $\angle PBE = \angle QCE$ . Show that the quadrilateral  $ABCD$  is cyclic.
- B1. (IMO Shortlist 2000) Let  $O$  be the circumcenter and  $H$  the orthocenter of an acute triangle  $ABC$ . Show that there exist points  $D, E$ , and  $F$  on sides  $BC, CA$ , and  $AB$  respectively such that  $OD + DH = OE + EH = OF + FH$  and the lines  $AD, BE$ , and  $CF$  are concurrent.
- B2. (IMO Shortlist 2012) In an acute triangle  $ABC$  the points  $D, E$  and  $F$  are the feet of the altitudes through  $A, B$  and  $C$  respectively. The incenters of the triangles  $AEF$  and  $BDF$  are  $I_1$  and  $I_2$  respectively; the circumcenters of the triangles  $ACI_1$  and  $BCI_2$  are  $O_1$  and  $O_2$  respectively. Prove that  $I_1I_2$  and  $O_1O_2$  are parallel.

- B3. (IMO Shortlist 2005) Let  $ABCD$  be a parallelogram. A variable line  $g$  through the vertex  $A$  intersects the rays  $BC$  and  $DC$  at the points  $X$  and  $Y$ , respectively. Let  $K$  and  $L$  be the  $A$ -excenters of the triangles  $ABX$  and  $ADY$ . Show that the angle  $\angle KCL$  is independent of the line  $g$ .

- B4. (Tuymaada MO 2012) Point  $P$  is taken in the interior of the triangle  $ABC$ , so that

$$\angle PAB = \angle PCB = \frac{1}{4}(\angle A + \angle C).$$

Let  $L$  be the foot of the angle bisector of  $\angle B$ . The line  $PL$  meets the circumcircle of  $\triangle APC$  at point  $Q$ . Prove that  $QB$  is the angle bisector of  $\angle AQC$ .

- B5. (Japan MO 2009) Let  $\Gamma$  be the circumcircle of a triangle  $ABC$ . A circle with center  $O$  touches to line segment  $BC$  at  $P$  and touches the arc  $BC$  of  $\Gamma$  which doesn't have  $A$  at  $Q$ . If  $\angle BAO = \angle CAO$ , then prove that  $\angle PAO = \angle QAO$ .
- B6. (Bulgarian TST 2004) The points  $P$  and  $Q$  lie on the diagonals  $AC$  and  $BD$ , respectively, of a quadrilateral  $ABCD$  such that  $\frac{AP}{AC} + \frac{BQ}{BD} = 1$ . The line  $PQ$  meets the sides  $AD$  and  $BC$  at points  $M$  and  $N$ . Prove that the circumcircles of the triangles  $AMP$ ,  $BNQ$ ,  $DMQ$ , and  $CNP$  are concurrent.
- B7. (Chinese TST 2002) Circles  $\omega_1$  and  $\omega_2$  intersect at points  $A$  and  $B$ . Points  $C$  and  $D$  are on circles  $\omega_1$  and  $\omega_2$ , respectively, such that lines  $AC$  and  $AD$  are tangent to circles  $\omega_2$  and  $\omega_1$ , respectively. Let  $I_1$  and  $I_2$  be the incenters of triangles  $ABC$  and  $ABD$ , respectively. Segments  $I_1I_2$  and  $AB$  intersect at  $E$ . Prove that  $\frac{1}{AE} = \frac{1}{AC} + \frac{1}{AD}$ .
- C1. (Russia 2012) The point  $E$  is the midpoint of the segment connecting the orthocenter of the scalene triangle  $ABC$  and the point  $A$ . The incircle of triangle  $ABC$  is tangent to  $AB$  and  $AC$  at points  $C'$  and  $B'$ , respectively. Prove that point  $F$ , the point symmetric to point  $E$  with respect to line  $B'C'$ , lies on the line that passes through both the circumcenter and the incenter of triangle  $ABC$ .
- C2. (Chinese TST 2005) Let  $\omega$  be the circumcircle of  $\triangle ABC$ .  $P$  is an interior point of  $\triangle ABC$ .  $A_1, B_1, C_1$  are the intersections of  $AP, BP, CP$  respectively and  $A_2, B_2, C_2$  are the symmetrical points of  $A_1, B_1, C_1$  with respect to the midpoints of side  $BC, CA, AB$ . Show that the circumcircle of  $\triangle A_2B_2C_2$  passes through the orthocentre of  $\triangle ABC$ .
- C3. (USA TST 2005) Let  $ABC$  be an acute scalene triangle with  $O$  as its circumcenter. Point  $P$  lies inside triangle  $ABC$  with  $\angle PAB = \angle PBC$  and  $\angle PAC = \angle PCB$ . Point  $Q$  lies on line  $BC$  with  $QA = QP$ . Prove that  $\angle AQP = 2\angle OQB$ .
- C4. (USA TST 2006) Let  $ABC$  be a triangle. Triangles  $PAB$  and  $QAC$  are constructed outside of triangle  $ABC$  such that  $AP = AB$  and  $AQ = AC$  and  $\angle BAP = \angle CAQ$ . Segments  $BQ$  and  $CP$  meet at  $R$ . Let  $O$  be the circumcenter of triangle  $BCR$ . Prove that  $AO \perp PQ$ .

- C5. (RMM 2012) Let  $ABC$  be a triangle and let  $I$  and  $O$  denote its incentre and circumcentre respectively. Let  $\omega_A$  be the circle through  $B$  and  $C$  which is tangent to the incircle of the triangle  $ABC$ ; the circles  $\omega_B$  and  $\omega_C$  are defined similarly. The circles  $\omega_B$  and  $\omega_C$  meet at a point  $A'$  distinct from  $A$ ; the points  $B'$  and  $C'$  are defined similarly. Prove that the lines  $AA'$ ,  $BB'$  and  $CC'$  are concurrent at a point on the line  $IO$ .
- C6. (RMM 2011) A triangle  $ABC$  is inscribed in a circle  $\omega$ . A variable line  $\ell$  chosen parallel to  $BC$  meets segments  $AB$ ,  $AC$  at points  $D$ ,  $E$  respectively, and meets  $\omega$  at points  $K$ ,  $L$  (where  $D$  lies between  $K$  and  $E$ ). Circle  $\gamma_1$  is tangent to the segments  $KD$  and  $BD$  and also tangent to  $\omega$ , while circle  $\gamma_2$  is tangent to the segments  $LE$  and  $CE$  and also tangent to  $\omega$ . Determine the locus, as  $\ell$  varies, of the meeting point of the common inner tangents to  $\gamma_1$  and  $\gamma_2$ .