Note 1: The Problem with Spin Operators and the Jordan–Wigner Transformation

1. The Problem with Spin Operators

Let us define the spin raising and lowering operators:

$$S_i^+ = S_i^x + iS_i^y, \tag{1}$$
$$S_i^- = S_i^x - iS_i^y. \tag{2}$$

We cannot get rid of S^z, though. This is not a trouble with the physical Hamiltonian itself, but rather it is an artifact of the $\mathfrak{su}(2)$ spin algebra. Since

$$[S_i^+, S_i^-] = 2S_i^z, \tag{3}$$

the moment a Heisenberg equation of motion (EOM) is to be solved, S^z pops out.

Like say, the typical XY Hamiltonian can be represented as:

$$\begin{aligned} H_{xy} &= J\sum_{\langle i,j\rangle}\left(S_i^xS_j^x + S_i^yS_j^y\right) \\ &= -\frac{J}{2}\sum_{\langle i,j\rangle}\left(S_i^+S_j^- + S_i^-S_j^+\right). \end{aligned} \tag{4}$$

Trying to solve for time dependence (in the Heisenberg picture):

$$[S_{k,H}^+, H_{xy}] = -\frac{J}{2}\sum_{\langle i,j\rangle}[S_k^+, S_i^+S_j^- + S_i^-S_j^+] \quad \text{— } (\dagger) \tag{5}$$

where S_i^z appears again. (Conventionally, it is said that $\mathfrak{su}(2)$ algebra does not close without S^z.)

What is the trouble with this? Notice that to solve for (†), we need to know the time dependence of S^z itself, which in turn requires $[S_k^z, H_{xy}]$ to be computed. This outputs S^+ and S^- again! S^z ends up acting like an interaction term that couples together a hierarchy of differential equations, making the problem complex.

Also, Hamiltonians must be Hermitian! To allow for this, each string of operators in it must be accompanied by its Hermitian conjugate (H.c.). This is due to $(S^+)^\dagger = S^-$ and vice versa.

What do these spin operators act like? Notice that:

$$\{S_i^+, S_i^-\} = S_i^+S_i^- + S_i^-S_i^+ = 1. \tag{6}$$

2. Hard-Core Bosons and the Jordan–Wigner Transformation

But for different sites ($i \neq j$):

$$[S_i^+, S_j^-] = 0. \tag{7}$$

So, they act like fermionic creation/annihilation operators on the same site, but like bosonic operators on different sites — **hard-core bosons**!

The Pauli Exclusion Principle (PEP) is not satisfied globally. Mapping $S_i^- \to c_i$ directly fails to map to standard bosonic or fermionic operators.

Since $(S_i^+)^2 = (S_i^-)^2 = 0$, the constraint demands an on-site repulsion term in the Hamiltonian preventing two bosons on the same site. (Apparently interacting boson problems are notoriously difficult to solve — find out why.)

Why do we want such a mapping? There exist standard methods to diagonalize and quantize free fermionic/bosonic systems, namely Fourier transforms.

Jordan–Wigner Transformation (1D Chain)

Let us assume a 1D chain of spin-1/2 particles. Our fundamental question here is: *Can we construct fermionic operators out of purely spin-like matrices?*

The representation of the ladder operators $S_i^\pm$ at a particular site i on the chain, represented in the full Hilbert space of the problem, is a tensor product:

$$S_i^\pm = I \otimes I \otimes \cdots \otimes \sigma^\pm \otimes \cdots \otimes I \quad (i\text{-th position}), \tag{8}$$

with N operators for N spins in the chain.

How can we then force two commuting operators $S_i^\pm$ and $S_j^\pm$ ($i \neq j$), operating on different tensor spaces, to anti-commute? We require:

$$S_i^\pm S_j^\pm = -S_j^\pm S_i^\pm. \tag{9}$$

If our new fermionic operators modify the operator at site j to carry/possess an S^z term from another site i, then the ladder operators at site i will anti-commute with the operator at site j. Of course, we require this to work for all sites simultaneously.

3. Constructing the Transformation

Personal Conjecturing: What will it take for the transform to work for every pair? We want to map:

$$S_i^+ \longrightarrow c_i^\dagger, \tag{10}$$

$$S_i^- \longrightarrow c_i, \tag{11}$$

such that c_i and $c_i^\dagger$ describe spinless fermion annihilation/creation operators at the i-th site:

$$\{c_i, c_j\} = 0, \quad \{c_i^\dagger, c_j^\dagger\} = 0, \quad \{c_i, c_j^\dagger\} = \delta_{ij}. \tag{12}$$

We also need to account for the underlying spin algebra constraints:

1. $[S_i^\pm, S_j^\pm] = 0 \quad \forall\, i \neq j$ (Bosonic off-site)
2. $[S_i^+, S_i^-] = 2S_i^z$ ($\mathfrak{su}(2)$ quirk)
3. $\{S_i^-, S_i^+\} = 1 \quad \forall\, i$ (Fermionic on-site, already satisfied)
4. $\{S_i^\pm, S_i^\pm\} = 0$ (Fermionic on-site, partially satisfied since $(S_i^\pm)^2 = 0$)

Ordering and Asymmetry: Since $S_i^\dagger S_j^\dagger = S_j^\dagger S_i^\dagger$ off-site, we require asymmetry in the mapping to spinless fermions. Why? Otherwise i and j will give two (-1) phase factors that cancel out to retain the bosonic behavior off-site. In other words, we must order the spins $(1, 2, 3, \ldots, N)$, with a later spin being affected by earlier spins in the map.

Assume c_j carries components other than S_j^- to account for all sites $l < j$:

$$c_j = K_j S_j^-, \tag{13}$$

where K_j is a modifier from sites $l < j$, and is not modified by the j-th site itself.

Useful relation: $\{AB, C\} = A[B, C] + \{A, C\}B$.

4. The Jordan–Wigner String Solution

Conditions on K_j:

- $\{K_j, S_i^-\} = 0 \quad \forall\, i < j$ (c_j contains modifier for i-th site)
- $[K_j, S_i^-] = 0 \quad \forall\, i > j$ (K_j must commute with all S_i^- for $i > j$)

Relevant Commutators:

$$[S_i^z, S_i^\pm] = \pm S_i^\pm \implies S_i^z S_i^\pm - S_i^\pm S_i^z = \pm S_i^\pm, \tag{14}$$

$$S_i^+ S_i^- - S_i^- S_i^+ = 2S_i^z. \tag{15}$$

Jordan–Wigner Map: The non-local string factor K_j is given by:

$$K_j = \prod_{l<j} \exp\left(i\pi S_l^+ S_l^-\right) = \prod_{l<j} \exp\left(i\pi n_l\right). \tag{16}$$

Thus, the Jordan–Wigner transformation reads:

$$c_j = \left[\prod_{l<j} \exp\left(i\pi S_l^+ S_l^-\right)\right] S_j^-, \tag{17}$$

$$c_j^\dagger = \left[\prod_{l<j} \exp\left(-i\pi S_l^+ S_l^-\right)\right] S_j^+. \tag{18}$$

Phase Factor Analysis: Note that $n_l \to 1$ if $|\uparrow\rangle_l$ and 0 if $|\downarrow\rangle_l$. The phase operator satisfies $\exp(i\pi n_l) = \exp(-i\pi n_l)$, making it Hermitian:

$$\exp(i\pi n_l) = 1 - 2n_l = 1 - 2S_l^+ S_l^-. \tag{19}$$

Using $1 - 2S_l^+ S_l^- = 1 - S_l^+ S_l^- - (2S_l^z + S_l^- S_l^+)$, this simplifies to:

$$1 - 2S_l^+ S_l^- = -2S_l^z = -\sigma_l^z. \tag{20}$$

The local S_l^z component stays as the core building block of the string phase!

Note: It appears that Jordan–Wigner spinless fermions are domain wall excitations!
(Now draw parallels to the Kitaev honeycomb method!)