

# Impact of Local Time on St. Patrick Geomagnetic Storm response at mid-low latitudes

Castellanos-Velazco C. I.<sup>1,3</sup> Corona-Romero P.<sup>2,3</sup> Sergeeva M.<sup>2,3</sup>  
Orrala-[] I. D Orrala-Legorreta I.D.<sup>1,3</sup>, Gonzalez-Esparza J. A.<sup>2,3</sup>

<sup>1</sup>Universidad Nacional Autonoma de Mexico, Posgrado en Ciencias de la Tierra

<sup>2</sup> Laboratorio Nacional de Clima Espacial

<sup>3</sup> Instituto de Geofisica

## Abstract

This study analyzes magnetic fluctuations associated with Prompt Penetration Electric Fields (PPEFs) using data from four mid-low latitude observatories separated by 60°–90° in longitude, covering different local time sectors. We focus on the St. Patrick’s Day geomagnetic storm of 17 March 2015. Using Fourier analysis, we reconstruct the magnetic signatures of ionospheric currents. Our results demonstrate that these ionospheric magnetic responses depend strongly on longitudinal location during the storm’s main phase and are particularly influenced by partial ring current activity.

**Keywords:** Regional Space weather, Magnetic fluctuations, Prompt-Penetration Electric Fields

## 1 Introduction

A geomagnetic storm (GS) is a temporary disturbance of Earth’s magnetosphere caused by the injection of energy and charged particles into the system, leading to a significant reduction in the geomagnetic field intensity. These GS are typically triggered by solar phenomena such as Coronal Mass Ejections (CMEs) or high-speed solar wind (HSSW) streams. Due to their potential impacts on technological systems—including disruptions to communications, navigation, and power grids—GSs are among the most extensively studied phenomena in space weather research.

During the Main Phase (MP) of a GS, energy and charged particles are transported into Earth’s inner magnetosphere. Since the magnetosphere is dynamically coupled with the ionosphere, particularly at high latitudes [C.T Russell, 2016], the ionosphere exhibits pronounced responses to these perturbations. One critical ionospheric effect is the generation of storm-time electric fields, which can be manifested at different latitudes:

- Prompt Penetration Electric Fields (PPEF) [Nishida, 1968a,b]
- Disturbed Dynamo Electric Fields (DDEF) [Blanc and Richmond, 1980]

The concept of PPEF was first introduced by Nishida [1968a,b], who observed periodic magnetic fluctuations between 4 hours and 30 minutes in ground magnetometers located around the auroral oval and the dip equator, both showing nearly identical fluctuations. Later, Vasyliunas [1970], Jaggi and Wolf [1973] developed the first model explaining the penetration of convective electric fields ( $\mathbf{E}_C$ ) during disturbed periods, propagating through Region 1 Field-Aligned Currents (R1-FAC). During quiet periods,  $\mathbf{E}_C$  is shielded by a polarized electric field induced by Region 2 Field-Aligned Currents (R2-FAC). However, during the MP of a GS, magnetic reconnection leads to an almost immediate intensification of R1-FAC activity due to the injection of solar wind particles and energy. Since R2-FAC depends on ring current activity to strengthen, it initially fails the shielding of  $\mathbf{E}_C$ , allowing its transfer to the ionosphere through R1-FAC. This process continues until R2-FAC intensifies its activity though the intensification of ring current, allowing R2-FAC balancing R1-FAC activity, which can take from tens of minutes to several hours [Wei et al., 2015]. During this period of imbalance,  $\mathbf{E}_C$  penetrates into the mid- and low-latitude ionosphere, being that the reason for be called Penetration Electric fields. Furthermore, Kelley et al.

[1979], Wei et al. [2015] showed that PPEF depends on the southward orientation of the interplanetary magnetic field (IMF) component  $B_Z$  and can also be triggered by sharp directional changes in  $B_Z$ .

DDEF on the other hand, are associated with an ionospheric current driven by the injection of charged particles into the auroral ionosphere. This injection, produces a Joule heating and leads to the circulation of equatorward thermospheric winds which drag charged particles (protons) Blanc and Richmond [1980]. With the dragged protons, it gets to a charge separation which induces a Pedersen current in the same direction of neutral wind flow. Near the equator, the Coriolis effect produces proton accumulation and westward circulation occur around the dip equator, generating a poleward electric field. Consequently, a second Pedersen current is induced in the poleward direction, opposing the first current. At mid-latitudes ( $\sim 40^\circ$ ), the above commented phenomena, drives Hall currents, which are interrupted at the terminators (limit of day-side with night-side) and get a closure or circuit at adjacent latitudes. This process results in vortex-shaped currents known as the Disturbed Dynamo, characterized by directions opposite to the Solar Quiet (SQ) current. For this reason, the disturbed dynamo is also referred to as the anti-SQ current.

Both PPEF and DDEF were studied by Fejer et al. [1983], who observed their effects using incoherent scatter radar at Jicamarca latitude. Later, Amory-Mazaudier [1982] detected DDEF above Saint-Santin latitude, with results consistent with the disturbed dynamo model. On the other hand, Le Huy and Amory-Mazaudier [2005] modeled by the first time the magnetic field associated with ionospheric disturbances, assuming that, for mid-low latitudes, they are mainly driven by DDEF and PPEF. They considered that during the MP, PPEF magnetic effects are dominant, whereas during the recovery phase (RP), regional magnetic field variations are mainly driven by DDEF. Following this initial approach, several studies [Le Huy and Amory-Mazaudier, 2008, Zaka et al., 2009] have focused on analyzing local responses at different latitudes and longitudes to analyze the different local responses associate to DDEF phenomena. In Liu et al. [2013], a statistical study was conducted to understand ionospheric responses as a function of local sectors during GS onset. Later, Younas et al. [2020] proposed using Fourier analysis to isolate magnetic fluctuations related to DDEF and PPEF based on their quasi-periodic behavior, identifying periods of approximately 24 hours for DDEF and less than 4 hours for PPEF.

In case of Mexico, regional space weather has a relatively recent history. On 2014, it was form the Mexican Space Weather Service (SCIESMEX), leading to combined efforts to understand the regional space weather phenomena, in the middle of a global perspective Gonzalez-Esparza et al. [2017]. From this teamwork, studies like Corona-Romero et al. [2018, 2017] has been focused on local geomagnetic response during GS periods, and the computing of a local K index specific for Mexico. On the other hand, Romero-Hernandez et al. [2017], Sergeeva et al. [2017], Sergeeva et al. [2020] found important ionospheric activity related to geomagnetic storms. In Castellanos-Velazco et al. [2024], the main focus was analyzing the local geomagnetic response related to PPEF and DDEF phenomena, during 20 strong GS, from 2003 to 2018, where one of the main conclusions were the need of comparing the geomagnetic response in the center of Mexico with different locations at similar geomagnetic latitudes.

In this paper, we analyze the PPEF magnetic signatures at different longitude sectors, looking for the sector where these fluctuations are more intense. We compare the results with the Asymmetric Ring Current activity in order to analyze it's influence on PPEF strength and so, on local geomagnetic response. We selected the GS of march 17<sup>th</sup>, 2015 which is also known as the St. Patrick's event. This GS is known to be the most intense GS during the solar cycle 24<sup>th</sup>, reaching -274 nT during its MP.

## 2 St. Patrick Geomagnetic Storm

The geomagnetic storm (GS) analyzed in this work is the intense event that occurred on March 17, 2015, widely known as the St. Patrick's Day storm. It is considered the most intense storm of solar cycle 24 Wu et al. [2016]. The storm was triggered by a coronal mass ejection (CME) associated with a C9.1 solar flare and a subsequent dimming observed on March 15. This CME, recorded as a partial halo by the SOHO/LASCO coronagraphs, had an estimated speed of approximately 660 km/s.

Figure 1 provides an overview of the interplanetary conditions that led to the St. Patrick's Day storm. A more detailed description about the trigger event is provided by Wu et al. [2016].

Panel (a) shows the total interplanetary magnetic field,  $B_T$  (black line), and its north-south component,  $B_Z$  (red line). Panels (b) and (c) presents solar wind properties, including the flow speed  $V_X$  (positive in anti sunward direction, black line), proton density  $n_P$  (orange line) and proton temperature  $T_P$  (blue line). In panel (b) also shows the driven Interplanetary Electric Field  $E_Y$ , calculated as  $-V_X \times B_Z$ . Finally, panel (e) displays the geomagnetic response through the SYMH index (green line).

The key milestones of the event are marked by three vertical dashed lines in Figure 1. The first line (left) indicates the arrival of an interplanetary shock (IPS) at 04:30 UT at the L1 point. The IPS signature is clear from the step-like increases in  $B_T$ ,  $V_X$  and  $n_P$ . This shock compressed the magnetosphere, causing a Sudden Storm Commencement (SSC) visible as a sharp increase in the SYMH index between 04:30 and 07:00 UT, highlighted by light gray shading.

The region following the shock, the CME sheath, is identified by highly fluctuating  $B_T$  and  $B_Z$ , enhanced  $n_P$  and rising  $T_P$ . A period of southward  $B_Z$  between 06:00 and 09:00 UT initiated the what we called, the first stage of main phase (MP), driving the SYMH index down to its first minimum of -80 nT.

The second dashed line ( 14:00 UT) marks the arrival of the magnetic cloud, identified by Wu et al. [2016] as the main driver of the GS. Its passage is identified by a smooth, strong rotation of the magnetic field, a drop in temperature, and so, a corresponding low plasma beta. A sustained and intense southward  $B_Z$  component (reaching -40 nT) within the cloud marked the beginning of the second (and more intense) stage of the MP. The SYMH index decreased sharply, reaching its global minimum of -234 nT at 22:47 UT. The entire main phase period is shaded in dark gray in panel (e).

The third dashed line ( 23:00 UT) indicates the trailing edge of the magnetic cloud and the entry into a high-speed solar wind (HSSW) stream. While the high solar wind speed persists, the geomagnetic activity subsides as  $B_Z$  turns northward and the  $E_Y$  drops close to zero, ending the geoeffective driving of the storm.

## 3 Procedure

### 3.1 Geomagnetic Data

The geomagnetic data analyzed in this paper were downloaded from the INTERMAGNET public database (*International Real-Time Magnetic Observatory Network* [INTERMAGNET, 2021]), corresponding to the observatories (OBs) listed in Table 1. All OBs are located within a narrow range of magnetic latitude ( $21^\circ$ – $33^\circ$ ) and are longitudinally separated by  $60^\circ$ – $90^\circ$  from each other. The closest OB is KAK, while the farthest is TEO from JAI. However, this spacing was tolerated since KAK is the closest OB to the longitude of  $165^\circ$  E.

Table 1: List of observatories considered for this study. Each OB is separated longitudinally at  $70 - 90^\circ$  apart.

| Observatory | IAGA code | Country/<br>Territory | Latitude   |          | Longitude  |          |
|-------------|-----------|-----------------------|------------|----------|------------|----------|
|             |           |                       | Geographic | Magnetic | Geographic | Magnetic |
| Guimar      | GUI       | Spain                 | 28.317 N   | 33.21 N  | 16.433 W   | 61.1 E   |
| Jaipur      | JAI       | India                 | 26.917 N   | 18.42 N  | 75.8 E     | 150.47 E |
| Kakioka     | KAK       | Japan                 | 36.232 N   | 27.8 N   | 140.186 E  | 149.73 W |
| Teoloyucan  | TEO       | Mexico                | 19.747 N   | 27.81 N  | 99.182 W   | 28.4 W   |

To study how the local time (LT) sector affects the geomagnetic response during the main phase (MP) of a geomagnetic storm (GS), we focused on data from the four observatories (OB) listed in Table 1. Our aim was to analyze the GS response at these OB during a time window (TW) of maximum partial ring current (PRC) activity, from 14:07 to 01:00 UT (11 hours), covering the second stage of the MP and the first few hours of the recovery phase (RP).




Figure 1: Interplanetary measurements. Panel (a): interplanetary magnetic field total component (black line) and z component (red line). Panel (b):  $V_X$  component of solar wind velocity (black line) and Interplanetary Electric Field (blue line). Panel (c): Proton Temperature (blue line) and Proton Density (orange line). Panel (d): SYMH (green line) and ASYMH index (orange line), highlighting the initial phase (light gray shade), main phase 1 (medium gray shade) and main phase 2 (dark gray shade).




Figure 2: Geographic position of the four magnetic OB, distributed at  $70 - 90^\circ$  from each other

We therefore considered four local TW sectors during which the MP of the GS was active. These sectors are:

- morning sector, 06:01 - 12:00 LT.
- Afternoon sector, 12:01 - 18:00 LT.
- night sector, 18:01 - 00:00 LT.
- After midnight sector, 00:01 - 06:00 LT

Considering that the analyzed TW lasts approximately 11 hours and the local times at which it begins and ends, each OB is located within approximately two of these sectors during the TW. For this reason, we divided the TW into two subsets (TW1 and TW2), which for each OB cover two of the mentioned LT sectors (covering most of the sectors for GUI and KAK).

In Table 2, we present a brief description of the time windows during which each OB was located in TW1 and TW2. It can be seen that TEO was located in the morning and afternoon sectors, while JAI was located in the night and post-midnight sectors, placing these two at almost longitudinally opposite locations. GUI and KAK, on the other hand, are slightly shifted from the afternoon/night and post-midnight/morning sectors, respectively. This is consistent with the fact that they are not exactly  $90^\circ \pm 10^\circ$  apart from the adjacent OBs.

Table 2: Characteristics of the St. Patrick's Day event. Column 1 displays TW1 and TW2 in UT spanning March 17 and the first hour of March 18. Columns 2 and 3 present the minimum (SYMH) and maximum (ASYH) values observed during the main phase, respectively. Columns 4–7 show the local time sectors for each observatory during TW1 and TW2.

| GS TW             | Min SYMH | Max ASYH | Local TW      |               |               |               |
|-------------------|----------|----------|---------------|---------------|---------------|---------------|
| UT                | [nT]     | [nT]     | GUI           | JAI           | KAK           | TEO           |
| TW1 13:01 - 19:00 | -234     | 273      | 11:01 - 17:00 | 18:01 - 00:00 | 22:01 - 04:00 | 06:01 - 12:00 |
| TW2 19:00 - 01:00 | -234     | 174      | 17:00 - 23:00 | 00:00 - 06:00 | 04:00 - 10:00 | 12:00 - 18:00 |

### 3.2 Magnetic fluctuations of ionospheric origin

It is well accepted that [Rishbeth and Garriott, 1969, Knecht and B.M, 1985, Gjerloev, 2012, van de Kamp, 2013] local recordings  $H_{loc}$  of a ground magnetometer can be considered a linear combination of different geomagnetic sources:

$$H_{loc} = H_{SQ} + H_0 + H_{MT} + H_{others} + H_I, \quad (1)$$

where  $H_{SQ}$  represents the diurnal variation and  $H_0$  is the main field during a month window.  $H_{MT}$  represents the variations induced by the intensification of magnetospheric currents during the GS. In the case of  $H_{others}$ , it corresponds to magnetic sources less significant, like (and no limited to)) magnetotellurics, crustal magnetization, etc; which can be neglected for not being considered as relevant compared to the others mentioned. Finally,  $H_I$  corresponds to the magnetic variations driven by ionospheric current activity during a GS. Thus, the local geomagnetic response during a GS is given by:

$$\Delta H = H_{loc} - (H_{SQ} + H_0). \quad (2)$$

We provide a detailed explanation of how we model the regular variations in Appendix A.

From Le Huy and Amory-Mazaudier [2005], Amory-Mazaudier et al. [2017], Younas et al. [2020], for mid-low magnetic latitudes, with  $\lambda$  as the geomagnetic latitude of each OB, we can approximate  $H_{MT}$  as:

$$H_{MT} \approx SYMH \cdot \cos(\lambda), \quad (3)$$

Thus,  $H_I$  can be approximated as:

$$H_I = H_{DDEF} + H_{PPEF} \approx H - (H_{SQ} + H_0 + SYMH - H \cdot \cos(\lambda)), \quad (4)$$

where  $H_{DDEF}$  and  $H_{PPEF}$  represent the magnetic fields associated with the disturbed dynamo electric fields and the prompt penetration electric fields, respectively.

In order to isolate  $H_{DDEF}$  and  $H_{PPEF}$  from each other, we proceed under the assumption that both ionospheric currents drive quasi-periodic fluctuations [Nishida, 1968a, Blanc and Richmond, 1980, Younas et al., 2020]. Hence, we begin with a Fourier analysis, applying passband filters to extract fluctuations with frequencies around  $11.5\mu\text{Hz}$  ( 24-hour periods) to reconstruct  $H_{DDEF}$ . For  $H_{PPEF}$ , we apply high-pass filters for frequencies higher than  $69.5\mu\text{Hz}$  (corresponding to periods  $\leq 4$  hours).

### 3.3 Total Electron Content

The overall ionosphere dynamics in each considered region were characterized by the vertical Total Electron Content (TEC) variations in the vicinity of the chosen magnetic OB. TEC values were derived from GNSS measurements using TayAbsTEC method Yasyukevich et al. [2015a,b], Mylnikova et al. [2015]. In particular, data from UCOE, LPAL, NPGJ and TSK2 receivers was used to obtain TEC values for TEO, GUI, JAI and KAK regions, respectively.

However, the open data obtained for this work, doesn't have the best quality, even more considering the period of time. The quality of data is a sum of errors which depend on the satellite swarm, so as the receptor itself. We could smooth some of these errors but couldn't be avoided at all.

Considering that we are studying results from points in different longitudinal sectors (Mexico, Canarias Islands, India and Japan), there can be issues by comparing absolute TEC values. To deal with this issue, we instead analyze relative values, which is, measuring TEC percentage difference of TEC response with respect to the 27 days median TEC prior to the GS Vijaya Lekshmi et al. [2011], Sergeeva et al. [2017].

## 4 Results

In the following section, we focus on the results related to the geomagnetic response to the St. Patrick's GS for each OB listed in Table 1. Then we show the results of Equation 4 for isolating  $H_I$ , as well as the reconstruction of  $H_{PPEF}$  through Fourier analysis. Finally, we provide an analysis related to local




Figure 3: Top panels, local geomagnetic  $\Delta H$  (black line), ASYH (yellow line) and the magnetic contribution of ionospheric origin (red line). Mid panels, reconstruction of  $H_{PPEF}$ . Bottom panels, TEC (orange line) and median TEC (blue line). In the upper side of each panel, the Time Windows (TW) are highlighted with a blue and black solid line respectively and vertical dotted lines. Each gray shading represents each night hours in Local Time

time sector in which the GS developed as well as a comparison between each OB result.

In Figure 3, we show in top panels, the local index  $\Delta H$  (black line), the ASYH index (yellow line), and the computed ionospheric disturbance magnetic contribution  $H_I$  (red line) for each observatory. We focus on the period, TW, of high ASYH activity from 13:00 UT on March 17 to 01:00 UT the next day. As mentioned previously, this period is divided into two subsets (TW1 and TW2), highlighted by vertical dashed lines and two solid horizontal lines at the top of each panel (blue for TW1 and green for TW2). During these periods,  $H_I$  is primarily driven by  $H_{PPEF}$  fluctuations. The results of applying frequency band filters to the  $\text{FFT}(H_I)$ , compared with the ionospheric response are shown in mid panels, represented by red lines. Finally, in bottom panels, we show the TEC response (orange lines), compared with the median TEC (blue line).

We will focus on periods where ASYH exhibited sharp temporal enhancements, or ASYH pulses, at different universal times. Our aim is to observe how each observatory registered different local responses depending on the local time at which these pulses occurred. The first pulse (273 nT at 14:07 UT), which is the most intense, is our primary interest, but we identified three additional pulses. The second pulse (212 nT) occurs at 16:42 UT, while the third (150 nT) is located almost at the transition between TW1 and TW2, at 19:17 UT. Finally, we identify a fourth pulse (174 nT) around 23:00 UT, near the end of TW2.

In the case of Ionospheric response, we are focusing primarily on whether or not, we observe a positive or negative response, given the conditions of GNSS data used for the vertical TEC computation. As mentioned in the previous section, we are analyzing the relative variations of TEC response with respect to the 27 days median TEC. Considering that we are analyzing quite different longitude positions, the threshold we are considering to define disturbing conditions is above 30% Vijaya Lekshmi et al. [2011].

We begin by analyzing the magnetic data from GUI. During its local TW1, GUI was on the dayside (11:00-17:00 LT), while TW2 occurred mostly on the nightside. The transition between TW1 and TW2 is nearly centered on the dusk sector (18:00 LT). The peak ASYH enhancement (274 nT) occurs at 12:07 LT at GUI, nearly coinciding with the noon sector. A few hours later at 14:42 LT, there is another ASYH enhancement (212 nT).  $\Delta H$  shows a smooth decrease during TW1, consistent with the main phase, while  $H_I$  appears to exhibit a pattern opposite to ASYH. During TW2, ASYH shows significantly reduced activity, with two smaller enhancements at 17:17 LT and 22:00 LT. In this period,  $\Delta H$  maintains sustained minimum values around -212 nT until 21:00 LT, when it suddenly increases concurrently with an ASYH enhancement.  $H_{PPEF}$  fluctuation amplitudes within TW1 and TW2 (afternoon and night sectors, respectively) show peak amplitudes of -29.98 nT at 14:56 LT and 35.66 nT at 21:48 LT. Outside these peaks, we observe little contrast between  $H_{PPEF}$  amplitudes inside and outside the TW1 and TW2 periods. This increase is also evident in  $H_I$ . In TEC response, we observe an enhancement period around 16:00 LT, reaching an increment of around 55% of  $\text{TEC}_0$ , followed by a second enhancement of 58% almost at the end of TW2 and near midnight.

For JAI magnetic data, we observe that both time windows occur when JAI is on the night side. During TW1, the peak ASYH enhancement occurs at 19:07 LT, while both  $\Delta H$  and  $H_I$  show a more pronounced decrease compared to GUI.  $\Delta H$  reaches its minimum nearly simultaneously with the second ASYH pulse. During TW1,  $H_I$  appears to have a more clearly opposite pattern to ASYH, which changes completely during TW2. Within TW2 (post-midnight sector),  $\Delta H$  exhibits periods of pronounced decreases and increases, differing from the behavior at GUI. Around 04:00 LT, there is a sharp increase in both  $\Delta H$  and  $H_I$ , corresponding to the final ASYH enhancement in TW2. JAI observatory presents a higher contrast in  $H_{PPEF}$  fluctuations between inside and outside TW1 and TW2 (night and post-midnight sectors for JAI). The  $H_{PPEF}$  fluctuations show a sharp enhancement upon entering TW1, immediately after the dusk sector. During both TW1 and TW2,  $H_{PPEF}$  maintains values above  $|20|$  nT, with peaks of -41.32 nT at 18:55 LT (TW1) and 52.73 nT at 04:55 LT (TW2). TEC response on the other hand, does not present a significant difference until the end of TW2, on March 18. During this day, we observe a consistent negative response of around 70%.

At KAK, we observe during TW1 that  $\Delta H$  shows a significantly less pronounced decrease during ASYH activity compared to the previous observatories. The minimum  $\Delta H$  is reached at approximately 02:40 LT, temporally correlated with a small ASYH pulse.  $H_I$  also shows a pattern opposite to the

ASYH index, though this is only evident until 01:00 LT. After this time,  $H_I$  gradually appears to follow a pattern more similar to ASYH. During TW2,  $H_I$  maintains intense negative values between -90 nT and -150 nT. There is no apparent response in  $\Delta H$  during the final ASYH pulse beginning at 08:00 LT. Conversely,  $H_I$  shows a growth phase between the two time windows, reaching a maximum value shortly after dawn at 06:00 LT. For KAK  $H_{PPEF}$  amplitudes are less intense than at JAI, with peaks of 30.58 nT at 00:48 LT and -29.78 nT at 02:40 LT, both within TW1. Apart from the observed contrast in  $H_{PPEF}$  amplitudes upon entering TW1, we observe a decreasing amplitude trend transitioning from TW1 to TW2 toward the dayside. In TEC, we observe similar to JAI, a negative response of 50% the next day of MP.

For TEO, which represents the opposite case of JAI, we observe that both TW1 and TW2 occur during daytime hours (morning and afternoon sectors, respectively). In both time windows, we observe an almost linear descending trend in  $\Delta H$ , unlike the previous cases where the behavior changed between time windows.  $\Delta H$  reaches a minimum value at approximately 16:20 LT, consistent with the timing of the final ASYH pulse.  $H_I$ , on the other hand, shows a pattern more similar to ASYH behavior during TW1, with positive values that contrast with previous cases and peak values occurring approximately simultaneously with ASYH pulses. During TW2, however,  $H_I$  begins a decreasing phase, reaching a minimum at the same time as  $\Delta H$  and the final ASYH enhancement. We observe  $H_{PPEF}$  amplitude peaks of 35.12 nT at 15:57 LT and -34.63 nT at 16:40 LT during TW2. TEO also shows a noticeable contrast in  $H_{PPEF}$  amplitudes inside versus outside the TW1 and TW2 periods. Examining the TEC DIX, we observe two main peaks: the first occurring near the end of TW1 at 11:00 LT, and the second around 06:00 LT the following day. While the first peak can be related to high  $H_{PPEF}$  activity at the end of TW1, the second peak cannot be attributed to the same phenomenon. We specifically compare this with ASYH and  $\Delta H$  behavior in the vicinity of this peak's occurrence. In this region, we observe a positive response of 70% around 11:00 LT, near noon sector.

## 5 Discussion

From Figure 3, we observe significant variations in the results depending on the local time sector of each observatory during TW1 and TW2. More specifically, we note variations in the partial ring current (PRC) influence on the local response. When comparing the geomagnetic responses, we observe characteristic behaviors in  $\Delta H$  and consequently in  $H_I$ .

For GUI, JAI, and KAK, a decrease in  $\Delta H$  and consequently in  $H_I$  is observed during TW1, particularly at 14:07 UT (corresponding to 12:07, 19:07, and 23:07 LT, respectively). All three observatories exhibit a clear decreasing trend during TW1, with varying degrees of intensity. JAI shows the most pronounced decrease, while KAK displays the least pronounced. In contrast, TEO did not experience a decrease in  $\Delta H$  and  $H_I$ , but rather a smooth increase in their magnitudes during the sharp ASYH enhancement.

TW2 also shows notable differences among the observatories. At GUI,  $\Delta H$  maintains sustained minimum values throughout most of TW2, while JAI and KAK exhibit both decreasing and increasing  $\Delta H$  values.  $H_I$  shows an increasing trend at all three observatories until 23 UT (21, 04 and 08 LT). At TEO, after ASYH reaches its maximum at 07:07 LT, both  $\Delta H$  and  $H_I$  show a consistent decreasing trend throughout TW1 and TW2 until the final ASYH pulse at approximately 23:00-00:00 UT (16:00-17:00 LT).

An interesting observation is that during various periods of TW1 and TW2,  $H_I$  sometimes appears to behave oppositely to ASYH activity (e.g., during TW1 at JAI), while in other cases it follows ASYH fluctuations more closely (e.g., during TW1 at TEO). In some instances, these similar or opposite patterns are only present during ASYH pulses of activity.

To quantitatively confirm the contribution of the PRC to  $H_I$  and consequently to  $H_{PPEF}$ , we conducted two validation experiments focusing on TW1 (13:00-19:00 UT) and TW2 (19:00-01:00 UT), during which PRC activity occurs. The experiments were:

1. Analyze the correlation between  $H_I$  and ASYH.




Figure 4: Correlation Analysis ASYH and  $H_I$ , during the period 1300 - 1900 UT, which is consistent with the period of highest ASYH activity

2. Analyze the  $H_{PPEF}$  fluctuations distribution during MP, looking for the sectors where  $H_{PPEF}$  is more intense in general.

In the first experiment, we computed the Pearson correlation coefficient ( $\rho$ ) in two stages: first,  $\rho_1$  for ASYH vs  $H_I$ , and second,  $\rho_2$  for ASYH vs  $H_I$  after removing  $H_{PPEF}$  from  $H_I$  using a low-pass filter. The results for TW1 are shown in Figure 4, where red diamonds represent  $\rho_1$  (before removing  $H_{PPEF}$ ), and gray diamonds represent  $\rho_2$  (after removing  $H_{PPEF}$ ).

According to Makarov [2022], it is implied a strong anti-correlation between ASYH and SYMH indices. Therefore, a strongly positive or negative  $\rho$  coefficient can be interpreted as  $H_I$  (and consequently  $\Delta H$ ) being influenced by PRC activity, which is well described by the ASYH index. In this context, a strongly positive  $\rho$  implies a negative contribution to the local H component; in other words, the PRC may intensify the local geomagnetic response. Conversely, a strongly negative  $\rho$  suggests that the PRC may attenuate the local geomagnetic response.

Note that the two observatories showing strong correlation (or anti-correlation) are JAI and TEO. Interestingly, these were also the observatories closest to the dusk and dawn sectors, respectively, at the beginning of TW1. After removing  $H_{PPEF}$  fluctuations from  $H_I$ , the scattering increases at all observatories except GUI. In case of correlation during TW2, the coefficients are:  $\rho_{GUI} = 0.39$ ,  $\rho_{JAI} = 0.19$ ,  $\rho_{KAK} = -0.17$ , and  $\rho_{TEO} = -0.14$ . Given that ASYH activity is significantly weaker during TW2, this suggests that ASYH intensity is an important factor influencing  $\Delta H$ .

By measuring  $\rho$  during individual ASYH pulses, we find that the first pulse shows  $\rho$  values of [-0.81 (GUI), -0.91 (JAI), -0.70 (KAK), and 0.88 (TEO)]. For the second and third pulses, only JAI and TEO maintain correlations above 0.8 or below -0.8, while KAK shows values around 0.55. For the final ASYH pulse, only TEO exhibits a correlation slightly above 0.6 (or below -0.6). This supports the interpretation that  $H_{PPEF}$  magnitudes are strongly influenced by PRC activity, with a dependence on local time but also, the intensity of PRC activity can be another factor for this influence.

Following the idea of PRC influencing on  $H_{PPEF}$  fluctuation amplitudes, we carried out an analysis of  $H_{PPEF}$  distribution magnitudes, from march 17<sup>th</sup> to march 18<sup>th</sup>, shown in Figure 5. From top panels, we observe that  $H_{PPEF}$  magnitudes (blue bars) in each OB seems to approach to a Gaussian distribution (fitted with a red line). All OB present media  $\mu \sim 0$  nT and a very small skewness, seen at first glance. The standard deviation  $\sigma$  seen in each OB has a maximum difference goes from 5.2 to 8.15, being not so significant.

Of the four observatories, the KAK  $H_{PPEF}$  distribution is the narrowest and sharpest, characterized by the smallest standard deviation ( $\sigma$ ), the highest peak, and the shortest tails. This indicates that KAK exhibits the least dispersed data with the fewest extreme values and the most stable  $H_{PPEF}$  amplitudes throughout the 2-day period. The opposite extreme is observed at JAI, which has the flattest and widest distribution, consistent with the largest  $\sigma$ , indicating the greatest overall data variability. JAI also possesses the longest tails, suggesting a significantly higher likelihood of extreme  $H_{PPEF}$  values.

The observatories GUI and TEO exhibit intermediate characteristics. GUI shows the second-largest dispersion after JAI, indicating highly variable  $H_{PPEF}$  values. However, its tails are shorter than JAI's, meaning it has fewer extreme values than JAI despite its high variability. This high dispersion at GUI implies no strong systematic difference exists between the  $H_{PPEF}$  values in the specific time windows (TW1, TW2) and the entire 2-day period, a conclusion supported by the minor amplitude differences shown in Figure 3. In contrast, TEO's distribution is narrower than both GUI and JAI but exhibits a higher propensity for extreme values than both GUI and KAK. Consequently, the narrow distributions of KAK and TEO imply a more limited range of background  $H_{PPEF}$  amplitudes, making any deviations during the TW1 and TW2 windows more statistically significant relative to their low background variability.

Hence, we are interested in estimating the point at which  $H_{PPEF}$  can be considered as extreme, for each OB. We refer this point as *Threshold*. Thus, higher thresholds imply higher extreme values in general.

First, from the good symmetry in all distributions, we can consider an absolute value threshold. From this approach, we set each threshold as the 95% of each cumulative  $H_{PPEF}$  probability distribution, shown in bottom panels of Figure 5 (blue lines). Since we are considering  $-H_{PPEF}$ , we fit an accumulated half-Gaussian distribution (red lines). The threshold is indicated by the red dashed line for the 2 days period. We compare this threshold with the threshold of  $H_{PPEF}$  intensities within TW.

The highest couple of thresholds correspond to JAI (28.63 and 40.98 nT) followed by TEO (20.31 and 33.79). It is remarkable too how each threshold grew by considering only TW data. In JAI and TEO, the second threshold presents an increment of 43.13 and 66.37% respectively with respect to the first one. Meanwhile, in GUI and KAK, having significantly minor thresholds, the increment is about 35 and 45 % respectively. From here, we observe that the Threshold increment in KAK is not so different from JAI meanwhile TEO presents the highest threshold increment of the four OB. What this difference may indicate is how  $H_{PPEF}$  intensity is affected by the LT sector at the moment of the highest ASYH activity. This result is consistent with the results of  $\rho_{1,2}$  for these OB, which reinforces the idea of  $H_{PPEF}$  being influenced by PRC activity, indicating not just the impact of PRC activity but also on the proximity of JAI and TEO to one of the joint points of PRC - FAC during its highest activity.

Regarding the ionospheric response, we observe that the TEO and GUI regions (monitored by the UCOE and LPAL receivers) present a positive storm effect, while the JAI and KAK regions (monitored by the NPGJ and TSK2 receivers) present a negative effect. From Thomas and Venables [1966], Vijaya Lekshmi et al. [2011], it is understood that positive phases at mid-low latitudes are often initiated on the




Figure 5:  $H_{PPEF}$  distribution for each OB. Color dashed vertical lines represent the a threshold for  $H_{PPEF}$  to be considered as intense amplitudes

dayside during the main phase of the storm. The physical mechanism for this is complex and involves thermospheric winds,  $E \times B$  drift, and, most importantly for this study, the presence of eastward Prompt Penetration Electric Fields (PPEF). According to Vijaya Lekshmi et al. [2011], the influence of PPEF alone may not always induce large TEC enhancements; its effects are often combined with neutral wind circulation and  $E \times B$  drift to produce significant positive ionospheric storms.

The negative effects, on the other hand, are more related to regions that were on the nightside during the storm's main phase, which is the case for JAI and KAK in our event. These negative phases, however, typically manifest the following morning in local time. Studies like Fuller-Rowell et al. [1994], Prölss [1995] describe the mechanism responsible for these negative effects. It involves a storm-induced circulation that pushes molecular-rich air ( $[N_2]$ ,  $[O_2]$ ) from high latitudes toward lower latitudes. This increases the  $\frac{[N_2]}{[O]}$  ratio, which in turn enhances the loss rate of atomic oxygen ions ( $O^+$ ) through recombination. Although this circulation begins during the night, its effects peak the next morning when photo-ionization commences but is overwhelmed by the heightened recombination rate, resulting in an electron density depletion and a negative TEC storm.

## 6 conclusions

This project analyzed longitudinal magnetic fluctuations driven by Prompt Penetration Electric Fields (PPEF) during the intense St. Patrick's Day geomagnetic storm on March 17, 2015. To achieve this, we utilized geomagnetic data from ground-based observatories separated by 60–90° in longitude.

The study focused on isolating two key signatures: the magnetic component associated with the ionospheric current system ( $H_I$ ) and the fluctuations specifically linked to PPEF ( $H_{PPEF}$ ). The  $H_{PPEF}$  signal was reconstructed via high-pass filtering to retain fluctuations with periods less than four hours. Our analysis targeted the period of enhanced activity in the ASYH index, which reflects the asymmetric part of the magnetospheric ring current system and is linked to the Partial Ring Current (PRC).

We investigated the influence of the PRC on both  $H_I$  and  $H_{PPEF}$ . These observations were complemented by analyzing relative Vertical TEC variations, considering deviations greater than 30% from the 27-day median as significant ionospheric perturbations. We observed positive ionospheric storm effects at the TEO and GUI regions (55% and 70% TEC increase, respectively), while the JAI and KAK regions exhibited negative storm effects (70% and 50% TEC decrease). This pattern is consistent with the local time of each region during the storm's main phase onset.

By comparing  $H_I$  and  $H_{PPEF}$  with ASY-H activity across different local time sectors and integrating TEC observations, we reached the following conclusions:

- PRC activity is highly correlated with fluctuations in both  $H_I$  and  $H_{PPEF}$  during the most geoeffective period of the geomagnetic storm.
- The local geomagnetic response ( $\Delta H$ ) is either enhanced or attenuated depending on the observatory's proximity to the dawn or dusk sectors, where the PRC influence is strongest.
- High PRC activity significantly affects the magnitude of  $H_{PPEF}$ , particularly when an observatory is magnetically conjugate to the regions where the PRC connects to the ionosphere via Field-Aligned Currents (FACs).
- Conversely, when an observatory is located far from these PRC-FAC coupling regions (e.g., near the noon sector), the contribution to  $H_{PPEF}$  is less significant.
- The TEC enhancements at GUI and TEO are primarily driven by eastward PPEF combined with  $\mathbf{E} \times \mathbf{B}$  drift from neutral wind circulation.
- The TEC depletions at JAI and KAK are attributed to storm-induced changes in thermospheric composition and circulation. The distinct mechanisms observed between the dayside and nightside regions suggest the presence of a disturbed dynamo.

## 7 Future work

### A Processing magnetic data

It is crucial to eliminate regular variations from magnetic records to isolate magnetic contributions of interest. The main field  $H_0$  can be removed by modeling a monthly baseline. This baseline was approximated by computing an array of daily mode values for March 2015, following Gjerloev [2012]. These daily mode values were obtained from data between 00:00 and 03:00 LT ( $H_{0.3}$ ), then  $H_0$  was computed by a linear least squares fitting from these daily values. To exclude data from disturbed days for the fitting, we adapted the procedure outlined in van de Kamp [2013]. Specifically, we applied a moving hourly window to calculate the hourly interquartile range (IQR). Disturbed days were identified by setting a variation threshold where:

$$IQR_{med}^i > threshold = disturbed \quad (5)$$

Here,  $IQR_{med}^i$  is the median of each daily IQR values, so if more than 50% of IQR values for a given day, surpass a determined threshold, then that day is considered as disturbed day and depreciated for

the LS fitting.

The threshold was set, based on the criteria seen in Solari et al. [2017]. Basically, we analyze the monthly distribution of maximum IQR values ( $IQR_i^{max}$ ) within tri-hours IQR bins, by shifting another moving window and sorting those peaks, so  $IQR_1^{max} < IQR_2^{max} < \dots < IQR_n^{max}$ , in order to get a cumulative  $IQR^{max}$  distribution.

Hence, we have to compute the parameters of the General Pareto Distribution (GPD) Pickands [1975] which is a type of distribution used for modeling other distributions extreme values, and it is expressed as:

$$f(x) = \begin{cases} \frac{1}{\sigma} \left(1 + k \frac{x-u}{\sigma}\right)^{-1-\frac{1}{k}} & \text{if } k \neq 0 \\ \frac{1}{\sigma} \exp\left(-\frac{x-u}{\sigma}\right) & \text{if } k = 0 \end{cases} \quad (6)$$

where the three GPD parameters are shape  $k$ , scale  $\sigma$  and location  $u$  for the cumulative  $IQR_i^{max}$ . The last parameter,  $u$  is the one which can be considered as a threshold point [Hosking, 1990, Solari et al., 2017]. Then, for each  $IQR_i^{max}$ , we compute  $[k_i, \sigma_i, u_i]$ , then we apply a modified Anderson-Darling test. The threshold,  $u_t$ , is considered once the test gives the most optimum validation result.

To compute the diurnal variation, we identified what we refer to as local quiet days (LQD) for each station. Following the methodology in van de Kamp [2013], we considered the five days with the smallest daily IQR peaks as LQDs. We then computed the average of the  $H$  component data during these five quietest days:

$$\langle LDQ_H(t) \rangle = \frac{1}{5} \sum_{i=1}^5 H_i(t) - \langle H_i 0 : 3 \rangle \quad (7)$$

where we normalize the offset of each daily data by subtracting the average of the first four hours of magnetic data in local time, as indicated by the second term in Equation 7. Subsequently, we model the diurnal variation  $H_{SQ}$  by applying a low-pass filter to the Fourier Transform of the result from Equation 7, following the procedure in Equation (1) of van de Kamp [2013].

**Acknowledgements** The authors gratefully acknowledge the data and services provided by various institutions that made this work possible. Geomagnetic indices (ASY/SYM-H) were obtained from the International Service of Geomagnetic Indices (ISGI). Global geomagnetic data from the GUI, JAI, and KAK observatories were provided by INTERMAGNET, while local data from the TEO Observatory were sourced from LANCE-SciesMEX, REGMEX, and the Magnetic Service. GNSS data for the LPAL and TSK2 stations were accessed via the International GNSS Service (IGS), and data for the NPGJ station were provided by the NSF SAGE facility operated by EarthScope Consortium (under NSF award 1724509).

We extend our special thanks to the personnel of the Servicio de Geodesia Satelital, particularly Luis Salazar Tlaczani, and the TLALOCNet team for the maintenance of the UCOE station, data acquisition, and IT support. The GNSS data from this network, essential for TEC calculations, are supported by project PAPIIT IN111225 (E. Cabral-Cano).

Finally, this research was funded by the Universidad Nacional Autónoma de México (UNAM) and the Secretaría de Educación, Ciencia, Tecnología e Innovación (SECIHTI).

## References

- R. J. Strangeway C.T Russell, J. G. Luhman. *SPACE PHYSICS AN INTRODUCTION*. Cambridge university press. University Pringting House, Cambridge CB2 (BS, United Kingdom, 2016. ISBN 978-1-107-09882-4.
- Atsuhiko Nishida. Geomagnetic  $D_p$  2 fluctuations and associated magnetospheric phenomena. *Journal of Geophysical Research*, 73(5):1795–1803, March 1968a. doi: 10.1029/JA073i005p01795.
- Atsuhiko Nishida. Coherence of geomagnetic DP 2 fluctuations with interplanetary magnetic variations. *Journal of Geophysical Research*, 73(17):5549–5559, September 1968b. doi: 10.1029/JA073i017p05549.

- M. Blanc and A. D. Richmond. The ionospheric disturbance dynamo. *Journal of Geophysical Research*, 85(A4):1669–1686, April 1980. doi: 10.1029/JA085iA04p01669.
- Vytenis M. Vasyliunas. Mathematical models of magnetospheric convection and its coupling to the ionosphere. In B. M. McCormac, editor, *Particles and Fields in the Magnetosphere*, pages 60–71, Dordrecht, 1970. Springer Netherlands. ISBN 978-94-010-3284-1.
- R. K. Jaggi and Richard A. Wolf. Self-consistent calculation of the motion of a sheet of ions in the magnetosphere. *Journal of Geophysical Research*, 78:2852–2866, 1973. URL <https://api.semanticscholar.org/CorpusID:121315484>.
- Yong Wei, Biqiang Zhao, Guozhu Li, and Weixing Wan. Electric field penetration into earth’s ionosphere: a brief review for 2000–2013. *Science Bulletin*, 60(8):748–761, 2015. ISSN 2095-9273. doi: <https://doi.org/10.1007/s11434-015-0749-4>. URL <https://www.sciencedirect.com/science/article/pii/S2095927316304376>.
- M. C. Kelley, B. G. Fejer, and C. A. Gonzales. An explanation for anomalous equatorial ionospheric electric fields associated with a northward turning of the interplanetary magnetic field. *Geophysical Research Letters*, 6(4):301–304, April 1979. doi: 10.1029/GL006i004p00301.
- B. G. Fejer, M. F. Larsen, and D. T. Farley. Equatorial disturbance dynamo electric fields. *Geophysical Research Letters*, 10(7):537–540, 1983. doi: <https://doi.org/10.1029/GL010i007p00537>. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/GL010i007p00537>.
- Christine Amory-Mazaudier. Electric currents above saint-santin. i - data. *Journal of Geophysical Research*, 87:2459–2464, 04 1982. doi: 10.1029/JA087iA04p02459.
- Minh Le Huy and Christine Amory-Mazaudier. Magnetic signature of the ionospheric disturbance dynamo at equatorial latitudes: “ $D_{dyn}$ ”. *Journal of Geophysical Research (Space Physics)*, 110(A10): A10301, October 2005. doi: 10.1029/2004JA010578.
- Minh Le Huy and Christine Amory-Mazaudier. Planetary magnetic signature of the storm wind disturbance dynamo currents:  $D_{dyn}$ . *Journal of Geophysical Research (Space Physics)*, 113(A2): A02312, February 2008. doi: 10.1029/2007JA012686.
- K. Z. Zaka, A. T. Koba, P. Assamoi, O. K. Obrou, V. Doumbia, K. Boka, B. J. P. Adohi, and N. M. Mene. Latitudinal profile of the ionospheric disturbance dynamo magnetic signature: comparison with the DP2 magnetic disturbance. *Annales Geophysicae*, 27(9):3523–3536, September 2009. doi: 10.5194/angeo-27-3523-2009.
- J. Y. Liu, W. H. Yang, C. H. Lin, Y. I. Chen, and I. T. Lee. A statistical study on the characteristics of ionospheric storms in the equatorial ionization anomaly region: GPS-TEC observed over Taiwan. *Journal of Geophysical Research (Space Physics)*, 118(6):3856–3865, June 2013. doi: 10.1002/jgra.50366.
- W. Younas, C. Amory-Mazaudier, Majid Khan, and R. Fleury. Ionospheric and Magnetic Signatures of a Space Weather Event on 25-29 August 2018: CME and HSSWs. *Journal of Geophysical Research (Space Physics)*, 125(8):e27981, August 2020. doi: 10.1029/2020JA027981.
- J. A. Gonzalez-Esparza, V. De la Luz, P. Corona-Romero, J. C. Mejia-Ambriz, L. X. Gonzalez, M. A. Sergeeva, E. Romero-Hernandez, and E. Aguilar-Rodriguez. Mexican space weather service (sciesmex). *Space Weather*, 15(1):3–11, 2017. doi: <https://doi.org/10.1002/2016SW001496>. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016SW001496>.
- P. Corona-Romero, E. Sanchez-Garcia, M. Sergeeva, J.A. Gonzalez-Esparza, G. Cifuentes-Nava, E. Hernandez-Quintero, A. Caccavari, E. Aguilar-Rodriguez, J.C. Mejía-Ambriz, V. de-la-Luz, and L.X. Gonzalez. Kmex: the mexican geomagnetic k index. In *Proceedings of the II Pan-American Workshop on Geomagnetism*, pages 96–100, Basouras, Brazil, 2018.
- P. Corona-Romero, M. Sergeeva, J.A. Gonzalez-Esparza, G. Cifuentes-Nava, E. Hernandez-Quintero, and E. Caccavari, A. Sanchez-Garcia. Mexican geomagnetic k index. pages 1–6, 2017.

- E. Romero-Hernandez, J. A. Gonzalez-Esparza, M. Rodriguez-Martinez, M. A. Sergeeva, E. Aguilar-Rodriguez, J. C. Mejia-Ambriz, and V. De la Luz. Study of ionospheric disturbances over Mexico associated with transient space weather events. *Advances in Space Research*, 60(8):1838–1849, October 2017. doi: 10.1016/j.asr.2017.06.042.
- M. A. Sergeeva, O. A. Maltseva, J. A. Gonzalez-Esparza, V. De la Luz, and P. Corona-Romero. Features of TEC behaviour over the low-latitude North-American region during the period of medium solar activity. *Advances in Space Research*, 60(8):1594–1605, October 2017. doi: 10.1016/j.asr.2017.06.021.
- Maria Sergeeva, Aleksei Kalishin, Olga Maltseva, D. Blagoveshchensky, A. Gonzalez-Esparza, Pedro Corona-Romero, and Victor Gatica Acevedo. Possibility of signal reflection from the northern crest of eia: case study. pages 1–4, 03 2020. doi: 10.23919/EuCAP48036.2020.9135786.
- C.I. Castellanos-Velazco, P. Corona-Romero, J.A. González-Esparza, M.A. Sergeeva, A.L. Caccavari-Garza, and V.J. Gatica-Acevedo. Low latitude geomagnetic response associated with intense geomagnetic storms: Regional space weather in mexico. *Journal of Atmospheric and Solar-Terrestrial Physics*, 259:106237, 2024. ISSN 1364-6826. doi: <https://doi.org/10.1016/j.jastp.2024.106237>. URL <https://www.sciencedirect.com/science/article/pii/S1364682624000658>.
- Chin-Chun Wu, Kan Liou, Ronald P. Lepping, Lynn Hutting, Simon Plunkett, Russ A. Howard, and Dennis Socker. The first super geomagnetic storm of solar cycle 24: “The St. Patrick’s day event (17 March 2015)”. *Earth, Planets and Space*, 68(1):151, December 2016. doi: 10.1186/s40623-016-0525-y.
- INTERMAGNET. International real-time magnetic observatory network, 2021. URL <https://www.intermagnet.org/index-eng.php>.
- Henry Rishbeth and Owen K. Garriott. *Introduction to ionospheric physics*. 1969.
- D.J Knecht and . Shuman B.M. *Handboor of Geophysics and the Space Enviromnent, chaper 4, The Geomagnetic Field*. ed by A. S. Jursa, Boston: Air Force Geophysics Laborator, 1985.
- J. W. Gjerloev. The SuperMAG data processing technique. *Journal of Geophysical Research (Space Physics)*, 117(A9):A09213, September 2012. doi: 10.1029/2012JA017683.
- M. van de Kamp. Harmonic quiet-day curves as magnetometer baselines for ionospheric current analyses. *Geoscientific Instrumentation, Methods and Data Systems*, 2(2):289–304, November 2013. doi: 10.5194/gi-2-289-2013.
- C. Amory-Mazaudier, O. S. Bolaji, and V. Doubbia. On the historical origins of the CEJ, DP2, and Ddyn current systems and their roles in the predictions of ionospheric responses to geomagnetic storms at equatorial latitudes. *Journal of Geophysical Research (Space Physics)*, 122(7):7827–7833, July 2017. doi: 10.1002/2017JA024132.
- Yu. V. Yasyukevich, A. A. Mylnikova, and A. S. Polyakova. Estimating the total electron content absolute value from the GPS/GLONASS data. *Results in Physics*, 5:32–33, January 2015a. doi: 10.1016/j.rinp.2014.12.006.
- Yu. V. Yasyukevich, A. A. Mylnikova, V. E. Kunitsyn, and A. M. Padokhin. Influence of GPS/GLONASS differential code biases on the determination accuracy of the absolute total electron content in the ionosphere. *Geomagnetism and Aeronomy*, 55(6):763–769, November 2015b. doi: 10.1134/S001679321506016X.
- A. A. Mylnikova, Yu. V. Yasyukevich, V. E. Kunitsyn, and A. M. Padokhin. Variability of GPS/GLONASS differential code biases. *Results in Physics*, 5:9–10, January 2015. doi: 10.1016/j.rinp.2014.11.002.
- D. Vijaya Lekshmi, N. Balan, S. Tulasi Ram, and J. Y. Liu. Statistics of geomagnetic storms and ionospheric storms at low and mid latitudes in two solar cycles. *Journal of Geophysical Research (Space Physics)*, 116(A11):A11328, November 2011. doi: 10.1029/2011JA017042.
- Georgy Makarov. Geomagnetic indices ASY-H and SYM-H and their relation to interplanetary parameters Geomagnetic indices ASY-H and SYM-H and their relation to interplanetary parameters Geomagnetic indices ASY-H and SYM-H and their relation to interplanetary parameters. *Solar-Terrestrial Physics*, 8(4):36–43, December 2022. doi: 10.12737/stp-84202203.

- L. Thomas and F. H. Venables. The onset of the F-region disturbance at middle latitudes during magnetic storms. *Journal of Atmospheric and Terrestrial Physics*, 28(6-7):599–605, January 1966. doi: 10.1016/0021-9169(66)90073-0.
- T. J. Fuller-Rowell, M. V. Codrescu, R. J. Moffett, and S. Quegan. Response of the thermosphere and ionosphere to geomagnetic storms. *jgr*, 99(A3):3893–3914, March 1994. doi: 10.1029/93JA02015.
- G. W. Prölss. *Ionospheric F-region storms*, volume 2, pages 195–248. CRC Press, 1995. doi: 10.1201/9780203713297.
- Sebastián Solari, Marta Egüen, María José Polo, and Miguel A. Losada. Peaks over threshold (pot): A methodology for automatic threshold estimation using goodness of fit p-value. *Water Resources Research*, 53(4):2833–2849, 2017. doi: <https://doi.org/10.1002/2016WR019426>. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016WR019426>.
- James Pickands. Statistical Inference Using Extreme Order Statistics. *The Annals of Statistics*, 3(1):119 – 131, 1975. doi: 10.1214/aos/1176343003. URL <https://doi.org/10.1214/aos/1176343003>.
- J. R. M. Hosking. L-moments: Analysis and estimation of distributions using linear combinations of order statistics. *Journal of the Royal Statistical Society: Series B (Methodological)*, 52(1):105–124, 12 1990. ISSN 0035-9246. doi: 10.1111/j.2517-6161.1990.tb01775.x. URL <https://doi.org/10.1111/j.2517-6161.1990.tb01775.x>.