

# Bayesain Project

*Ammar Hasan*

*18 February 2019*

## Contents

<b>Introduction</b>	<b>2</b>
<b>Data Setup and Cleaning</b>	<b>2</b>
Data . . . . .	2
Data Exploration . . . . .	2
Correlation (Pairs Plot) . . . . .	2
Graphical Summaries . . . . .	3
Data Cleaning . . . . .	7
Indicators . . . . .	7
Weeks (Placebo) . . . . .	7
Sex . . . . .	7
<b>Models</b>	<b>7</b>
Linear Regression . . . . .	7
Basic Multiple . . . . .	7
Term Interactions . . . . .	7
Hierarchical . . . . .	7
Auto Regression . . . . .	7
AR(1) . . . . .	7
AR(2) . . . . .	7
Gaussain Process . . . . .	8
Modelling and Diagnostics . . . . .	8
Summaries . . . . .	8
<b>Conclusion</b>	<b>8</b>
<b>Appendix</b>	<b>8</b>
Abbreviations . . . . .	8
Code . . . . .	8
Data . . . . .	8

## Introduction

This report summaries the Bayesian analysis of the Reisby data-set. The Reisby data-set is based on a 5 week (first placebo) psychiatric study which investigates response of depressed patients to IMI. The Bayesian analysis will investigate how the drug affects depression.

## Data Setup and Cleaning

### Data

The data is loaded from a .Rdata file containing the Riesby data-set introduced in the introduction.

### Data Exploration


#### Correlation (Pairs Plot)

	hd	week	lnimi	lndmi	female	reactive_depression
hd	1.0000000	-0.3322674	-0.1165766	-0.3521130	-0.0624613	-0.0561811
week	-0.3322674	1.0000000	0.0357592	0.1239101	-0.0118326	0.0325663
lnimi	-0.1165766	0.0357592	1.0000000	0.2102979	0.0859277	-0.0365988
lndmi	-0.3521130	0.1239101	0.2102979	1.0000000	0.0945810	-0.1001591
female	-0.0624613	-0.0118326	0.0859277	0.0945810	1.0000000	0.1158473
reactive_depression	-0.0561811	0.0325663	-0.0365988	-0.1001591	0.1158473	1.0000000

The strongest correlations are negative and weak-moderate, and occur between Hamilton index with week and DMI, and rather obvious as the increase in blood concentration of the antidepressant would alleviate depression over weeks of treatment. Most other correlations are weak to very weak.


## Graphical Summaries

### Hamilton Scores




Most Hamilton scores are above 20 (i.e. moderate and severe depression dominates against mild and normal)

### Sex




The female test subjects are overwhelmingly higher than the males (nearly double!).

## Endogenous vs Reactive (Depression Type)



Most Depression cases in the test population are Endogenous (i.e. not a reaction to an environmental event).

## DMI and IMI Concentrations



Generally tight distribution \*especially around 25% to 75%) for both with little outliers with both distributions looking nearly identical sans the shift up with DMI. It seems that after being processed as DMI, the concentration of the antidepressant in the blood increases.

## Data Cleaning

Indicators

Weeks (Placebo)

Sex

## Models

### Linear Regression

Basic Multiple

Modelling and Diagnostics

Summaries

Term Interactions

Modelling and Diagnostics

Summaries

Not everyone was measured every week.

Hierarchical

Modelling and Diagnostics

Summaries

### Auto Regression

AR(1)

Modelling and Diagnostics

Summaries

AR(2)

Modelling and Diagnostics

Summaries

## Gaussain Process

## Modelling and Diagnostics

## Summaries

## Conclusion

## Appendix

### Abbreviations

IMI - antidepressant drug imipramine

DMI - desmethylimipramine (Processed IMI)

AR - Auto Regressive Models

AR(1) - Auto Regressive Models (1st Degree)

AR(2) - Auto Regressive Models (2nd Degree)

### Code

#### Data

#### Data Loading

```
load("Reisby.RData")
```

#### Correlation (Pairs Plot)

```
kable(cor(Reisby[, -1])) # no id
```

#### Hamilton Scores Histogram

```
hist(Reisby[,2], main = "Hamilton Index Scores Histogram",  
     xlab = "Hamilton Score")
```

#### Sex Histogram

```
# Replace x axis here with male/female axis  
  
hist(Reisby[,6], main = "Sex Histogram",  
     xlab = "Sex", xaxt = "n", ylim = c(0, 200))  
axis(1, at=0:1, labels=c("Male", "Female"))
```

#### Depression Type Histogram



```
hist(Reisby[,7], main = "Depression Type Histogram",
     xlab = "Depression Type", xaxt = "n", ylim = c(0, 150))
axis(1, at=0:1, labels=c("Endogenous", "Reactive"))
```

### DMI and IMI Boxplot

```
boxplot(Reisby[,4], Reisby[,5], main = "IMI and DMI Concentration Distributions",
        ylab = "Log Concentraion")
axis(1, at=0:2, labels=c("IMI", "IMI", "DMI")) # uses 3 labels due to strange bug
```