Appendix A

Source Codes

In this appendix, we have collected the source codes of a number of the most important numerical methods for fractional differential equations. This appendix is by no means meant to be comprehensive or complete in any sense: our goal is to provide the reader with a selection of generally applicable easy-to-use programs that should be helpful in a large number of potential applications.

Whereas most of the programs are written in FORTRAN (mainly, standard FORTRAN77), i.e. in a language for which compilers are generally available on almost every conceivable hardware environment, we have also included some Matlab [554], Mathematica [584] and R [555] codes for users who prefer these platforms.

A.1 The Adams-Bashforth-Moulton Method

We begin with the standard version of the Adams-Bashforth-Moulton method with a uniform mesh. The code is divided into a number of subroutines. The first of these computes the required weights.

Algorithm A.1. Computation of the weights of the method. This routine requires an external routine GAMMA (such as, e.g., the one provided in [146]) for the evaluation of Euler's Gamma function.

SUBROUTINE ABMW(ALPHA, APRED, ACORR, BCORR, CCORR, N, IERR)

С

C CALCULATION OF WEIGHTS FOR FRACTIONAL

ADAMS-BASHFORTH-MOULTON METHOD

C

```
INPUT VARIABLES
C
C
            - NUMBER OF GRID POINTS
C
      ALPHA - ORDER OF DIFFERENTIAL OPERATOR
C
С
      OUTPUT VARIABLES
C
            - PREDICTOR WEIGHTS
C
            - FUNDAMENTAL CORRECTOR WEIGHTS
      ACORR.
С
            - START-POINT CORRECTOR WEIGHTS
      BCORR
C
            - END-POINT CORRECTOR WEIGHT
      CCORR
С
      IERR
             - ERROR CODE
С
                0 : SUCCESSFUL EXIT
C
               12 : ALPHA < 0
С
      IMPLICIT NONE
C
                       N, IERR
      INTEGER
      DOUBLE PRECISION ALPHA
      DOUBLE PRECISION APRED(N), ACORR(N), BCORR(N), CCORR
C
      INTEGER
      DOUBLE PRECISION GMALP1, GMALP2
C
      DOUBLE PRECISION GAMMA
      EXTERNAL
                       GAMMA
C
      TERR = 0
      IF (ALPHA .LE. O.DO) IERR = 12
      IF (IERR .NE. O) RETURN
C
      GMALP1 = GAMMA(ALPHA + 1.DO)
      GMALP2 = GMALP1 * (ALPHA + 1.D0)
C
      CCORR = 1.DO / GMALP2
C
      DO 100 J = 1. N
         ACORR(J) = ((J+1)**(ALPHA+1.D0) + (J-1)**(ALPHA+1.D0)
                       - 2 * J**(ALPHA+1.DO)) * CCORR
         APRED(J) = (J**ALPHA - (J-1)**ALPHA) / GMALP1
         BCORR(J) = ((J-1)**(ALPHA+1.DO) - (J-1-ALPHA) * J**ALPHA)
                         * CCORR
 100
     CONTINUE
      RETURN
      END
```

The next piece of code demonstrates the application of the Adams-Bashforth-Moulton method, using the weights computed above, by means of a concrete example, namely the initial value problem

$$^{\mathsf{C}}D^{1.3}y(x) = x^{0.7}E_{1.1.7}(-x) + \exp(-2x) - [y(x)]^2, \quad y(0) = 1, \ y'(0) = -1,$$

where $E_{1,1.7}$ denotes the two-parameter Mittag-Leffler function with parameters 1 and 1.7. The exact solution of this equation is

$$y(x) = \exp(-x)$$
.

The following program solves this initial value problem with the Adams-Bashforth-Moulton method on the interval [0, 4]. The code required here for the evaluation of the Mittag-Leffler function that appears on the right-hand side of the differential equation, *i. e.* the routine ML2P, is given in Appendix A.4 below (see Algorithm A.8).

Algorithm A.2. Standard version of the Adams-Bashforth-Moulton method. The algorithm uses the routine ABMW from Algorithm A.1 to compute the weights of the method.

```
SUBROUTINE RHS(X, Y, F)
      IMPLICIT NONE
      DOUBLE PRECISION X, Y, F
      COMPUTE THE RIGHT-HAND SIDE OF THE DIFFERENTIAL EQUATION
C
C
      DOUBLE PRECISION ML2P
      EXTERNAL
                        ML2P
C
      F = X**0.7D0 * ML2P(1.D0, 1.7D0, -X) + EXP(-2.D0*X) - Y**2
      RETURN
      END
C
C
C
      DOUBLE PRECISION FUNCTION EXACT(X)
      IMPLICIT NONE
      DOUBLE PRECISION X
C
      COMPUTE THE EXACT SOLUTION
C
      OF THE DIFFERENTIAL EQUATION AT THE POINT X
C
      EXACT = EXP(-X)
      RETURN
      F.ND
```

C C SUBROUTINE ABMS (ALPHA, N, XMIN, XMAX, RHS, CORRIT, APRED, BPRED, ACORR, BCORR, CCORR, Y, YO, IERR) C FRACTIONAL ADAMS METHOD FOR SINGLE CAPUTO FDE OF ORDER ALPHA C IMPLICIT NONE C **EXTERNAL** RHS INTEGER N, CORRIT, IERR DOUBLE PRECISION ALPHA, XMIN, XMAX, APRED(N), BPRED(N), ACORR(N), BCORR(N), CCORR, YO(*), Y(N) INPUT VARIABLES (UNCHANGED ON OUTPUT) С ORDER OF DE, MUST BE > 0.0 ALPHA С NUMBER OF STEPS, MUST BE > 0 С XMIN LOWER BOUND OF INTERVAL WHERE SOLUTION IS SOUGHT С XMAX UPPER BOUND OF INTERVAL WHERE SOLUTION IS SOUGHT, С MUST BE > XMIN С RHS GIVEN SUBROUTINE THAT COMPUTES THE RHS OF THE DE С (SEE BELOW FOR DETAILS) С NUMBER OF REQUESTED CORRECTOR ITERATIONS С (MUST ALWAYS BE >= 1: С IT IS RECOMMENDED TO USE CORRIT = 1 С FOR ALPHA >= 1) С APRED, BPRED -PREDICTOR WEIGHTS AS PROVIDED BY WADAMS С ACORR, BCORR, CCORR - CORRECTOR WEIGHTS AS PROVIDED BY ABMW С INITIAL CONDITIONS: С YO(J) MEANS (J-1)ST DERIVATIVE OF Y AT XMIN С С OUTPUT VARIABLES C Y(J) DENOTES APPROXIMATE SOLUTION C AT POINT XMIN + J*(XMAX-XMIN)/N С ERROR FLAG: WILL BE O ON SUCCESSFUL TERMINATION С MUST HAVE THE FOLLOWING STRUCTURE: C THE SUBROUTINE RHS С SUBROUTINE RHS(X, Y, F) C DOUBLE PRECISION X, Y, F C INPUT VARIABLES: X, Y С OUTPUT VARIABLE: F = F(X,Y)C IT MUST BE DECLARED "EXTERNAL" IN THE CALLING SUBROUTINE. С C

DOUBLE PRECISION H, HALP, T, KFAC, WRK(3)

```
INTEGER.
                        J, K, M, CI
C
      TERR = 0
      IF (
               N .LE. 0)
                             IERR = 11
      IF ( ALPHA .LE. O.DO) IERR = 12
         (CORRIT .LE. 0)
                             IERR = 14
            XMAX .LE. XMIN) IERR = 15
      IF (IERR .NE. 0) RETURN
C
      Η
           = (XMAX - XMIN) / N
      HALP = H ** ALPHA
           = INT(ALPHA)
      IF (ALPHA .EQ. DBLE(M)) M = M - 1
C
      DO 1000 J = 1, N
         T = XMIN + J * H
C
С
         SET INITIAL CONDITION
         WRK(1) = YO(1)
         KFAC = 1.D0
         DO 50 K = 1. M
            KFAC = KFAC * K
            WRK(1) = WRK(1) + T**K / KFAC * YO(K+1)
 50
         CONTINUE
C
C
         PREDICTOR.
         CALL RHS(XMIN, YO(1), WRK(3))
         WRK(2) = WRK(1) + HALP * BPRED(J) * WRK(3)
         DO 70 K = 1, J-1
            CALL RHS(XMIN + (J-K)*H, Y(J-K), WRK(3))
            WRK(2) = WRK(2) + HALP * APRED(K) * WRK(3)
70
         CONTINUE
С
C
         CORRECTORS
         CALL RHS(XMIN, YO(1), WRK(3))
         WRK(1) = WRK(1) + HALP * BCORR(J) * WRK(3)
         DO 110 K = 1, J-1
            CALL RHS(XMIN + (J-K)*H, Y(J-K), WRK(3))
            WRK(1) = WRK(1) + HALP * ACORR(K) * WRK(3)
 110
         CONTINUE
         DO 120 CI = 1, CORRIT
            CALL RHS(T, WRK(2), WRK(3))
            WRK(2) = WRK(1) + HALP * CCORR * WRK(3)
 120
         CONTINUE
         Y(J) = WRK(2)
```

```
1000 CONTINUE
      RETURN
      END
C
С
     PROGRAM EXAMPL
      IMPLICIT NONE
С
      DOUBLE PRECISION YO(2), Y(40), XMIN, XMAX, ALPHA, X,
                       APRED(40), ACORR(40), BCORR(40), CCORR,
                       EXACT
      INTEGER
                       J, K, N, CORRIT, IERR
                       RHS, EXACT
      EXTERNAL
C
C
     DEFINE PARAMETERS OF EQUATION
      ALPHA = 1.3DO
      XMIN = O.DO
      XMAX = 4.DO
C
C
     DEFINE INITIAL CONDITIONS
      YO(1) = 1.D0
      YO(2) = -1.D0
C
C
     DEFINE PARAMETERS OF NUMERICAL METHOD
С
     MAXIMUM NUMBER OF GRID POINTS
      N = 40
C
      NUMBER OF CORRECTOR ITERATIONS ACCORDING TO RECOMMENDATION
      CORRIT = 1
C
C
      COMPUTE WEIGHTS
      CALL ABMW(ALPHA, APRED, ACORR, BCORR, CCORR, N, IERR)
      IF (IERR .NE. O) THEN
         WRITE(*,'(A,14)') 'ERROR IN COMPUTATION OF WEIGHTS: ', IERR
         STOP
      END IF
C
C
      SOLVE EQUATION NUMERICALLY WITH 10, 20, 40 GRID POINTS
      DO 100 J = 1, 3
         N = 5 * 2**J
         CALL ABMS (ALPHA, N, XMIN, XMAX, RHS, CORRIT,
                     APRED, APRED, ACORR, BCORR, CCORR, Y, YO, IERR)
         IF (IERR .NE. 0) THEN
            WRITE(*, '(A,I3,A,I3)') 'ERROR CODE', IERR,
```

C

C

C

C

Source Codes 341

```
' IN DIFFERENTIAL EQUATION SOLVER FOR N =', N
   &
       ELSE
                   '(A)') '=======;
          WRITE(*.
                   '(A,I3)') 'NUMBER OF GRID POINTS: ', N
          WRITE(*.
                   '(A)')
                       EXACT VALUE
   &
               , Х
                                      APPROX. VALUE
                                                          ERROR'
          DO 50 K = 1, N
              X = XMIN + (XMAX-XMIN) / DBLE(N) * DBLE(K)
             WRITE (*, '(F6.4,F13.6,F16.6,F12.6)')
                   X, EXACT(X), Y(K), EXACT(X) - Y(K)
   r
50
           CONTINUE
       END IF
100
    CONTINUE
    END
```

Next we provide the parallel version of the method as developed in [173]. It uses the OpenMP programming model [434] for the parallelization. The particular example implemented here solves the initial value problem

$${}^{\mathsf{C}}D^{\alpha}y(x) = -y(x), \qquad y(0) = 1, \quad y'(0) = 0,$$

on an interval [0, X]. The program should be called with three parameters, namely α (the order of the differential operator which must be in the interval (0, 2]; the second of the two initial conditions above is only used if $\alpha > 1$), the number of grid points, and X, the right end point of the interval of interest.

Algorithm A.3. OpenMP-based parallel version of the Adams-Bashforth-Moulton method. The algorithm uses the routine ABMW from Algorithm A.1 to compute the weights of the method.

```
SUBROUTINE ABMSP(ALPHA, N, XMIN, XMAX, RHS, CORRIT,
& APRED, BPRED, ACORR, BCORR, CCORR, Y, YO,
& WRK, IERR)

FRACTIONAL ADAMS METHOD FOR SINGLE CAPUTO FDE OF ORDER ALPHA
PARALLELIZED WITH OPENMP

IMPLICIT NONE

EXTERNAL RHS
INTEGER N, CORRIT, IERR, FNSHD
DOUBLE PRECISION ALPHA, XMIN, XMAX, APRED(N), BPRED(N),
```

```
ACORR(N), BCORR(N), CCORR, YO(*), Y(N),
     &
     &
                        WRK(N)
C
C
      INPUT VARIABLES (UNCHANGED ON OUTPUT)
C
                 ORDER OF DE. MUST BE > 0.0
C
                 NUMBER OF STEPS, MUST BE > 0
C
      XMIN
                 LOWER BOUND OF INTERVAL WHERE SOLUTION IS SOUGHT
С
                 UPPER BOUND OF INTERVAL WHERE SOLUTION IS SOUGHT,
      XMAX
C
                 MUST BE > XMIN
С
      R.HS
                 GIVEN SUBROUTINE THAT COMPUTES THE RHS OF THE DE
С
                  (SEE BELOW FOR DETAILS)
C
                 NUMBER OF REQUESTED CORRECTOR ITERATIONS
С
                  (MUST ALWAYS BE >= 1;
С
                  RECOMMENDATION: USE CORRIT = 1 FOR ALPHA >= 1)
С
     APRED, BPRED -
                     PREDICTOR WEIGHTS AS PROVIDED BY ABMW
С
      ACORR, BCORR, CCORR - CORRECTOR WEIGHTS AS PROVIDED BY ABMW
С
      Y0
                 INITIAL CONDITIONS;
С
                 YO(J) MEANS (J-1)ST DERIVATIVE OF Y AT XMIN
С
С
      OUTPUT VARIABLES
С
                 Y(J) DENOTES APPROXIMATE SOLUTION
С
                 AT POINT XMIN + J*(XMAX-XMIN)/N
С
      WRK
                 WORKSPACE
C
      IERR
                 ERROR FLAG: WILL BE O ON SUCCESSFUL TERMINATION
С
С
                              MUST HAVE THE FOLLOWING STRUCTURE:
      THE SUBROUTINE
                        RHS
С
            SUBROUTINE RHS(X, Y, F)
С
            DOUBLE PRECISION X, Y, F
С
            INPUT VARIABLES: X, Y
С
            OUTPUT VARIABLE: F = F(X,Y)
С
      IT MUST BE DECLARED "EXTERNAL" IN THE CALLING SUBROUTINE.
      DOUBLE PRECISION H, HALP, T, KFAC, ICND, PRD, CRR, F, WO
                        B, J, K, M, CI,
      INTEGER
      INTEGER
                        NBLKS. OMPNTH.
                        OMP_GET_NUM_THREADS, OMP_GET_THREAD_NUM
                        OMP_GET_NUM_THREADS, OMP_GET_THREAD_NUM
      EXTERNAL
C
      IERR = 0
               N .LE. 0)
                             IERR = 11
      IF ( ALPHA .LE. O.DO) IERR = 12
      IF (CORRIT .LE. 0)
                             IERR = 14
            XMAX .LE. XMIN) IERR = 15
      IF (IERR .NE. 0) RETURN
C
      Η
           = (XMAX - XMIN) / N
```

```
HALP = H ** ALPHA
           = INT(ALPHA)
      IF (ALPHA .EQ. DBLE(M)) M = M - 1
C
      CALL RHS(XMIN, YO(1), WO)
      WO = WO * HALP
      FNSHD = 0
C
C
! $0MP
         PARALLEL PRIVATE(I, J, K, ICND, KFAC, NBLKS,
                           T, PRD, CRR, CI, F, B)
! $0MP&
                   SHARED (OMPNTH, N, H, XMIN, YO, M, APRED,
! $0MP&
!$0MP&
                           ACORR, BCORR, CCORR, WRK, Y, CORRIT,
! $0MP&
                           HALP, FNSHD, WO)
C
      OMPNTH = OMP_GET_NUM_THREADS()
      NBLKS = CEILING(DBLE(N)/DBLE(OMPNTH))
C
      DO 1000 B = 0, NBLKS -1
C
! $0MP
         D0
C
         DO 2000 I = 1, OMPNTH
            J = B * OMPNTH + I
            IF (J .LE. N) THEN
               T = XMTN + .I * H
C
C
               SET INITIAL CONDITION
               ICND = YO(1)
               KFAC = 1.D0
               DO 50 K = 1, M
                   KFAC = KFAC * K
                   ICND = ICND + T**K / KFAC * YO(K+1)
 50
               CONTINUE
C
С
               PREDICTOR AND CORRECTOR: FIRST PART OF SUMMATION
               PRD = ICND + BPRED(J) * WO
               CRR = ICND + BCORR(J) * WO
               DO 70 K = J - B * OMPNTH, J - 1
                   PRD = PRD + APRED(K) * WRK(J-K)
                   CRR = CRR + ACORR(K) * WRK(J-K)
 70
               CONTINUE
               DO 80 K = I-1, 1, -1
                   DO WHILE (FNSHD .LT. J-K)
! $0MP
                      FLUSH (Y, WRK, FNSHD)
```

```
END DO
                   PRD = PRD + APRED(K) * WRK(J-K)
                   CRR = CRR + ACORR(K) * WRK(J-K)
 80
               CONTINUE
C
               CORRECTORS: PROPER CORRECTOR ITERATIONS
               DO 120 CI = 1, CORRIT
                   CALL RHS(T, PRD, F)
                   PRD = CRR + HALP * CCORR * F
 120
               CONTINUE
С
               Y(J) = PRD
               CALL RHS(T, Y(J), WRK(J))
               WRK(J) = WRK(J) * HALP
               FNSHD = J
! $0MP
               FLUSH (Y, WRK, FNSHD)
C
         ENDIF
2000
         CONTINUE
! $0MP
         END DO
 1000 CONTINUE
!$OMP END PARALLEL
      RETURN
      F.ND
C
С
      SUBROUTINE RHS (X, Y, F)
      IMPLICIT NONE
      DOUBLE PRECISION X, Y, F
      F = -1.D0 * Y
      RETURN
      END
C
С
C
      EXAMPLE MAIN PROGRAM
С
      PROGRAM PARFDE
C
      IMPLICIT NONE
C
      INTEGER MAXPTS
      PARAMETER (MAXPTS=1000000)
      INTEGER IARGC
```

```
C
      DOUBLE PRECISION YO(2), Y(MAXPTS), XMIN, XMAX, ALPHA, X,
                        APRED (MAXPTS), ACORR (MAXPTS), BCORR (MAXPTS),
     &
                        CCORR, WRK (MAXPTS)
      INTEGER
                        K. N. CORRIT, IERR, NPTS
      EXTERNAL
                        RHS, IARGC, GETARG
      CHARACTER *72
                        ARGV
C
      N = IARGC()
      IF (N .NE. 3) THEN
         WRITE (*, '(A)') 'WRONG NUMBER OF ARGUMENTS.'
         WRITE (*. '(A)') 'USAGE: PARFDE ALPHA NPTS XMAX'
         STOP
      ENDIF
C
      CALL GETARG (1, ARGV)
      READ (ARGV, *) ALPHA
      CALL GETARG (2, ARGV)
      READ (ARGV, *) NPTS
      IF (NPTS .GT. MAXPTS) THEN
         WRITE (*, '(A,I,A,I,A)') 'TOO MANY POINTS: ', NPTS,
                            '(MAX: ', MAXPTS, ')'
         STOP
      ENDIF
      CALL GETARG (3, ARGV)
      READ (ARGV, *) XMAX
C
C
      DEFINE PARAMETERS OF EQUATION
      XMIN = O.DO
C
C
      DEFINE INITIAL CONDITIONS
      YO(1) = 1.D0
      YO(2) = 0.D0
C
C
      DEFINE PARAMETERS OF NUMERICAL METHOD
      CORRIT = 1
C
C
      COMPUTE WEIGHTS
      CALL ABMW(ALPHA, APRED, ACORR, BCORR, CCORR, NPTS, IERR)
      IF (IERR .NE. O) THEN
         WRITE (*, '(A, I4)') 'ERROR IN COMPUTATION OF WEIGHTS: ',
     &
                              IERR
         STOP
      END IF
C
C
      SOLVE EQUATION NUMERICALLY
```

```
CALL ABMSP (ALPHA, NPTS, XMIN, XMAX, RHS, CORRIT, APRED,
                 APRED, ACORR, BCORR, CCORR, Y, YO, WRK, IERR)
      IF (IERR .NE. 0) THEN
                  '(A,I3,A,I3)') 'ERROR CODE', IERR,
                   ' IN DIFFERENTIAL EQUATION SOLVER FOR NPTS = ',
    &
                   NPTS
      ELSE
         WRITE(*, '(A,F14.6)') 'ALPHA = ', ALPHA
        WRITE(*, '(A,I14)')
                               'NPTS
                                     = '. NPTS
                 '(A,F14.6)') 'XMAX
                                      = ', XMAX
                 '(A)') '======;
         WRITE(*, '(A)')
                                    APPROX. VALUE?
    Хr.
                         X
        DO 50 K = 1, NPTS
            X = XMIN + (XMAX-XMIN) / DBLE(NPTS) * DBLE(K)
            WRITE (*, '(F16.8,F19.12)') X, Y(K)
50
         CONTINUE
      END IF
 100
     CONTINUE
C
     END
```

A.2 Lubich's Fractional Backward Differentiation Formulas

For the fractional backward differentiation formulas introduced by Lubich, we here list the Matlab implementations due to Weilbeer [576]. This implementation consists of three separate routines. The first of these routines computes the solution at the starting points of the grid.

Algorithm A.4. Matlab implementation of Lubich's fractional BDF: Starting points.

```
function[Y] = startpoints(C,S,a,y0,alpha,T)
 STARTPOINTS solves the nonlinear equation system for the first
%
    a values of the fractional differential equation
%
%
    (1)
           D^{\alpha} = f(t,y(t))
%
%
   using a simple Newton method with a start solution corresponding
%
    to the initial condition of (1).
%
%
    Input Data:
%
```

```
%
    C
         - Convolution weights vector
%
    S
         - Starting weights matrix
%
         - number of start weights
%
    alpha
           - Order of FDE alpha \in (0,1)
%
         - mesh
%
    y0
          - Initial conditions
%
%
    Output Data:
%
%
         - Solution at a-1 points
%
%
    Note:
%
%
    This is not a stand-alone function but a subroutine of FBDFP.m
h = T(2)-T(1);
Y = y0(1).*ones(a-1,1);
IC= fliplr(y0); % Initial conditions polynomial
syms y t;
% right-hand side diff w.r.t. y
fj = inline(char(diff(rhs(t,y,alpha),y)),'t','y');
% right-hand side
ff = inline(char(rhs(t,y,alpha)),'t','y');
B = 1/h^{alpha*}(rot90(hankel(fliplr(C(1:a)))) + S(1:a,1:a));
% Evaluation of the initial conditions polynomial
P = polyval(IC,T(2:a));
B = B(2:a,2:a);
Pp = B*P';
er = 1;
while er > 1e-8
   for k=1:a-1
      Fj(k) = fj(T(k+1),Y(k));
      Ff(k) = ff(T(k+1),Y(k))+Pp(k);
   X = (B - diag(Fj)) \setminus (B*Y - Ff');
   er = abs(max(X));
   Y = Y - X;
end
```

The second routine is devoted to the construction of the matrix that contains the weights of the method. Recall that a linear system of equations needs to be solved in this context; the parameter c of the subroutine allows the user to choose between various alternative numerical methods for the solution of this linear system.

Algorithm A.5. Matlab implementation of Lubich's fractional BDF: Computation of weights.

```
function[C, S, a, r] = ltsmp(N,alpha,p,c)
% LTSMP(N,alpha,p,c)
%
    LTSMP creates the p-th order lower triangular strip matrix
%
    corresponding to the discrete approximation of the fractional
    differential or integral operator D^alpha at N nodes.
%
%
    Input Data:
%
%
           - number of nodes of the discretization.
%
    alpha
          - order of the fractional integration / differentiation.
%
             alpha > 0 means integration,
%
             alpha < 0 means differentiation.
%
             abs(alpha) < 1.
%
   p

    order of approximation

%
             (between 1 and 6 for stability reasons).
%
    С
           - choice of numerical method for the
%
             calculation of the starting weights.
%
                   - solution of the equation system
%
                          using lu decomposition
%
        'lui'
                   - computation of the inverse using
%
                          lu decomposition of wcoeff followed by
%
                          matrix multiplication A^(-1)*rhs
%
        'gmres'
                   - gmres solution of the equation system
%
        'gmresi'
                   - inverse using gmres ...
%
        'gmresh'
                   - gmres solution of the equation system using
%
                          Householder orthogonalization
%
        'qr'
                   - solution of the equation system using
%
                          QR decomposition
%
        'svd'
                   - solution of the equation system using SVD
%
        'bpi'
                     inverse using Bjoerck-Pereyra Algorithm
%
                          (alpha has to be a unit fraction
%
                          for this choice)
%
%
    Output Data:
%
%
          - Vector with N convolution weights
```

```
%
               (omega_0,omega_1,...,omega_N-1)
%
    <S1>
           - Matrix with N*a starting weights:
%
              [w_0,1]
                       w_{0,2}
                               w_0,3 ... w_0,a ]
%
              [w_1, 1]
                       w_1,1
                               w_1,2 ... w_1,a ]
%
                                  ٦
%
              [ w_N-1,1
                                       w_N-1,a ]
%
           - Number of starting weights for each n.
    <a>>
%
           - average residual of the start weight computation
    <r>
%
              over all right hand sides.
% Error handling.
if nargin ~= 4
   error('Wrong number of input arguments.')
end
% Calculating length(a) and q (=gamma).
cnt = 1;
    = 0;
q
for k=0:p-1
   for l=0:ceil(1/abs(alpha)).*(p-1)
      qt = k+l*abs(alpha); % potential gamma value.
      de = find(chop(q,10)==chop(qt,10)); % Avoid double entries.
      if qt \le p-1 \& isempty(de) == 1
 cnt = cnt+1;
 q(cnt) = qt;
      end
   end
end
% Sort feasible gamma entries in increasing order.
q = sort(q);
[nil, a] = size(q);
                             % Counting number of starting weights.
% Convolution weights (via automatic differentiation).
U = zeros(6,N);
                [ 1
U(1:6,1:7) =
                          -1
                                0
                                       0
                                              0
                                                     0
                                                          0;
                 3/2
                          -2
                               1/2
                                       0
                                              0
                                                     0
                                                          0:
                11/6
                          -3
                               3/2
                                     -1/3
                                              0
                                                     0
                                                          0;
                25/12
                          -4
                                3
                                     -4/3
                                             1/4
                                                     0
                                                          0;
                                             5/4
                                                  -1/5
               137/60
                         -5
                                5
                                    -10/3
                                                          0;
               147/60
                         -6
                              15/2
                                    -20/3
                                            15/4
                                                   -6/5
                                                         1/6];
C(1) = U(p,1)^(-alpha);
for s=1:N-1
   tmp = zeros(1,p+1);
   if s < p+2
```

```
for v=0:s-1
         tmp(v+1) = (alpha*(v-s)-v)*C(v+1)*U(p,s-v+1);
      and
      C(s+1) = 1/(s*U(p,1)) * sum(tmp);
   else
      for v=s-(p+1):s-1
         tmp(v-(s-(p+1))+1) = (alpha*(v-s)-v)*C(v+1)*U(p,s-v+1);
      C(s+1) = 1/(s*U(p,1)) * sum(tmp);
   end
end
% Starting weights
if p = 1
            % if p=1 no additional starting weights are needed
   % Exact fractional integrals
   s(1,1:a) = gamma(1+q(1,1:a))./gamma(1+q(1,1:a)+alpha);
  % Right-hand sides
  for k=1:a
      T(1:N) = (1:N).^q(1,k);
      L(1:N-1) = (1:N-1).^{(q(1,k)+alpha)};
      Y = conv(C,T);
      rhs(2:N,k) = s(1,k)'*L(1:N-1)' - Y(1:N-1)';
   end
  % Coefficient Matrix
   for k=1:a
      Wcoeff(k,:) = (0:a-1).^q(1,k);
   end
   switch lower(c)
      case 'lu'
      % 1) Solution of the linear equation systems
           using the \ operator
      S(1:N,:) = (Wcoeff\rhs(1:N,:)')';
      case 'lui'
      \% 2) Solution of the linear equation systems
           using the inverse
      Wcoeffinv = inv(Wcoeff);
      S(1:N,:) = (Wcoeffinv*rhs(1:N,:)')';
      case 'gmres'
      % 3) Solution using gmres at each step
```

```
%
     on the actual system.
for k=1:N
   [S(:,k), flag,relres,iter]
       = gmres(Wcoeff,rhs(k,:)',a+1,1e-16,1);
end
S = S':
case 'gmresi'
% 4) Computing the inverse of Wcoeff using gmres
for k=1:a
   unitVector = zeros(a,1);
   unitVector(k,1) = 1;
   [Wcoeffinv(:,k), flag]
      = gmres(Wcoeff,unitVector,a+1,1e-16,1);
end
for k=1:N
   S(k,:) = (Wcoeffinv*rhs(k,:)')';
end
case 'gmresh'
% 5) Solution using gmreshouseholder at each step
     on the actual system.
for k=1:N
   S(:,k)
      = gmresh(Wcoeff,rhs(k,:)',zeros(a,1),1e-32,a);
end
S = S';
case 'qr'
% 6) Solution using QR decomposition.
[Q R] = qr(Wcoeff);
S(1:N,:) = (R\setminus(Q'*rhs(1:N,:)'))';
case 'svd'
% 7) Solution using SVD.
S = zeros(N,a);
[Uv Sv Vv] = svd(Wcoeff);
for 1=1:N
   for k=1:a
      S(1,:) = S(1,:)
                 + (Uv(:,k)'*rhs(1,:)'*Vv(:,k)./Sv(k,k))';
   end
end
case 'bpi'
\% 8) Calculating the inverse using the Bjoerck-Pereyra
```

```
algorithm (alpha has to be a unit fraction)
      [num, num2] = rat(alpha);
      if abs(num) \sim 1
         fprintf(1,
             '\n alpha is not a unit fraction. Cannot use bpi\n');
      else
         X = (0:a-1).^(abs(alpha));
         % Calculating the inverse
         Wcoeffinv = bpif(X,'p');
         % Solving for the N right-hand sides
         S(1:N,:) = (Wcoeffinv*rhs(1:N,:)')';
      end
      otherwise
      fprintf(1,
         '\n Wrong choice for parameter c.\n');
   end
   % residual tester
   Residual(1:N,1) = norm(Wcoeff*S(1:N,:)'- rhs(1:N,:)');
   r = sum(abs(Residual))/length(Residual);
else
   a = 1;
   S = zeros(N,1);
   r = 0;
end
fprintf(1,'\n The average residual is %g.\n',r);
```

Finally we give the main part of the algorithm that actually computes the required approximate solutions, using the two auxiliary routines listed above.

Algorithm A.6. Matlab implementation of Lubich's fractional BDF: Main part.

```
function[Y] = fbdfp(N,alpha,y0,b,p,in,ie,c)

% FBDFP(N,alpha,y0,b,p,in,ie,c)

% Fractional Backward Difference Formula of order p.

% FBDFP solves the fractional order differential equation (FDE)

%

(1) D^alpha * y(t) = f(t,y(t)) ,alpha \in (0,1)

%

at N equispaced mesh points on the interval [0,b].
```

```
The method has a theoretical convergence order of p. The
    parameters 'in' (iteration number) and 'ie' (iteration error)
    determine the maximal number of iterations or the minimal
%
    change of two consecutive iterations at each mesh points.
%
    The parameter c determines the numerical method used to generate
%
    the discretized fractional differential operator D^alpha.
%
    The right-hand side of (1) has to be stored in an external
%
    matlab function in the file rhs.m.
%
%
    Input Data:
%
%
           - Number of mesh points.
%
    alpha
          - Order of FDE alpha \in (0,1).
%
           - Right interval end (0 assumed to be left start).
    b
%
    v0
           - Starting values y^(k)(0) = c_k as vector
$
             y0 = [c_0, c_1, ..., c_k].
%
           - desired convergence order (p = 1-6).
   р
%
   in
          - Maximal number of iteration at each mesh point.
%
   ie
           - Minimal change of two consecutive iteration steps.
%
    С
           - choice of numerical method for the calculation
%
             of the starting weights
%
%
          - externally stored right-hand side f(t,y(t)) of FDE.
%
%
    Output Data:
%
%
    Υ
           - Solution at the N mesh points.
%
%
    See also FBDFPS, LTSMP, LTSMPS, RHS, STARTPOINTS.
if nargin ~= 8
   error('Wrong number of input arguments.')
end
Y
       = zeros(N,1);
                                        % Solution.
Yt
       = zeros(N,1);
                                        % Auxiliary variables.
Z
       = zeros(N,1);
                                        % Initial Conditions.
Y(1,1) = y0(1);
IC
       = fliplr(y0);
T = linspace(0,b,N);
                                        % Equidistant mesh points.
h = T(2)-T(1);
                                        % Step size.
[C S a] = ltsmp(N,-alpha,p,c);
                                        % Calculating lower
                                        % triangular strip matrix.
```

```
if p = 1
  Sp = startpoints(C,S,a,y0,alpha,T); % Startpoints calculation.
  Y(2:a) = Sp(1:a-1);
                                        % Setting start values.
end
for k=1+a:N
  K = zeros(1,k);
  Ym = Y(1:k);
   ICm(1:k) = polyval(IC,T(1:k));
  K(1:k) = C(k:-1:1);
  K(1:a) = K(1:a) + S(k,1:a);
  St = 1/C(1).*K*Ym;
  A = 1/C(1).*K*ICm';
  % First guess using previous solution point
  Z(k) = (h^{(alpha)})/C(1)*rhs(T(k),Y(k-1),alpha) - St + A;
  er = 5;
                                 % auxiliary variables for iteration.
   cnt = 1;
  while er >= ie & cnt < in
                                 % Fixed point iteration.
      Yt(k) = (h^(alpha))/C(1)*rhs(T(k),Z(k),alpha) - St + A;
            = abs(Yt(k)-Z(k));
      Z(k)
            = Yt(k);
      cnt
            = cnt+1;
   end
  Y(k) = Z(k);
   if imag(Y(k)) = 0
                                 % termination criterion.
      Y(k:N) = 0;
      break;
   end
```

end

A.3 Time-fractional Diffusion Equations

Our last differential equation solver is a finite difference method for a timefractional diffusion equation

$${}^{\mathtt{C}}D_t^{\alpha}y(x,t) + \Phi(x,t)\frac{\partial^2}{\partial x^2}y(x,t) = f(x,t), \qquad x \in [a,b], \quad t \in [0,T],$$

with $0 < \alpha < 1$, the initial condition

$$y(x,0) = y_0(x)$$

and mixed Dirichlet-Neumann boundary conditions

$$g_1(t)y(a,t) + h_1(t)\frac{\partial}{\partial x}y(a,t) = r_1(t),$$

$$g_2(t)y(b,t) + h_2(t)\frac{\partial}{\partial x}y(b,t) = r_2(t)$$

as discussed in Subsection 3.2.2. This algorithm is given in a form suitable for direct use within the Mathematica software package [584]. It is based on the approximation developed in Eqs. (2.1.10) and (2.1.11) for the fractional derivative with respect to t, using a stepsize of $\Delta t = T/n_t$, and on a standard centered difference approximation with step size $\Delta x = (b-a)/n_x$ for the second-order derivative with respect to x, where n_t and n_x are preassigned positive integers.

The algorithm requires that $\Phi(a,t) \neq 0$ and $\Phi(b,t) \neq 0$ for all $t \in [0,T]$. The algorithm computes (n_x+1) -dimensional vectors $\mathbf{u}[0]$, $\mathbf{u}[1], \ldots, \mathbf{u}[n_t]$ (in Mathematica notation) with the interpretation that the component $\mathbf{u}[k][[j]]$ is the approximation of the exact solution $y((j-1)\cdot\Delta_x,k\cdot\Delta_t)$ where $k\in\{0,1,\ldots n_t\}$ and $j\in\{1,2,\ldots,n_x+1\}$.

Algorithm A.7. Mathematica implementation of a backward differentiation method for the solution of the time-fractional diffusion equation in one space dimension.

- (* Load package for handling of tridiagonal linear systems *)
 << LinearAlgebra'Tridiagonal'</pre>
- (* Initialize parameters of the equation (given data) *)

```
(* boundaries of intervals *)
(* values are to be understood as examples *)
a = 0;
b = 1;
T = 1;
(* functions in boundary conditions *)
(* values are to be understood as examples *)
r1[t_] := 0;
r2[t_{-}] := 1;
g1[t_] := 1;
g2[t_] := 1;
h1[t_] := 0;
h2[t_] := 0;
(* function in initial condition *)
(* value is to be understood as example *)
u0[x_] := x;
(* functions in the differential equation itself *)
(* values are to be understood as examples *)
phi[x_{,}t_{]} := -1;
f[x_{-}, t_{-}] := 0;
(* order of fractional differential operator with respect to time *)
alpha = 0.5;
Clear[weight];
(* initialize parameters of the numerical scheme:
   number of steps in coordinate directions *)
(* values for nx and nt are to be understood as examples *)
nx = 67;
nt = 80:
deltax = (b - a)/nx;
deltat = T/nt;
(* auxiliaries *)
ga = Gamma[2 - alpha];
(* define weights of fractional BDF *)
weight[j_, k_] := weight[j, k] = Which[j == 0, 1,
    j == k, k^{-1}(-alpha) (1 - alpha - k) + (k - 1)^{1} - alpha),
    True, (j - 1)^(1 - alpha) -
     2 j^{(1 - alpha)} + (j + 1)^{(1 - alpha)}/ga
```

```
(* main part of algorithm *)
(* assign initial values *)
u[0] = Table[u0[a + (j - 1) deltax], {j, 1, nx + 1}];
(* time step loop *)
Do[
 tk = k deltat;
 (* construction of tridiagonal matrix of linear system
    for each time step *)
 lower =
 Append[Table[phi[a + j deltax, tk], {j, 1, nx - 1}],
           -h2[tk]] / deltax^2;
main = Append[
   Prepend[Table[-2 phi[a + j deltax, tk] / deltax^2 +
      weight[0, k] / deltat^alpha, \{j, 1, nx - 1\}],
    g1[tk] / deltax +
     h1[tk] (-1/deltax^2 + 1/(2 ga deltat^alpha phi[a, tk] ))],
   g2[tk] / deltax +
    h2[tk] (1/deltax^2 - 1/(2 ga deltat^alpha phi[b, tk] ))];
upper = Prepend[Table[phi[a + j deltax, tk], {j, 1, nx - 1}],
    h1[tk]]/deltax^2;
 (* construction of vector on right-hand side of system *)
 rhs = Prepend[
   Append [
    Table[f[a + j deltax, tk]
           + u[0][[j]] (1 - alpha) /ga/tk^alpha
           - Sum[weight[mu, k] u[k - mu][[j]], {mu, 1, k}] /
                 deltat^alpha, {j, 2, nx}],
    r2[tk]/deltax -
     0.5 h2[tk] / phi[b, tk] (
       f[b, tk] + u0[b] tk^(-alpha) (1 - alpha)/ga
        - Sum[weight[mu, k] u[k - mu][[nx + 1]], {mu, 1, k}]/
         deltat^alpha)],
    r1[tk]/deltax +
    0.5 h1[tk] / phi[a, tk] (
      f[a, tk] + u0[a] tk ^(-alpha) (1 - alpha)/ga
       - Sum [weight[mu, k] u[k - mu][[1]], {mu, 1, k}]
           / deltat^alpha)];
 (* approximate solution at present time step is obtained
    by solving this tridiagonal system *)
```

```
u[k] = TridiagonalSolve[lower, main, upper, rhs]
, {k, 1, nt}]
```

A.4 Computation of the Mittag-Leffler Function

In this section, we give a fast routine for the computation of the twoparameter Mittag-Leffler function $E_{\alpha,\beta}(x)$ for $\alpha \geq 0$, $\beta > 0$ and real values of x. The routine has been developed to run fast; its accuracy is very high only for $x \leq 1$. If highly accurate results are required for x > 1 or for complex arguments x then we recommend to use the routine of Gorenflo et al. [247, 248]. A Matlab version of such a code is available [455].

Algorithm A.8. FORTRAN77 implementation of a routine for the fast evaluation of two-parameter Mittag-Leffler functions. This routine requires an external routine GAMMA (such as, e.g., the one provided in [146]) for the evaluation of Euler's Gamma function.

```
DOUBLE PRECISION FUNCTION ML2P(ALPHA, BETA, X)
C
      IMPLICIT NONE
      DOUBLE PRECISION ALPHA, BETA, X
C
      FAST COMPUTATION OF TWO-PARAMETER MITTAG-LEFFLER FUNCTION
C
С
C
      PARAMETERS
C
      ALPHA - INPUT - FIRST PARAMETER OF MITTAG-LEFFLER FUNCTION
С
              INPUT - SECOND PARAMETER OF MITTAG-LEFFLER FUNCTION
C
            - INPUT - ARGUMENT OF THE MITTAG-LEFFLER FUNCTION
C
      INTEGER
                       K, IERR
      DOUBLE PRECISION TMP, NEW, GMARG, GAMMA
                       GAMMA
      EXTERNAL
C
      TMP = 1.DO / GAMMA(BETA)
      K = 1
 100
      GMARG = ALPHA * K + BETA
            X**K / GAMMA(GMARG)
      NEW =
      TMP = TMP + NEW
      K = K + 1
      IF (GMARG .LT. 5.DO) GOTO 100
      IF (K .LT. 100
                      .AND.
                             ABS(NEW) * 1.D16 .GT. TMP) GOTO 100
```

```
C ML2P = TMP
C RETURN
END
```

A.5 Monte Carlo Simulation of CTRW

We conclude this appendix with two implementations (in R and in Matlab) of a Monte Carlo program to simulate CTRWs according to the algorithm described in section 7.3. The program below generates and plots a single realization of a CTRW with a given number of jumps and durations. Even if this is a very simple algorithm, it consists of three parts. The first part is the generator of independent and indentically distributed Mittag-Leffler deviates according to equation 7.3.2. Then, Lévy α -stable deviates are generated following equation 7.3.1. Finally, cumulative sums give the position coordinates and the epochs and positions are plotted as a function of the epochs. This routine can be easily modified with suitable external cycles to generate many realizations up to a given time t and estimate the probability density $f_{X(t)}(x,t)$ from the histogram of realized positions X(t). This was explicitly done in reference [231].

Algorithm A.9. Matlab implementation for the Monte Carlo simulation of CTRW.

```
%Plot of a single CTRW realization

%Generation of Mittag-Leffler deviates
%See Fulger, Scalas, Germano 2008 and references therein

n=100; %number of points
gammat=1; %scale parameter
beta=0.99; %ML parameter

u1=rand(n,1); %uniform deviates
v1=rand(n,1); %uniform deviates
%Generation of symmetric alpha stable deviates

tau=-gammat*log(u1).*(sin(beta*pi).
/tan(beta*pi*v1)-cos(beta*pi)).^(1/beta);
```

```
gammax=1; %scale parameter
alpha=1.95; %Levy parameter
u2=rand(n,1); %uniform deviates
v2=rand(n,1); %uniform deviates
phi=pi*(v2-0.5);
xi=gammax*(sin(alpha*phi)./cos(phi)).
*(-log(u2).*cos(phi)./cos((1-alpha)*phi)).^(1-1/alpha);
%Random walk
x=cumsum(xi');
x=[0 x];
%Epochs
t=cumsum(tau');
t=[0 t];
stairs(t,x) %plots ctrw
Algorithm A.10. R implementation for the Monte Carlo simulation of
CTRW.
# Plot of a single CTRW realization
# Generation of Mittag-Leffler deviates
# See Fulger, Scalas, Germano 2008 and references therein
n <- 10000 #number of points
gammat <- 1 #scale parameter
beta <- 0.95 #ML parameter
u1 <- runif(n) #uniform deviates
v1 <- runif(n) #uniform deviates
tau \leftarrow -gammat * log(u1) * (sin(beta * pi)/tan(beta * pi * v1) -
cos(beta * pi))^(1/beta)
# Generation of symmetric alpha-stable deviates
# See Fulger, Scalas, Germano 2008 and references therein
gammax <- 1 #scale parameter
alpha <- 1.95 #Levy parameter
```

```
u2 <- runif(n)
v2 <- runif(n)
phi <- pi*(v2 - 0.5)

xi <- gammax *(sin(alpha*phi)/cos(phi))*
(-log(u2)*cos(phi)/cos((1-alpha)*phi))^(1-1/alpha)

# histogram of xi
# hist(xi)

# Random walk

x <- cumsum(xi)

# Epochs

t <- cumsum(tau)

# Stairplot
plot(t,x,type="s")</pre>
```

This page intentionally left blank

Bibliography

- [1] Abdeljawad, T. and Baleanu, D. (2011a). Caputo q-fractional initial value problems and a q-analogue Mittag-Leffler function, Commun. Nonlin. Sci. Numer. Sim. 16, pp. 4682–4688.
- [2] Abdeljawad, T. and Baleanu, D. (2011b). Fractional differences and integration by parts, J. Comput. Anal. Appl. 13, pp. 574-582.
- [3] Abel, N. H. (1823). Solution de quelques problèmes á l'aide d'integrales définies, Werke 1, pp. 10–14.
- [4] Abel, N. H. (1881). Solution de quelques problèmes à l'aide d'intégrales dèfinies, in *Oeuvres Complètes*, Vol. 1 (Grøndahl, Christiania), pp. 16–18.
- [5] Abramowitz, M. and Stegun, I. A. (1965). Handbook of Mathematical Functions (Dover Publ., New York).
- [6] Adamchick, V. S. (1997). On Stirling numbers and Euler sums, J. Comput. Appl. Math. 79, pp. 119–130.
- [7] Adolfsson, K., Enelund, M. and Larsson, S. (2004). Adaptive discretization of fractional order viscoelasticity using sparse time history, *Comput. Meth. Appl. Mech. Engin.* 193, pp. 4567–4590.
- [8] Adomian, G. (1989). Nonlinear Stochastic Systems Theory and Applications to Physics (Kluwer, Dordrecht).
- [9] Adomian, G. (1994). Solving Frontier Problems of Physics: The Decomposition Method (Kluwer, Dordrecht).
- [10] Agarwal, R. P., Benchohra, M. and Hamani, S. (2010a). A survey on existence results for boundary value problems of nonlinear fractional differential equations and inclusions, *Acta Appl. Math.* 109, pp. 973–1033.
- [11] Agarwal, R. P., Bohner, M., O'Regan, D. and Peterson, A. (2002). Dynamic equations on time scales: A survey, J. Comput. Appl. Math. 141, pp. 1–26.
- [12] Agarwal, R. P., Lakshmikantham, V. and Nieto, J. J. (2010b). On the concept of solution for fractional differential equations with uncertainty, Nonlinear Anal.: Theory, Meth. Appl. 72, pp. 2859–2862.
- [13] Agarwal, R. P., O'Regan, D. and Stanek, S. (2010c). Positive solutions for Dirichlet problems of singular nonlinear fractional differential equations, J. Math. Anal. App. 371, pp. 57–68.

- [14] Agarwal, R. P., Zhou, Y. and He, Y. (2010d). Existence of fractional neutral functional differential equations, Comp. Math. Appl. 59, pp. 1095–1100.
- [15] Agrawal, O. P. (2001). A new Lagrangian equation of motion for fractionally damped systems, J. Appl. Mech. 68, pp. 339–341.
- [16] Agrawal, O. P. (2002a). Formulation of Euler-Lagrange equations for fractional variational problems, J. Math. Anal. Appl. 52, pp. 1247–1253.
- [17] Agrawal, O. P. (2002b). Solution for a fractional diffusion-wave equation defined in a bounded domain, *Nonlinear Dyn.* 29, pp. 145–155.
- [18] Agrawal, O. P. (2004). A general formulation and solution scheme for fractional optimal control problems, *Nonlinear Dyn.* 38, pp. 323–337.
- [19] Agrawal, O. P. (2006). Fractional variational calculus and the transversality conditions, J. Phys. A.: Math. Gen. 39, pp. 10375–10384.
- [20] Agrawal, O. P. (2007a). Fractional variational calculus in terms of Riesz fractional derivatives, J. Phys. A: Math. Theor. 40, pp. 6287–6303.
- [21] Agrawal, O. P. (2007b). Generalized Euler-Lagrange equations and transversality conditions for FVPs in terms of the Caputo derivative, J. Vib. Contr. 13, pp. 1217–1237.
- [22] Agrawal, O. P. (2008). A formulation and numerical scheme for fractional optimal control problems, J. Vib. Contr. 14, pp. 1291–1299.
- [23] Agrawal, O. P. (2010). Generalized variational problems and Euler-Lagrange equations, Comput. Math. Appl. 59, pp. 1852–1864.
- [24] Agrawal, O. P. and Baleanu, D. (2007). A Hamiltonian formulation and a direct numerical scheme for fractional optimal control problems, J. Vib. Contr. 13, pp. 1269–1281.
- [25] Agrawal, O. P., Defterli, O. and Baleanu, D. (2010). Fractional optimal control problems with several state and control variables, J. Vib. Contr. 16, pp. 1967–1976.
- [26] Agrawal, O. P. and Gregory, J. (2003). $O(h^2)$ global pointwise algorithms for delay quadratic problems in the calculus of variations, SIAM J. Numer. Anal. 41, pp. 1773–1784.
- [27] Agrawal, O. P., Gregory, J. and Spector, K. P. (1997). A Bliss-type multiplier rule for constrained variational problems with time delay, J. Math. Anal. Appl. 20, pp. 702–711.
- [28] Agrawal, O. P., Muslih, S. I. and Baleanu, D. (2011). Generalized variational calculus in terms of multi-parameters fractional derivatives, Commun. Nonlin. Sci. Numer. Sim. 16, pp. 4756–4767.
- [29] Ahmad, B. and Nieto, J. J. (2009). Existence results for a coupled system of nonlinear fractional differential equations with three-point boundary conditions, Comp. Math. Appl. 58, pp. 1838–1843.
- [30] Ahmed, E. and Elgazzar, A. S. (2007). On fractional order differential equations model for nonlocal epidemics, *Phys. A* 379, pp. 607–614.
- [31] Aigner, M. (1975). Kombinatorik I (Springer, Berlin).
- [32] Aitchison, J. and Brown, J. A. (1957). The Lognormal Distribution (Cambridge Univ. Press, Cambridge).
- [33] Alaimia, M. R. and Tatar, N. E. (2005). Blow up for the wave equation with a fractional damping, J. Appl. Anal. 11, pp. 133–144.

- [34] Almeida, R. and Torres, D. F. (2009). Calculus of variations with fractional derivatives and fractional integrals, Appl. Math. Lett. 22, pp. 1816–1820.
- [35] Ames, W. F. (1992). Numerical Methods for Partial Differential Equations (Acad. Press, Boston).
- [36] Anderson, D. R. and Zafer, A. (2009). Nonlinear oscillation of second-order dynamic equations on time scales, Appl. Math. Lett. 22, pp. 1591–1597.
- [37] Apostol, T. M. (1957). Mathematical Analysis: A Modern Approach to Advanced Calculus (Addison-Wesley, London).
- [38] Applebaum, D. (2009). Lévy Processes and Stochastic Calculus (Cambridge Univ. Press, Cambridge).
- [39] Arena, P., Caponetto, R., Fortuna, L. and Porto, D. (2000). Nonlinear Noninteger Order Circuits and Systems: Theory and Applications (World Scientific, Singapore).
- [40] Artin, E. (1964). The Gamma Function (Holt, Rinehart and Winston, New York).
- [41] Atanacković, T. M., Budinčević, M. and Pilipović, S. (2005). On a fractional distributed-order oscillator, J. Phys. A: Math. Gen. 38, pp. 6703–6713.
- [42] Atanacković, T. M., Konjik, S. and Pilipović, S. (2008). Variational problems with fractional derivatives: Euler-Lagrange equations, J. Phys. A: Math. Theor. 41, pp. 095201–095212.
- [43] Atici, F. M. and Eloe, P. W. (2007). A transform method in discrete fractional calculus, Int. J. Differ. Equ. 2, pp. 165–176.
- [44] Atici, F. M. and Eloe, P. W. (2009). Initial value problem in discrete fractional calculus, Proc. Amer. Math. Soc. 137, pp. 981–989.
- [45] Audounet, J., Giovangigli, V. and Roquejoffre, J. M. (1998). A threshold phenomenon in the propagation of a point source initiated flame, *Phys. D* 121, pp. 295–316.
- [46] Bachelier, L. (1900). Théorie de la spéculation, Annales Scientifiques de l'École Normale Supérieure 3, pp. 21–86.
- [47] Bagley, R. L. and Torvik, P. J. (2000a). On the existence of the order domain and the solution of distributed order equations – Part I, *Internat. J. Appl. Math.* 2, pp. 865–882.
- [48] Bagley, R. L. and Torvik, P. J. (2000b). On the existence of the order domain and the solution of distributed order equations – Part II, *Internat. J. Appl. Math.* 2, pp. 965–987.
- [49] Baillie, R. T. and King, M. L. (1996). Fractional differencing and long memory processes, J. Econometrics 73, pp. 1–3.
- [50] Balachandran, K. and Trujillo, J. J. (2010). The nonlocal Cauchy problem for nonlinear fractional integrodifferential equations in Banach spaces, Nonlinear Anal.: Theory, Meth. Appl. 72, pp. 4587–4593.
- [51] Balakrishnan, V. (1985). Anomalous diffusion in one dimension, Phys. A 132, pp. 569–580.
- [52] Baleanu, D. (1999). Killing-Yano tensors and Nambu tensors, Nuovo Cimento B 114, pp. 1065–1072.
- [53] Baleanu, D. (2004). Constrained systems and Riemann-Liouville fractional derivative, in A. Le Méhauté, J. A. Tenreiro Machado, J. C. Trigeassou

- and J. Sabatier (eds.), Proceedings of the 1st IFAC Workshop on Fractional Differentiation and its Applications (ENSEIRB, Bordeaux), pp. 597–602.
- [54] Baleanu, D. (2005a). About fractional calculus of singular Lagrangians, J. Adv. Comput. Intell., Intel. Inform. 9, pp. 395–398.
- [55] Baleanu, D. (2005b). Constrained systems and Riemann-Liouville fractional derivatives, in A. Le Méhauté, J. A. Tenreiro Machado, J. C. Trigeassou and J. Sabatier (eds.), Fractional Differentiation and its Applications (Ubooks, Neusäß), pp. 69–80.
- [56] Baleanu, D. (2006). Fractional Hamiltonian analysis of irregular systems, Signal Process. 86, pp. 2632–2636.
- [57] Baleanu, D. (2008a). Fractional constrained systems and Caputo derivatives, J. Comput. Nonlin. Dyn. 3, pp. 021102–021105.
- [58] Baleanu, D. (2008b). New applications of fractional variational principles, Rep. Math. Phys. 61, pp. 199–206.
- [59] Baleanu, D. (2009a). About fractional quantization and fractional variational principles, Commun. Nonlin. Sci. Numer. Sim. 14, pp. 2520–2523.
- [60] Baleanu, D. (2009b). Fractional variational principles in action, Phys. Scr. T136, p. 014006.
- [61] Baleanu, D. and Agrawal, O. P. (2006). Fractional Hamilton formalism within Caputo's derivative, Czech. J. Phys. 56, pp. 1087–1092.
- [62] Baleanu, D. and Avkar, T. (2004). Lagrangians with linear velocities within Riemann-Liouville fractional derivatives, *Nuovo Cimento B* 119, pp. 73–79.
- [63] Baleanu, D., Defterli, O. and Agrawal, O. P. (2009a). A central difference numerical scheme for fractional optimal control problems, J. Vibr. Contr. 15, pp. 583–597.
- [64] Baleanu, D., Golmankhaneh, A. K. and Golmankhaneh, A. K. (2009b). The dual action of fractional multi time Hamilton equations, *Int. J. Theor. Phys.* 48, pp. 2558–2569.
- [65] Baleanu, D., Golmankhaneh, A. K. and Golmankhaneh, A. K. (2009c). Fractional Nambu mechanics, Int. J. Theor. Phys. 48, pp. 1044–1052.
- [66] Baleanu, D., Guvenc, Z. B. and Tenreiro Machado, J. A. (2010a). New Trends in Nanotechnology and Fractional Calculus Applications (Springer, Heidelberg).
- [67] Baleanu, D., Jarad, F. and Muslih, S. (2006). Difference discrete and fractional variational principles, Facta Univ.: Phys. Chem. Tech. 4, pp. 175–183.
- [68] Baleanu, D., Maraaba, T. and Jarad, F. (2008a). Fractional variational principles with delay, J. Phys. A: Math. Theor. 41, p. 315403.
- [69] Baleanu, D. and Muslih, S. I. (2005a). About fractional supersymmetric quantum mechanics, Czech. J. Phys. 55, pp. 1063–1066.
- [70] Baleanu, D. and Muslih, S. I. (2005b). Lagrangian formulation of classical fields within Riemann-Liouville fractional derivatives, *Phys. Scr.* 72, pp. 119–121.
- [71] Baleanu, D. and Muslih, S. I. (2007). Fractional Euler-Lagrange and fractional Hamilton equations for super symmetric classical model, Fractals 15, pp. 379–383.

- [72] Baleanu, D., Muslih, S. I. and Rabei, E. M. (2008b). On fractional Euler-Lagrange and Hamilton equations and the fractional generalization of total time derivative, *Nonlinear Dyn.* 53, pp. 67–74.
- [73] Baleanu, D., Muslih, S. I., Rabei, E. M., Golmankhaneh, A. K. and Golmankhaneh, A. K. (2010b). On fractional dynamics on the extended phase space, J. Comput. Nonlin. Dyn. 5, pp. 041011-041014.
- [74] Baleanu, D., Muslih, S. I., Rabei, E. M., Golmankhaneh, A. K. and Golmankhaneh, A. K. (2011). On fractional Hamiltonian systems possessing first-class constraints within Caputo derivatives, *Rom. Rep. Phys.* 63, pp. 3–8.
- [75] Baleanu, D., Mustafa, O. G. and Agarwal, R. P. (2010c). Asymptotically linear solutions for some linear fractional differential equations, *Abstr. Appl. Anal.* 2010, p. 865139.
- [76] Baleanu, D., Sadati, S. J., Ranjbar, A., Ghaderi, R., Abdeljawad, T. and Jarad, F. (2010d). Razumikhin stability theorem for fractional systems with delay, Abstr. Appl. Anal. ID124812, pp. 1–9.
- [77] Baleanu, D. and Trujillo, J. J. (2008). On exact solutions of a class of fractional Euler-Lagrange equations, Nonlinear Dyn. 52, pp. 331–335.
- [78] Baleanu, D. and Trujillo, J. J. (2010). A new method of finding the fractional Euler-Lagrange and Hamilton equations within Caputo fractional derivatives, Commun. Nonlin. Sci. Numer. Sim. 15, pp. 1111–1115.
- [79] Baleanu, D. and Vacaru, S. (2011a). Constant curvature coefficients and exact solutions in fractional gravity and geometric mechanics, Centr. Europ. J. Phys. 9, pp. 1267–1279.
- [80] Baleanu, D. and Vacaru, S. (2011b). Fedosov quantization of fractional Lagrange spaces, Int. J. Theor. Phys. 50, pp. 233–243.
- [81] Baleanu, D. and Vacaru, S. (2011c). Fractional almost Kähler-Lagrange geometry, Nonlinear Dyn. 64, pp. 365–373.
- [82] Baleanu, D. and Vacaru, S. (2011d). Fractional curve flows and solitonic hierarchies in gravity and geometric mechanics, J. Math. Phys. (in press).
- [83] Bastos, N. R., Ferreira, R. A. and Torres, D. F. (2011). Necessary optimality conditions for fractional difference problems of the calculus of variations, *Discr. Contin. Dyn. Syst.*, Ser. A 29, pp. 417–437.
- [84] Beghin, L. and Orsingher, E. (2009). Fractional Poisson process and related random motions, *Electron. J. Probab.* 14, pp. 1790–1826.
- [85] Ben-Avraham, D. and Havlin, S. (2005). Diffusion and Reactions in Fractals and Disordered Systems (Cambridge Univ. Press, Cambridge).
- [86] Benson, D. A. (1998). The Fractional Advection-Dispersion Equation: Development and Application, Ph.D. thesis, University of Nevada at Reno.
- [87] Benson, D. A., Wheatcraft, S. W. and Meerschaert, M. M. (2000). Application of a fractional advection-dispersion equation, Water Resour. Res. 36, pp. 1403–1412.
- [88] Bering, K. (2000). A note on non-locality and Ostrogradski's construction, Tech. rep., preprint:hep-th/0007192.
- [89] Bertoin, J. (1998). Lévy Processes (Cambridge Univ. Press, Cambridge).

- [90] Bertram, W. (2004). An empirical investigation of Australian Stock Exchange data, Phys. A 341, pp. 533–546.
- [91] Bhalekar, S. and Daftardar-Gejji, V. (2010). Fractional ordered Liu system with time-delay, Commun. Nonlin. Sci. Numer. Sim. 15, pp. 2178–2191.
- [92] Bhalekar, S., Daftardar-Gejji, V., Baleanu, D. and Magin, R. (2011). Fractional Bloch equation with delay, Comput. Math. Applic. 61, pp. 1355–1365.
- [93] Bingham, N. H., Goldie, C. M. and Teugels, J. L. (1987). Regular Variation (Cambridge Univ. Press, Cambridge).
- [94] Birk, C. and Song, C. (2009). A continued-fraction approach for transient diffusion in unbounded medium, Comput. Methods Appl. Mech. Eng. 198, pp. 2576–2590.
- [95] Birk, C. and Song, C. (2010). A local high-order doubly asymptotic open boundary for diffusion in a semi-infinite layer, J. Comput. Phys. 229, pp. 6156–6179.
- [96] Bisquert, J. (2005). Interpretation of a fractional diffusion equation with nonconserved probability density in terms of experimental systems with trapping or recombination, *Phys. Rev. E* **72**, pp. 011109–1–011109–6.
- [97] Biswas, R. J. and Sen, S. (2011). Fractional optimal control problems with specified final time, J. Comput. Nonlin. Dyn. 6, pp. 021009.1–021009.6.
- [98] Björck, Å. and Pereyra, V. (1970). Solution of Vandermonde systems of equations, Math. Comp. 24, pp. 893–903.
- [99] Blank, L. (1997). Numerical treatment of differential equations of fractional order, Nonlinear World 4, pp. 473–490.
- [100] Bochner, S. (1932). Vorlesungen über Fouriersche Integrale (Akademische Verlagsgesellschaft, Leipzig).
- [101] Bohner, M. and Guseinov, G. S. (2004). Partial differentiation on time scales, Dyn. Sys. Appl. 13, pp. 351–379.
- [102] Bohner, M. and Peterson, A. (2001). Dynamic equations on time scales (Birkhäuser, Basel).
- [103] Bolster, D., Benson, D. A., Le Borgne, T. and Dentz, M. (2010). Anomalous mixing and reaction induced by superdiffusive nonlocal transport, *Phys. Rev. E* 82, pp. 021119–021123.
- [104] Bonilla, B., Kilbas, A. A. and Trujillo, J. J. (2003). Cálculo Fraccionario y Ecuaciones Diferenciales Fraccionarias (UNED, Madrid).
- [105] Bonilla, B., Rivero, M., Rodríguez-Germá, L. and Trujillo, J. J. (2007). Fractional differential equations as alternative models to the nonlinear fractional differential equations, Appl. Math. Comp. 187, pp. 79–88.
- [106] Bottazzi, G. and Secchi, A. (2003). Why are distributions of firm growth rates tent shaped? *Economics Letters* **80**, pp. 415–420.
- [107] Bottazzi, G. and Secchi, A. (2006). Explaining the distribution of firm growth rates, The RAND Journal of Economics 37, pp. 235–256.
- [108] Bounaim, A., Holm, S., Chen, W. and Odegard, A. (2004). Sensitivity of the ultrasonic CARI technique for breast tumor detection using a FETD scheme, *Ultrasonics* 42, pp. 919–25.
- [109] Braß, H. (1995). On the principle of avoiding the singularity in quadrature, Z. Angew. Math. Mech. 75, pp. S617–S618.

- [110] Brunner, H. and van der Houwen, P. J. (1986). The Numerical Solution of Volterra Equations (North-Holland, Amsterdam).
- [111] Butzer, P. L. and Hauss, M. (1991a). On Stirling functions of the second kind, Stud. Appl. Math. 84, pp. 71–91.
- [112] Butzer, P. L. and Hauss, M. (1991b). Stirling functions of first and second kind: Some new applications, in S. Baron and D. Leviatan (eds.), Approximation, Interpolation and Summability (AMS, Providence), pp. 89–108.
- [113] Butzer, P. L. and Hauss, M. (1999). Application of sampling theory to combinatorical analysis, Stirling numbers, special functions and the Riemann zeta functions, in J. R. Higgins and R. L. Stens (eds.), Sampling Theory in Fourier Analysis: Advanced Topics (Oxford Univ. Press, Oxford), pp. 1–37.
- [114] Butzer, P. L., Hauss, M. and Schmidt, M. (1989). Factorial functions and Stirling numbers of fractional orders, *Results Math.* 16, pp. 16–48.
- [115] Butzer, P. L. and Jansche, S. (1997). A direct approach to the Mellin transform, J. Fourier Anal. Appl. 3, pp. 325–375.
- [116] Butzer, P. L., Kilbas, A. A. and Trujillo, J. J. (2002a). Compositions of Hadamard-type fractional integration operators and the semigroup property, J. Math. Anal. Appl. 269, pp. 387–400.
- [117] Butzer, P. L., Kilbas, A. A. and Trujillo, J. J. (2002b). Fractional calculus in the Mellin setting and Hadamard-type fractional integrals, J. Math. Anal. Appl. 269, pp. 1–27.
- [118] Butzer, P. L., Kilbas, A. A. and Trujillo, J. J. (2002c). Mellin transform analysis and integration by parts for Hadamard-type fractional integrals, J. Math. Anal. Appl. 270, pp. 1–15.
- [119] Butzer, P. L., Kilbas, A. A. and Trujillo, J. J. (2003a). Generalized Stirling functions of second kind and representations of fractional order differences via derivatives, J. Diff. Equat. Appl. 9, pp. 503–533.
- [120] Butzer, P. L., Kilbas, A. A. and Trujillo, J. J. (2003b). Stirling functions of the second kind in the setting of difference and fractional calculus, *Numer. Funct. Anal. Optim.* 24, pp. 673–711.
- [121] Butzer, P. L., Markett, C. and Schmidt, M. (1991). Stirling numbers, central factorial numbers, and representations of the Riemann zeta function, Results Math. 19, pp. 257–275.
- [122] Butzer, P. L., Rodríguez-Germá, L., Kilbas, A. A. and Trujillo, J. J. (2007). Stirling functions of the first kind in the setting of fractional calculus, J. Diff. Equat. Appl. 13, pp. 683–721.
- [123] Butzer, P. L., Schmeisser, G. and Stens, R. L. (2001). An introduction to sampling analysis, in F. Marvasti (ed.), *Nonuniform Sampling: Theory and Practice* (Kluwer Acad. Publ., New York), pp. 17–121.
- [124] Butzer, P. L., Splettstoesser, W. and Stens, R. L. (1988). The sampling theorem and linear prediction in signal analysis, *Jahresber. Dtsch. Math.-Ver.* 90, pp. 1–70.
- [125] Cabré, X. and Roquejoffre, J. M. (2009). Front propagation in Fisher-KPP equations with fractional diffusion, C. R. (Math.) Acad. Sci. Paris 347, pp. 1361–1366.

- [126] Caffarelli, L. and Silvestre, L. (2007). An extension problem related to the fractional Laplacian, Comm. Part. Diff. Equ. 32, pp. 1245–1260.
- [127] Caffarelli, L. and Vázquez, J. L. (2011). Asymptotic behaviour of a porous medium equation with fractional diffusion, *Discret. Contin. Dyn. Syst.* 29, pp. 1393–1404.
- [128] Caniego, F. J., Espejo, R., Martín, M. A. and San José Martínez, F. (2005). Multifractal scaling of soil spatial variability, *Ecological Model.* 182, pp. 291–303.
- [129] Cao, X., Burrage, K. and Abdullah, F. (2006). A variable stepsize implementation for fractional differential equations, Tech. rep., University of Queensland, Brisbane.
- [130] Caponetto, R., Dongola, G. and Fortuna, L. (2010). Fractional Order Systems: Modeling and Control Applications (World Scientific, Singapore).
- [131] Caputo, M. (1966). Linear models of dissipation whose q is almost frequency independent, Annals of Geophysics 19, pp. 383–393.
- [132] Caputo, M. (1969). Elasticità e Dissipazione (Zanichelli, Bologna).
- [133] Caputo, M. (1995). Mean fractional-order-derivatives: Differential equations and filters, Ann. Univ. Ferrara Sez. VII (N. S.) 41, pp. 73–84.
- [134] Caputo, M. (2001). Distributed order differential equations modelling dielectric induction and diffusion, Fract. Calc. Appl. Anal. 4, pp. 421–442.
- [135] Caputo, M. (2003). Diffusion with space memory modelled with distributed order space fractional differential equations, Annals of Geophysics 46, pp. 223–234.
- [136] Caputo, M. and Mainardi, F. (1971). A new dissipation model based on memory mechanism, Pure Appl. Geophys. 91, pp. 134–147.
- [137] Castro, A., Córdoba, D., Gancedo, F. and Orive, R. (2009). Incompressible flow in porous media with fractional diffusion, C. R. (Math.) Acad. Sci. Paris 347, pp. 1361–1366.
- [138] Çinlar, E. (1975). Introduction to Stochastic Processes (Prentice-Hall, Englewood Cliffs).
- [139] Chambers, J. M., Mallows, C. L. and Stuck, B. W. (1976). A method for simulating stable random variables, J. Amer. Statist. Ass. 71, pp. 340–344.
- [140] Champeney, D. C. (1987). A Handbook of Fourier Theorems (Cambridge Univ. Press, Cambridge).
- [141] Chatterjee, A. (2005). Statistical origins of fractional derivatives in viscoelasticity, J. Sound Vibration 284, pp. 1239–1245.
- [142] Cheang, G. H. and Chiarella, C. (2011). A modern view on Merton's jumpdiffusion model, University of Technology, Sydney, Quantitative Finance Research Centre, Research paper 287.
- [143] Chelluri, R., Richmond, L. B. and Temme, N. M. (2000). Asymptotic estimates for generalized Stirling numbers, Analysis 20, pp. 1–13.
- [144] Chen, Y. Q. and Moore, K. L. (2002). Analytical stability bound for a class of delayed fractional-order dynamic systems, *Nonlinear Dyn.* 29, pp. 191–200.
- [145] Chen, W., Baleanu, D. and Tenreiro Machado, J. A. (eds.) (2010). Frac-

- tional differentiation and its applications, Comput. Math. Applic. 59, pp. 1585–1896.
- [146] Cody, W. J. and Stoltz, L. (1989). GAMMA, in SPECFUN 2.5, URL http://www.netlib.org/specfun/gamma.
- [147] Compte, A. (1996). Stochastic foundations of fractional dynamics, Phys. Rev. E 53, pp. 4191–4193.
- [148] Comtet, L. (1974). Advanced Combinatorics: The Art of Finite and Infinite Expansion (Reidel, Dordrecht).
- [149] Copson, E. T. (1943). Some applications of Marcel Riesz's integral of fractional order, Proc. Roy. Soc. Edinburgh A 61, pp. 260–272.
- [150] Corcino, R. B. and Hsu, L. C. (2001). Combinatorial and statistical applications of generalized Stirling numbers, J. Math. Res. Exposition 21, pp. 337–343.
- [151] Cramér, H. (1930). On the Mathematical Theory of Risk (Skandia Jubilee Volume, Stockholm).
- [152] Cresson, J. (2007). Fractional embedding of differential operators and Lagrangian systems, J. Math. Phys. 48, pp. 033504–033537.
- [153] Cresson, J. (2010). Inverse problem of fractional calculus of variations for partial differential equations, *Commun. Nonlin. Sci. Numer. Sim.* 15, pp. 987–996.
- [154] Cresson, J. and Inizan, P. (2009). About fractional Hamiltonian systems, Phys. Scr. T136, p. 014007.
- [155] Cuesta, E. (2010). A variable step size numerical method based on fractional type quadratures for linear integro-differential equations, Adv. Engineer. Softw. 41, pp. 64–69.
- [156] Cuesta, E., Lubich, C. and Palencia, C. (2006). Convolution quadrature time discretization of fractional diffusion-wave equations, *Math. Comput.* 75, pp. 673–696.
- [157] Cuesta, E. and Palencia, C. (2003). A numerical method for an integrodifferential equation with memory in Banach spaces: Qualitative properties, SIAM J. Numer. Anal. 41, pp. 1232–1241.
- [158] Dahlquist, G. (1956). Convergence and stability in the numerical integration of ordinary differential equations, *Math. Scand.* 4, pp. 33–53.
- [159] Dahlquist, G. (1959). Stability and error bounds in the numerical integration of ordinary differential equations, Tekn. Högskol. Handl. 130, Royal Inst. of Technology, Stockholm.
- [160] Dahlquist, G. (1978). On accuracy and unconditional stability of linear multistep methods for second order differential equations, BIT 18, pp. 133–136.
- [161] Das, S. (2008). Functional Fractional Calculus for System Identification and Controls (Springer, Berlin).
- [162] Davis, P. J. and Rabinowitz, P. (1984). Methods of Numerical Integration (Acad. Press, Orlando).
- [163] de Boor, C. (1978). A Practical Guide to Splines (Springer, New York).
- [164] de Donder, T. (1935). Théorie invariantive de calculus de variations (Gauthier-Villars, Paris).

- [165] de Finetti, B. (1975). Theory of Probability, Vol. II (Wiley, New York).
- [166] Deng, W. H. (2007). Short memory principle and a predictor-corrector approach for fractional differential equations, J. Comput. Appl. Math. 206, pp. 174–188.
- [167] Diethelm, K. (1997a). An algorithm for the numerical solution of differential equations of fractional order, *Elec. Transact. Numer. Anal.* 5, pp. 1–6.
- [168] Diethelm, K. (1997b). Generalized compound quadrature formulae for finite-part integrals, IMA J. Numer. Anal. 17, pp. 479–493.
- [169] Diethelm, K. (2003). Efficient solution of multi-term fractional differential equations using P(EC)^mE methods, Computing 71, pp. 305–319.
- [170] Diethelm, K. (2008a). An analysis of some nonclassical methods for the numerical approximation of Caputo-type fractional derivatives, *Numer. Al*gorithms 47, pp. 361–390.
- [171] Diethelm, K. (2008b). Multi-term fractional differential equations, multiorder fractional differential systems and their numerical solution, J. Europ. Syst. Autom. 42, pp. 665–676.
- [172] Diethelm, K. (2010). The Analysis of Fractional Differential Equations (Springer, Heidelberg).
- [173] Diethelm, K. (2011). An efficient parallel algorithm for the numerical solution of fractional differential equations, Fract. Calc. Appl. Anal. 14, pp. 475–490.
- [174] Diethelm, K., Ford, J. M., Ford, N. J. and Weilbeer, M. (2005a). A comparison of backward differentiation approaches for ordinary and partial differential equations of fractional order, in A. Le Méhauté, J. A. Tenreiro Machado, J. C. Trigeassou and J. Sabatier (eds.), Fractional Differentiation and its Applications (Ubooks, Neusäß), pp. 557–569.
- [175] Diethelm, K., Ford, J. M., Ford, N. J. and Weilbeer, M. (2006). Pitfalls in fast numerical solvers for fractional differential equations, J. Comput. Appl. Math. 186, pp. 482–503.
- [176] Diethelm, K. and Ford, N. J. (2001). Numerical solution methods for distributed order differential equations, Fract. Calc. Appl. Anal. 4, pp. 531– 542.
- [177] Diethelm, K. and Ford, N. J. (2002a). Analysis of fractional differential equations, J. Math. Anal. Appl. 265, pp. 229–248.
- [178] Diethelm, K. and Ford, N. J. (2002b). Numerical solution of the Bagley-Torvik equation, BIT 42, pp. 490–507.
- [179] Diethelm, K. and Ford, N. J. (2004). Multi-order fractional differential equations and their numerical solution, Appl. Math. Comput. 154, pp. 621– 640.
- [180] Diethelm, K. and Ford, N. J. (2009). Numerical analysis for distributedorder differential equations, J. Comput. Appl. Math. 225, pp. 96–104.
- [181] Diethelm, K. and Ford, N. J. (2011). Volterra integral equations and fractional calculus: Do neighbouring solutions intersect? J. Integr. Equ. Appl. (in press).
- [182] Diethelm, K., Ford, N. J. and Freed, A. D. (2002). A predictor-corrector

- approach for the numerical solution of fractional differential equations, Nonlinear Dyn. 29, pp. 3–22.
- [183] Diethelm, K., Ford, N. J. and Freed, A. D. (2004). Detailed error analysis for a fractional Adams method, *Numer. Algorithms* 36, pp. 31–52.
- [184] Diethelm, K., Ford, N. J., Freed, A. D. and Luchko, Y. (2005b). Algorithms for the fractional calculus: A selection of numerical methods, *Comput. Methods Appl. Mech. Eng.* 194, pp. 743–773.
- [185] Diethelm, K. and Freed, A. D. (1999a). The FracPECE subroutine for the numerical solution of differential equations of fractional order, in S. Heinzel and T. Plesser (eds.), Forschung und wissenschaftliches Rechnen 1998, no. 52 in GWDG-Berichte (Gesellschaft für wissenschaftliche Datenverarbeitung, Göttingen), pp. 57–71.
- [186] Diethelm, K. and Freed, A. D. (1999b). On the solution of nonlinear fractional differential equations used in the modeling of viscoplasticity, in F. Keil, W. Mackens, H. Voß and J. Werther (eds.), Scientific Computing in Chemical Engineering II Computational Fluid Dynamics, Reaction Engineering, and Molecular Properties (Springer, Heidelberg), pp. 217–224.
- [187] Diethelm, K. and Weilbeer, M. (2004a). Initial-boundary value problems for time-fractional diffusion-wave equations and their numerical solution, in A. Le Méhauté, J. A. Tenreiro Machado, J. C. Trigeassou and J. Sabatier (eds.), Proceedings of the 1st IFAC Workshop on Fractional Differentiation and its Applications (ENSEIRB, Bordeaux), pp. 551–557.
- [188] Diethelm, K. and Weilbeer, M. (2004b). A numerical approach for Joulin's model of a point source initiated flame, Fract. Calc. Appl. Anal. 7, pp. 191–212.
- [189] Doetsch, G. (1937). Theorie und Anwendung der Laplace-Transformation (Springer, Berlin).
- [190] Dong, J. P. and Xu, M. Y. (2007). Some solutions to the space fractional Schrödinger equation using momentum representation method, J. Math. Phys. 48, p. 072105.
- [191] Driver, R. D. (1977). Ordinary and Delay Differential Equations (Springer, Berlin).
- [192] Duarte Ortigueira, M. (2011). Fractional Calculus for Scientists and Engineers (Springer, Berlin).
- [193] Duarte Ortigueira, M. and Tenreiro Machado, J. A. (eds.) (2003b). Fractional signal processing and applications, Signal Process. 83.
- [194] Duarte Ortigueira, M. and Tenreiro Machado, J. A. (eds.) (2006a). Fractional calculus applications in signals and systems, Signal Process. 86.
- [195] Duarte Ortigueira, M., Vinagre, B., Tenreiro Machado, J. A. and Trujillo, J. J. (eds.) (2011). Advances in fractional signals and systems, Signal Process. 91.
- [196] Dzherbashyan, M. M. (1966). Integral Transforms and Representations of Functions in the Complex Plane (Nauka, Moscow).
- [197] Dzherbashyan, M. M. (1970). A boundary value problem for a Sturm-Liouville type differential operator of fractional order, *Izv. Akad. Nauk Arm. SSR*, Ser. Mat. 5, pp. 71–96.

- [198] Edwards, J. T., Ford, N. J. and Simpson, A. C. (2002). The numerical solution of linear multi-term fractional differential equations: Systems of equations, J. Comput. Appl. Math. 148, pp. 401–418.
- [199] Edwards, J. T., Roberts, J. A. and Ford, N. J. (1997). A comparison of Adomian's decomposition method and Runge-Kutta methods for approximate solution of some predator prey model equations, Numerical Analysis Report 309, Manchester Centre for Computational Mathematics.
- [200] Einstein, A. (1905). Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen, Annalen der Physik 17, pp. 549–560.
- [201] El-Nabulsi, R. A. and Torres, D. F. M. (2008). Fractional action-like variational problems, J. Math. Phys. 49, pp. 053521–053527.
- [202] Elliott, D. (1993). An asymptotic analysis of two algorithms for certain Hadamard finite-part integrals, IMA J. Numer. Anal. 13, pp. 445–462.
- [203] Engheta, N. (1996). On fractional calculus and fractional multipoles in electromagnetism, *IEEE Trans. Antennas and Propagation* 44, pp. 554– 566.
- [204] Engheta, N. (1998). Fractional curl operator in electromagnetics, Microwave and Optical Technology Letters 17, pp. 86–91.
- [205] Engle, R. and Russel, J. (1997). Forecasting the frequency of changes in quoted foreign exchange prices with the autoregressive conditional duration model, *Journal of Empirical Finance* 4, pp. 187–212.
- [206] Engle, R. and Russel, J. (1998). Autoregressive conditional duration: A new model for irregularly spaced transaction data, *Econometrica* 66, pp. 1127–1162.
- [207] Erdélyi, A. (1940). Transformation of hypergeometric integrals by means of fractional integration by parts, Quart. J. Math. 11, pp. 293–303.
- [208] Erdélyi, A. (1965). An application of fractional integrals, J. d'Analyse Math. 14, pp. 113–126.
- [209] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G. (1953). Higher Transcendental Functions, Vol. I (McGraw-Hill, New York).
- [210] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G. (1954). Tables of Integral Transforms, Vol. II (McGraw-Hill, New York).
- 211] Euler, L. (1738). De progressionibus transcentibus, sev quarum termini algebraice dari nequeunt, Comment. Acad. Sci. Imperiales Petropolitanae 1, pp. 38–57.
- [212] Faddeev, L. and Jackiw, R. (1988). Hamiltonian reduction of unconstrained and constrained systems, Phys. Rev. Lett. 60, pp. 1692–1694.
- [213] Feller, W. (1943). On a general class of "contagious" distributions, Ann. Math. Stat. 14, pp. 389–400.
- [214] Feller, W. (1971). Introduction to Probability Theory and its Applications, Vol. II (Wiley, New York).
- [215] Fischer, H. (2010). A History of the Central Limit Theorem: From Classical to Modern Probability Theory (Springer, New York).
- [216] Fitzgerald, W. J. and Leung, C. M. (1996). A critical review of the Adomian method, Tech. Rep. CUED/F-INFENG/TR. 277, University of Cambridge.

- [217] Flajolet, P. and Prodinger, H. (1999). On Stirling numbers for complex arguments and Hankel contours, SIAM J. Discrete Math. 12, pp. 155–159.
- [218] Flomenbom, O. and Klafter, J. (2005). Closed-form solutions for continuous time random walks on finite chains, *Phys. Rev. Lett.* 95, p. 098105.
- [219] Flomenbom, O. and Silbey, R. J. (2007). Path PDFs for semi-Markovian random walks, Phys. Rev. E 76, p. 041101.
- [220] Fogedby, H. C. (1994). Langevin equations for continuous time Lévy flights, Phys. Rev. E 50, pp. 1657–1660.
- [221] Ford, N. J. and Connolly, J. A. (2006). Comparison of numerical methods for fractional differential equations, *Comm. Pure Appl. Anal.* 5, pp. 289– 307.
- [222] Ford, N. J. and Connolly, J. A. (2009). Systems-based decomposition schemes for the approximate solution of multi-term fractional differential equations, J. Comput. Appl. Math. 229, pp. 382–391.
- [223] Ford, N. J. and Simpson, A. C. (2000). The approximate solution of fractional differential equations of order greater than 1, in M. Deville and R. Owens (eds.), Proceedings of the 16th IMACS World Congress on Scientific Computation, Applied Mathematics, and Simulation (Lausanne), paper 213-1.
- [224] Ford, N. J. and Simpson, A. C. (2001). The numerical solution of fractional differential equations: Speed versus accuracy, *Numer. Algorithms* 26, pp. 333–346.
- [225] Ford, N. J., Xiao, J. and Yan, Y. (2011). A finite element method for time fractional partial differential equations, Fract. Calc. Appl. Anal. 14, pp. 454–474.
- [226] Fourier, J. (1888). Théorie Analytique de la Chaleur (Gauthier-Villars, Paris).
- [227] Frederico, G. S. and Torres, D. F. (2010). Fractional Noether's theorem in the Riesz-Caputo sense, Appl. Math. Comput. 217, pp. 1023–1033.
- [228] Fujita, Y. (1990a). Cauchy problems of fractional order and stable processes, Japan J. Appl. Math. 7, pp. 459–476.
- [229] Fujita, Y. (1990b). Integro-differential equation which interpolates the heat equation and the wave equation. I, Osaka J. Math. 27, pp. 309–321.
- [230] Fujita, Y. (1990c). Integro-differential equation which interpolates the heat equation and the wave equation. II, Osaka J. Math. 27, pp. 797–804.
- [231] Fulger, D., Scalas, E. and Germano, G. (2008). Monte Carlo simulation of uncoupled continuous-time random walks yielding a stochastic solution of the space-time fractional diffusion equation, *Phys. Rev. E* 77, pp. 021122.1– 021122.7.
- [232] Gaffeo, E., Delli Gatti, D., Di Guilmi, D., Giulioni, G., Gallegati, M. and Palestrini, A. (2004). Business cycle fluctuations and firms' size distribution dynamics, Adv. Complex Syst. 7, pp. 223–240.
- [233] Gaume, L. A. and Barbon, J. L. F. (2001). Nonlinear vacuum phenomena in noncommutative QED, Int. J. Mod. Phys. A 16, pp. 1123–1146.
- [234] Germano, G., Politi, M., Scalas, E. and Schilling, R. L. (2009). Stochastic

- calculus for uncoupled continuous-time random walks, $Phys.\ Rev.\ E$ **79**, pp. 066102–066113.
- [235] Gibrat, R. (1931). Les Inégalités Economiques, Application: Aux Inégalités des Richesses, à la Concentrations des Entreprises, aux Populations des Villes, aux Statistiques des Familles, etc., d'une Loi Nouvelle, al Loi de l'Effet Proportionnel (Librairie du Recueil Sirey, Paris).
- [236] Gitman, D. M. and Tytin, I. V. (1990). Quantization of fields with constraints (Springer, Berlin).
- [237] Gnedenko, B. V. and Kolmogorov, A. N. (1954). Limit Distributions for Sums of Independent Random Variables (Addison-Wesley, Cambridge, MA).
- [238] Goldfain, E. (2010). Fractional dynamics and the TeV regime of field theory, Commun. Nonlin. Sci. Numer. Sim. 13, pp. 666-676.
- [239] Golmankhaneh, A. K. (2009). Relativistic scalar fields for non-conservative systems, Phys. Scr. T136, p. 014008.
- [240] Golmankhaneh, A. K., Golmankhaneh, A. K., Baleanu, D. and Baleanu, M. C. (2010). Hamiltonian structure of fractional first order Lagrangian, Int. J. Theor. Phys. 49, 2, pp. 365–375.
- [241] Golmankhaneh, A. K., Golmankhaneh, A. K., Baleanu, D. and Baleanu, M. C. (2011). Fractional odd-dimensional mechanics, Adv. Difference Equ. ID526472, pp. 1–12.
- [242] Gomis, J., Kamimura, K. and Llosa, J. (2001). Hamiltonian formalism for space-time noncommutative theories, Phys. Rev. D 63, p. 045003.
- [243] Gomis, J., Kamimura, K. and Ramirez, T. (2004). Physical degrees of freedom of non-local theories, Nucl. Phys. B 696, pp. 263–291.
- [244] Gomis, J. and Mehen, T. (2000). Space-time noncommutative field theories and unitarity, Nucl. Phys. B 591, pp. 265–276.
- 245] Gorenflo, R. (1965). Computation of an integral transform of Abel's type in the presence of noise by quadratic programming, in *Proceedings of IFIP Congress* 1965, Vol. II (New York), pp. 575–576.
- [246] Gorenflo, R. and Abdel-Rehim, E. A. A. (2004). From power laws to fractional diffusion: The direct way, Vietnam Journal of Mathematics 32 (SI), pp. 65–75.
- [247] Gorenflo, R., Loutchko, J. and Luchko, Y. (2002). Computation of the Mittag-Leffler function $E_{\alpha,\beta}(z)$ and its derivative, Fract. Calc. Appl. Anal. 5, pp. 491–518.
- [248] Gorenflo, R., Loutchko, J. and Luchko, Y. (2003). Corrections to the paper "Computation of the Mittag-Leffler function $E_{\alpha,\beta}(z)$ and its derivative", Fract. Calc. Appl. Anal. 6, pp. 111–112.
- [249] Gorenflo, R. and Vessella, S. (1991). Abel Integral Equations (Springer, Berlin).
- [250] Graham, I. G. (1982). Galerkin methods for second kind integral equations with singularities, Math. Comp. 39, pp. 519–533.
- [251] Graham, R. L., Knuth, D. E. and Patashnik, O. (1994). Concrete Mathematics: A Foundation for Computer Science (Addison-Wesley, Boston).

- [252] Grünwald, A. K. (1867). Über "begrenzte" Derivationen und deren Anwendung, Z. Math. Phys. 12, pp. 441–480.
- [253] Hackbusch, W. (1995). Integral Equations: Theory and Numerical Treatment (Birkhäuser, Basel).
- [254] Hadamard, J. (1892). Essai sur l'étude des fonctions donnés par leur développement de Taylor, J. Math. Pures Appl. 8, pp. 101–186.
- [255] Hadamard, J. (1952). Lectures on Cauchy's Problem in Linear Partial Differential Equations (Dover Publ., New York), reprint.
- [256] Hairer, E., Lubich, C. and Schlichte, M. (1988). Fast numerical solution of weakly singular Volterra equations, J. Comput. Appl. Math. 23, pp. 87–98.
- [257] Hairer, E., Nørsett, S. P. and Wanner, G. (1993). Solving Ordinary Differential Equations I: Nonstiff Problems (Springer, Berlin).
- [258] Hairer, E. and Wanner, G. (1996). Solving Ordinary Differential Equations II: Stiff and Differential-algebraic Problems (Springer, Berlin).
- [259] Hardy, G. H. (1917). On some properties of integrals of fractional order, Messenger Math. 47, pp. 145–150.
- [260] Hardy, G. H. and Littlewood, J. E. (1917). Some properties of fractional integrals, Proc. London Math. Soc. 24, pp. 37–43.
- [261] Hardy, G. H. and Littlewood, J. E. (1928). Some properties of fractional integrals I, Math. Z. 27, pp. 565–570.
- [262] Hardy, G. H. and Littlewood, J. E. (1931). Some properties of fractional integrals II, Proc. London Math. Soc. 34, pp. 403–410.
- [263] Hartley, T. T. and Lorenzo, C. F. (1999). Fractional system identification: An approach using continuous order-distributions, NASA Technical Memorandum 1999-209640, NASA Glenn Research Center, Cleveland.
- [264] Hauss, M. (1989). Über die Theorie der fraktionierten Stirlingzahlen und deren Anwendung, Diploma thesis, Lehrstuhl A für Mathematik, RWTH Aachen.
- [265] Hauss, M. (1995). Verallgemeinerte Stirling-, Bernoulli- und Euler-Zahlen, deren Anwendungen und schnell konvergente Reihen fuer Zeta-Funktionen, Ph.D. thesis, RWTH Aachen.
- [266] He, J. H. (1998). Approximate solution of non linear differential equations with convolution product nonlinearities, Computer Meth. Appl. Mech. Eng. 167, pp. 69–73.
- [267] He, J. H. (2007). Variational iteration method. Some recent results and new interpretations, J. Comput. Appl. Math. 207, pp. 3–17.
- [268] Heaviside, O. (1892). Electrical Papers (Macmillan, London).
- [269] Heaviside, O. (1893). On operators in physical mathematics, Proc. Roy. Soc. 52, pp. 504–510.
- [270] Henneaux, M. and Teitelboim, C. (1993). Quantization of Gauge Systems (Princeton Univ. Press, Princeton).
- [271] Hernández, E. and Balachandran, K. (2010). Existence results for abstract degenerate neutral functional differential equations, *Bull. Aust. Math. Soc.* 81, pp. 329–342.
- [272] Herrmann, R. (2010). Common aspects of q-deformed Lie algebras and fractional calculus, Phys. A 389, pp. 4613–4622.

- [273] Herrmann, R. (2011). Fractional Calculus, an Introduction for Physicists (World Sci., New Jersey).
- [274] Herzallah, M. A. and Baleanu, D. (2009). Fractional-order Euler-Lagrange equations and formulation of Hamiltonian equations, *Nonlinear Dyn.* 58, pp. 385–391.
- [275] Herzallah, M. A. and Baleanu, D. (2011). Fractional-order variational calculus with generalized boundary conditions, Adv. Difference Equ. ID357580, pp. 1–9.
- [276] Herzallah, M. A., Muslih, S. I., Baleanu, D. and Rabei, E. M. (2011). Hamilton-Jacobi and fractional like action with time scaling, *Nonlinear Dyn.* (in press).
- [277] Heymans, N. and Podlubny, I. (2005). Physical interpretation of initial conditions for fractional differential equations with Riemann-Liouville fractional derivatives, *Rheol. Acta* 45, pp. 765–771.
- [278] Higgins, J. R. (1996). Sampling Theory in Fourier and Signal Analysis: Foundations (Clarendon Press, Oxford).
- [279] Higham, N. J. (1987). Error analysis of the Björck-Pereyra algorithms for solving Vandermonde systems, *Numer. Math.* 50, pp. 613–632.
- [280] Hilfer, R. (2000). Applications of Fractional Calculus in Physics (World Scientific, Singapore).
- [281] Hilfer, R. (2008). Threefold introduction to fractional derivatives, in R. Klages, G. Radons and I. M. Sokolov (eds.), Anomalous Transport: Foundations and Applications (Wiley-VCH, Weinheim), pp. 17–74.
- [282] Hilfer, R. and Anton, L. (1995). Fractional master equations and fractal time random walks, Phys. Rev. E 51, pp. R848–R851.
- [283] Hilger, S. (1988). Ein Maßkettenkalkül mit Anwendung auf Zentrumsmannigfaltigkeiten, Ph.D. thesis, Universität Würzburg.
- [284] Hsu, L. C. and Shiue, P. J. (1998). A unified approach to generalized Stirling numbers, Adv. Appl. Math. 20, pp. 366–384.
- [285] Hull, J. C. (2005). Options, Futures and Other Derivatives (Prentice Hall, Upper Saddle River).
- [286] Iserles, A. (1996). A First Course in the Numerical Analysis of Differential Equations (Cambridge Univ. Press, Cambridge).
- [287] Jacob, N. (2005). Pseudo-differential Operators and Markov Processes, Vol. III (Imperial College Press, London).
- [288] Jacod, J. and Shiryaev, A. (2002). *Limit Theorems for Stochastic Processes* (Springer, New York).
- [289] Janicki, A. and Weron, A. (1994). Simulation and Chaotic Behavior of α-Stable Stochastic Processes (Marcel Dekker, New York).
- [290] Janssen, J. and Manca, R. (2007). Semi-Markov Risk Models for Finance, Insurance and Reliability (Springer, New York).
- [291] Jarad, F., Abdeljawad, T. and Baleanu, D. (2010a). Fractional variational optimal control problems with delayed arguments, *Nonlinear Dyn.* 62, pp. 609–614.
- [292] Jarad, F., Abdeljawad, T. and Baleanu, D. (2010b). Fractional variational principles with delay within Caputo derivatives, Rep. Math. Phys. 65, pp. 17–28.

- [293] Jarad, F., Abdeljawad, T. and Baleanu, D. (2011). Higher order fractional variational optimal control problems with delayed arguments, ArXiv:1105.5148v1 [math.DS].
- [294] Jayakumar, K. (2003). Mittag-Leffler process, Math. Comput. Modelling 37, pp. 1427–1434.
- [295] Jelicic, Z. D. and Petrovacki, N. (2009). Optimality conditions and a solution scheme for fractional optimal control problems, Struct. Multidisc. Optim. 38, pp. 571–581.
- [296] Jordan, C. (1950). Calculus of Finite Differences (Chelsea Publ. Comp., New York).
- [297] Joulin, G. (1985). Point-source initiation of lean spherical flames of light reactants: An asymptotic theory, Combustion Science and Technology 43, pp. 99–113.
- [298] Jumarie, G. (2007). Lagrangian mechanics of fractional order, Hamilton-Jacobi fractional PDE and Taylor's series of nondifferentiable functions, Chaos Soliton Frac. 32, pp. 969–987.
- [299] Jumarie, G. (2009). From Lagrangian mechanics fractal in space to space fractal Schrödinger's equation via fractional Taylor's series, Chaos Soliton Frac. 41, pp. 1590–1604.
- [300] Kalia, R. N. (ed.) (1993). Recent Advances in Fractional Calculus (Global Publ. Co., Sauk Rapids).
- [301] Kanatchikov, I. V. (1999). De Donder-Weyl theory and a hypercomplex extension of quantum mechanics to field theory, *Rep. Math. Phys.* 43, pp. 157–170.
- [302] Kaneko, H., Noren, R. and Xu, Y. (1992). Numerical solutions for weakly singular Hammerstein equations and their superconvergence, J. Int. Equat. Appl. 4, pp. 391–406.
- [303] Kaymakcalan, B. (1993). Existence and comparison results for dynamic systems on time scales, J. Math. Anal. Appl. 172, pp. 243–255.
- [304] Kern, J. (1974). Lagrange geometry, Arch. Math. 25, pp. 438–443.
- [305] Kilbas, A. A. (2005). Fractional calculus of the generalized Wright function, Fract. Calc. Appl. Anal. 8, pp. 113–126.
- [306] Kilbas, A. A., Pierantozzi, T., Trujillo, J. J. and Vázquez, L. (2004). On the solution of fractional evolution equations, J. Phys. A: Math. Gen. 37, pp. 3271–3282.
- [307] Kilbas, A. A. and Saigo, M. (2004). H-Transforms: Theory and Applications (Chapman & Hall, Boca Raton).
- [308] Kilbas, A. A., Saigo, M. and Trujillo, J. J. (2002). On the generalized Wright function, Fract. Calc. Appl. Anal. 5, pp. 437–460.
- [309] Kilbas, A. A., Srivastava, H. M. and Trujillo, J. J. (2006). Theory and Applications of Fractional Differential Equations (Elsevier, Amsterdam).
- [310] Kilbas, A. A. and Trujillo, J. J. (2001). Differential equations of fractional order: Methods, results and problems. I, Appl. Anal. 78, pp. 153–192.
- [311] Kilbas, A. A. and Trujillo, J. J. (2002). Differential equations of fractional order: Methods, results and problems. II, *Appl. Anal.* **81**, pp. 435–493.
- [312] Kirane, M. and Laskri, Y. (2005). Nonexistence of global solutions to a

- hyperbolic equation with a space-time fractional damping, *Appl. Math. Comput.* **167**, pp. 1304–1310.
- [313] Kirane, M., Medve, M. and Tatar, N. (2011). Semilinear Volterra integrodifferential problems with fractional derivatives in the nonlinearities, Abstr. Appl. Anal. 2011, pp. 1248–1251.
- [314] Kiryakova, V. S. (1994). Generalized Fractional Calculus and Applications (Longman, Harlow).
- [315] Klafter, J., Lim, S. C. and Metzler, R. (eds.) (2011). Fractional Dynamics in Physics: Recent Advances (World Scientific, Singapore).
- [316] Klafter, J., Shlesinger, M. F. and Zumofen, G. (1996). Beyond Brownian motion, Phys. Today 49, pp. 33–39.
- [317] Klimek, M. (2001). Fractional sequential mechanics: models with symmetric fractional derivative, Czech. J. Phys. 51, pp. 1348–1354.
- [318] Klimek, M. (2002). Lagrangean and Hamiltonian fractional sequential mechanics, Czech. J. Phys. 52, pp. 1247–1253.
- [319] Knabner, P. and Angermann, L. (2003). Numerical Methods for Elliptic and Parabolic Partial Differential Equations (Springer, Berlin).
- [320] Kober, H. (1940). On fractional integrals and derivatives, Quart. J. Math. 11, pp. 193–211.
- [321] Koeller, R. C. (1986). Polynomial operators, Stieltjes convolution, and fractional calculus in hereditary mechanics, *Acta Mech.* **58**, pp. 251–264.
- [322] Konotop, V. V. and Vázquez, L. (1994). Nonlinear Random Waves (World Scientific, Singapore).
- [323] Kozubowski, T. J. (1998). Mixture representation of Linnik distribution revisited, Stat. Probab. Lett. 38, pp. 157–160.
- [324] Kozubowski, T. J. (2000a). Computer simulation of geometric stable distributions, J. Comput. Appl. Math. 116, pp. 221–229.
- [325] Kozubowski, T. J. (2000b). Exponential mixture representation of geometric stable distributions, *Annals of the Institute of Statistical Mathematics* **52**, pp. 231–238.
- [326] Kozubowski, T. J. and Rachev, S. T. (1999). Univariate geometric stable distributions, J. Comput. Anal. Appl. 1, pp. 177–217.
- [327] Lakshmikantham, V., Leela, S. and Devi, J. V. (2009). Theory of Fractional Dynamic Systems (Cambridge Sci. Publ., Cambridge).
- [328] Lang, W. (2000). On generalizations of the Stirling number triangles, J. Integer Seq. 3, pp. 1–19.
- [329] Laplace, P. S. (1812). Théorie Analytique des Probabilités (Courcier, Paris).
- [330] Larsson, S. and Thomée, V. (2003). Partial Differential Equations with Numerical Methods (Springer, Berlin).
- [331] Laskin, N. (2000). Fractional quantum mechanics, Phys. Rev. E 62, pp. 3135–3145.
- [332] Laskin, N. (2003). Fractional Poisson process, Commun. Nonlinear Sci. Numer. Simul. 8, pp. 201–213.
- [333] Laurent, H. (1884). Sur le calcul des dérivées a indices quelconques, Nouv. Ann. Math. 3, pp. 240–252.
- [334] Le Méhauté, A., Tenreiro Machado, J. A., Trigeassou, J. C. and Sabatier,

- J. (eds.) (2005). Fractional Differentiation and its Applications (Ubooks, Neusäß).
- [335] Lederman, C., Roquejoffre, J. M. and Wolanski, N. (2002). Mathematical justification of a nonlinear integro-differential equation for the propagation of spherical flames, C. R., Math., Acad. Sci. Paris 334, pp. 569–574.
- [336] LePage, W. R. (1980). Complex Variables and the Laplace Transform for Engineers (Dover Publ., New York).
- [337] Letnikov, A. V. (1868a). On historical development of differentiation with an arbitrary index, Mat. Sb. 3, pp. 85–112, in Russian.
- [338] Letnikov, A. V. (1868b). Theory of differentiation with an arbitrary index, Mat. Sb. 3, pp. 1–66, (in Russian).
- [339] Letnikov, A. V. (1872). An explanation of the main propositions of differentiation theory with an arbitrary index, Mat. Sb. 6, pp. 413–445, in Russian.
- [340] Lévy, P. P. (1954). Théorie de l'addition des variables aléatoires (Gauthier-Villars, Paris).
- [341] Li, C. and Chen, G. (2004). Chaos in the fractional order chen system and its control, Chaos, Solitons and Fractals 22, pp. 549–554.
- [342] Li, J. R. (2010). A fast time stepping method for evaluating fractional integrals, SIAM J. Sci. Comput. 31, pp. 4696–4714.
- [343] Li, Y. M. and Xu, C. J. (2007). Finite difference/spectral approximations for the time-fractional diffusion equation, J. Comput. Phys. 225, pp. 1533– 1552.
- [344] Liang, S. and Jeffrey, D. J. (2009). Comparison of homotopy analysis method and homotopy perturbation method through an evolution equation, Commun. Nonlin. Sci. Numer. Sim. 14, pp. 4057–4064.
- [345] Liao, S. J. (2003). Beyond Perturbation: Introduction to the Homotopy Analysis Method (Chapman & Hall, Boca Raton).
- [346] Lim, S. C. and Muniandy, S. V. (2004). Stochastic quantization of nonlocal fields, Phys. Lett. A 324, pp. 396–405.
- [347] Liouville, J. (1832). Mémoire sur quelques questions de géométrie et de mécanique, et sur un nouveau genre de calcul pour résoudre ces questions, J. de l'Ecole Roy. Polytéchn. 13, pp. 1–69.
- [348] Liouville, J. (1835). Sur le changement de la variable indépendante dans calcul des différentielles à indices quelconques, J. Ecole Polytech. 15, pp. 17–54.
- [349] Llosa, J. and Vives, J. (1994). Hamiltonian-formalism for nonlocal Lagrangians, J. Math. Phys. 35, pp. 2856–2877.
- [350] Loeb, D. E. (1992). A generalization of the Stirling numbers, Discrete Math. 103, pp. 325–376.
- [351] Love, E. R. (1971). Fractional derivatives of imaginary order, J. London Math. Soc. 2, pp. 241–259.
- [352] Lu, J. F. and Hanyga, A. (2005). Wave field simulation for heterogeneous porous media with singular memory drag force, J. Comput. Phys. 208, pp. 651–674.

- [353] Lubich, C. (1983a). On the stability of linear multistep methods for Volterra convolution equations, IMA J. Numer. Anal. 3, pp. 439–465.
- [354] Lubich, C. (1983b). Runge-Kutta theory for Volterra and Abel integral equations of the second kind, Math. Comput. 41, pp. 87–102.
- [355] Lubich, C. (1985). Fractional linear multistep methods for Abel-Volterra integral equations of the second kind, Math. Comp. 45, pp. 463–469.
- [356] Lubich, C. (1986). Discretized fractional calculus, SIAM J. Math. Anal. 17, pp. 704–719.
- [357] Lubich, C. (1988a). Convolution quadrature and discretized operational calculus. I, Numer. Math. 52, pp. 129–145.
- [358] Lubich, C. (1988b). Convolution quadrature and discretized operational calculus. II, Numer. Math. 52, pp. 413–425.
- [359] Luchko, Y. F. (2012). Operational Method in Fractional Calculus (Springer, Berlin), (in press).
- [360] Luchko, Y. F., Rivero, M., Trujillo, J. J. and Velasco, M. P. (2010). Fractional models, non-locality, and complex systems, Comp. Math. Appl. 59, pp. 1048–1056.
- [361] Lundberg, P. (1903). Approximerad Framställning av Sannolikehetsfunktionen. Återförsäkering av Kollektivrisker (Almqvist & Wicksell, Uppsala).
- [362] Luo, A. C. and Afraimovich, V. S. (eds.) (2011). Long-range Interactions, Stochasticity and Fractional Dynamics (Springer, Berlin).
- [363] M. López-Fernández, C. L. and Schadle, A. (2008). Adaptive, fast, and oblivious convolution in evolution equations, SIAM J. Sci. Comput. 30, pp. 1015–1037.
- [364] Magdziarz, M. and Weron, K. (2006). Anomalous diffusion schemes underlying the Cole-Cole relaxation: The role of the inverse α -stable subordinator, *Phys. A* **367**, pp. 1–6.
- [365] Magin, R. (2006). Fractional Calculus in Bioengineering (Begell House, Redding).
- [366] Magin, R., Abdullah, O., Baleanu, D. and Zhou, X. H. (2008). Anomalous diffusion expressed through fractional order differential operators in the Bloch-Torrey equation, J. Magn. Res. 190, pp. 255–270.
- [367] Magin, R., Duarte Ortigueira, M., Podlubny, I. and Trujillo, J. J. (2011). On the fractional signals and systems, Signal Process. 91, pp. 350–371.
- [368] Magin, R., Feng, X. and Baleanu, D. (2009). Solving the fractional order Bloch equation, Concep. Magn. Res. 34A, pp. 16 –23.
- [369] Mainardi, F. (1997). Fractional calculus: Some basic problems in continuum and statistical mechanics, in A. Carpinteri and F. Mainardi (eds.), Fractals and Fractional Calculus in Continuum Mechanics (Springer, Wien), pp. 291–348.
- [370] Mainardi, F. (2010). Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models (Imperial College Press, London).
- [371] Mainardi, F. (2012). An historical perspective on fractional calculus in linear viscoelasticity, http://arxiv.org/abs/1007.2959 (in press).
- [372] Mainardi, F., Gorenflo, R. and Scalas, E. (2004). A fractional generalization

- of the Poisson process, Vietnam Journal of Mathematics **32** (SI), pp. 53–64.
- [373] Mainardi, F., Luchko, Y. and Pagnini, G. (2001). The fundamental solution of the space-time fractional diffusion equation, Fract. Calc. Appl. Anal. 4, pp. 153–192.
- [374] Mainardi, F., Raberto, M., Gorenflo, R. and Scalas, E. (2000). Fractional calculus and continuous-time finance: The waiting-time distribution, *Phys.* A 287, pp. 468–481.
- [375] Malinowska, A. B. and Torres, D. F. M. (2010). Generalized natural boundary conditions for fractional variational problems in terms of the Caputo derivative, *Comput. Math. Appl.* 59, pp. 3110–3116.
- [376] Maravall, D. (1956). Nuevos tipos de ecuaciones integrales e integrodiferenciales. Nuevos fenómenos de oscilaciones, Rev. R. Acad. Cienc. Exactas, Fis. Nat. 1, pp. 1–151.
- [377] Maravall, D. (1971). Ecuaciones diferenciales lineales no enteras y oscilaciones fraccionarias, Rev. R. Acad. Cienc. Exactas, Fis. Nat. 65, pp. 245– 258.
- [378] Markett, C. (1994). Triple sums and the Riemann zeta function, J. Number Theory 48, pp. 113–132.
- [379] Martinez, C. and Sanz, M. (2001). The Theory of Fractional Powers of Operators (Elsevier, Amsterdam).
- [380] Masoliver, J., Montero, M., Perelló, J. and Weiss, G. H. (2006). The continuous time random walk formalism in financial markets, *Journal of Economic Behavior and Organization* 61, pp. 577–591.
- [381] Mathai, A. M. (1993). A Handbook of Generalized Special Functions for Statistical and Physical Sciences (Clarendon Press, Oxford).
- [382] Mathai, A. M. and Haubold, H. J. (2008). Special Functions for Applied Scientists (Springer, Berlin).
- [383] Mathai, A. M. and Saxena, R. K. (1978). The H-Function with Applications in Statistics and other Disciplines (Wiley & Sons, New York).
- [384] Matignon, D. and Montseny, G. (eds.) (1998). Fractional Differential Systems: Models, Methods, and Applications (SMAI, Paris).
- [385] McBride, A. C. (1979). Fractional Calculus and Integral Transforms of Generalized Functions (Pitman, London).
- [386] McLean, W. and Mustapha, K. (2007). A second-order accurate numerical method for a fractional wave equation, *Numer. Math.* 105, pp. 481–510.
- [387] Medicis, A. and Leroux, R. (1995). Generalized Stirling numbers, convolution formulae and p, q-analogues, Canad. J. Math. 47, pp. 474–499.
- [388] Meerschaert, M. M., Nane, E. and Vellaisamy, P. (2011). The fractional Poisson process and the inverse stable subordinator, *Elect. J. Probab.* 16, pp. 1600–1620.
- [389] Meerschaert, M. M. and Scalas, E. (2006). Coupled continuous time random walks in finance, Phys. A 370, pp. 114–118.
- [390] Meerschaert, M. M. and Scheffler, H. P. (2001). Limit Distributions for Sums of Independent Random Vectors: Heavy Tails in Theory and Practice (Wiley, New York).

- [391] Meerschaert, M. M., Scheffler, H.-P. and Tadjeran, C. (2006). Finite difference methods for two-dimensional fractional dispersion equation, J. Comput. Phys. 211, pp. 249–261.
- [392] Merton, R. C. (1976). Option pricing when underlying stock returns are discontinuous, *Journal of Financial Economics* 3, pp. 125–144.
- [393] Metzler, R., Barkai, E. and Klafter, J. (1999). Anomalous diffusion and relaxation close to thermal equilibrium: A fractional Fokker-Planck equation approach, *Phys. Rev. Lett.* 82, pp. 3563–3567.
- [394] Metzler, R., Glöckle, W. G. and Nonnenmacher, T. F. (1994). Fractional model equation for anomalous diffusion, Phys. A: Stat. Mech. Appl. 211, pp. 13–24.
- [395] Metzler, R. and Klafter, J. (2000). The random walk guide to anomalous diffusion: A fractional dynamics approach, *Phys. Rep.* 339, pp. 1–77.
- [396] Metzler, R. and Klafter, J. (2004). The restaurant at the end of the random walk: Recent developments in the description of anomalous transport by fractional dynamics, J. of Phys. A: Math. General 37, pp. R161–R208.
- [397] Metzler, R. and Nonnenmacher, T. F. (1997). Fractional diffusion: exact representations of spectral functions, J. Phys. A: Math. Gen. 30, pp. 1089– 1093.
- [398] Mikosch, T. (1999). Elementary Stochastic Calculus With Finance in View (World Scientific, Singapore).
- [399] Miller, K. S. and Ross, B. (1989). Fractional difference calculus, in H. M. Srivastava and S. Owa (eds.), Univalent Functions, Fractional Calculus, and Their Applications (Ellis Horwood, Chichester), pp. 139–152.
- [400] Miller, K. S. and Ross, B. (1993). An Introduction to the Fractional Calculus and Fractional Differential Equations (J. Wiley & Sons, New York).
- [401] Miller, R. K. and Feldstein, A. (1971). Smoothness of solutions of Volterra integral equations with weakly singular kernels, SIAM J. Math. Anal. 2, pp. 242–258.
- [402] Miron, R. and Anastasiei, M. (1994). The Geometry of Lagrange Spaces: Theory and Applications (Kluwer, Dordrecht).
- [403] Mittag-Leffler, G. M. (1903). Sur la nouvelle fonction E_{α} , C. R. Acad. Sci. Paris 137, pp. 554–558.
- [404] Mittag-Leffler, G. M. (1904). Sopra la funzione $E_{\alpha}(x)$, Rend. Accad. Lincei 13, pp. 3–5.
- [405] Mittag-Leffler, G. M. (1905). Sur la représentation analytique d'une branche uniforme d'une fonction monogene. Cinquième note, Acta Math. 29, pp. 101–181.
- [406] Momani, S. and Odibat, Z. (2007). Homotopy perturbation method for nonlinear partial differential equations of fractional order, *Phys. Lett. A* 365, pp. 345–350.
- [407] Monje, C. A., Chen, Y., Vinagre, B., Xue, D. and Feliu, V. (2010). Fractional-order Systems and Controls: Fundamentals and Applications (Springer, New York).
- [408] Montroll, E. W. and Scher, H. (1973). Random walks on lattices IV:

- Continuous-time random walks and influence of absorbing boundary conditions, *J. Stat. Phys.* **9**, pp. 101–135.
- [409] Montroll, E. W. and Weiss, G. H. (1965). Random walks on lattices II, J. Math. Phys. 6, pp. 167–181.
- [410] Montseny, G., Audounet, J. and Matignon, D. (2000). Diffusive representation for pseudo-differentially damped nonlinear systems, in A. Isidori, F. Lamnabhi-Lagarrigue and W. Respondek (eds.), *Nonlinear Control, Vol.* 2 (Springer, Berlin), pp. 163–182.
- [411] Mozyrska, D. and Torres, D. F. M. (2010). Minimal modified energy control for fractional linear control systems with the Caputo derivative, *Carpat. J. Math.* 26, pp. 210–221.
- [412] Musiela, M. and Rutkowski, M. (2002). Martingale Methods in Financial Modelling (Springer, New York).
- [413] Muslih, S. I., Agrawal, O. P. and Baleanu, D. (2010a). Fractional Schrödinger equation and its solution, Int. J. Theor. Phys. 49, pp. 1746– 1752.
- [414] Muslih, S. I. and Baleanu, D. (2005). Hamiltonian formulation of systems with linear velocities within Riemann-Liouville fractional derivatives, J. Math. Anal. Appl. 304, pp. 599–606.
- [415] Muslih, S. I. and Baleanu, D. (2007). Fractional Euler-Lagrange equations of motion in fractional space, J. Vib. Contr. 13, pp. 1209–1216.
- [416] Muslih, S. I., Baleanu, D. and Rabei, E. M. (2007). Fractional Hamilton's equations of motion in fractional time, Cent. Eur. J. Phys. 5, pp. 549–557.
- [417] Muslih, S. I., Saddallah, M., Baleanu, D. and Rabei, E. (2010b). Lagrangian formulation of Maxwell's field in fractional d-dimensional space-time, Rom. J. Phys. 55, pp. 659–663.
- [418] Naber, M. (2004). Time fractional Schrödinger equation, J. Math. Phys. 45, pp. 3339–3352.
- [419] Nakhushev, A. M. (2003). Fractional Calculus and its Applications (Fizmatlit, Moscow).
- [420] Nambu, Y. (1973). Generalized Hamiltonian dynamics, Phys. Rev. D 7, pp. 2405–2412.
- [421] Nasiri, S. Y., Sobouti, Y. and Taati, F. (2006). Phase space quantum mechanics-direct, J. Math. Phys. 47, p. 092106.
- [422] Nesterenko, V. V. (1989). Singular Lagrangians with higher derivatives, J. Phys. A: Math. Gen. 22, pp. 1673–1687.
- [423] Nielsen, N. (1965). Handbuch der Theorie der Gammafunktionen (Chelsea Publ. Comp., New York).
- [424] Nieto, J. J. (2010). Maximum principles for fractional differential equations derived from Mittag-Leffler functions, Appl. Math. Letters 23, pp. 1248– 1251.
- [425] Nigmatulin, R. I. (1979). Spatial averaging in the mechanics of heterogeneous and dispersed systems, Int. J. of Multiphase Flow 5, pp. 353–385.
- [426] Nigmatullin, R. R. (2005). Theory of dielectric relaxation in non-crystalline solids: From a set of micromotions to the averaged collective motion in the mesoscale region, *Phys. B: Phys. Cond. Matt.* 358, pp. 201–215.

- [427] Nigmatullin, R. R., Arbuzov, A. A., Salehli, F., Gis, A., Bayrak, I. and Catalgil-Giz, H. (2007). The first experimental confirmation of the fractional kinetics containing the complex power-law exponents: Dielectric measurements of polymerization reaction, *Phys. B: Phys. Cond. Matt.* 388, pp. 418–434.
- [428] Nigmatullin, R. R. and Baleanu, D. (2010). Is it possible to derive Newtoninan equation of motion with memory? J. Theor. Phys. 49, pp. 701–708.
- [429] Nigmatullin, R. R. and Le Méhauté, A. (2005). Is there geometrical/physical meaning of the fractional integral with complex exponent? J. Non-Cryst. Sol. 351, pp. 2888–2899.
- [430] Nolan, J. P. (2012). Stable Distributions: Models for Heavy Tailed Data (Birkhäuser, Boston).
- [431] Odibat, Z., Momani, S. and Erturk, V. S. (2008). Generalized differential transform method: Application to differential equations of fractional order, *Appl. Math. Comput.* 197, pp. 467–477.
- [432] Oldham, K. B. (1969). A new approach to the solution electrochemical problems involving diffusion, Anal. Chem. 41, pp. 1904–1910.
- [433] Oldham, K. B. and Spanier, J. (1974). The Fractional Calculus (Acad. Press, New York).
- [434] OpenMP Architecture Review Board (2010). The OpenMP API specification for parallel programming, URL http://www.openmp.org.
- [435] O'Shaughnessy, L. (1918). Problem # 433, Amer. Math. Monthly 25, pp. 172–173.
- [436] Osler, T. J. (1970). Leibniz rule for fractional derivatives generalized and an application to infinite series, SIAM J. Appl. Math. 18, pp. 658–674.
- [437] Oustaloup, A. (1981). Fractional order sinusoidal oscillators: Optimization and their use in highly linear fm modulation, *IEEE Transactions on Cir*cuits and Systems 28, pp. 1007–1009.
- [438] Oustaloup, A. (1983). Systèmes asservis linéaires d'ordre fractionnaire: Théorie et pratique (Masson, Paris).
- [439] Oustaloup, A. (1991). La Commande CRONE: Commande Robuste d'Ordre Non Entier (Hermès, Paris).
- [440] Oustaloup, A. (1995). La Dérivation Non Entière. Théorie, Synthèse et Applications (Hermès Sci., Paris).
- [441] Oustaloup, A., Cois, O., Lanusse, P., Melchior, P., Moreau, X. and Sabatier, J. (2006). The CRONE approach: Theoretical developments and major applications, in *Proceedings of the 2nd IFAC Workshop on Fractional Dif*ferentiation and its Applications (Porto), pp. ID10.3182/20060719–3–PT– 4902.00059.
- [442] Oustaloup, A. and Mathieu, B. (1999). La commande CRONE: du scalaire au multivariable (Hermés, Paris).
- [443] Oustaloup, A., Sabatier, J. and Lanusse, P. (1999). From fractal robustness to CRONE control, Fract. Calc. Appl. Anal. 2, pp. 1–30.
- [444] Ozdemir, N., Agrawal, O. P., Karadeniz, D. and Iskender, B. B. (2009). Fractional optimal control problem of an axis-symmetric diffusion-wave propagation, *Phys. Scr.* T136, p. 014024.

- [445] Pachepsky, Y., Benson, D. A. and Rawls, W. (2000). Simulating scale-dependent solute transport in soils with the fractional advective-dispersive equation, Soil Sci. Soc. Am. J. 64, pp. 1234–1243.
- [446] Pakes, A. G. (1998). Mixture representations for symmetric generalized Linnik laws, Stat. Probab. Lett. 37, pp. 213–221.
- [447] Perron, O. (1919a). Über Integration von gewöhnlichen Differentialgleichungen durch Reihen, Sitz.-Ber. Heidelberger Akademie der Wissenschaften A: Math.-Nat. Klasse Abhandlung 2.
- [448] Perron, O. (1919b). Über Integration von gewöhnlichen Differentialgleichungen durch Reihen, II, Sitz.-Ber. Heidelberger Akademie der Wissenschaften A: Math.-Nat. Klasse Abhandlung 8.
- [449] Perron, O. (1919c). Über Integration von gewöhnlichen Differentialgleichungen durch Reihen, III, Sitz.-Ber. Heidelberger Akademie der Wissenschaften A: Math.-Nat. Klasse Abhandlung 12.
- [450] Perron, O. (1920). Über Integration partieller Differentialgleichungen durch Reihen, Sitz.-Ber. Heidelberger Akademie der Wissenschaften A: Math.-Nat. Klasse Abhandlung 9.
- [451] Picozzi, S. and West, B. J. (2002). Fractional Langevin model of memory in financial time series, *Phys. Rev. E* **66**, pp. 046118–046129.
- [452] Platonov, M. L. (1997). The relations between generalized Stirling numbers constructed on different basis, Combinatorial and Asymptotic Analysis. Krasnoyarsk. Gos. Univ. 2, pp. 142–152.
- [453] Podlubny, I. (1999). Fractional Differential Equations (Acad. Press, San Diego).
- [454] Podlubny, I. (2000). Matrix approach to discrete fractional calculus, Fract. Calc. Appl. Anal. 3, pp. 359–386.
- [455] Podlubny, I. (2009). Mittag-Leffler function, URL http://www.mathworks.com/matlabcentral/fileexchange/8738-mittag-leffler-function.
- [456] Podlubny, I., Petras, I., Vinagre, B., Chen, Y. Q., O'Leary, P. and Dorcak, L. (2003). Realization of fractional order controllers, *Acta Montan. Slovaca* 8, pp. 233–235.
- [457] Post, E. L. (1919). Discussion of the solution of $(d/dx)^{1/2}y = y/x$ (problem \sharp 433), Amer. Math. Monthly **26**, pp. 37–39.
- [458] Prudnikov, A. P., Brychkov, Y. A. and Marichev, O. I. (1986). Integrals and Series, Vols. I, II and III (Gordon and Breach, New York).
- [459] Pskhu, A. V. (2005). Partial Differential Equations of Fractional Order (Nauka, Moscow).
- [460] Quirós, F., Rossi, J. D. and Vázquez, J. L. (2002). Complete blow-up and thermal avalanche for heat equations with nonlinear boundary conditions, Comm. Part. Diff. Equ. 27, pp. 395–424.
- [461] Rabei, E. M. and Ababenh, B. S. (2008). Hamilton-Jacobi fractional mechanics, J. Math. Anal. Appl. 344, pp. 799–805.
- [462] Rabei, E. M., Ajlounid, A. W. and Gassib, H. B. (2006). Quantization of Brownian motion, Int. J. Theor. Phys. 45, pp. 1619–1629.
- [463] Rabei, E. M., Alhalholy, T. S. and Rousan, A. A. (2004a). Potentials of arbitrary forces with fractional derivatives, J. Mod. Phys. 19, pp. 3083– 3092.

- [464] Rabei, E. M., Alhalholy, T. S. and Taani, A. A. (2004b). On Hamiltonian formulation of non-conservative systems, *Turkish J. Phys.* 28, pp. 213–221.
- [465] Rabei, E. M., Almayteh, I., Muslih, S. I. and Baleanu, D. (2008). Hamilton-Jacobi formulation of systems within Caputo's fractional derivative, *Phys. Scr.* 77, p. 015101.
- [466] Rabei, E. M., Altarazi, I. A., Muslih, S. I. and Baleanu, D. (2009a). Fractional WKB approximation, Nonlinear Dyn. 57, pp. 171–175.
- [467] Rabei, E. M., Muslih, S. I. and Baleanu, D. (2010). Quantization of fractional systems using WKB approximation, Commun. Nonlin. Sci. Numer. Sim. 15, pp. 807–811.
- [468] Rabei, E. M., Nawafleh, K. I., Hijjawi, R. S., Muslih, S. I. and Baleanu, D. (2007). The Hamilton formalism with fractional derivatives, J. Math. Anal. Appl. 327, pp. 891–897.
- [469] Rabei, E. M., Rawashdeh, I. M., Muslih, S. I. and Baleanu, D. (2011). Hamilton-Jacobi formulation for systems in terms of Riesz's fractional derivatives, Int. J. Theor. Phys. 50, pp. 1569–1576.
- [470] Rabei, E. M., Tarawneh, D. M., Muslih, S. I. and Baleanu, D. (2009b). Heisenberg's equations of motion with fractional derivatives, J. Vibr. Contr. 13, pp. 1239–1247.
- [471] Raberto, M., Scalas, E. and Mainardi, F. (2002). Waiting-times and returns in high-frequency financial data: An empirical study, *Phys. A* 314, pp. 749– 755.
- [472] Rabinowitz, P. (1979). On avoiding the singularity in the numerical integration of proper integrals, *BIT* **19**, pp. 104–110.
- [473] Rabotnov, Y. N. (1969). Creep Problems in Structural Members (North-Holland, Amsterdam).
- [474] Rabotnov, Y. N. (1973). On the Use of Singular Operators in the Theory of Viscoelasticity (Lecture Notes for the CISM course on Rheology, Bologna).
- [475] Répaci, A. (1990). Nonlinear dynamical systems: On the accuracy of Adomian's decomposition method, Appl. Math. Lett. 3, 4, pp. 35–39.
- [476] Repin, O. N. and Saichev, A. I. (2000). Fractional Poisson law, Radiophys. Quantum Electron. 43, pp. 738–741.
- [477] Richmond, R. and Merlini, D. (1997). Stirling numbers for complex arguments, SIAM J. Discrete Math. 10, pp. 73–82.
- [478] Riemann, B. (1876). Versuch einer allgemeinen Auffasung der Integration und Differentiation, in *Gesammelte Werke* (Teubner, Leipzig), pp. 331–344.
- [479] Riesz, M. (1936). L'intégrale de Riemann-Liouville et solution invariantive du probléme de cauchy pour l'equation des ondes, C. R. Congrès Intern. Math. 2, pp. 44–45.
- [480] Riesz, M. (1949). L'intégrale de Riemann-Liouville et le problème de Cauchy, Acta Math. 81, pp. 1–223.
- [481] Riewe, F. (1996). Nonconservative Lagrangian and Hamiltonian mechanics, Phys. Rev. E 53, pp. 1890–1899.
- [482] Riewe, F. (1997). Mechanics with fractional derivatives, Phys. Rev. E 55, pp. 3581–3592.
- [483] Riordan, J. (1979). Combinatorial Identities (Krieger Publ., Huntington).

- [484] Rivero, M., Rodríguez-Germá, L., Trujillo, J. J. and Velasco, M. P. (2010). Fractional operators and some special functions, *Comp. Math. Appl.* 59, pp. 1822–1834.
- [485] Rivero, M., Trujillo, J. J., Vázquez, L. and Velasco, M. P. (2011). Fractional dynamic of populations, Appl. Math. Comp. 218, pp. 1089–1095.
- [486] Rivero, M., Trujillo, J. J. and Velasco, M. P. (2009). On deterministic fractional models, in D. Baleanu, Z. B. Güvenç and J. A. Tenrreiro (eds.), New Trends in Nanotechnology and Fractional Calculus Applications (Springer-Verlag, Berlin), pp. 123–150.
- [487] Rodríguez-Germá, L., Trujillo, J. J., Vázquez, L. and Velasco, M. P. (2008). Fractional calculus framework to avoid singularities of differential equations, Fract. Calc. Appl. Anal. 11, pp. 431–441.
- [488] Roman, H. E. and Alemany, P. A. (1994). Continuous-time random walks and the fractional diffusion equation, J. Phys. A 27, pp. 3407–3410.
- [489] Ross, B. (ed.) (1974). Fractional Calculus and its Applications: Proceedings of the International Conference, New Haven, June 1974 (Springer, New York).
- [490] Ross, B. (1977). The development of fractional calculus 1695–1900, Hist. Math. 4, pp. 75–89.
- [491] Rossikhin, Y. A. and Shitikova, M. V. (2007). Comparative analysis of viscoelastic models involving fractional derivatives of different orders, *Fract. Calc. Appl. Anal.* 10, pp. 111–121.
- [492] Rucinski, A. and Voigt, B. (1990). A local limit theorem for generalized Stirling numbers, Rev. Roumaine Math. Pures Appl. 35, pp. 161–172.
- [493] Saad, Y. (2003). Iterative Methods for Sparse Linear Systems (SIAM, Philadelphia).
- [494] Saad, Y. and Schultz, M. H. (1986). GMRES: A generalized minimal residual algorithm for solving nonsymmetric linear systems, SIAM J. Sci. Statist. Comput. 7, pp. 856–869.
- [495] Sabatier, J., Agrawal, O. P. and Tenreiro Machado, J. A. (eds.) (2007). Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering (Springer, Berlin).
- [496] Sabatier, J., Melchior, P., Tenreiro Machado, J. A. and Vinagre, B. (eds.) (2008). Fractional order systems: Applications in modelling, identification and control, J. Europ. Syst. Autom. 42, pp. 622–1051.
- [497] Sadati, S. J., Baleanu, D., Ranjbar, A., Ghaderi, R. and Abdeljawad, T. (2010). Mittag-Leffler stability theorem for fractional nonlinear systems with delay, Abstr. Appl. Anal. 2010, pp. 108651–108657.
- [498] Saichev, A. I. and Zaslavsky, G. M. (1997). Fractional kinetic equations: solutions and applications, Chaos 7, pp. 753–764.
- [499] Samko, S. (2002). Hypersingular Integrals and Their Applications (Taylor and Francis, New York).
- [500] Samko, S. G. (1967). Solution of generalized Abel equation by means of an equation with Cauchy kernel, Soviet Math. Dokl. 6, pp. 1259–1262, in Russian.

- [501] Samko, S. G., Kilbas, A. A. and Marichev, O. I. (1993). Fractional Integrals and Derivatives. Theory and Applications (Gordon and Breach, Yverdon).
- [502] Samorodnitsky, G. and Taqqu, M. S. (1994). Stable non-Gaussian Random Processes: Stochastic Models with Infinite Variance (Chapman & Hall, London).
- [503] San José Martínez, F., Pachepsky, Y. A. and Rawls, W. J. (2007). Fractional advective-dispersive equation as a model of solute transport in porous media, in J. Sabatier, O. P. Agrawal and J. A. Tenreiro Machado (eds.), Advances in Fractional Calculus Theoretical Developments and Applications in Physics and Engineering (Springer, Berlin), pp. 199–212.
- [504] San José Martínez, F., Pachepsky, Y. A. and Rawls, W. J. (2010). Modelling solute transport in soil columns using advective-dispersive equations with fractional spatial derivatives, Adv. Engineer. Soft. 41, pp. 4–8.
- [505] Sanz-Serna, J. M. (1988). A numerical method for a partial integrodifferential equation, SIAM J. Numer. Anal. 25, pp. 319–327.
- [506] Sato, K.-I. (1999). Lévy Processes and Infinitely Divisible Distributions (Cambridge Univ. Press, Cambridge).
- [507] Scalas, E. (2006). The application of continuous-time random walks in finance and economics, *Phys. A* 362, pp. 225–239.
- [508] Scalas, E. (2011). A class of CTRWs: Compound fractional Poisson processes, in R. Metzler (ed.), Fractional Dynamics, chap. 15 (World Scientific, Singapore), pp. 351–372.
- [509] Scalas, E., Gorenflo, R. and Mainardi, F. (2000). Fractional calculus and continuous-time finance, Phys. A 284, pp. 376–384.
- [510] Scalas, E., Gorenflo, R. and Mainardi, F. (2004a). Uncoupled continuoustime random walks: Solutions and limiting behavior of the master equation, *Phys. Rev. E* 69, p. 011107.
- [511] Scalas, E., Gorenflo, R., Mainardi, F. and Meerschaert, M. M. (2004b). Speculative option valuation and the fractional diffusion equation, in A. Le Méhauté, J. A. Tenreiro Machado, J. C. Trigeassou and J. Sabatier (eds.), Proceedings of the 1st IFAC Workshop on Fractional Differentiation and its Applications (ENSEIRB, Bordeaux).
- [512] Scher, H. and Lax, M. (1973a). Stochastic transport in a disordered solid I. Theory, Phys. Rev. B 7, pp. 4491–4502.
- [513] Scher, H. and Lax, M. (1973b). Stochastic trasport in a disordered solid II. Impurity conduction, Phys. Rev. B 7, pp. 4506–4519.
- [514] Schiesser, W. E. (1991). The Numerical Method of Lines (Acad. Press, San Diego).
- [515] Schmidt, A. and Gaul, L. (2006). On a critique of a numerical scheme for the calculation of fractionally damped dynamical systems, *Mech. Res.* Commun. 33, pp. 99–107.
- [516] Schoutens, W. (2003). Lévy Processes in Finance (Wiley, New York).
- [517] Schumer, R., Meerschaert, M. M. and Baeumer, B. (2009). Fractional advection-dispersion equations for modeling transport at the earth surface, J. Geophysical Research 114, pp. F00A07-F00A21.
- [518] Schwartz, L. M. (1950). Théorie des Distributions (Hermann, Paris).

- [519] Seiberg, N., Susskind, L. and Toumbas, T. (2000). Space/time noncommutativity and causality, J. H. Energ. Phys. 6, p. 021.
- [520] Shlesinger, M. F. (1996). Random processes, in G. L. Trigg (ed.), Encyclopedia of Applied Physics, Vol. 16 (VCH, New York), pp. 45–70.
- [521] Shlesinger, M. F., Zaslavsky, G. M. and Klafter, J. (1993). Strange kinetics, Nature 363, pp. 31–37.
- [522] Singh, S. J. and Chatterjee, A. (2006). Galerkin projections and finite elements for fractional order derivatives, *Nonlinear Dyn.* 45, pp. 183–206.
- [523] Singh-Chandel, R. S. (1977). Generalized Stirling numbers and polynomials, Publ. Inst. Math., Beograd (N. S.) 22, pp. 43–48.
- [524] Sinha, V. P. and Dhawan, G. K. (1969). On generalized Stirling numbers and polynomials, Riv. Math. Univ. Parma 10, pp. 95–100.
- [525] Sixdeniers, J. M., Penson, K. A. and Solomon, A. L. (2001). Extended Bell and Stirling numbers from hypergeometric exponentiation, J. Integer Seq. 4, pp. 1–11.
- [526] Sneddon, I. N. (1957). Elements of Partial Differential Equations (McGraw-Hill, New York).
- [527] Sneddon, I. N. (1966). Mixed boundary value problems in potential theory (North-Holland, Amsterdam).
- [528] Sneddon, I. N. (1979). The use of operators of fractional integration in applied mathematics, (Polish Acad. Sci., Warszawa-Poznan).
- [529] Sokolov, I. M., Chechkin, A. V. and Klafter, J. (2004). Distributed order fractional kinetics, Acta Physica Polonica B 35, pp. 1323–1341.
- [530] Srivastava, H. M. (1968). On an extension of the Mittag-Leffler function, Yokohama Math. J. 16, pp. 77–88.
- [531] Srivastava, H. M. and Buschman, R. G. (1977). Convolution Integral Equations with Special Function Kernels (Wiley & Sons, New York).
- [532] Srivastava, H. M., Gupta, K. C. and Goyal, S. P. (1982). The H-Functions of One and Two Variables with Applications (South Asian Publ., New Delhi and Madras).
- [533] Srivastava, H. M. and Kashyap, B. R. (1982). Special Functions in Queuing Theory and Related Stochastic Processes (Academic Press, New York).
- [534] Stanislavsky, A. A. (2004). Probability interpretation of the integral of fractional order, *Theor. Math. Phys.* 138, pp. 418–431.
- [535] Stanislavsky, A. A. (2006). Hamiltonian formalism of fractional systems, Eur. Phys. J. B 49, pp. 93–101.
- [536] Stenger, F. (1976). Approximation via Whittaker's cardinal function, J. Approx. Theory 17, pp. 222–240.
- [537] Stirling, J. (1730). Methodus Differentialis, sive Tractatus de Summatione et Interpolatione Serium Infinitarum (Gul. Bowyer, London).
- [538] Su, H., Chen, W., Li, C. and Chen, Y. (2010). Fractional differential models for anomalous diffusion, Phys. A: Stat. Mech. Appl. 389, pp. 2719–2724.
- [539] Sun, H. G., Chen, W. and Chen, Y. Q. (2009). Variable-order fractional differential operators in anomalous diffusion modeling, *Phys. A* 388, pp. 4586–4592.

- [540] Tarasov, V. E. (2005). Fractional hydrodynamics equations for fractal media, Anal. Phys. 318, pp. 286–307.
- [541] Tarasov, V. E. (2006). Fractional statistical mechanics, Chaos 16, pp. 033108-033114.
- [542] Tarasov, V. E. (2010). Fractional dynamics of relativistic particle, Int. J. Theor. Phys. 49, pp. 293–303.
- [543] Tarasov, V. E. (2011). Fractional Dynamics: Application of Fractional Calculus to Dynamics of Particles, Fields and Media (Springer, Berlin).
- [544] Tarasov, V. E. and Zaslavsky, G. M. (2006). Nonholonomic constraints with fractional derivatives, J. Phys. A: Math. Gen. 39, pp. 9797–9815.
- [545] Tarasov, V. E. and Zaslavsky, G. M. (2008). Conservation laws and Hamilton's equations for systems with long-range interaction and memory, Commun. Nonlin. Sci. Numer. Sim. 13, pp. 1860–1878.
- [546] Tatar, N. (2010). Existence results for an evolution problem with fractional nonlocal conditions, Comput. Math. Appl. 60, pp. 2971–2982.
- [547] Tatar, N. E. (2011). Mild solutions for a problem involving fractional derivatives in the nonlinearity and in the non-local conditions, Adv. Diff. Equ. ID2011:18, pp. 1–12.
- [548] Tatari, M. and Dehghan, M. (2007). On the convergence of He's variational iteration method, J. Comput. Appl. Math. 207, pp. 121–128.
- [549] Tenreiro Machado, J. A. (2001). Discrete-time fractional-order controllers, J. Fract. Calc. Appl. Anal. 4, pp. 47–66.
- [550] Tenreiro Machado, J. A. (ed.) (2002). Fractional order systems, Nonlinear Dyn. 29.
- [551] Tenreiro Machado, J. A. (2003). A probabilistic interpretation of the fractional-order differentiation, Fract. Calc. Appl. Anal. 6, pp. 73–80.
- [552] Tenreiro Machado, J. A. (2009). Fractional derivatives: probability interpretation and frequency response of rational approximations, Comm. Nonlin. Sci. Num. Simul. 14, pp. 3492–3497.
- [553] Tenreiro Machado, J. A., Kiryakova, V. and Mainardi, F. (2011). Recent history of fractional calculus, *Commun. Nonlin. Sci. Numer. Sim.* 16, pp. 1140–1153.
- [554] The MathWorks, Inc. (2011). Matlab, URL http://www.mathworks.com/ products/matlab/.
- [555] The R Project for Statistical Computing (2011). R, URL http://www.r-project.org/.
- [556] Toader, S. and Toader, G. (1998). Generalized Stirling numbers, Studia Univ. Babes-Bolyai Math. 33, pp. 50–53.
- [557] Tomowski, Z., Hilfer, R. and Srivastava, H. M. (2010). Fractional and operational calculus with generalized fractional derivative operators and Mittag-Leffler type functions, *Int. Trans. Spec. Func.* 21, pp. 797–814.
- [558] Torvik, P. J. and Bagley, R. L. (1984). On the appearance of the fractional derivative in the behavior of real materials, J. Appl. Mech. 51, pp. 294–298.
- [559] Tricaud, C. and Chen, Y. Q. (2009). Time-optimal control of fractional dynamic systems, Proceedings of the IEEE Conference on Decision and Control, pp. 5027–5032.

- [560] Trujillo, J. J. and Baleanu, D. (eds.) (2011). Fractional models and their applications, Advances in Difference Equations.
- [561] Uchaikin, V. V. (2008). Method of Fractional Derivatives (Artishok Press, Ulyanovsk).
- [562] Umarov, S. and Gorenflo, R. (2005). Cauchy and non-local multi-point problems for distributed order pseudo-differential equations, Part I, J. Anal. Appl. 24, pp. 449–466.
- [563] Usero, D. and Vázquez, L. (2003). Fractional derivative. A new formulation for damped systems, in L. Vázquez, R. S. MacKay and M. P. Zorzano (eds.), *Localization and Energy Transfer in Nonlinear Systems* (World Scientific, Singapore), pp. 296–303.
- [564] Vacaru, S. (2008). Finsler and Lagrange geometries in Einstein and string gravity, Int. J. Geom. Methods. Mod. Phys. 5, pp. 473–511.
- [565] Vacaru, S. (2010a). Fractional dynamics from Einstein gravity, general solutions, and black holes, Tech. rep., preprint:hep-th/1004.0628.
- [566] Vacaru, S. (2010b). Fractional nonholonomic Ricci flows, Tech. rep., preprint:hep-th/1004.0625.
- [567] Vacaru, S., Stavrinos, P., Gaburov, E. and Gonţa, D. (2006). Clifford and Riemann-Finsler structures in geometric mechanics and gravity, Preprint:gr-qc/050802 URL http://www.mathem.pub.ro/dgds/mono/ va-t.pdf.
- [568] Vainikko, G. and Uba, P. (1981). A piecewise polynomial approximation to the solution of an integral equation with weakly singular kernel, J. Austral. Math. Soc., Ser. B 22, pp. 431–438.
- [569] Vasilév, V. V. and Simak, L. A. (2008). Fractional Calculus and Approximation Methods in Modeling of Dynamic Systems (Academic Press, Kiev).
- [570] Vázquez, L. (2003). Fractional diffusion equations with internal degrees of freedom, J. Comp. Math. 21, pp. 491–494.
- [571] Vázquez, L. and Vilela Mendes, R. (2003). Fractionally coupled solutions of the diffusion equation, Appl. Math. Comp. 141, pp. 125–130.
- [572] Wagner, C. G. (1996). Generalized Stirling and Lah numbers, Discrete Math. 160, pp. 199–218.
- [573] Walker, H. F. (1988). Implementation of the GMRES method using Householder transformations, SIAM J. Sci. Statist. Comput. 9, pp. 152–163.
- [574] Wang, J. and Zhou, Y. (2011). A class of fractional evolution equations and optimal controls, Nonlinear Anal.: Real World Appl. 12, pp. 262–272.
- [575] Wang, S. W. and Xu, M. Y. (2007). Generalized fractional Schrödinger equation with space-time fractional derivatives, J. Math. Phys. 48, p. 043502.
- [576] Weilbeer, M. (2006). Efficient Numerical Methods for Fractional Differential Equations and their Analytical Background, Ph.D. thesis, Technische Universität Braunschweig.
- [577] Weyl, H. (1917). Bemerkungen zum Begriff des Differentialquotienten gebrochener Ordnung, Zürich Naturf. Ges. 62, pp. 296–302.
- [578] Weyl, H. (1935). Geodesic fields in the calculus of variation for multiple integrals, Ann. Math. 36, pp. 607–629.

- [579] Whittaker, E. T. (1915). On the functions which are represented by the expansions of the interpolation-theory, *Proc. Roy. Soc. Edinburgh* 35, pp. 181–194.
- [580] Whittaker, J. M. (1935). Interpolatory Function Theory (Cambridge Univ. Press, Cambridge).
- [581] Widder, D. V. (1941). The Laplace Transform (Princeton Univ. Press, Princeton).
- [582] Williams, D. (1991). Probability with Martingales (Cambridge Univ. Press, Cambridge).
- [583] Wilmott, P. (1998). Derivatives: The Theory and Practice of Financial Engineering (Wiley, New York).
- [584] Wolfram Research, Inc. (2011). Mathematica, URL http://www.wolfram.com/mathematica/.
- [585] Wu, J. (2006). Lower bounds for an integral involving fractional Laplacians and the generalized Navier-Stokes equations in Besov spaces, *Comm. Math. Phys.* 263, pp. 803–831.
- [586] Wyss, W. (1986). The fractional diffusion equation, J. Math. Phys 27, pp. 2782–2795.
- [587] Wyss, W. and Schneider, W. R. (1989). Fractional diffusion and wave equations, J. Math. Phys 30, pp. 569–580.
- [588] Xue, D., Chen, Y. and Atherton, D. P. (2008). Linear Feedback Control: Analysis and Design with MATLAB (Society for Industrial Mathematics, Philadelphia).
- [589] Ye, H. and Ding, Y. (2009a). A fractional-order differential equation model of HIV infection of CD4+ T-cells, Math. Comput. Model. 50, pp. 386–392.
- [590] Ye, H. and Ding, Y. (2009b). Nonlinear dynamics and chaos in a fractional-order HIV model, Math. Probl. Eng. ID378614, pp. 1–12.
- [591] Yuan, L. and Agrawal, O. P. (2002). A numerical scheme for dynamic systems containing fractional derivatives, J. Vibration Acoustics 124, pp. 321–324.
- [592] Yuste, S. B. and Acedo, L. (2005). An explicit finite difference method and a new von Neumann type stability analysis for fractional diffusion equations, SIAM J. Numer. Anal. 42, pp. 1862–1874.
- [593] Zaslavsky, G. M. (1994). Fractional kinetic equation for Hamiltonian chaos, Phys. D 76, pp. 110–122.
- [594] Zaslavsky, G. M. (2002). Chaos, fractional kinetics, and anomalous transport, Phys. Rep. 371, pp. 461–580.
- [595] Zemanian, A. H. (1968). Generalized Integral Transformations (Wiley, New York).
- [596] Zhang, C. (1994). Quelques études en théorie des équations fonctionnelles et en analyse combinatoire, Ph.D. thesis, Université Louis Pasteur Strasbourg.
- [597] Zhang, X., Lv, M., Crawford, J. and Young, I. M. (2007). The impact of boundary on the fractional advection—dispersion equation for solute transport in soil: Defining the fractional dispersive flux with the Caputo derivatives, Adv. Water Resour. 30, pp. 1205–1217.

- [598] Zhang, Y. (2009). Finite difference approximations for space-time fractional partial differential equation, J. Numer. Math. 17, pp. 319–326.
- [599] Zoia, A., Rosso, A. and Kardar, M. (2007). Fractional Laplacian in bounded domains, Phys. Rev. E 76, pp. 021116–021126.
- [600] Zolotarev, V. M. (1986). One-dimensional Stable Distributions (Am. Math. Soc., Providence).
- [601] Zygmund, A. (1934). Some points in the theory of trigonometric and power series, Trans. Amer. Math. Soc. 36, pp. 586-617.
- [602] Zygmund, A. (1945). A theorem on fractional derivatives, Duke Math J. 3, pp. 455–464.

This page intentionally left blank

Index

 α -stable Lévy Process, 316 α -stable distribution, 299, 314, 315 β -stable subordinator, 305, 316

absolute continuity of measures, 328 Adams-Bashforth method, 62, 112, 335 Adams-Bashforth-Moulton method, 63, 100, 108, 335 Adams-Moulton method, 60, 112, 335 Adomian decomposition, 69

ask limit order, 317

backward differentiation formula, 57, 346
Bagley-Torvik equation, 77
Basset equation, 77
Bell function, 181
best ask, 318
best bid, 317
Beta function, 7
bid limit order, 317
bid-ask spread, 318
binomial coefficients, 7

Caputo fractional derivative, 16, 19, 43
Cauchy problem for the space-time fractional diffusion equation, 311 characteristic polynomial, 50 collocation method, 136

complex powers and Stirling functions, 182 compound fractional Poisson process, 310, 316 confluent hypergeometric function, 8 consistency, 66 continuous-double auction, 317 continuous-time random walks, 289, 290, 306, 311, 314, 315 convergence, 66 convolution operator, 3, 4 convolution quadrature, 52, 53 convolution relation, 182 convolution weights, 52–54 corrector, 62 counting process, 291

decoupled continuous-time random walks, 306 delay, 246, 281 diffusion-wave equation, 83 Digamma function, 6, 149, 236 distributed-order equations, 82

economic growth model, 333
equivalent martingale measure, 328
Euler Gamma function, 5
Euler-Lagrange equations
discrete fractional, 253
fractional, 241–265

Faddeev-Jackiw formalism, 263

finite difference operator, 23, 39, 191 generalized fractional, 191, 206, 217, 234 Grünwald-Letnikov, 39, 49 left-sided, 39 right-sided, 39 finite memory principle, 94 Finsler geometry, 255 Fourier convolution, 2 Fourier transform, 2, 3 fractional Poisson process, 315 fractional compound Poisson process, 314 fractional convolution quadrature, 54, 66 fractional diffusion, 289, 299, 311 fractional diffusion equation, 315 fractional Euler-Lagrange equations, 241 - 265fractional Hamiltonian dynamics, 266 - 287fractional integration by parts, 15, 22 fractional Poisson process, 303, 305, 310, 314 g-Liouville fractional derivative, 37 q-Liouville fractional integral, 36 q-Riemann-Liouville fractional derivative, 37 q-Riemann-Liouville fractional integral, 36 Gauss hypergeometric function, 8 generalized differential transform method, 72

Hadamard fractional derivative, 31, 185, 229 Hadamard fractional integral, 30, 185 Hadamard type fractional derivative, 31, 186 Hadamard type fractional integral, 31, 186 Hamiltonian dynamics fractional, 266–287 homotopy analysis method, 72 homotopy perturbation method, 72 infinitely divisible distribution, 290, 305, 313, 314 insurance ruin, 332 intra-day hedging, 325 inverse β stable subordinator, 316 Laplace operator, 3 Laplace transform, 4, 5 left-sided difference operator, 39 linear multistep method, 50, 66, 104 Liouville fractional derivative, 20, 38, 169, 174, 213, 224 Liouville fractional integral, 20, 172 log-normal distribution, 330 log-price, 321 log-return, 321 logarithmic return, 321 Marchaud fractional derivative, 23, 40, 170, 175 market order, 318 martingale, 328 generalized exponential function, 24 maturity, 326 generalized hypergeometric series, 9 method of lines horizontal, 127 generating function for linear multistep methods, 51 vertical, 124 for Stirling functions, 145, 181, 201 mid-price, 318 for Stirling numbers, 141 Mittag-Leffler function, 9, 10, 358 Monte Carlo simulations, 332 Grünwald-Letnikov finite difference, 39, 49 multi-order fractional differential left-sided, 39 system, 78 right-sided, 39 multi-term fractional differential Grünwald-Letnikov fractional equation, 78 derivative, 39, 40, 48, 49 multiple sum representation, 151, 152 Index 399

Nambu mechanics, 276

option pricing, 325

plain-vanilla European call, 325
Pochhammer symbol, 5, 8
polygamma function, 7
predictor, 61
predictor-corrector method, 62, 335
probability of ruin, 332
product rectangle formula, 44
product trapezoidal formula
for Riemann-Liouville derivatives,
47
for Riemann-Liouville integrals, 46

random walk, 291
recurrence relation for Stirling
functions, 143, 149, 166, 167, 176,
199, 211
representation of fractional difference
operators by Stirling functions,
216, 234
representation of fractional operators
by Stirling functions, 186, 189, 229
representation of Stirling functions,
146, 158, 162, 168–172, 174, 175,
179, 193, 196, 203, 206, 213, 224,
226
Riemann zeta function, 152
Riemann-Liouville fractional

sampling theorem and applications, 225

derivative, 10, 11, 42, 146, 229

right-sided difference operator, 39

10, 42, 161, 229

Rothe's method, 127

risk-free interest rate, 324

Riemann-Liouville fractional integral,

Schrödinger equations, 260 semi-Markov process, 311 semigroup property, 33, 37, 39 short memory principle, 94 space-time fractional diffusion equation, 311, 315 spectral method, 137 stability, 66 stable distribution, 295–297, 313, 314 starting quadrature, 54 Stirling functions and complex powers, 182 applications of, 216, 219, 221, 222, 232, 236 asymptotic behavior, 180 connections between — of first and second kind, 223 differentiability, 164, 185 generalized orthogonality, 227 generating function, 145, 181, 201 of the first kind for both parameters, 153, 164, 166 for the first parameter, 143, 145, 151, 227, 230, 233 for the second parameter, 163, 223 of the second kind for both parameters, 205, 213 for the first parameter, 168, 172, 173, 176, 179, 185, 223, 226 for the second parameter, 191, 193, 199, 216, 217, 219 recurrence relation, 143, 149, 166, 167, 176, 199, 211 representation of, 146, 158, 162, 168–172, 174, 175, 179, 193, 196, 203, 206, 213, 224, 226 representation of fractional difference operators by —, 216 representation of fractional operators by —, 186, 229,

Stirling numbers
generating function, 141
of the first kind, 141
of the second kind, 141, 191
strike price, 326
supersymmetry, 278

234

terminal value problem, 74 tick-by-tick return, 322 tick-by-tick trade, 317

variable-order equations, 82

variational iteration method, 71

Yuan-Agrawal method, 72

zero-coupon bond, 324

Advisory Board

Valentin Afraimovich

San Luis Potosi University, IICO-UASLP Av. Karakorum 1470, Lomas 4a Seccion San Luis Potosi, SLP 78210, Mexico Email: valentin@cactus.iico.uaslp.mx

Maurice Courbage

Universite Paris 7-Denis Diderot/L.P.T.M.C. Tour 24-14.5eme etage, 4, Place Jussieu 75251, Paris Cedex 05, France Email: courbage@ccr.jussieu.fr

Ben-Jacob Eshel

School of Physics and Astronomy, Tel Aviv University, 69978 Tel Aviv, Israel Email: eshel@tamar.tau.ac.il

Bernold Fiedler

Freie Universität Berlin, Institut für Mathematik I, Arnimallee 2-6 14195 Berlin, Germany E-mail: fiedler@math.fu-berlin.de

James A. Glazier

Biocomplexity Institute
Department of Physics
Indiana University, Bloomington
Bloomington, IN 47405-7105, USA
Email: glazier@indiana.edu

Nail Ibragimov

Department of Mathematics, IHN, Blekinge Institute of Technology, S-371 79 Karlskrona, Sweden Email: nib@bth.se

Anatoly Neishtadt

Space Research Institute Russian Academy of Sciences Profsoyuznaya 84/32 Moscow 117997, Russia E-mail: aneishta@iki.rssi.ru

Leonid Shilnikov

Research Institute for Applied Mathematics & Cybernetics 10 Ul'yanov Street Nizhny Novgorod 603005, Russia Email: diffequ@unn.ac.ru

Michael Shlesinger

Office of Naval Research, Physics and Chemistry Department 800 North Quincy Str, Room 503 Arlington, VA 22217-5660, USA Email: shlesim@onr.navy.mil

Dietrich Stauffer

Institute for Theoretical Physics University of Cologne, Zülpicher Straße 77 D-50937 Köln, Germany Email: stauffer@thp.uni-koeln.de

Jian Qiao Sun

Department of Mechanical Engineering, University of Delaware, Newark DE 19716, USA

Email: sun@me.udel.edu

Dimitry Treschev

Department of Mechanics and Mathematics Moscow State University, Leninskie Gogy Moscow 119899, Russia

E-mail: dtresch@mech.math.msu.su

Vladimir V. Uchaikin

Ulyanovsk State University L.Tolstoy Str., 42, Ulyanovsk 432700 Russia

Email: uchaikin@sv.uven.ru

Angelo Vulpiani

Department of Physics, University La Sapinza, P.le A.Moro 2 00185 Roma, Italy E-mail: Angelo.Vulpiani@roma1.infn.it

Pei Yu

Department of Applied Mathematics The University of Western Ontario, London Ontario N6A 5B7, Canada Email: pyu@pyu1.apmaths.uwo.ca