

Soluções

• Derivadas parciais

1. (a) $\lim_{t \rightarrow 0} g(t) = y^2 + 3$; $\lim_{t \rightarrow 0} h(t) = 2xy$. (b) $\lim_{t \rightarrow 0} g(t) = 8x + y^3$; $\lim_{t \rightarrow 0} h(t) = 3xy^2$.
2. (a) $f_x = 3x^2y + 14x$; $f_y = x^3 - 6y^2$;
(b) $f_x = \frac{(3-7y)y}{(7x+y)^2}$; $f_y = \frac{y^2 + 14xy - 3x}{(7x+y)^2}$;
(c) $f_x = ye^{xy} \cos(1 + e^{xy})$;
 $f_y = xe^{xy} \cos(1 + e^{xy})$;
(d) $f_x = 6x^2(x^3 - y^2)$; $f_y = -4y(x^3 - y^2)$;
(e) $f_x = e^y + y \cos x$; $f_y = xe^y + \sin x$;
(f) $f_s = e^s (\ln(st) + 1/s)$; $f_t = \frac{e^s}{t}$;
(g) $f_x = e^x [\ln(y^2 + 3x) + 3/(y^2 + 3x)]$;
 $f_y = 2ye^x/(y^2 + 3x)$;
(h) $f_x = \cos \frac{x}{y} - \frac{x}{y} \sin \frac{x}{y}$; $f_y = \frac{x^2}{y^2} \sin \frac{x}{y}$;
(i) $f_x = 4x^3 + 6y$; $f_y = 3y^2 + 6x$;
(j) $f_x = 2y^3 e^{2xy^3}$; $f_y = 6xy^2 e^{2xy^3}$;
(k) $f_x = e^{\sqrt{xy}} \left(1 + \frac{xy}{2\sqrt{xy}}\right) = e^{\sqrt{xy}} \left(1 + \frac{\sqrt{xy}}{2}\right)$;
 $f_y = \frac{x^2}{2\sqrt{xy}} e^{\sqrt{xy}}$;
(l) $f_x = yx^{y-1}$; $f_y = x^y \ln x$;
(m) $f_r = \frac{2\pi}{T}$; $f_T = -\frac{2\pi r}{T^2}$;
(n) $f_x = (1 + xy)e^{xy} \sin yz$;
 $f_y = xe^{xy} (x \sin yz + z \cos yz)$;
 $f_z = xye^{xy} \cos yz$;
- (o) $f_s = 2s \cos(2tu)$; $f_t = -2us^2 \sin(2tu)$;
 $f_u = -2ts^2 \sin(2tu)$;
(p) $f_x = 2z$; $f_y = 0$; $f_z = 2(x + z)$;
(q) $f_x = (1 + xyz) yze^{xyz}$;
 $f_y = (1 + xyz) xze^{xyz}$;
 $f_z = (1 + xyz) xye^{xyz}$;
(r) $f_x = \frac{1}{1 + x + y^2 + z^3}$;
 $f_y = \frac{2y}{1 + x + y^2 + z^3}$;
 $f_z = \frac{3z^2}{1 + x + y^2 + z^3}$;
(s) $f_r = -2r \cos(r^2)$; $f_u = 1$; $f_v = 1$;
(t) $f_x = e^x (\sin(x + y) + \cos(x + y))$;
 $f_y = e^x \cos(x + y) + 3 \sin(z - 3y)$;
 $f_z = -\sin(z - 3y)$;
(u) $f_m = \frac{v^2}{r}$; $f_v = \frac{2mv}{r}$; $f_r = -\frac{mv^2}{r^2}$;
(v) $f_x = \frac{yx^{y-1}}{e^z + xy}$; $f_y = \frac{x^y \ln x}{e^z + xy}$; $f_z = \frac{e^z}{e^z + xy}$.
3. (a) $w_{xy} = (w_x)_y = 4y^3 - 12xy^2 = w_{yx}$;
(b) $w_{xy} = (w_x)_y = -6x^2 e^{-2y} + 2y^{-3} \sin x = w_{yx}$;
4. $w_{xyz} = (w_x)_{yz} = (6xy^3z + 2y^4z^2)_{yz} = (18xy^2z + 8y^3z^2)_z = 18xy^2 + 16y^3z$.
5. $\frac{\partial^3 w}{\partial x \partial y \partial z} = 1$.
6. $w_{rrs} = w_{rsr} = w_{srr} = 36r^2 s^2 t - 6st^2 e^{rt}$.
7. $\frac{\partial v}{\partial t} = -\frac{1}{2} t^{-\frac{3}{2}} e^{-\frac{x^2}{4t}} + t^{-\frac{1}{2}} \frac{x^2}{4t^2} e^{-\frac{x^2}{4t}} = \frac{1}{2} t^{-\frac{1}{2}} e^{-\frac{x^2}{4t}} \left(t^{-1} + \frac{x^2}{2t^2}\right)$;
 $\frac{\partial^2 v}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial v}{\partial x}\right) = \frac{\partial}{\partial x} \left(-\frac{x}{2t} t^{-\frac{1}{2}} e^{-\frac{x^2}{4t}}\right) = \frac{\partial}{\partial x} \left(-\frac{x}{2} t^{-\frac{3}{2}} e^{-\frac{x^2}{4t}}\right) = \frac{\partial v}{\partial t}$.
8. (a) $\frac{\partial^2 f}{\partial x^2} = k^2 e^{kx} \cos(ky) = -\frac{\partial^2 f}{\partial y^2}$ (c) $\frac{\partial^2 f}{\partial y^2} = f$; $\frac{\partial^2 f}{\partial x^2} = -f$
(b) $\frac{\partial^2 f}{\partial x^2} = 6y = -\frac{\partial^2 f}{\partial y^2}$

9. $\lambda = -1$.

10. $\frac{\partial^2 w}{\partial x^2} = -\cos(x-y) + \frac{1}{(x+y)^2} = \frac{\partial^2 w}{\partial y^2}$.

11. (a) $\frac{\partial T}{\partial x}(1, 2) = 40x(x^2 + y^2)|_{(1,2)} = 200$; (b) $\frac{\partial T}{\partial y}(1, 2) = 40y(x^2 + y^2)|_{(1,2)} = 400$.

12. (a) $\frac{\partial V}{\partial x}(2, -1, 1) = -\frac{200x}{(x^2 + y^2 + z^2)^2} \Big|_{(2,-1,1)} = -\frac{100}{9}$;

(b) $\frac{\partial V}{\partial y}(2, -1, 1) = -\frac{200y}{(x^2 + y^2 + z^2)^2} \Big|_{(2,-1,1)} = \frac{50}{9}$;

(c) $\frac{\partial V}{\partial z}(2, -1, 1) = -\frac{200z}{(x^2 + y^2 + z^2)^2} \Big|_{(2,-1,1)} = -\frac{50}{9}$;

• Planos tangentes e diferenciais

13. (a) $-4x - 8y + z + 8 = 0$.

(d) $-x - y + z = 0$.

(b) $-6x - 4y + z + 5 = 0$.

(e) $\frac{3}{2}x - \frac{3}{4}y + z - \frac{5}{2} = 0$.

(c) $-e^3 y + z + e^3 = 0$.

(f) $-x - y + z = 0$.

14. (a) $dz = (2x + 3y)dx + (3x - 2y)dy$;

(Observação: dado (x, y) , dz é uma função de dx e dy ; em notação completa, escrevemos $dz = dz_{(x,y)}(dx, dy)$).

(b) $(x, y) = (2, 3)$; $(x + \Delta x, y + \Delta y) = (2.05, 2.96)$;

$dx = \Delta x = 2.05 - 2 = 0.05$; $dy = \Delta y = 2.96 - 3 = -0.04$;

$dz = (2x + 3y)|_{(2,3)} \times 0.05 + (3x - 2y)|_{(2,3)} \times (-0.04) = 0.65$;

$\Delta z = f(2.05, 2.96) - f(2, 3) = 0.6449 \simeq dz$.

15. $(x, y) = (1, 2)$; $(x + \Delta x, y + \Delta y) = (1.05, 2.1)$;

$dx = \Delta x = 1.05 - 1 = 0.05$; $dy = \Delta y = 2.1 - 2 = 0.1$;

$dz = (10x)|_{(1,2)} \times 0.05 + (2y)|_{(1,2)} \times 0.1 = 0.9$;

$\Delta z = f(1.05, 2.1) - f(1, 2) = 0.9225 \simeq dz$.

16. Utilize diferenciais para calcular um valor aproximado de

(a) Seja $f(x, y) = \sqrt{9x^2 + y^2}$. Assim,

$$\sqrt{9(1.95)^2 + (8.1)^2} = f(1.95, 8.1) = f(2 + \Delta x, 8 + \Delta y)$$

com $(\Delta x, \Delta y) = (-0.05, 0.1)$. Usando diferenciais, sabemos que

$$f(2 + \Delta x, 8 + \Delta y) \simeq f(2, 8) + df_{(2,8)}(\Delta x, \Delta y)$$

onde

$$df_{(x,y)}(\Delta x, \Delta y) = \frac{9x \Delta x}{\sqrt{9x^2 + y^2}} + \frac{y \Delta y}{\sqrt{9x^2 + y^2}}.$$

Assim,

$$\sqrt{9(1.95)^2 + (8.1)^2} \simeq \sqrt{9 \times 2^2 + 8^2} + \frac{9 \times 2 \times (-0.05)}{\sqrt{9 \times 2^2 + 8^2}} + \frac{8 \times 0.1}{\sqrt{9 \times 2^2 + 8^2}} = 9.99$$

(b) Definindo $f(x, y) = \sqrt{x} e^y$, temos

$$df_{(x,y)}(\Delta x, \Delta y) = \frac{e^y \Delta x}{2\sqrt{x}} + \sqrt{x} e^y \Delta y$$

e

$$\sqrt{99} e^{0.02} = f(99, 0.02) \simeq f(100, 0) + df_{(100,0)}(\Delta x, \Delta y)$$

com $(\Delta x, \Delta y) = (-1, 0.02)$. Assim,

$$\sqrt{99} e^{0.02} \simeq 10 + \frac{-1}{20} + 10 \times 0.02 = 10.15.$$

(c) Para $f(x, y) = x^2 - y \ln \frac{y}{x}$, temos

$$df_{(x,y)}(\Delta x, \Delta y) = \left(2x + \frac{y}{x}\right) \Delta x + \left(-\ln \frac{y}{x} - 1\right) \Delta y.$$

Assim, com $(x, y) = (1, 1)$ e $(\Delta x, \Delta y) = (-0.02, 0.01)$, podemos escrever

$$(0.98)^2 - 1.01 \ln \frac{1.01}{0.98} = f(0.98, 1.01) \simeq f(1, 1) + df_{(1,1)}(-0.02, 0.01) = 1 - 0.07 = 0.93.$$

(d) Considere-se $f(x, y) = x^{1/3}y^{1/2}$, $(x, y) = (27, 36)$ e $(\Delta x, \Delta y) = (-0.02, 0.04)$. Temos

$$df_{(x,y)}(\Delta x, \Delta y) = \frac{1}{3}x^{-2/3}y^{1/2}\Delta x + \frac{1}{2}x^{1/3}y^{-1/2}\Delta y$$

e

$$26.98^{1/3} \times 36.04^{1/2} = f(26.98, 36.04) \simeq f(27, 36) + df_{(27,36)}(-0.02, 0.04) = 18 + 0.0056 = 18.0056.$$

17. (a) $dw = (3x^2 - 2xy)dx + (-x^2 + 6y)dy$;

(b) $dw = xe^{xy}(2 + xy)dx + \left(x^3e^{xy} - \frac{2}{y^3}\right)dy$;

(c) $dz = e^x (\cos(xy) - y \operatorname{sen}(xy))dx - x e^x \operatorname{sen}(xy)dy$.

(d) $dw = 2x \ln(y^2 + z^2)dx + \frac{2x^2y}{y^2 + z^2}dy + \frac{2x^2z}{y^2 + z^2}dz$;

(e) $dw = \frac{yz(y+z)}{(x+y+z)^2}dx + \frac{xz(x+z)}{(x+y+z)^2}dy + \frac{xy(x+y)}{(x+y+z)^2}dz$;

(f) $dw = (2xz - z^2t)dx + 4t^3dy + (x^2 - 2xzt)dz + (12yt^2 - xz^2)dt$.

18. (a) $dz = \frac{1}{x-3y} \Big|_{(7,2)} \times (-0.1) - \frac{3}{x-3y} \Big|_{(7,2)} \times 0.06 = -0.28 \simeq \Delta z = \ln(6.9 - 3 \times 2.06) - \ln 1$

(b) $dw = y^2 \operatorname{sen} \pi z \Big|_{(4,5,4)} \times (-0.01) + 2xy \operatorname{sen} \pi z \Big|_{(4,5,4)} \times (-0.02) + \pi xy^2 \cos \pi z \Big|_{(4,5,4)} \times 0.03 = 9.4248 \simeq \Delta w$

(c) $dw = \frac{-x}{\sqrt{20-x^2-7y^2}} \Big|_{(2,1)} \times (-0.05) + \frac{-7y}{\sqrt{20-x^2-7y^2}} \Big|_{(2,1)} \times 0.08 = -0.1533 \simeq \Delta w$

19. A área de um retângulo é dada por $A(x, y) = xy$, onde x e y representam o comprimento da base e a altura do retângulo, respetivamente.

Pretende-se estimar o valor máximo de $|A(10 + \Delta x, 5 + \Delta y) - A(10, 5)|$ quando $|\Delta x| \leq 0.1$ e $|\Delta y| \leq 0.1$.

Usando diferenciais, tem-se $dA = y\Delta x + x\Delta y$ e

$$|A(10 + \Delta x, 5 + \Delta y) - A(10, 5)| \simeq |5\Delta x + 10\Delta y| \leq 5 \times 0.1 + 10 \times 0.1 = 1.5 \text{ cm}^2.$$

20. Temos $|\Delta x| \leq 0.2$, $|\Delta y| \leq 0.2$, $|\Delta z| \leq 0.2$ e $dV = yz\Delta x + xz\Delta y + xy\Delta z$. Assim,

$$\begin{aligned} |\Delta V| &= |V(75 + \Delta x, 60 + \Delta y, 40 + \Delta z) - V(75, 60, 40)| \\ &\simeq |60 \times 40 \times \Delta x + 75 \times 40 \times \Delta y + 75 \times 60 \times \Delta z| \leq 1980 \text{ cm}^3. \end{aligned}$$

• Derivadas de funções compostas

21. $\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial s} = -2xy \operatorname{sen}(x^2y)3s^2t^2 - x^2 \operatorname{sen}(x^2y)2s = -2xs \operatorname{sen}(x^2y)(3yst^2 + x)$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial t} = -2xy \operatorname{sen}(x^2y)2s^3t - x^2 \operatorname{sen}(x^2y) \left(-\frac{1}{t^2}\right) = -x \operatorname{sen}(x^2y) \left(4ys^3t - \frac{x}{t^2}\right)$$

22. $\frac{\partial w}{\partial x} = \frac{\partial w}{\partial u} \cdot \frac{\partial u}{\partial x} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial x} = 6ux^2 \operatorname{sen} v + u^2y^2 \cos v$

$$\frac{\partial w}{\partial y} = \frac{\partial w}{\partial u} \cdot \frac{\partial u}{\partial y} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial y} = -12uy^2 \operatorname{sen} v + 2u^2xy \cos v$$

23. $\frac{\partial w}{\partial r} = \frac{\partial w}{\partial u} \cdot \frac{\partial u}{\partial r} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial r} = \frac{ue^{-s}}{\sqrt{u^2 + v^2}} - \frac{vs^2e^{-r}}{\sqrt{u^2 + v^2}}$

$$\frac{\partial w}{\partial s} = \frac{\partial w}{\partial u} \cdot \frac{\partial u}{\partial s} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial s} = \frac{-ur e^{-s}}{\sqrt{u^2 + v^2}} + \frac{2vs e^{-r}}{\sqrt{u^2 + v^2}}$$

$$24. \frac{\partial z}{\partial x} = \frac{\partial w}{\partial r} \cdot \frac{\partial r}{\partial x} + \frac{\partial w}{\partial s} \cdot \frac{\partial s}{\partial x} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial x} = \frac{\cos y}{v} + \frac{y \cos x}{v} - \frac{2x(r+s)e^{-y}}{v^2}$$

$$\frac{\partial z}{\partial y} = \frac{\partial w}{\partial r} \cdot \frac{\partial r}{\partial y} + \frac{\partial w}{\partial s} \cdot \frac{\partial s}{\partial y} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial y} = \frac{-x \operatorname{sen} y}{v} + \frac{\operatorname{sen} x}{v} + \frac{x^2(r+s)e^{-y}}{v^2}$$

$$25. (a) \frac{dw}{dt} = \frac{\partial w}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial w}{\partial y} \cdot \frac{dy}{dt} = -\frac{3x^2}{(1+t)^2} - \frac{3y^2}{(1+t)^2};$$

$$(b) \frac{dw}{dt} = \frac{\partial w}{\partial u} \cdot \frac{du}{dt} + \frac{\partial w}{\partial v} \cdot \frac{dv}{dt} = \frac{2e^{2t}}{u+v} + \frac{3t^2-2t}{u+v};$$

$$(c) \frac{dw}{dt} = \frac{\partial w}{\partial r} \cdot \frac{dr}{dt} + \frac{\partial w}{\partial s} \cdot \frac{ds}{dt} + \frac{\partial w}{\partial t} \cdot \frac{dt}{dt} = 2r \cos t + v \operatorname{sen} t - 4s;$$

$$(d) \frac{dw}{dt} = \frac{\partial w}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial w}{\partial y} \cdot \frac{dy}{dt} + \frac{\partial w}{\partial z} \cdot \frac{dz}{dt} = 4xy^3z^4 + 9x^2y^2z^4 + 20x^2y^3z^3.$$

$$26. \frac{\partial z}{\partial t} = xy^2.$$

$$\frac{dz}{dt} = [t + \ln(e^t + t^2)] e^{2t} + t \left(1 + \frac{e^t + 2t}{e^t + t^2} \right) e^{2t} + 2t [t + \ln(e^t + t^2)] e^{2t}.$$

Sugestão: fazer a composição $z = t [t + \ln(e^t + t^2)] e^{2t}$ e usar as regras de derivação para funções de uma variável.

$$27. \frac{d^2u}{dt^2} = e^{\sin t - 2t^3} [-\sin t - 12t + (\cos t - 6t^2)^2]$$

Sugestão: fazer a composição $u = e^{\sin t - 2t^3}$ e usar as regras de derivação para funções de uma variável.

$$28. \frac{\partial u}{\partial s} = [4x^3yr e^t + 2(x^4 + 2yz^3)rs e^{-t} + 3y^2z^2r^2 \operatorname{sen} t]_{(r=2, s=1, t=0)} = 2^7 + 2^6.$$

$$29. T = \frac{1}{c}pV; \quad \frac{dT}{dt} = \frac{\partial T}{\partial p} \cdot \frac{dp}{dt} + \frac{\partial T}{\partial V} \cdot \frac{dV}{dt} = \frac{1}{c}V \frac{dp}{dt} + \frac{1}{c}p \frac{dV}{dt} = \frac{1}{c} \left(V \frac{dp}{dt} + p \frac{dV}{dt} \right).$$

$$T(t) = \frac{1}{c}p(t)V(t); \quad T'(t) = \frac{1}{c}(p'(t)V(t) + p(t)V'(t)).$$

$$30. \frac{\partial w}{\partial x} = \frac{dw}{du} \frac{\partial u}{\partial x} = 2x \frac{dw}{du} \quad \text{e} \quad \frac{\partial w}{\partial y} = \frac{dw}{du} \frac{\partial u}{\partial y} = 2y \frac{dw}{du}.$$

$$31. \frac{\partial w}{\partial s} = \frac{\partial w}{\partial u} \cdot \frac{\partial u}{\partial s} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial s} = 2s \frac{\partial w}{\partial u} - 2s \frac{\partial w}{\partial v}; \quad \frac{\partial w}{\partial t} = \frac{\partial w}{\partial u} \cdot \frac{\partial u}{\partial t} + \frac{\partial w}{\partial v} \cdot \frac{\partial v}{\partial t} = -2t \frac{\partial w}{\partial u} + 2t \frac{\partial w}{\partial v};$$

$$32. \text{Fazendo } u = x - y, \text{ vem } \frac{\partial z}{\partial x} = \frac{dz}{du} \cdot \frac{\partial u}{\partial x} = \frac{dz}{du} \quad \text{e} \quad \frac{\partial z}{\partial y} = \frac{dz}{du} \cdot \frac{\partial u}{\partial y} = -\frac{dz}{du}.$$

• Derivada da função implícita

$$33. (a) \text{ Seja } F(x, y) = 2x^3 + x^2y + y^3 - 1.$$

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{6x^2 + 2xy}{x^2 + 3y^2}.$$

$$(b) \text{ Seja } F(x, y) = 6x + \sqrt{xy} - 3y + 4.$$

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{6 + \frac{y}{2\sqrt{xy}}}{\frac{x}{2\sqrt{xy}} - 3} = -\frac{12\sqrt{xy} + y}{x - 6\sqrt{xy}}.$$

$$(c) \frac{dy}{dx} = -\frac{3x^2 - 6y}{3y^2 - 6x}.$$

$$(d) \frac{dy}{dx} = -\frac{2xy^2 + 1}{2x^2y - 6y^2}.$$

$$34. \frac{dy}{dx}(x) = -\frac{(1+y-x^2+\ln y)_x}{(1+y-x^2+\ln y)_y} = -\frac{-2x}{1+\frac{1}{y}} = \frac{2xy}{y+1}; \quad \frac{dy}{dx}(\sqrt{2}) = \frac{2\sqrt{2}}{2} = \sqrt{2}.$$

$$35. (a) \text{ Seja } F(x, y, z) = 2xz^3 - 3yz^2 + x^2y^2 + 4z.$$

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{2x^2y^2}{6xz^2 - 6yz + 4};$$

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{-3z^2 + 2x^2y}{6xz^2 - 6yz + 4}.$$

$$\frac{\partial z}{\partial x} = -\frac{e^{yz} - 2yze^{xz} + 3ye^{xy}}{xye^{yz} - 2xye^{xz} + 3e^{xy}};$$

$$\frac{\partial z}{\partial y} = -\frac{xze^{yz} - 2e^{xz} + 3xe^{xy}}{xye^{yz} - 2xye^{xz} + 3e^{xy}}.$$

$$(c) \frac{\partial z}{\partial x} = -\frac{2xy - yz \operatorname{sen}(xyz)}{2z - xy \operatorname{sen}(xyz)};$$

$$\frac{\partial z}{\partial y} = -\frac{x^2 - xz \operatorname{sen}(xyz)}{2z - xy \operatorname{sen}(xyz)}.$$

$$(d) \frac{\partial z}{\partial x} = -\frac{z^2 + 4xy}{2xz - 4y^2};$$

$$\frac{\partial z}{\partial y} = -\frac{2x^2 - 8yz + 3}{2xz - 4y^2}.$$