

Some Regularity Results for the Obstacle Problem

Semester project

Florian Noah Grün
École Polytechnique Fédérale de Lausanne, Switzerland

AMCV Professor Maria Colombo
Supervisor Xavier Fernández-Real

09.06.2022

Outline

1. Introduction to the obstacle problem
2. Regularity of solution
3. The free boundary
 - 3.1 Classification of blow-ups
 - 3.2 Regularity
 - 3.3 Singular points

Outline

1. Introduction to the obstacle problem
2. Regularity of solution
3. The free boundary
 - 3.1 Classification of blow-ups
 - 3.2 Regularity
 - 3.3 Singular points

Introduction to the obstacle problem

$$\text{minimize } \int_{\Omega} \frac{1}{2} |\nabla v(x)|^2 dx \quad \text{s.t. } v \geq \varphi \text{ and } v|_{\partial\Omega} = g \quad (\text{OP})$$

$$\text{minimize } \int_{\Omega} \left(\frac{1}{2} |\nabla u|^2 + fu \right) dx \quad \text{s.t. } u \geq 0, u|_{\partial\Omega} = g - \varphi, f := -\Delta\varphi \quad (\text{ZOP})$$

- Under reasonable assumptions existence and uniqueness of minimizer
- Domain has two parts $\{u = 0\}$ and $\{u > 0\}$
- How does $\partial\{u > 0\}$ look like? Smooth?
- Blackboard-Sketch of basic setup

Introduction to the obstacle problem

$$\text{minimize } \int_{\Omega} \frac{1}{2} |\nabla v(x)|^2 dx \quad \text{s.t. } v \geq \varphi \text{ and } v|_{\partial\Omega} = g \quad (\text{OP})$$

$$\text{minimize } \int_{\Omega} \left(\frac{1}{2} |\nabla u|^2 + fu \right) dx \quad \text{s.t. } u \geq 0, u|_{\partial\Omega} = g - \varphi, f := -\Delta\varphi \quad (\text{ZOP})$$

- Under reasonable assumptions existence and uniqueness of minimizer
- Domain has two parts $\{u = 0\}$ and $\{u > 0\}$
- How does $\partial\{u > 0\}$ look like? Smooth?
- Blackboard-Sketch of basic setup

Introduction to the obstacle problem

$$\text{minimize } \int_{\Omega} \frac{1}{2} |\nabla v(x)|^2 dx \quad \text{s.t. } v \geq \varphi \text{ and } v|_{\partial\Omega} = g \quad (\text{OP})$$

$$\text{minimize } \int_{\Omega} \left(\frac{1}{2} |\nabla u|^2 + fu \right) dx \quad \text{s.t. } u \geq 0, u|_{\partial\Omega} = g - \varphi, f := -\Delta\varphi \quad (\text{ZOP})$$

- Under reasonable assumptions existence and uniqueness of minimizer
- Domain has two parts $\{u = 0\}$ and $\{u > 0\}$
- How does $\partial\{u > 0\}$ look like? Smooth?
- Blackboard-Sketch of basic setup

Introduction to the obstacle problem

$$\text{minimize } \int_{\Omega} \frac{1}{2} |\nabla v(x)|^2 dx \quad \text{s.t. } v \geq \varphi \text{ and } v|_{\partial\Omega} = g \quad (\text{OP})$$

$$\text{minimize } \int_{\Omega} \left(\frac{1}{2} |\nabla u|^2 + fu \right) dx \quad \text{s.t. } u \geq 0, u|_{\partial\Omega} = g - \varphi, f := -\Delta\varphi \quad (\text{ZOP})$$

- Under reasonable assumptions existence and uniqueness of minimizer
- Domain has two parts $\{u = 0\}$ and $\{u > 0\}$
- How does $\partial\{u > 0\}$ look like? Smooth?
- Blackboard-Sketch of basic setup

Introduction to the obstacle problem

$$\text{minimize } \int_{\Omega} \frac{1}{2} |\nabla v(x)|^2 dx \quad \text{s.t. } v \geq \varphi \text{ and } v|_{\partial\Omega} = g \quad (\text{OP})$$

$$\text{minimize } \int_{\Omega} \left(\frac{1}{2} |\nabla u|^2 + fu \right) dx \quad \text{s.t. } u \geq 0, u|_{\partial\Omega} = g - \varphi, f := -\Delta\varphi \quad (\text{ZOP})$$

- Under reasonable assumptions existence and uniqueness of minimizer
- Domain has two parts $\{u = 0\}$ and $\{u > 0\}$
- How does $\partial\{u > 0\}$ look like? Smooth?
- Blackboard-Sketch of basic setup

Regularity of solution

- EL equation: $\Delta u = f \chi_{\{u>0\}} \Leftrightarrow \begin{cases} \Delta u = f & \text{in } \{u > 0\}, \\ u = 0 & \text{on } \partial\{u > 0\}, \\ \nabla u = 0 & \text{on } \partial\{u > 0\}. \end{cases}$
- Directly $u \in C^{1,1-\epsilon}$ by Schauder estimate
- $0 < cr^2 < \sup_{B_r(x_0)} u \leq Cr^2$ for any FBP x_0 (Harnack's inequality and scaling)
- Nondegeneracy needs extra assumption: $f \geq c_0 > 0$
- Quadratic growth & Schauder estimates give $C^{1,1}$ regularity of u .
- Schauder estimate: solution is two derivatives smoother than RHS

Regularity of solution

- EL equation: $\Delta u = f \chi_{\{u>0\}} \Leftrightarrow \begin{cases} \Delta u = f & \text{in } \{u > 0\}, \\ u = 0 & \text{on } \partial\{u > 0\}, \\ \nabla u = 0 & \text{on } \partial\{u > 0\}. \end{cases}$
- Directly $u \in C^{1,1-\epsilon}$ by Schauder estimate
- $0 < cr^2 < \sup_{B_r(x_0)} u \leq Cr^2$ for any FBP x_0 (Harnack's inequality and scaling)
- Nondegeneracy needs extra assumption: $f \geq c_0 > 0$
- Quadratic growth & Schauder estimates give $C^{1,1}$ regularity of u .
- Schauder estimate: solution is two derivatives smoother than RHS

Regularity of solution

- EL equation: $\Delta u = f \chi_{\{u>0\}} \Leftrightarrow \begin{cases} \Delta u = f & \text{in } \{u > 0\}, \\ u = 0 & \text{on } \partial\{u > 0\}, \\ \nabla u = 0 & \text{on } \partial\{u > 0\}. \end{cases}$
- Directly $u \in C^{1,1-\epsilon}$ by Schauder estimate
- $0 < cr^2 < \sup_{B_r(x_0)} u \leq Cr^2$ for any FBP x_0 (Harnack's inequality and scaling)
- Nondegeneracy needs extra assumption: $f \geq c_0 > 0$
- Quadratic growth & Schauder estimates give $C^{1,1}$ regularity of u .
- Schauder estimate: solution is two derivatives smoother than RHS

Regularity of solution

- EL equation: $\Delta u = f \chi_{\{u>0\}} \Leftrightarrow \begin{cases} \Delta u = f & \text{in } \{u > 0\}, \\ u = 0 & \text{on } \partial\{u > 0\}, \\ \nabla u = 0 & \text{on } \partial\{u > 0\}. \end{cases}$
- Directly $u \in C^{1,1-\epsilon}$ by Schauder estimate
- $0 < cr^2 < \sup_{B_r(x_0)} u \leq Cr^2$ for any FBP x_0 (Harnack's inequality and scaling)
- Nondegeneracy needs extra assumption: $f \geq c_0 > 0$
- Quadratic growth & Schauder estimates give $C^{1,1}$ regularity of u .
- Schauder estimate: solution is two derivatives smoother than RHS

Free Boundary: Classification of blow-ups

- Our setting:
$$\begin{cases} u & \in C^{1,1}(B_1), \\ u & \geq 0, \\ \Delta u & = \chi_{\{u>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$

- Rescaling ("zooming in") $u_r(x) := \frac{u(x_0+rx)}{r^2} \longrightarrow u_0$ in $C_{loc}^1(\mathbb{R}^n)$

- How does the blow-up (if it exists) look?

- Global problem:
$$\begin{cases} u_0 & \in C_{loc}^{1,1}(\mathbb{R}^n), \\ u_0 & \geq 0, \\ \Delta u_0 & = \chi_{\{u_0>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$

Free Boundary: Classification of blow-ups

- Our setting:
$$\begin{cases} u & \in C^{1,1}(B_1), \\ u & \geq 0, \\ \Delta u & = \chi_{\{u>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$
- Rescaling ("zooming in") $u_r(x) := \frac{u(x_0+rx)}{r^2} \longrightarrow \mathbf{u_0}$ in $C_{loc}^1(\mathbb{R}^n)$

• How does the blow-up (if it exists) look?

- Global problem:
$$\begin{cases} u_0 & \in C_{loc}^{1,1}(\mathbb{R}^n), \\ u_0 & \geq 0, \\ \Delta u_0 & = \chi_{\{u_0>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$

Free Boundary: Classification of blow-ups

- Our setting:
$$\begin{cases} u & \in C^{1,1}(B_1), \\ u & \geq 0, \\ \Delta u & = \chi_{\{u>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$
- Rescaling ("zooming in") $u_r(x) := \frac{u(x_0+rx)}{r^2} \longrightarrow u_0$ in $C_{loc}^1(\mathbb{R}^n)$
- How does the blow-up (if it exists) look?

- Global problem:
$$\begin{cases} u_0 & \in C_{loc}^{1,1}(\mathbb{R}^n), \\ u_0 & \geq 0, \\ \Delta u_0 & = \chi_{\{u_0>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$

Free Boundary: Classification of blow-ups

- Our setting:
$$\begin{cases} u & \in C^{1,1}(B_1), \\ u & \geq 0, \\ \Delta u & = \chi_{\{u>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$
- Rescaling ("zooming in") $u_r(x) := \frac{u(x_0+rx)}{r^2} \longrightarrow u_0$ in $C_{loc}^1(\mathbb{R}^n)$
- How does the blow-up (if it exists) look?
- Global problem:
$$\begin{cases} u_0 & \in C_{loc}^{1,1}(\mathbb{R}^n), \\ u_0 & \geq 0, \\ \Delta u_0 & = \chi_{\{u_0>0\}}, \\ 0 & \text{is FBP.} \end{cases}$$

Free Boundary: Classification of blow-ups

- Blow-ups are 2-homogeneous and convex!
- Two types of blow-ups:
 1. $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$ if $\{u_0 = 0\}$ has non-empty interior (half-space)
 2. $u_0(x) = x^T A x$ $\{u_0 = 0\}$ has empty interior (line)


Free Boundary: Classification of blow-ups

- Blow-ups are 2-homogeneous and convex!
- Two types of blow-ups:
 1. $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$ if $\{u_0 = 0\}$ has non-empty interior (half-space)
 2. $u_0(x) = x^T A x$ $\{u_0 = 0\}$ has empty interior (line)


Free Boundary: Classification of blow-ups

- Blow-ups are 2-homogeneous and convex!
- Two types of blow-ups:
 1. $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$ if $\{u_0 = 0\}$ has non-empty interior (half-space)
 2. $u_0(x) = x^T A x$ if $\{u_0 = 0\}$ has empty interior (line)

$$u_0(x) = \frac{1}{2}(x \cdot e)_+^2$$




$$u_0(x) = \frac{1}{2}x_1^2$$




Free Boundary: Classification of blow-ups

- Blow-ups are 2-homogeneous and convex!
- Two types of blow-ups:
 1. $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$ if $\{u_0 = 0\}$ has non-empty interior (half-space)
 2. $u_0(x) = x^T A x$ if $\{u_0 = 0\}$ has empty interior (line)


$$u_0(x) = \frac{1}{2}(x \cdot e)_+^2$$



$$u_0(x) = \frac{1}{2}x_1^2$$



Free Boundary: Classification of blow-ups



Free Boundary: Regularity

- Regular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} > 0$
- u has a blow-up $u_0 = \frac{1}{2}(x \cdot e)_+^2$ for some $e \in \mathbb{S}^{n-1}$.
- u_0 and its derivative are "close" to u_{r_0} for some r_0 .
- The free boundary of the scaled version u_{r_0} around 0 is contained in a strip.
- There exists a cone (not a full half space) of directions τ , in where u_{r_0} is non-decreasing. (transferring monotonicity from u_0 to u_{r_0})
- Free boundary of u_{r_0} is Lipschitz (interior and exterior cone condition)
 \implies free boundary of u is also locally Lipschitz.

Free Boundary: Regularity

- Regular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} > 0$
- u has a blow-up $u_0 = \frac{1}{2}(x \cdot e)_+^2$ for some $e \in \mathbb{S}^{n-1}$.
- u_0 and its derivative are "close" to u_{r_0} for some r_0 .
- The free boundary of the scaled version u_{r_0} around 0 is contained in a strip.
- There exists a cone (not a full half space) of directions τ , in where u_{r_0} is non-decreasing. (transferring monotonicity from u_0 to u_{r_0})
- Free boundary of u_{r_0} is Lipschitz (interior and exterior cone condition)
 \implies free boundary of u is also locally Lipschitz.

Free Boundary: Regularity

- Regular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} > 0$
- u has a blow-up $u_0 = \frac{1}{2}(x \cdot e)_+^2$ for some $e \in \mathbb{S}^{n-1}$.
- u_0 and its derivative are "close" to u_{r_0} for some r_0 .
- The free boundary of the scaled version u_{r_0} around 0 is contained in a strip.
- There exists a cone (not a full half space) of directions τ , in where u_{r_0} is non-decreasing. (transferring monotonicity from u_0 to u_{r_0})
- Free boundary of u_{r_0} is Lipschitz (interior and exterior cone condition)
 \implies free boundary of u is also locally Lipschitz.

Free Boundary: Regularity

- Regular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} > 0$
- u has a blow-up $u_0 = \frac{1}{2}(x \cdot e)_+^2$ for some $e \in \mathbb{S}^{n-1}$.
- u_0 and its derivative are "close" to u_{r_0} for some r_0 .
- The free boundary of the scaled version u_{r_0} around 0 is contained in a strip.
- There exists a cone (not a full half space) of directions τ , in where u_{r_0} is non-decreasing. (transferring monotonicity from u_0 to u_{r_0})
- Free boundary of u_{r_0} is Lipschitz (interior and exterior cone condition)
 \implies free boundary of u is also locally Lipschitz.

Free Boundary: Regularity

- Regular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} > 0$
- u has a blow-up $u_0 = \frac{1}{2}(x \cdot e)_+^2$ for some $e \in \mathbb{S}^{n-1}$.
- u_0 and its derivative are "close" to u_{r_0} for some r_0 .
- The free boundary of the scaled version u_{r_0} around 0 is contained in a strip.
- There exists a cone (not a full half space) of directions τ , in where u_{r_0} is non-decreasing. (transferring monotonicity from u_0 to u_{r_0})
- Free boundary of u_{r_0} is Lipschitz (interior and exterior cone condition)
 \implies free boundary of u is also locally Lipschitz.

Free Boundary: Regularity

- Regular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} > 0$
- u has a blow-up $u_0 = \frac{1}{2}(x \cdot e)_+^2$ for some $e \in \mathbb{S}^{n-1}$.
- u_0 and its derivative are "close" to u_{r_0} for some r_0 .
- The free boundary of the scaled version u_{r_0} around 0 is contained in a strip.
- There exists a cone (not a full half space) of directions τ , in where u_{r_0} is non-decreasing. (transferring monotonicity from u_0 to u_{r_0})
- Free boundary of u_{r_0} is Lipschitz (interior and exterior cone condition)
 \implies free boundary of u is also locally Lipschitz.


Free Boundary: Regularity

- Boundary Harnack:

$$\Omega \text{ is Lip, } w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{0,\alpha}} \leq C.$$

- Higher order Boundary Harnack:

$$\Omega \text{ is } C^{k,\alpha}, w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{k,\alpha}} \leq C$$



- Boundary Harnack gives $C^{1,\alpha}$ regularity around the origin.
- Higher order Boundary Harnack gives C^∞ regularity around the origin.
- FB has no corners!

Free Boundary: Regularity


- Boundary Harnack:

$$\Omega \text{ is Lip, } w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{0,\alpha}} \leq C.$$

- Higher order Boundary Harnack:

$$\Omega \text{ is } C^{k,\alpha}, w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{k,\alpha}} \leq C$$

- Boundary Harnack gives $C^{1,\alpha}$ regularity around the origin.
- Higher order Boundary Harnack gives C^∞ regularity around the origin.
- FB has no corners!



Free Boundary: Regularity


- Boundary Harnack:

$$\Omega \text{ is Lip, } w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{0,\alpha}} \leq C.$$

- Higher order Boundary Harnack:

$$\Omega \text{ is } C^{k,\alpha}, w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{k,\alpha}} \leq C$$

- Boundary Harnack gives $C^{1,\alpha}$ regularity around the origin.
- Higher order Boundary Harnack gives C^∞ regularity around the origin.
- FB has no corners!



Free Boundary: Regularity


- Boundary Harnack:

$$\Omega \text{ is Lip, } w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{0,\alpha}} \leq C.$$

- Higher order Boundary Harnack:

$$\Omega \text{ is } C^{k,\alpha}, w_1, w_2 \text{ harmonic} \implies \left\| \frac{w_1}{w_2} \right\|_{C^{k,\alpha}} \leq C$$

- Boundary Harnack gives $C^{1,\alpha}$ regularity around the origin.
- Higher order Boundary Harnack gives C^∞ regularity around the origin.
- FB has no corners!



Free Boundary: Singular points

- Singular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} = 0$
- Caffarelli's dichotomy (1977):
 1. regular points: $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$
 2. singular points: $u_0(x) = \frac{1}{2}x^T A x$
- Blow-up at singular points is unique (Monneau monotonicity formula)
- Continuous dependence of blow-ups
- Caffarelli [Caf98]: The singular set is locally contained in a C^1 $(n-1)$ dimensional manifold.
- Monneau [Mon03]: Generically (for "almost every" boundary data) the free boundary has no singular points in \mathbb{R}^2 .

Free Boundary: Singular points

- Singular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} = 0$
- Caffarelli's dichotomy (1977):
 1. regular points: $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$
 2. singular points: $u_0(x) = \frac{1}{2}x^T A x$
- Blow-up at singular points is unique (Monneau monotonicity formula)
- Continuous dependence of blow-ups
- Caffarelli [Caf98]: The singular set is locally contained in a C^1 $(n-1)$ dimensional manifold.
- Monneau [Mon03]: Generically (for "almost every" boundary data) the free boundary has no singular points in \mathbb{R}^2 .

Free Boundary: Singular points

- Singular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} = 0$
- Caffarelli's dichotomy (1977):
 1. regular points: $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$
 2. singular points: $u_0(x) = \frac{1}{2}x^T A x$
- Blow-up at singular points is unique (Monneau monotonicity formula)
- Continuous dependence of blow-ups
- Caffarelli [Caf98]: The singular set is locally contained in a C^1 $(n-1)$ dimensional manifold.
- Monneau [Mon03]: Generically (for "almost every" boundary data) the free boundary has no singular points in \mathbb{R}^2 .

Free Boundary: Singular points

- Singular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} = 0$
- Caffarelli's dichotomy (1977):
 1. regular points: $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$
 2. singular points: $u_0(x) = \frac{1}{2}x^T A x$
- Blow-up at singular points is unique (Monneau monotonicity formula)
- Continuous dependence of blow-ups
- Caffarelli [Caf98]: The singular set is locally contained in a C^1 $(n-1)$ dimensional manifold.
- Monneau [Mon03]: Generically (for "almost every" boundary data) the free boundary has no singular points in \mathbb{R}^2 .

Free Boundary: Singular points

- Singular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} = 0$
- Caffarelli's dichotomy (1977):
 1. regular points: $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$
 2. singular points: $u_0(x) = \frac{1}{2}x^T A x$
- Blow-up at singular points is unique (Monneau monotonicity formula)
- Continuous dependence of blow-ups
- Caffarelli [Caf98]: The singular set is locally contained in a C^1 $(n-1)$ dimensional manifold.
- Monneau [Mon03]: Generically (for "almost every" boundary data) the free boundary has no singular points in \mathbb{R}^2 .

Free Boundary: Singular points

- Singular FBP: $\limsup_{r \rightarrow 0} \frac{|\{u=0\} \cap B_r(x_0)|}{|B_r|} = 0$
- Caffarelli's dichotomy (1977):
 1. regular points: $u_0(x) = \frac{1}{2}(x \cdot e)_+^2$
 2. singular points: $u_0(x) = \frac{1}{2}x^T A x$
- Blow-up at singular points is unique (Monneau monotonicity formula)
- Continuous dependence of blow-ups
- Caffarelli [Caf98]: The singular set is locally contained in a C^1 $(n-1)$ dimensional manifold.
- Monneau [Mon03]: Generically (for "almost every" boundary data) the free boundary has no singular points in \mathbb{R}^2 .

Thank you for listening!



H.W. Alt and D. Phillips.

A free boundary problem for semilinear elliptic equations.

J. Reine Angew. Math., 368:63–107, 1986.



L.A. Caffarelli.

The obstacle problem revisited.

J. Fourier Anal. Appl., 4:383–402, 1998.



Xavier Fernández-Real and Xavier Ros-Oton.

Regularity Theory for Elliptic PDE.

2022.



R. Monneau.

On the number of singularities for the obstacle problem in two dimensions.

The Journal of Geometric Analysis, 13(2), 2003.



Arshak Petrosyan, Henrik Shahgholian, and Nina Uraltseva.

Regularity of Free Boundaries in Obstacle-Type Problems, volume 136 of *Graduate Studies in Mathematics*.

American Mathematical Society, Providence, RI, 2012.



Joel Spruck.

Uniqueness in a diffusion model of population biology.

Comm. Partial Differential Equations, 8:1605–1620, 1983.



Georg Weiss.

A homogeneity improvement approach to the obstacle problem.

Inventiones Mathematicae, 138:23–50, 10 1999.